

EXTRACTA MATHEMATICAE Vol. 28, Núm. 2, 157–167 (2013)

On the Approximate Solution of D'Alembert Type Equation Originating from Number Theory

B. BOUIKHALENE¹, E. ELQORACHI², A. CHARIFI¹

¹*Sultan Moulay Slimane University, Polydisciplinaire Faculty, Beni-Mellal, Morocco*

²*Ibn Zohr University, Faculty of Sciences, Agadir, Morocco*

bbouikhalene@yahoo.fr, elqorachi@yahoo.fr, charifi2000@yahoo.fr

Presented by David Yost

Received December 2, 2012

Abstract: We solve the functional equation

$$E(\alpha) : f(x_1x_2 + \alpha y_1y_2, x_1y_2 + x_2y_1) + f(x_1x_2 - \alpha y_1y_2, x_2y_1 - x_1y_2) = 2f(x_1, y_1)f(x_2, y_2),$$

where $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$, $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ and α is a real parameter, on the monoid \mathbb{R}^2 . Also we investigate the stability of this equation in the following setting:

$$|f(x_1x_2 + \alpha y_1y_2, x_1y_2 + x_2y_1) + f(x_1x_2 - \alpha y_1y_2, x_2y_1 - x_1y_2) - 2f(x_1, y_1)f(x_2, y_2)| \\ \leq \min\{\varphi(x_1), \psi(y_1), \phi(x_2), \zeta(y_2)\}.$$

From this result, we obtain the superstability of this equation.

Key words: D'Alembert functional equation, monoid \mathbb{R}^2 , multiplicative function, stability, superstability.

AMS *Subject Class.* (2010): 47D09, 22D10, 39B82.

1. INTRODUCTION

For any $\alpha \in \mathbb{R}$, Berrone and Dieulefait [5] equipped \mathbb{R}^2 with the multiplication rule \cdot_α , defined by

$$(x_1, y_1) \cdot_\alpha (x_2, y_2) = (x_1x_2 + \alpha y_1y_2, x_1y_2 + x_2y_1), \quad (x_1, y_1), (x_2, y_2) \in \mathbb{R}^2.$$

For $\alpha = -1$, the multiplication is the usual product of complex numbers in $\mathbb{C} = \mathbb{R}^2$. The rule makes \mathbb{R}^2 into a commutative monoid with neutral element $(1, 0)$ and $\sigma(x, y) = (x, -y)$ (complex conjugation) as an involution.

Berrone and Dieulefait [5, Theorem 1] studied the homomorphisms $m : (\mathbb{R}^2, \cdot_\alpha) \rightarrow (\mathbb{R}, \cdot)$, i.e., the multiplicative, real-valued functions on the monoid $(\mathbb{R}^2, \cdot_\alpha)$. We extend their investigations by finding the bigger set of all multiplicative, complex-valued functions $M : (\mathbb{R}^2, \cdot_\alpha) \rightarrow (\mathbb{C}, \cdot)$. Combining

this information with Davison's work [9] about D'Alembert's functional equation on monoids, we obtain an explicit description of the solutions $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ of D'Alembert's functional equation

$$E(\alpha) : f(a \cdot_{\alpha} b) + f(a \cdot_{\alpha} \sigma(b)) = 2f(a)f(b), \quad a, b \in \mathbb{R}^2,$$

on the monoid $(\mathbb{R}^2, \cdot_{\alpha})$. The description falls into three different cases, according to whether $\alpha > 0$ or $\alpha < 0$. The equation $E(\alpha)$ is a common generalization of many functional equations of type D'Alembert

$$f(ab) + f(a\sigma(b)) = 2f(a)f(b), \quad a, b \in \mathbb{R}^2 \quad (1.1)$$

on the monoid \mathbb{R}^2 , like, e.g.,

1) If $\alpha = 0$,

$$E(0) : f(x_1x_2, x_1y_2 + x_2y_1) + f(x_1x_2, x_2y_1 - x_1y_2) = 2f(x_1, y_1)f(x_2, y_2),$$

for all $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$. Setting $x_1 = x_2 = 1$ and $F(y) = f(1, y)$ for any $y \in \mathbb{R}$ respectively $y_1 = y_2 = 0$ and $m(x) = f(x, 0)$ for any $x \in \mathbb{R}$ in $E(0)$, we get the classical D'Alembert functional equation

$$F(y_1 + y_2) + F(y_1 - y_2) = 2F(y_1)F(y_2), \quad y_1, y_2 \in \mathbb{R} \quad (1.2)$$

on \mathbb{R} (see [1], [4], [15] and [23]) respectively the classical Cauchy equation

$$m(x_1x_2) = m(x_1)m(x_2), \quad x_1, x_2 \in \mathbb{R} \quad (1.3)$$

on \mathbb{R} . We call m a multiplicative function on \mathbb{R} (see[1]).

2) If $\alpha = -1$,

$$\begin{aligned} E(-1) : f(x_1x_2 - y_1y_2, x_1y_2 + x_2y_1) + f(x_1x_2 + y_1y_2, x_2y_1 - x_1y_2) \\ = 2f(x_1, y_1)f(x_2, y_2), \end{aligned}$$

$(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$. The equation $E(-1)$ is in connection with the identity

$$\begin{aligned} (x_1x_2 - y_1y_2)^2 + (x_1y_2 + x_2y_1)^2 + (x_1x_2 + y_1y_2)^2 + (x_2y_1 - x_1y_2)^2 \\ = 2(x_1^2 + y_1^2)(x_2^2 + y_2^2) \end{aligned} \quad (1.4)$$

for any $x_1, x_2, y_1, y_2 \in \mathbb{R}$.

3) If $\alpha \neq 1$ is a square free integer and $\mathbb{Q}(\sqrt{\alpha}) = \{x + y\sqrt{\alpha} : x, y \in \mathbb{Q}\}$ is the quadratic monoid equipped with the multiplicative rule

$$(x_1 + y_1\sqrt{\alpha})(x_2 + y_2\sqrt{\alpha}) = (x_1x_2 + \alpha y_1y_2) + (x_1y_2 + x_2y_1)\sqrt{\alpha}, \quad (1.5)$$

then $E(\alpha)$ reduces to D'Alembert functional equation (1.1) on the monoid $\mathbb{Q}(\sqrt{\alpha})$. In [9] Davison solved the D'Alembert functional equation with involution on a monoid A : any solution $f : A \rightarrow \mathbb{C}$ has the general form $f = \frac{M+M\circ\sigma}{2}$, where $M : A \rightarrow \mathbb{C}$ is a multiplicative function.

In 1940, Ulam [22] gave a talk before the Mathematics Club of the University of Wisconsin in which he discussed a number of unsolved problems. Among these was the following question concerning the stability of homomorphisms.

QUESTION 1.1. Let $(G_1, *)$ be a group and let (G_1, \diamond, d) be a metric group with the metric d . Given $\varepsilon > 0$, does there exist $\delta(\varepsilon) > 0$ such that if a mapping $h : G_1 \rightarrow G_2$ satisfies the inequality $d(h(x * y), h(x) \diamond h(y)) < \delta$ for all $x, y \in G_1$, then there is a homomorphism $H : G_1 \rightarrow G_2$ with $d(h(x), H(x)) < \delta(\varepsilon)$ for all $x \in G_1$?

In 1941, Hyers [12] answered this question for the case where G_1 and G_2 are Banach spaces. In 1978, Rassias [20] provided a generalization of Hyer's theorem which allows the Cauchy difference to be unbounded. The interested reader may refer to the book by Hyers, Isac, Rassias [13] for an in depth account on the subject of stability of functional equations. In 1982, Rassias [19] solved the Ulam problem by involving a product of powers of norms. Since then, the stability problems of various functional equations ha been investigated by many authors (see [10], [11] and [14]). In [3] and [7] Baker et al. and Bourgin respectively, introduced the notion that by now is frequently referred to as superstability or Baker's stability: if a function f satisfies the stability inequality $|E_1(f) - E_2(f)| \leq \varepsilon$, then either f is bounded or $E_1(f) = E_2(f)$. The superstability of D'Alembert's functional equation $f(x + y) + f(x - y) = 2f(x)f(y)$ was investigated by Baker [4] and Cholewa [8]. Badora and Ger [2], and Kim ([16], [17] and [18]) proved its superstability under the condition $|f(x + y) + f(x - y) - 2f(x)f(y)| \leq \varphi(x)$ or $\varphi(y)$. In a previuos work, Bouikhalene et al. [6] investigated the superstability of the cosine functional equation on the Heisenberg group. Following this investigation we study the superstability of the functional equation $E(\alpha)$ on the monoid $(\mathbb{R}^2, \cdot_\alpha)$. Also we say that a function $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ is of approximate a cosine type function,

if there is $\delta > 0$ such that

$$|f(a \cdot_{\alpha} b) + f(a \cdot_{\alpha} i(b)) - 2f(a)f(b)| < \delta, \quad a, b \in \mathbb{R}^2. \quad (1.6)$$

In the case where $\delta = 0$, f satisfies the functional equation $E(\alpha)$. We call f a cosine type function on \mathbb{R}^2 . The paper is organized as follows: In the first section after this introduction we solve the functional equation $E(\alpha)$. In the second section we study the superstability equation $E(\alpha)$.

2. SOLUTION OF EQUATION $E(\alpha)$

According to [9] we drive the following lemma.

LEMMA 2.1. *The solution $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ of $E(\alpha)$ is of the form*

$$f = \frac{M + M \circ \sigma}{2},$$

where $M : (\mathbb{R}^2, \cdot_{\alpha}) \rightarrow (\mathbb{C}, \cdot)$ is a multiplicative function.

By extending Berrone-Dieulefait's result [5] to complex-valued multiplicative functions, we get the following lemmas.

LEMMA 2.2. *The multiplicative functions $M : (\mathbb{R}^2, \cdot_1) \rightarrow (\mathbb{C}, \cdot)$ are the functions*

$$M(x, y) = m_1(x + y)m_2(x - y), \quad x, y \in \mathbb{R},$$

where $m_1, m_2 : \mathbb{R} \rightarrow \mathbb{C}$ are multiplicative functions.

LEMMA 2.3. *The multiplicative functions $M : (\mathbb{R}^2, \cdot_0) \rightarrow (\mathbb{C}, \cdot)$ are the trivial function $M = 1$ and $M(0, y) = 0$ for any $y \in \mathbb{R}$ and $M(x, y) = m(x)\gamma(\frac{y}{x})$ for any $(x, y) \in \mathbb{R}^2$, with $x \neq 0$, where $m : \mathbb{R} \rightarrow \mathbb{C}$ is a multiplicative function and $\gamma : (\mathbb{R}, +) \rightarrow \mathbb{C}$ is an arbitrary character.*

LEMMA 2.4. *The multiplicative functions $M : (\mathbb{C}, \cdot_{-1}) \rightarrow (\mathbb{C}, \cdot)$ are the trivial functions $M = 0$ and $M = 1$ and*

$$M(z) = \begin{cases} \tilde{m}(|z|)\Gamma(\exp(i\theta)), & \text{for } z = |z|\exp(i\theta) \neq 0 \\ 0, & \text{for } z = 0. \end{cases}$$

where $\tilde{m} : (\mathbb{R}^+, \cdot) \rightarrow \mathbb{C}^*$ and $\Gamma : \{\exp(i\theta), \theta \in \mathbb{R}\} \rightarrow \mathbb{C}^*$ are arbitrary characters.

Proof. When $\alpha = -1$, the multiplicative rule \cdot_{-1} becomes the usual product numbers in \mathbb{C} . By using the polar decomposition $z = |z| \exp(i\theta)$ for any $z \in \mathbb{C}^*$ where $\theta = \arg(z)$, we get

$$M(|z_1||z_2|) = M(|z_1|)M(|z_2|), \quad z_1, z_2 \in \mathbb{C}^* \tag{2.1}$$

and

$$M(\exp(i(\theta_1 + \theta_2))) = M(\exp(i\theta_1))M(\exp(i\theta_2)), \quad \theta_1, \theta_2 \in \mathbb{R}. \tag{2.2}$$

By letting $\tilde{m}(|z|) = M(|z|)$, for any $z \in \mathbb{C}^*$, and $\Gamma(\exp(i\theta)) = M(\exp(i\theta))$ for any $\theta \in \mathbb{R}$ it follows that $\tilde{m} : (\mathbb{R}^+, \cdot) \rightarrow \mathbb{C}^*$ and $\Gamma : \{\exp(i\theta), \theta \in \mathbb{R}\} \rightarrow \mathbb{C}^*$ are characters. If $z = 0$, we set $M(z) = 0$. ■

In the next corollary we give the set of all multiplicative complex-valued functions $M : (\mathbb{R}^2, \cdot_\alpha) \rightarrow \mathbb{C}$.

COROLLARY 2.5. *The multiplicative functions $M : (\mathbb{R}^2, \cdot_\alpha) \rightarrow (\mathbb{C}, \cdot)$ are given by the following list:*

I) *If $\alpha > 0$, then*

$$M(x, y) = m_1(x + y\sqrt{\alpha})m_2(x - y\sqrt{\alpha}), \quad (x, y) \in \mathbb{R}^2.$$

II) *If $\alpha = 0$, then*

- a) $M(x, y) = 1$, for any $(x, y) \in \mathbb{R}^2$.
- b) $M(0, y) = 0$, for any $y \in \mathbb{R}$.
- c) $M(x, y) = m(x)\gamma(\frac{y}{x})$, for any $(x, y) \in \mathbb{R}^2$ with $x \neq 0$.

III) *If $\alpha < 0$, then*

- a) $M(x, y) = 0$, for any $(x, y) \in \mathbb{R}^2$.
- b) $M(x, y) = 1$, for any $(x, y) \in \mathbb{R}^2$.
- c) $M(x, y) = \begin{cases} \tilde{m}(\sqrt{x^2 - \alpha y^2})\Gamma(\arg(x + iy)), & \text{for } (x, y) \neq (0, 0) \\ 0, & \text{for } (x, y) = (0, 0). \end{cases}$

where $m_1, m_2, m : \mathbb{R} \rightarrow \mathbb{C}$ are multiplicative functions, and $\tilde{m} : (\mathbb{R}^+, \cdot) \rightarrow \mathbb{C}^*$, $\Gamma : \{\exp(i\theta), \theta \in \mathbb{R}\} \rightarrow \mathbb{C}^*$ and $\gamma : (\mathbb{R}, +) \rightarrow \mathbb{C}$ are arbitrary characters.

The next theorem is the main result of this section.

THEOREM 2.6. *The set of solutions of the functional equation $E(\alpha)$ consists of the following three cases:*

A) *If $\alpha > 0$, then*

$$f(x, y) = \frac{m_1(x)m_2(y)}{2} \{m_1(y\sqrt{\alpha})m_2(-y\sqrt{\alpha}) + m_1(-y\sqrt{\alpha})m_2(y\sqrt{\alpha})\},$$

for any $(x, y) \in \mathbb{R}^2$.

B) *If $\alpha = 0$, then*

a) $f(x, y) = 1$, for any $(x, y) \in \mathbb{R}^2$.

b) $f(0, y) = 0$, for any $y \in \mathbb{R}$.

c) $f(x, y) = \frac{m(x)}{2} \{ \gamma(\frac{y}{x}) + \gamma(\frac{-y}{x}) \}$, $(x, y) \in \mathbb{R}^2$, $x \neq 0$.

C) *If $\alpha < 0$, then $f(0, 0) = 0$ and*

$$f(x, y) = \frac{\tilde{m}(\sqrt{x^2 - \alpha y^2})}{2} \{ \Gamma(\arg(x + iy)) \}, (x, y) \in \mathbb{R}^2 \setminus (0, 0),$$

where $m_1, m_2, m : \mathbb{R} \rightarrow \mathbb{C}$ are multiplicative functions, and $\tilde{m} : (\mathbb{R}^+, \cdot) \rightarrow \mathbb{C}^*$, $\Gamma : \{ \exp(i\theta), \theta \in \mathbb{R} \} \rightarrow \mathbb{C}^*$ and $\gamma : \mathbb{R} \rightarrow \mathbb{C}$ are arbitrary characters.

Proof. According to Lemma 2.1 and Corollary 2.5 we get the proof of theorem. ■

3. SUPERSTABILITY OF EQUATION $E(\alpha)$

In the next theorem we establish the stability of $E(\alpha)$.

THEOREM 3.1. *Let $\varphi, \psi, \phi, \zeta : \mathbb{R} \rightarrow [0, +\infty[$ be functions and let $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ be a function such that*

$$\begin{aligned} & |f(x_1x_2 + \alpha y_1y_2, x_1y_2 + x_2y_1) + f(x_1x_2 - \alpha y_1y_2, x_2y_1 - x_1y_2) \\ & - 2f(x_1, y_1)f(x_2, y_2)| \leq \min\{\varphi(x_1), \psi(y_1), \phi(x_2), \zeta(y_2)\} \end{aligned} \quad (3.1)$$

for all $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$ and α is a real parameter. Then either f is bounded or f satisfies the functional equation

$$E(\alpha) : f(x_1x_2 + \alpha y_1y_2, x_1y_2 + x_2y_1) + f(x_1x_2 - \alpha y_1y_2, x_2y_1 - x_1y_2) = 2f(x_1, y_1)f(x_2, y_2)$$

for all $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$.

Proof. For all $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$ and α a real parameter we get from the inequality (3.1) that

$$\begin{aligned} &|f(x_1x_2 + \alpha y_1y_2, x_1y_2 + x_2y_1) + f(x_1x_2 - \alpha y_1y_2, x_2y_1 - x_1y_2) \\ &\quad - 2f(x_1, y_1)f(x_2, y_2)| \qquad (3.2) \\ &\leq \varphi(x_1) \text{ or } \psi(y_1). \end{aligned}$$

Since f is unbounded then we can choose a sequence $(x_n, y_n)_{n \geq 3}$ in \mathbb{R}^2 such that $f(x_n, y_n) \neq 0$ and $\lim_{n \rightarrow +\infty} |f(x_n, y_n)| = +\infty$. Taking $(x_2, y_2) = (x_n, y_n)$ in (3.2) we obtain

$$\begin{aligned} &|f(x_1x_n + \alpha y_1y_n, x_1y_n + x_ny_1) + f(x_1x_n - \alpha y_1y_n, x_ny_1 - x_1y_n) \\ &\quad - 2f(x_1, y_1)f(x_n, y_n)| \\ &\leq \varphi(x_1) \text{ or } \psi(y_1) \end{aligned}$$

and

$$\begin{aligned} &\left| \frac{f(x_1x_n + \alpha y_1y_n, x_1y_n + x_ny_1) + f(x_1x_n - \alpha y_1y_n, x_ny_1 - x_1y_n)}{2f(x_n, y_n)} - f(x_1, y_1) \right| \\ &\leq \frac{\varphi(x_1)}{2|f(x_n, y_n)|} \text{ or } \frac{\psi(y_1)}{2|f(x_n, y_n)|}. \end{aligned}$$

That is we get

$$\begin{aligned} &f(x_1, y_1) \\ &= \lim_{n \rightarrow +\infty} \frac{f(x_1x_n + \alpha y_1y_n, x_1y_n + x_ny_1) + f(x_1x_n - \alpha y_1y_n, x_ny_1 - x_1y_n)}{2f(x_n, y_n)}. \end{aligned} \tag{3.3}$$

Setting $X_n = x_2x_n + \alpha y_2y_n$, $Y_n = x_2y_n + x_ny_2$, $\tilde{X}_n = x_2x_n - \alpha y_2y_n$, $\tilde{Y}_n = x_2y_n - x_ny_2$. For any $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$ it follows that

$$\begin{aligned}
& |f((x_1x_2 + \alpha y_1y_2)x_n + \alpha(x_1y_2 + x_2y_1)y_n, \\
& \quad (x_1x_2 + \alpha y_1y_2)y_n + x_n(x_1y_2 + x_2y_1)) \\
& + f((x_1x_2 + \alpha y_1y_2)x_n - \alpha(x_1y_2 + x_2y_1)y_n, \\
& \quad x_n(x_1y_2 + x_2y_1) - (x_1x_2 + \alpha y_1y_2)y_n) \\
& - 2f(x_1, y_1)f(x_2x_n + \alpha y_2y_n, x_2y_n + x_ny_2) \\
& + f((x_1x_2 - \alpha y_1y_2)x_n + \alpha(x_2y_1 - x_1y_2)y_n, \\
& \quad (x_1x_2 - \alpha y_1y_2)y_n + x_n(x_2y_1 - x_1y_2)) \\
& + f((x_1x_2 - \alpha y_1y_2)x_n - \alpha(x_2y_1 - x_1y_2)y_n, \\
& \quad x_n(x_2y_1 - x_1y_2) - (x_1x_2 - \alpha y_1y_2)y_n) \\
& - 2f(x_1, y_1)f(x_2x_n - \alpha y_2y_n, x_2y_n - x_ny_2)| \\
\leq & |f((x_1x_2 + \alpha y_1y_2)x_n + \alpha(x_1y_2 + x_2y_1)y_n, \\
& \quad (x_1x_2 + \alpha y_1y_2)y_n + x_n(x_1y_2 + x_2y_1)) \\
& + f((x_1x_2 - \alpha y_1y_2)x_n - \alpha(x_2y_1 - x_1y_2)y_n, \\
& \quad x_n(x_2y_1 - x_1y_2) - (x_1x_2 - \alpha y_1y_2)y_n) \\
& - 2f(x_1, y_1)f(x_2x_n + \alpha y_2y_n, x_2y_n + x_ny_2)| \\
& + |f((x_1x_2 - \alpha y_1y_2)x_n + \alpha(x_2y_1 - x_1y_2)y_n, \\
& \quad (x_1x_2 - \alpha y_1y_2)y_n + x_n(x_2y_1 - x_1y_2)) \\
& + f((x_1x_2 + \alpha y_1y_2)x_n - \alpha(x_1y_2 + x_2y_1)y_n, \\
& \quad x_n(x_1y_2 + x_2y_1) - (x_1x_2 + \alpha y_1y_2)y_n) \\
& - 2f(x_1, y_1)f(x_2x_n - \alpha y_2y_n, x_2y_n - x_ny_2)| \\
= & |f(x_1X_n + \alpha y_1Y_n, x_1Y_n + X_ny_1) + f(x_1X_n - \alpha y_1Y_n, X_ny_1 - x_1Y_n) \\
& \quad - 2f(x_1, y_1)f(X_n, Y_n)| \\
& + |f(x_1\tilde{X}_n + \alpha y_1\tilde{Y}_n, x_1\tilde{Y}_n + \tilde{X}_ny_1) + f(x_1\tilde{X}_n - \alpha y_1\tilde{Y}_n, \tilde{X}_ny_1 - x_1\tilde{Y}_n) \\
& \quad - 2f(x_1, y_1)f(\tilde{X}_n, \tilde{Y}_n)| \\
\leq & 2\varphi(x_1) \text{ or } 2\psi(y_1).
\end{aligned}$$

So that

$$\begin{aligned}
 & \left| \frac{f((x_1x_2 + \alpha y_1y_2)x_n + \alpha(x_1y_2 + x_2y_1)y_n, (x_1x_2 + \alpha y_1y_2)y_n + x_n(x_1y_2 + x_2y_1))}{f(x_n, y_n)} \right. \\
 & + \frac{f((x_1x_2 + \alpha y_1y_2)x_n - \alpha(x_1y_2 + x_2y_1)y_n, x_n(x_1y_2 + x_2y_1) - (x_1x_2 + \alpha y_1y_2)y_n)}{f(x_n, y_n)} \\
 & + \frac{f((x_1x_2 - \alpha y_1y_2)x_n + \alpha(x_2y_1 - x_1y_2)y_n, x_n(x_2y_1 - x_1y_2) + (x_1x_2 - \alpha y_1y_2)y_n)}{f(x_n, y_n)} \\
 & + \frac{f((x_1x_2 - \alpha y_1y_2)x_n - \alpha(x_2y_1 - x_1y_2)y_n, x_n(x_2y_1 - x_1y_2) - (x_1x_2 - \alpha y_1y_2)y_n)}{f(x_n, y_n)} \\
 & \left. - 2f(x_1, y_1) \left\{ \frac{f(x_2x_n + \alpha y_2y_n, x_2y_n + x_ny_2) + f(x_2x_n - \alpha y_2y_n, x_2y_n - x_ny_2)}{f(x_n, y_n)} \right\} \right| \\
 & \leq 2 \frac{\varphi(x_1)}{|f(x_n, y_n)|} \text{ or } 2 \frac{\psi(y_1)}{|f(x_n, y_n)|}.
 \end{aligned}$$

for any $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$. Since $|f(x_n, y_n)| \rightarrow +\infty$ as $n \rightarrow +\infty$ we get that f satisfies $E(\alpha)$. ■

By letting $\min\{\varphi(x_1), \psi(y_1), \phi(x_2), \zeta(y_2)\} = \delta$ we get the Baker's stability ([3], [4]) for the functional equation $E(\alpha)$.

COROLLARY 3.2. *Let $\delta > 0$ and let $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ be a function such that*

$$\begin{aligned}
 & |f(x_1x_2 + \alpha y_1y_2, x_1y_2 + x_2y_1) + f(x_1x_2 - \alpha y_1y_2, x_2y_1 - x_1y_2) \\
 & - 2f(x_1, y_1)f(x_2, y_2)| \leq \delta
 \end{aligned}$$

for all $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$ and α is a real parameter. Then either f is bounded and $|f(x, y)| \leq \frac{1+\sqrt{1+2\delta}}{2}$ for all $(x, y) \in \mathbb{R}^2$ or f satisfies the functional equation $E(\alpha)$.

REFERENCES

- [1] J. ACZÉL, J. DHOMBRES, “Functional Equations in Several Variables”, Encyclopedia of Mathematics and its Applications 31, Cambridge University Press, Cambridge, 1989.
- [2] R. BADORA, R. GER, On some trigonometric functional inequalities, in “Functional Equation-Results and Advances”, Adv. Math. (Dordr.) 3, Kluwer Acad. Publ., Dordrecht, 2002, 3–15.
- [3] J. BAKER, J. LAWRENCE, F. ZORZITTO, The stability of the equation $f(x+y) = f(x)f(y)$, *Proc. Amer. Math. Soc.* **74** (2) (1979), 242–246.
- [4] J. BAKER, The stability of the cosine equation, *Proc. Amer. Math. Soc.* **80** (3) (1980), 411–416.
- [5] L. R. BERRONE, L. V. DIEULEFAIT, A functional equation related to the product in a quadratic number field, *Aequationes Math.* **81** (1-2) (2011), 167–175.
- [6] B. BOUIKHALENE, E. ELQORACHI, J. M. RASSIAS, The superstability of d’Alembert’s functional equation on the Heisenberg group, *Appl. Math. Lett.* **23** (1) (2010), 105–109.
- [7] D. G. BOURGIN, Approximately isometric and multiplicative transformations on continuous function rings, *Duke. Math. J.* **16** (2) (1949), 385–397.
- [8] P. W. CHOLEWA, The stability of sine equation, *Proc. Amer. Math. Soc.* **88** (4) (1983), 631–634.
- [9] T. M. K. DAVISON, D’Alembert’s functional equation on topological monoids, *Publ. Math. Debrecen* **75** (1-2) (2009), 41–66.
- [10] Z. GAJDA, On stability of additive mappings, *Internat. J. Math. Math. Sci.* **14** (3) (1991), 431–434.
- [11] P. GĂVRUTA, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, *J. Math. Anal. Appl.* **184** (3) (1994), 431–436.
- [12] D. H. HYERS, On the stability of the linear functional equation, *Proc. Nat. Acad. Sci. U.S.A.* **27** (1941), 222–224.
- [13] D. H. HYERS, G. ISAC, TH. M. RASSIAS, “Stability of Functional Equations in Several Variables”, Progress in Nonlinear Differential Equations and their Applications 34, Birkhäuser Boston, Inc, Boston, 1998.
- [14] S. M. JUNG, J. H. BAE, Some functional equations originating from number theory, *Proc. Indian Acad. Sci. Math. Sci.* **113** (2) (2003), 91–98.
- [15] PL. KANNAPPAN, On the functional equation $f(x+y) + f(x-y) = 2f(x)f(y)$, *Amer. Math. Monthly* **72** (1965), 374–377.
- [16] G. H. KIM, On the stability of trigonometric functional equations, *Adv. Difference Equ.* (2007), Art. ID 90405, 10 pp.
- [17] G. H. KIM, A stability of the generalized sine functional equations, *J. Math. Anal. Appl.* **331** (2) (2007), 886–894.
- [18] G. H. KIM, On the stability of generalized D’Alembert and Jensen functional equations, *Int. J. Math. Math. Sci.* **2006**, Article ID 43185, (2006), 1–12.
- [19] J. M. RASSIAS, On approximation of approximately linear mapping by linear

- mappings, *J. Funct. Anal.* **46** (1) (1982), 126–130.
- [20] TH. M. RASSIAS, On the stability of the linear mapping in Banach spaces, *Proc. Amer. Math. Soc.* **72** (2) (1978), 297–300.
- [21] L. SZÉKELYHIDI, On a theorem of Baker, Lawrence and Zorzitto, *Proc. Amer. Math. Soc.* **84** (1) (1982), 95–96.
- [22] S. M. ULAM, “A Collection of Mathematical Problems”, Interscience Tracts in Pure and Applied Mathematics 8, Interscience Publishers, New York-London, 1960.
- [23] W. H. WILSON, On certain related functional equations, *Bull. Amer. Math. Soc.* **26** (7) (1920), 300–312.