

TRABAJOS ORIGINALES

Vascular flora and phytogeographical links of the Carabaya Mountains, Peru

Flora vascular y conexiones fitogeográficas de las montañas Carabaya, Perú

Paúl Gonzáles¹, Blanca León^{1,2,3}, Asunción Cano^{1,2*} and Peter M. Jørgensen⁴

1 Universidad Nacional Mayor de San Marcos, Museo de Historia Natural, Departamento de Dicotiledóneas. Av. Arenales 1256, Lima-14, Perú.

2 Universidad Nacional Mayor de San Marcos, Facultad de Ciencias Biológicas, Instituto de Investigación de Ciencias Biológicas Antonio Raimondi.

3 Department of Geography and the Environment, University of Texas at Austin, Austin, TX 78712-0530, USA.

4 Missouri Botanical Garden-St. Louis, MO 63166, USA

* Corresponding author

E-mail Paúl Gonzáles: pgonzalesarce@hotmail.com

E-mail Asunción Cano: acano@unmsm.edu.pe

E-mail Blanca León: leon@austin.utexas.edu

E-mail Peter M. Jørgensen: peter.jorgensen@mobot.org

Abstract

Studies of floristic composition and plant species richness in tropical mountains support their recognition as areas of high biological diversity, and therefore of their importance for plant conservation. Here, we present data on the flora of the high Andes of eight sites centered in the Carabaya mountains, and also provide a floristic comparison with nine other floras within Peru and northern Bolivia. The study area includes 506 species of vascular plants, grouped in 203 genera and 66 families. The highest species richness was found in two families: Asteraceae and Poaceae, which collectively encompass 37% of all species. Other important families were Caryophyllaceae, Fabaceae, Malvaceae, Brassicaceae, Caprifoliaceae, Gentianaceae, Plantaginaceae and Cyperaceae. The most diverse genera were *Senecio*, *Calamagrostis*, *Poa* and *Nototriche*. Perennial herbs were the dominant growth form. The vascular flora of the Carabaya Mountains is closely related to those of other regions of southern Peru. Also, more than half of all vascular plants registered for the Carabaya Mountain occur in the Andean region of Bolivia, which shows the undoubted geophysical and phytogeographical connection of the Carabaya and the Bolivian Apolobamba Mountains. This study also shows that there is still a need for more extensive plant collecting and future exploration, since the Carabaya, as other parts of Peru's high Andes are subject of dramatic change that may threaten these plant populations.

Keywords: High Andean flora; Peru; floristic composition; taxonomy.

Resumen

Los estudios sobre la composición florística y riqueza de especies en montañas tropicales apoyan su reconocimiento como áreas de alta diversidad biológica, y, por tanto, de su importancia para la conservación. En este trabajo presentamos datos sobre la flora altoandina de ocho sitios localizados en la Cordillera de Carabaya, proveemos también una comparación florística con otros nueve lugares tanto en Perú como en el norte de Bolivia. El área de estudio incluye 506 especies de plantas vasculares, reconocidas en 203 géneros y 66 familias. Las tasas más altas de riqueza de especies se hallan en dos familias: Asteraceae y Poaceae, que colectivamente abarcan el 37% de todas las especies. Otras familias importantes fueron Caryophyllaceae, Fabaceae, Malvaceae, Brassicaceae, Caprifoliaceae, Gentianaceae, Plantaginaceae y Cyperaceae. Los géneros más diversos fueron *Senecio*, *Calamagrostis*, *Poa* y *Nototriche*. La forma de crecimiento predominante fueron las hierbas perennes. La flora vascular de la Cordillera Carabaya está muy relacionada con otras regiones del sur de Perú. Además, más de la mitad de todas las plantas vasculares registradas para la Cordillera Carabaya se encuentran en la región andina de Bolivia, lo que demuestra la indudable conexión geofísica y fitogeográfica entre las cordilleras Carabaya y Apolobamba de Bolivia. Este estudio también demuestra la necesidad de una extensa colección botánica y futura exploración, desde que Carabaya, como otras partes de los altos Andes del Perú, están sujetos a cambios dramáticos que amenazan las poblaciones de esas plantas.

Palabras clave: flora altoandina; Perú; composición florística; taxonomía.

Citación:

Gonzáles P., B. León, A. Cano & P.M. Jørgensen. 2018. Vascular flora and phytogeographical links of the Carabaya Mountains, Peru. *Revista peruana de biología* 25(3): 191 - 210 (Agosto 2018). doi: <http://dx.doi.org/10.15381/rpb.v25i3.15228>

Presentado: 26/01/2018

Aceptado: 03/06/2018

Publicado online: 25/09/2018

Declaration of authorship

The authors state that all participated in the development of the work. PG, BL, AC and PJ: did sampling, drafting of the manuscript, and data analysis. PG and BL: did interpretation and preparation of the final version.

The authors declare that they have no conflict of interest.

Journal home page: <http://revistasinvestigacion.unmsm.edu.pe/index.php/rpb/index>

© Los autores. Este artículo es publicado por la Revista Peruana de Biología de la Facultad de Ciencias Biológicas, Universidad Nacional Mayor de San Marcos. Este es un artículo de acceso abierto, distribuido bajo los términos de la Licencia Creative Commons Atribución-NoComercial-CompartirIgual 4.0 Internacional. (<http://creativecommons.org/licenses/by-nc-sa/4.0/>), que permite el uso no comercial, distribución y reproducción en cualquier medio, siempre que la obra original sea debidamente citadas. Para uso comercial, por favor póngase en contacto con editor.revperubiol@gmail.com.

Introduction

The high Andes are places of concern due to the foreseeable effects of climate change (Markham et al. 1993, Beniston 1994, Thompson et al. 2006, Conde & Saldaña 2007, Pauli et al. 2003, 2007, Young 2014). In Peru, high Andean sites cover close to 16% of the total country area, and they are highly vulnerable to landscape changes that may affect its vegetation and components (Rodríguez & Young 2000, Young 2011). The urgency for the development of conservation strategies for sites and plant species is widely recognized; however, it faces challenges in relation to the dynamism and complexity of environmental changes and their interactions with human influences (Markham et al. 1993, Fort 2015, Kohler et al. 2014).

The Peruvian high Andes (above 3500 m) includes an interesting native vascular flora estimated to consist of more than 2000 plant species (see Jørgensen et al. 2011), of which nearly 32% of them are endemic to Peru. This region has recently seen an increase in taxonomical and ecological studies, accompanied by plant recording and exploration (e.g. Ballard and Iltis 2012, Al-Shehbaz et al. 2013, 2015a, 2015b, González et al. 2015, González & Cano 2016, Montesinos-Tubée et al. 2015, Linares et al. 2015, Ospina et al. 2016, Sylvester et al. 2016a, 2016b). The completion of the Bolivian catalog (Jørgensen et al. 2014) also provides needed data for a floristic comparison to understand plant richness in the Andes, and floristic connections. Here, we present a compilation of the vascular plant flora of a mountain range in southern Peru: the Carabaya Mountains. We asked how taxonomic richness of our study area compares to other parts of the Andes, particularly to neighboring sites in Peru, and northern Bolivia. The Carabaya is geographically linked to Bolivia through the Apolobamba Mountains, and so we asked if richness and composition at high elevations are similar. We also explore the potential implications of our findings for plant conservation based on the state of knowledge of the high Andean flora.

Material and methods

Study area.- The study area encompasses the high Andes between 4000–5300 m of the Carabaya Cordillera, along the eastern Andes in Puno, between 13°50'–15°05' S and 69°58'–70°58' W, crossing the provinces of Carabaya, Sandia and San Antonio de Putina (Fig. 1). The Carabaya also includes the largest tropical glacier system: Quelccaya, an area that has contributed to our understanding of climate change (Thompson et al. 2003, 2006, 2013). The Carabaya Mountains extends into Bolivian territory, and in the Department of La Paz, the northeastern Andes are known as the Apolobamba Mountains (Argollo et al. 1987, Argollo & Iriondo 2008). We obtained field data from eight sites (Fig. 1).

In the Carabaya we recognize four vegetation formations: puna grasslands; rock vegetation dominated by chasmophytes; high Andean wetlands; river shores and lakes, and periglacial areas including cryoturbated soils and subnival pumices (Weberbauer 1945, Galán de Mera et al. 2014, Montesinos-Tubée et al. 2015a).

Collection and species identification.- We compiled all species names, habitats, and distribution information initially based on specimens collected by the authors for the study area. For each species from the Carabaya, we gathered information on

collectors and collection number. However, for a small group of species that had only been recorded by photo or by authors' field notes, and lack a herbarium voucher, we cited other collections made in neighboring areas. All collections from the study area were identified using taxonomical keys and descriptions available from the botanical literature (Macbride 1936, Tovar 1993, and others). A few reported species were excluded, for example a photo of *Stangea rhizantha* did not match the species (Linares et al. 2012), and *Valeriana globularis* (Linares et al. 2012) was replaced by its correct spelling, *Valeriana globularioides* Graebn.

We also compiled all records for eight floras at high elevations, three sites (Parinacochas, Juli and Tambo Ichuna) were located above 3100 m elevation on the Pacific Basin, and the remaining five sites (from north to south: Cordillera Blanca, Concepcion, Bofedales, Apacheta and Vilcanota) were located above 4000 m. Species considered as national endemics were recorded from León et al. (2007), Al-Shehbaz et al. (2013, 2015a, 2015b), González et al. (2015), Montesinos-Tubée et al. (2015b), and González and Cano (2016).

For the flora of the Department of La Paz in northern Bolivia, we downloaded the species records from the Bolivia Catalog database (<http://www.tropicos.org/ProjectAdvSearch.aspx?projectid=13&langid=66>).

Specimen data were also obtained from online sources of several institutions (e.g., Field Museum Neotropical specimens [<http://fm1.fieldmuseum.org/vrrc/>], the Tropicos® database of Missouri Botanical Garden [<http://www.tropicos.org/>], New York Botanical Garden digital herbarium [<http://sciweb>].




Figure 1. Map of the study area. (Puno, Peru). Locality (1) Corani, (2) Macusani, (3) Ajoyani, (4) Cruzero, (5) Patambuco, (6) Sandia, (7) Cuyocuyo, (8) Ananea.

nybg.org/Science2/vii2.asp]), the digital herbarium portion of Atrium [http://atrium.andesamazon.org/digital_herbarium.php], JSTOR [https://plants.jstor.org/]. We also included or reviewed collections of the following herbaria B, BM, CUZ, F, G, GH, HUSA, HUT, K, LIL, LP, MO, MOL, NY, S, US and USM (Index Herbariorum). The Angiosperm Phylogeny Group IV (2016) classification system was used for the management of the angiosperm taxa.

Floristic comparison.— Our database of all taxa was compared with checklists of nine studies in the high Andes. These nine sites encompass about seven degrees of latitude (8°50' – 17°26' S), and represent both Pacific and Amazon basins. Within these sites there are a variety of habitats from open grasslands, rocky places, shrublands and wetlands. Four floras represent mostly wetlands and surrounding areas (Flores et al. 2005, Arteta et al. 2006, Roque & Ramírez 2008, Valencia et al. 2013), two floras represent sites located in periglacial areas with processes of cryoturbation (Cano et al. 2010, 2011), and four floras represent large or detailed surveys in the puna ecosystem (Tupayachi 2005, Montesinos-Tubée 2011, 2012, Jørgensen et al. 2014).

Results

We recorded over 10000 plant collections for all nine sites in the Peruvian Andes. This sampling effort still represents a low collection density index with a CDI (specimens/100 km²) of nearly 5. For southern Peru, and particularly for Puno (up to 4000 m), the largest plant collections were made by Augusto Weberbauer, César Vargas, Hugh Weddell, Franz Meyen, Jaroslav Soukup, Francis Pennell, Werner Rauh, Wilibald Lechler, Alfredo Tupayachi, Oscar Tovar and Dora Stafford, all of them made during the 19th and 20th centuries.

The highest elevations (above 4000 m) of the Carabaya Mountains harbors 506 species of vascular plants (pteridophytes, Gymnosperms, Eudicots and Monocots) grouped in 203 genera and 66 families (Table 1; Fig. 2 and Appendix 1). Eudicots were the largest group (with 68.2% of the total registered), and including 69.5% of genera and 71.9% of all species; followed by Monocots with 16.7%, 22.7% and 23.7% respectively for all three main taxonomic categories. Ferns and lycophytes were scarcely represented with only nine families (13.6%), fifteen genera (7.4%) and twenty-one species (4.2%). For these elevations, we found only one species of gymnosperm (*Ephedra rupestris*).

Asteraceae was the most diverse family, with 38 genera and 110 species, followed by Poaceae (21/78), Caryophyllaceae (10/24), Fabaceae (6/23), Malvaceae (3/21), Brassicaceae (11/21), Caprifoliaceae (3/17), Gentianaceae (3/16), Plantaginaceae (4/13) and Cyperaceae (7/12). These ten families include 52% of all genera and 66% of all species registered for Carabaya. Asteraceae and Poaceae together account for 29.1% of the total genera and 37.2% of all species (Table 2). Another 10 families have six to ten species, containing 16.2% of the total taxa registered. Sixty-six percent of the remaining families include five or less number of species: 24 families have between two and five species, containing 13.2% of total taxa, while 22 families are represented by a single species.

The most diverse genera were *Senecio* (28 species), *Calamagrostis* (21), *Poa* (21), *Nototriche* (19), *Gentianella* (13), *Valeriana* (13), *Werneria* (10), *Lupinus* and *Astragalus* with nine species each, *Festuca* and *Plantago* with eight species each, *Mniodes* and

Ranunculus with seven each; *Geranium*, *Neobartsia*, *Oxalis*, *Perezia*, *Pycnophyllum*, and *Xenophyllum* with six each; *Cerastium*, *Hypochaeris*, *Sisyrinchium* and *Viola* with five each (Table 3). The remaining, 62 genera (30.5%), include between two and four species, while 117 genera (57.6%) only have one species.

Perennial herbs are the dominant life form with 467 (92.3%) species, these include aquatics, palustrine and terrestrial forms. While woody life forms include 20 species (4%) of subshrubs, 15 (3%) shrubs, three (0.6%) cactus, and one tree (0.2%) (Appendix 1).

Comparison of Carabaya flora with nine sites, including

Table 1. Number of families, genera and species by taxa recorded in the study area.

Taxa	Families	Genera	Species
Eudicots	45	141	364
Monocots	11	46	120
Gymnosperms	1	1	1
Pteridophytes	9	15	21
Total	66	203	506

Table 2. Families with more genera and species in the Cordillera Carabaya.

Families	Genera	Species
Asteraceae	38	110
Poaceae	21	78
Caryophyllaceae	10	24
Fabaceae	6	23
Malvaceae	3	21
Brassicaceae	11	21
Caprifoliaceae	3	17
Gentianaceae	3	16
Plantaginaceae	4	13
Cyperaceae	7	12

Table 3. Genera with more species in the Cordillera Carabaya.

Genera	Species	Genera	Species
<i>Senecio</i>	28	<i>Mniodes</i>	7
<i>Calamagrostis</i>	21	<i>Pycnophyllum</i>	6
<i>Nototriche</i>	19	<i>Bartsia</i>	6
<i>Poa</i>	19	<i>Perezia</i>	6
<i>Gentianella</i>	13	<i>Geranium</i>	6
<i>Valeriana</i>	13	<i>Oxalis</i>	6
<i>Werneria</i>	10	<i>Lachemilla</i>	6
<i>Lupinus</i>	9	<i>Xenophyllum</i>	6
<i>Astragalus</i>	9	<i>Cerastium</i>	5
<i>Plantago</i>	8	<i>Hypochaeris</i>	5
<i>Festuca</i>	8	<i>Sisyrinchium</i>	5
<i>Ranunculus</i>	7	<i>Viola</i>	5



Figure 2. a) *Hypochaeris echegarayi* (Asteraceae); b) *Werneria heteroloba* (Asteraceae); c) *Stangea paulae* (Caprifoliaceae); d) *Gentiana sedifolia* (Gentianaceae); e) *Nototriche staffordiae* (Malvaceae); f) *Myrosmodes paludosa* (Orchidaceae); g) *Poa apiculata* (Poaceae); h) *Salpichroa amoena* (Solanaceae).

northern Bolivia reveals that Asteraceae and Poaceae remain the most species-rich families. All sites, except Parinacochas, have most of its genera and species (over 50%) included in the same ten families identified in Table 4. Other sites representing specific habitats (periglacial areas, and wetlands), also have the same species-rich families as those found in Carabaya (see also Table 2). The richness of all eight sites of Carabaya is as high as those found in northern Bolivia and Moquegua in southwestern Peru.

Over 91 % of species recorded for Carabaya have an altitudinal range above 4000 m (Append. 1). Plant endemism in the Department of Puno reaches 185 species, and most species are found at the upper limits of our study area (Append. 1).

Discussion

The high Andes is an important biogeographical region, distributed as an archipelago along the mountain range. Its plant species richness is high compared to other mountain ranges, and where past geological changes has allowed expansion and contraction, which may explain current plant composition, diversity and endemism (Sklenář et al. 2013, Hughes & Atchison 2015). Most recent studies have shown the northern Andes, particularly the paramo, as a place of high plant species richness (Sklenář & Balslev 2005, 2007, Cuesta et al. 2017). Our data also demonstrates that other parts of the upper tropical central Andes are as important in terms of plant composition and diversity.

Over 50% of floristic records for the study area are new, particularly for the department of Puno (Brako & Zarucchi 1993, Ulloa Ulloa et al. 2004, Linares et al. 2012). For the flora at high elevations (≥4000 m) of Puno, our results have tripled (20 more species) the known number of endemic species. It is

worth noting that this department has the highest percentage of rarity (65%) compared to other departments (León et al. 2007). The value of the CDI is lower than the national value of 34 recorded almost 30 years ago (see Toledo & Sosa 1993), and it demonstrates the need for further botanizing. This need is further complicated by pressures of global changes, creating important challenges for plant conservation (Young 2014).

The flora of the Carabaya, at the family level, includes two families (Asteraceae and Poaceae) that contribute significantly to its composition with over a third of species. The predominance of both families is a common pattern in the High Andean flora (Gentry 1993, Cuesta et al. 2017), where these two families can include 30 to 60% of all species in an area (Table 4). In addition, this trend of family oligarchy extends in our study to 18 other families, among them Fabaceae, Caryophyllaceae, Brassicaceae and Malvaceae, which has been shown to be species rich throughout the Puna (Cuesta et al. 2017). Most families, however, are species poor, and they usually represent taxa that originated in temperate areas. Probable explanations of species richness of those few families may include geological history, altitudinal range and different evolutionary paths of mountain ranges within the Andes.

A total of 506 species of vascular plants growing above 4000 m in the Carabaya shows that our floristic results are consistent with those found for other equivalent areas in the tropical high Andes. Species richness in our study area is like those found for Peru's jalca (Sánchez 1996). For the Andean highlands of northern Bolivia (Department of La Paz) reported by Jørgensen (2014), total species (737) number is larger than for our study area, which can be explained by representing a flora of a bigger

Table 4. Comparing the number of species (E) and genera (G) of the principal families in different Andean localities (¹Flores et al. 2005, ²Arteta et al. 2006, ³Roque & Ramírez 2008, ⁴Valencia et al. 2013, ⁵Tupayachi 2005, ⁶Cano et al. 2011, ⁷Cano et al. 2010, ⁸Montesinos–Tubée 2011, 2012, ⁹Jørgensen et al. 2014), with respect to what was found in the study area.

	Concepción ¹ 4350–4550 m		Bahía de Juli ² 3830–4200 m		Parinacochas ³ 3100–3500 m		Bofedales ⁴ 4200–4800 m		Vilcanota ⁵ >4000		Apacheta ⁶ >4500 m		Cordillera Blanca ⁷ >4500 m		Tambo-Ichuña ⁸ 3400–4850 m		Carabaya >4000 m		Bolivia ⁹ >4000 m	
Familias	G	E	G	E	G	E	G	E	G	E	G	E	G	E	G	E	G	E	G	E
Asteraceae	15	24	27	39	30	43	15	22	15	40	20	51	15	40	52	127	38	110	50	189
Poaceae	10	25	12	18	16	26	7	23	9	25	6	26	11	31	28	46	21	78	30	129
Brassicaceae	1	1	5	7	3	4	1	1	2	2	5	11	6	15	13	24	11	21	14	35
Caryophyllaceae	4	4	5	5	3	3	4	5	3	3	4	8	5	7	10	18	10	24	12	50
Fabaceae	1	3	5	8	8	14	2	2	0	0	2	4	1	1	9	18	6	23	13	63
Malvaceae	1	2	2	3	2	2	2	2	2	5	1	7	1	4	4	19	3	21	4	28
Plantaginaceae	1	3	1	3	1	1	2	3	3	3	0	0	0	0	2	12	4	13	5	13
Apiaceae	1	1	4	5	4	4	5	7	1	2	3	4	3	3	8	11	6	9	9	18
Cyperaceae	2	2	2	5	5	5	8	10	1	2	0	0	0	0	8	8	7	12	10	28
Caprifoliaceae	0	0	0	0	1	1	1	1	1	8	3	6	3	7	3	9	3	17	5	13
Others familias	28	35	74	92	106	122	28	50	32	34	16	17	20	28	134	215	94	178	120	271
% species by 10 families	56	65	46	50	41	46	63	60	54	73	73	87	69	79	51	58	54	65	56	68
Total	64	100	137	185	179	225	75	126	69	124	60	134	65	136	271	507	203	506	272	837

area with more extensive botanizing.

As proposed by Sklenář et al. (2013), floristic similarities in the Andes are higher in areas of geographical proximity. The undoubted geophysical connection of the Carabaya and the Bolivian Apolobamba Mountains, may suggest a close floristic relationship with Bolivia's high Andean flora. We found nearly half of all vascular plants (262 species) registered for the Carabaya Mountain also occur in the Andean region of Bolivia. At the family level, eight families Asteraceae, Poaceae, Fabaceae, Caryophyllaceae, Brassicaceae, Malvaceae, Cyperaceae and Apiaceae are part of the most diverse families in both study areas (Tabla 4). However, there are thirteen families (Araceae, Araliaceae, Basellaceae, Bromeliaceae, Calyceraceae, Ericaceae, Dennstaedtiaceae, Frankeniaceae, Polygalaceae, Portulacaceae, Ruppiaceae, Schoepfiaceae and Selaginellaceae) that have not yet been collected in the Carabaya. While another four families (Asparagaceae, Escalloniaceae, Convolvulaceae and Berberidaceae) have not yet been recorded for the Bolivian flora at similar altitudes

There are also 35 genera and 244 species in Carabaya not registered for Bolivia, the largest of these differences are present in genera such as *Senecio* (with 15 species), *Nototriche* (13), *Valeriana*, *Calamagrostis* and *Gentianella* (with 9 species each), *Poa* (7), *Xenophyllum* (6), *Lupinus*, *Neobartsia* and *Pycnophyllum* (with 5 species each). Most of these species (186) are endemic to Peru, and they occupy the highest part of the study area.

A comparison with the species composition of Cordillera Vilcanota, located to the north of our study area, reveals that 46% of the species registered there have not yet been found in Carabaya. Studies in Vilcanota have emphasized areas covered by *Polylepis* forest that were not included for Carabaya. Still, excluding those species of *Polylepis* forests, a third of species of the flora of Vilcanota have not been recorded in Carabaya.

At a more local scale, we found that Carabaya has similar number of taxa as those reported for an upper basin in Moquegua (Montesinos-Tubée 2011, 2012; Table 4) on the Pacific Basin. Seven plant families: Asteraceae, Poaceae, Brassicaceae, Caryophyllaceae, Fabaceae, Malvaceae and Plantaginaceae are part of the most diverse families in both study areas (Tabla 4). Two families Asteraceae and Poaceae have also equal representation for both sites, with 22%, and 15% for Carabaya, and 25% and 9% for Tambo-Ichuña respectively. However, some differences at the family level are present due to the altitudinal extent of our study (4000–5300 m) vs. those of Montesinos-Tubée (3400–4850 m). For example Cyperaceae and Caprifoliaceae are among the top ten most diverse families in the Carabaya, while Solanaceae and Cactaceae are for Moquegua (Montesinos-Tubée 2011, 2012). These differences may also reflect floristic differences between western and southern ranges.

The top ten most diverse families have also been reported by Cano et al. (2010, 2011) for periglacial sites with cryoturbated soils in Ancash (Cordillera Blanca) and Ayacucho-Huancavelica (Apacheta), reaching 79 and 87% respectively (Table 4). However, these families decrease their diversity at lower elevation (Flores et al. 2005, Arteta et al. 2006, Roque & Ramírez 2008), or when the richness is restricted to specific vegetation formations such as wetlands (Ruthsatz 2012, Valencia et al. 2013).

For the flora of Carabaya, the most abundant genera in

terms of species richness were *Senecio*, *Calamagrostis*, *Nototriche* and *Poa*; this taxonomic composition towards a few highly diverse genera has been explained due to the high degree of a recent local speciation driven by geographic isolation between high alpine 'continental islands' or complexes (Sklenář & Balslev 2005, Sklenář et al. 2013). In addition, these same genera are the most abundant in terms of vegetation cover (Cuesta et al. 2017).

A perennial life form is obviously the dominant plant type in the high Andes. This life form includes most species in plant lineages with rapid diversification and radiation in the Andes, such as *Lupinus*, *Nototriche*, *Pycnophyllum*, *Senecio* and others. Hughes and Atchison (2015) suggest "that perenniality could have played a general role as a key adaptation, enabling lineages to take advantage of ecological opportunities associated with the recent availability of alpine habitats to rapidly diversify" as was observed in *Lupinus* (Drummond et al. 2012). In this study, half of the Carabaya flora has longer life cycles as "Perennials" represented by herbaceous forms. However, a more detailed exam might allow recognition of other types such as rosettes, mats, and cushions.

Because of the availability of data from local, regional and country floras further comparisons are now feasible. The similarities in floristic composition and richness confirm the importance of the tropical Andes as global hotspot of plant diversity. Differences in flora sizes, especially between Carabaya and its closest neighboring sites, demonstrate the need of extending botanical exploration in Peru, and for the detailed recording in different types of habitats. Plant collections are still needed in many areas to fully understand the patterns of species diversity, and characteristics of their populations. As it can be found in the results of this study, several species have not been found since their discovery (e.g., *Alsine rupestris* Muschl., *Nototriche cupuliforme* Krapov. and *Nototriche erinacea* A.W. Hill). The high number of species (>91%) with its lower limits above 4000 m underscores the importance of the high Andes for conservation. This is even more important today in the Peruvian high Andes, where climatic conditions and human induced changes play an important role for impacting them (see Young 2014). Threats originate directly by overlapping competing human activities such as mining, fire, agriculture, landslides, erosion and other land uses. In addition, climate change associated to these anthropic influences are leading to the fragmentation of habitats and therefore, the loss of species survival and diversity, and possible mountaintop extinctions. Finally, it should be mentioned that many taxonomic studies are still pending to fully understand the biodiversity of the fascinating Andean highlands.

Acknowledgements

We thank Daniel Montesinos for providing valuable comments on the manuscript. We also thank Kenneth R. Young for his comments and critical review of this manuscript. Curators of all the herbaria cited are thanked for making available their plant collections. We also acknowledge all the people who made the botanical collections in the studied region.

Literature cited

- Al-Shehbaz I.A., A. Cano, H. Trinidad & E. Navarro. 2013. New Species of *Brayopsis*, *Descurainia*, *Draba*, *Neuontobotrys* and *Weberbaueria* (Brassicaceae) from Peru. *Kew Bulletin* 68 (2):219–231. doi: <https://doi.org/10.1007/s12225-013-9447-z>
- Al-Shehbaz I.A., P. Gonzáles & A. Cano. 2015a. *Englerocharis blanca-leoniae* (Brassicaceae), a new species from Puno, Peru. *Harvard Papers in Botany* 20(1):1–4. doi: <https://doi.org/10.3100/hpib.v20iss1.2015.n1>
- Al-Shehbaz I.A., P. Gonzáles & A. Cano. 2015b. *Weberbaueria incisa* (Brassicaceae), a new species from southern Peru. *Novon* 24(1):6–8. doi: <https://doi.org/10.3417/2015003>
- Angiosperm Phylogeny Group IV. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181:1–20. doi: <https://doi.org/10.1111/j.1095-8339.2009.00996.x>
- Argollo J., M. Fornari, G. Herail, V. Miranda & G. Viscarra. 1987. Estratigrafía de los depósitos glaciares en la Cordillera de Apolobamba (Bolivia) y su asociación con mineralizaciones auríferas. *Décimo Congreso Geol. Argentino, San Miguel de Tucumán, Actas II*:67–69.
- Argollo J. & M.H. Iriondo. 2008. El cuaternario de Bolivia y regiones vecinas. Museo Provincial de Ciencias Naturales Florentino Ameghino. Santa Fe, Argentina.
- Arteta M., M. Corrales, C. Dávalos, A. Delgado, F. Sinca, L. Hernani & J. Bojórquez. 2006. Plantas vasculares de la bahía de Juli, lago Titicaca, Perú. *Ecología Aplicada* 5(1–2):29–36.
- Ballard H.E. & H.H. Iltis. 2012. *Viola lilliputana* sp. nov. (*Viola* sect. *Andinium*, Violaceae), one of the world's smallest violets, from the Andes of Peru. *Brittonia* 64(4):353–358. doi: <https://doi.org/10.1007/s12228-012-9238-0>
- Beniston M. 1994. Mountain environments in changing climates. Routledge, London.
- Bridson D. & L. Forman. 1992. *Herbarium Handbook*. Royal Botanical Gardens, Kew.
- Brako L. & J. Zarucchi. 1993. Catalogue of the flowering plants and gymnosperms of Peru. *Monographs in Systematic Botany from the Missouri Botanical Garden* 45:1–1286.
- Cano A., W. Mendoza, S. Castillo, M. Morales, M.I. La Torre, H. Aponte, A. Delgado, N. Valencia & N. Vega. 2010. Flora y vegetación de suelos crioturbados y habitats asociados en la Cordillera Blanca, Ancash, Peru. *Revista Peruana de Biología* 17(1):095–103. doi: <https://doi.org/10.15381/rpb.v17i1.56>
- Cano A., A. Delgado, W. Mendoza, H. Trinidad, P. Gonzáles, M.I. La Torre, M. Chanco, H. Aponte, J. Roque, N. Valencia & E. Navarro. 2011. Flora y vegetación de suelos crioturbados y habitats asociados en los alrededores del Abra Apacheta, Ayacucho–Huancavelica (Perú). *Revista Peruana de Biología* 18(2):169–178. doi: <https://doi.org/10.15381/rpb.v18i2.224>
- Cerrate E. 1969. Manera de preparar plantas para el herbario. Museo de Historia Natural, Serie de Divulgación, N°1.
- Conde-Álvarez C. & S. Saldaña–Zorrilla. 2007. Cambio climático en América Latina y el Caribe: Impactos, vulnerabilidad y adaptación. *Revista Ambiente y Desarrollo* 23 (2):23–30.
- Cuesta F., P. Muriel, L. D. Llambí, S. Halloy, N. Aguirre, S. Beck, J. Carilla, R. I. Meneses, S. Cuello, A. Grau, L. E. Gámez, J. Irazábal, J. Jácome, R. Jaramillo, L. Ramírez, N. Samaniego, D. Suárez-Duque, N. Thompson, A. Tupayachi, P. Viñas, K. Yager, M. T. Becerra, H. Pauli & W. D. Gosling. 2017. Latitudinal and altitudinal patterns of plant community diversity on mountain summits across the tropical Andes. *Ecography* 40:1–14. doi: <https://doi.org/10.1111/ecog.02567>
- Drummond C. S., R. J. Eastwood, S. T. S. Miotto, & C. E. Hughes. 2012. Multiple continental radiations and correlates of diversification in *Lupinus* (Leguminosae): Testing for Key innovation with incomplete taxon sampling. *Systematic Biology* 61(3): 443–460. doi: <https://doi.org/10.1093/sysbio/syr126>
- Flores M., J. Alegría, & A. Granda. 2005. Diversidad florística asociada a las lagunas andinas Pomacocha y Habascocha, Junín, Perú. *Revista Peruana de Biología* 12(1): 125–134. Doi: <http://dx.doi.org/10.15381/rpb.v12i1.2366>
- Fort M. 2015. Impact of climate change on mountain environment dynamics. *Journal of Alpine Research / Revue de géographie alpine* 103-2. doi: <http://dx.doi.org/10.4000/rga.2877>
- Galán de Mera A., B. Del Monte, E.M. Mendoza, E. Linares, J. Campos, & J.A. Vicente. 2014. Las comunidades vegetales relacionadas con los procesos criogénicos en los Andes peruanos. *Phytocoenologia* 44(1–2):121–161. doi: <http://dx.doi.org/10.1127/0340-269X/2014/0044-0576>
- Gentry A.H. 1993. Overview of Peruvian Flora. In: Brako L, Zarucchi J, Catalogue of the Flowering Plants and Gymnosperms of Peru. *Monographs in Systematic Botany from the Missouri Botanical Garden* 45:1–1286.
- Gonzáles P., E. Navarro, M. Chanco & A. Cano. 2015. *Nototriche carabayensis* (Malvaceae), una especie nueva de los Andes de Perú. *Darwiniana* 3(1):1–6.
- Gonzáles P., A. Cano, I. Al-Shehbaz, D.W. Ramírez, E. Navarro, H. Trinidad & M. Cueva. 2016. Doce nuevos registros de plantas vasculares para los Andes de Perú. *Arnaldoa* 23(1): 159–170.
- Gonzáles P. & A. Cano. 2016. Two new species of *Viola* (Violaceae) named in honor preceding Peruvian botanists. *Phytotaxa* 283(1): 083–090. doi: <http://dx.doi.org/10.11646/phytotaxa.283.1.6>
- Hughes C.E. & G.W. Atchison. 2015. The ubiquity of alpine plant radiations: from the Andes to the Hengduan Mountains. *New Phytologist* 207:275–282. doi: <http://dx.doi.org/10.1111/nph.13230>
- Jørgensen P.M., C. Ulloa Ulloa, B. León, S. León-Yáñez, S.G. Beck, M. Nee, J.L. Zarucchi, M. Celis, R. Bernal & R. Grads-tein. 2011. Regional patterns of vascular plant diversity and endemism. 192–203. IN: S.K. Herzog, R. Martinez, P.M. Jørgensen & H. Tiessen (Eds.). *Climate Change and Biodiversity in the Tropical Andes*. SCOPE, MacArthur Foundation & IAI.
- Jørgensen P.M., M.H. Nee & S.G. Beck. (eds.) 2014. *Catálogo de las plantas vasculares de Bolivia*, *Monogr. Syst. Bot. Missouri Bot. Gard.* 127(1–2): i–viii, 1–1744. Missouri Botanical Garden Press, St. Louis.
- Kohler T., A. Wehrli, M. Jurek. (eds.). 2014. *Mountains and climate change: A global concern*. Sustainable Mountain Development Series. Bern, Switzerland, Centre for Development and Environment (CDE), Swiss Agency for Development and Cooperation (SDC) and Geographica Bernensia.
- Macbride J.F. et al. 1936 and next. *Flora of Peru*. Field Museum of Natural History, Botanical Series, Chicago.
- León B., J. Roque, C. Ulloa Ulloa, N. Pitman, P.M. Jørgensen & A. Cano. 2007. El libro rojo de las especies endémicas del Perú. *Revista Peruana de Biología, Número especial* 13(2):1–971.
- Linares E., J.E. Campos & A. Galán de Mera. 2012. Nuevas adiciones a la flora del Perú, VI. *Arnaldoa* 19(1):37–45.
- Linares E., J.E. Campos, J.A. Vicente & A. Galán de Mera. 2015. *Adesmia schickendantzii* (Fabaceae, subgen. *Acanthadesmia*), novedad para la flora del Perú. *Acta botánica malacitana* (40):206–208.
- Markham A., N. Dudley & S. Stolton. 1993. *Some like it hot: climate change, biodiversity and the survival of species*. WWF–International, Gland.
- Montesinos-Tubée D.B. 2011. Diversidad florística de la cuenca alta del río Tambo–Ichuña (Moquegua, Perú). *Revista Peruana de Biología* 18(1):119–132. doi: <http://dx.doi.org/10.15381/rpb.v18i1.156>
- Montesinos-Tubée D.B. 2012. Lista anotada de nuevas adiciones para la flora andina de Moquegua, Perú. *Revista Peruana de Biología* 19(3):303–312. doi: <http://dx.doi.org/10.15381/rpb.v19i3.1045>
- Montesinos-Tubée D.B., A.M. Cleef & K.V. Sýkora. 2015a. The Puna vegetation of Moquegua, South Peru: Chasmophytes, grasslands and *Puya raimondii* stands. *Phytocoenologia* 45(4):365–397. doi: <http://dx.doi.org/10.1127/phyto/2015/0006>
- Montesinos-Tubée D.B., P. Gonzáles & E. Navarro. 2015b. *Senecio canoi* (Compositae), a new species of the Andes of Peru. *Anales Jardín Botánico de Madrid* 72(2):1–4. doi: <http://dx.doi.org/10.3989/ajbm.2409>
- Ospina J.C., S.P. Sylvester & M.D.P.V. Sylvester. 2016. Multivariate

- analysis and taxonomic delimitation within the *Festuca setifolia* complex (Poaceae) and a new species from the Central Andes. *Systematic Botany* 41(3):727-746. doi: <http://dx.doi.org/10.1600/036364416X692398>
- Pauli H., M. Gottfried & G. Grabherr. 2003. Effects of climate change on the alpine and nival vegetation of the Alps. *Journal of Mountain Ecology* 7 (Suppl.):9-12.
- Pauli H., M. Gottfried, K. Reiter, C. Klettner & G. Grabherr. 2007. Signal of range expansions and contractions of vascular plants in the high Alps: Observations (1994-2004) at The GLORIA Master Site Schrankage, Tyrol, Austria. *Global Change Biology* 13:147-156. doi: <http://dx.doi.org/10.1111/j.1365-2486.2006.01282.x>
- Ruthsatz B. 2012. Vegetación y ecología de los bofedales altoandinos de Bolivia. *Phytocoenologia* 42:133-179. doi: <https://doi.org/10.1127/0340-269X/2012/0042-0535>
- Roque J. & E.K. Ramírez. 2008. Flora vascular y vegetación de la laguna Parinacochas y alrededores (Ayacucho, Perú). *Revista Peruana de Biología* 15(1):105-110. doi: <http://dx.doi.org/10.15381/rpb.v15i1.1677>
- Sánchez V.I. 1996. Aspectos florísticos de la jalca y alternativas de manejo sustentable. En: *Anales del Simposio Estrategias para Bioconservación en el Norte del Perú*. Arnaldo Ed. Esp. 4(2): 25-62.
- Sklenář P. & H. Balslev. 2005. Superpáramo plant species diversity and phytogeography in Ecuador. *Flora - Morphology, Distribution, Functional Ecology of Plants*, 200(5):416-433. doi: <http://dx.doi.org/10.1016/j.flora.2004.12.006>
- Sklenář P. & H. Balslev. 2007. Geographic flora elements in the Ecuadorian superpáramo. *Flora - Morphology, Distribution, Functional Ecology of Plants*, 202(1):50-61. doi: <http://dx.doi.org/10.1016/j.flora.2006.03.002>
- Sklenář P., I. Hedberg & A. M. Cleef. 2014. Island biogeography of tropical alpine floras. *Journal of Biogeography* 41:287-297. doi: <http://dx.doi.org/10.1111/jbi.12212>
- Sylvester S.P., D. Quandt, L. Ammann & M. Kessler. 2016. The world's smallest Campanulaceae: *Lysipomia mitsyae*, sp. nov. *Taxon* 65(2):305-314. doi: <http://dx.doi.org/10.12705/652.7>
- Sylvester S.P., R.J. Soreng, P. M. Peterson & M.D.P.V. Sylvester. 2016. An updated checklist and key to the open-panicled species of *Poa* L. (Poaceae) in Peru including three new species, *Poa ramoniana*, *Poa tayacajensis*, and *Poa urubambensis*. *PhytoKeys* 65:57-90. doi: <http://dx.doi.org/10.3897/phytokeys.65.7024>
- Thompson L.G., E. Mosley-Thompson, M.E. Davis, P.N. Lin, K. Henderson & T.A. Mashiotta. 2003. Tropical glacier and ice core evidence of climate change on annual to millennial time scales. *Climate Change* 59:137-155. doi: <https://doi.org/10.1023/A:1024472313775>
- Thompson L.G., E. Mosley-Thompson, H. Brecher, M. Davis, B. León, D. Les, L. Ping-Nan, M. Tracy & K. Mountain. 2006. Abrupt tropical climate change: Past and present. *Proceedings of the National Academy of Sciences* 103(28):10536-10543. doi: <https://doi.org/10.1073/pnas.0603900103>
- Thompson L.G., E. Mosley-Thompson, M.E. Davis, V.S. Zagorodnov; I.M. Howat, V.N. Mikhalenko & P.N. Lin. 2013. Annually resolved ice core records of tropical climate variability over past ~1800 years. *Science* 340(6135):945-950. doi: <http://dx.doi.org/10.1126/science.1234210>
- Toledo V.M. & V. Sosa. 1993. Floristics in Latin America and the Caribbean: An evaluation of the numbers of plant collections and botanists. *Taxon* 42:355-364.
- Tovar O. 1993. Las gramíneas (Poaceae) del Perú. *Ruizia* 13:9-474.
- Tupayachi A. 2005. Flora de la Cordillera Vilcanota. *Arnaldo* 12(1-2):126-144.
- Ulloa Ulloa C., J.L. Zarucchi & B. León. 2004. Diez años de adiciones a la flora del Perú. *Arnaldo* (edición especial):7-242.
- Valencia N., A. Cano, A. Delgado, H. Trinidad & P. Gonzáles. 2013. Composición y cobertura de la vegetación de los bofedales en un macrotrasecto este-oeste, en los Andes centrales del Perú pp. 278-293 In A. Alonso, F. Dallmeier & G. Servat. (Ed.), *Monitoreo de la biodiversidad: lecciones de un megaproyecto transandino*. Smithsonian Institution Scholarly Press. USA.
- Weberbauer A. 1945. *El Mundo Vegetal de los Andes Peruanos*. Ministerio de Agricultura, Lima. Lumen S.A.
- Young K. R. 2011. Introduction to Andean Geographies. pp. 128-137. In: S. K. Herzog et al. *Climate Change and biodiversity in the tropical Andes*. IAI.
- Young K. R. 2014. Ecology of land cover change in glaciated tropical mountains. *Revista peruana de Biología* 21:259-270. doi: <https://doi.org/10.15381/rpb.v21i3.10900>
- Young K. R., C. Ulloa Ulloa, J. Luteyn & S. Knapp. 2002. Plant evolution and endemism in Andean South America: An Introduction. *Botanical review* 68(1):4-21. DOI: [https://doi.org/10.1663/0006-8101\(2002\)068\[0004:PEAEIA\]2.0.CO;2](https://doi.org/10.1663/0006-8101(2002)068[0004:PEAEIA]2.0.CO;2)

Appendix 1. Checklist of vascular flora of the Cordillera Carabaya, indicating growth form (F) (h: herb, s: subshrub, a: shrub, ab: tree, c: cactoide), endemic species (E) (x: reported in León et al. (2007); x* no collections in other countries but not previously reported as endemic to Peru), Elev.: elevational record of these collection(s); Voucher: PG (Paúl Gonzáles), FP (Francis Pennell), JS (Jaroslav Soukup), CV (César Vargas), OT (Oscar Tovar), AW (Augusto Weberbauer), MW (Maximilian Weigend), fl: not registered herbarium specimen, known only as photograph from study area; []: collections within brackets come from neighboring areas.

family	Species	F	E	Elev.	Voucher
Pteridophytes					
Aspleniaceae	<i>Asplenium gilliesii</i> Hook.	h	-	>4000	fl
	<i>Asplenium monanthes</i> L.	h	-	>4000	Boeke JD. 2972
	<i>Asplenium triphyllum</i> C. Presl	h	-	4499	PG 3816
Dryopteridaceae	<i>Elaphoglossum matthewsii</i> (Fée) T. Moore	h	-	>4000	PG 3025
	<i>Polystichum cochleatum</i> (Klotzsch) Hieron.	h	-	>4000	fl
Isoetaceae	<i>Isoetes andina</i> Hook.	h	-	>4000	León 5316
	<i>Isoetes lechleri</i> Mett.	h	-	>4000	Lechler 1937
Lycopodiaceae	<i>Phlegmariurus crassus</i> (Humb. & Bonpl. ex Willd.) B. Øllg.	h	-	>4000	fl
Ophioglossaceae	<i>Ophioglossum crotalophoroides</i> Walter	h	-	>4000	fl
Polypodiaceae	<i>Campyloneurum amphostenon</i> (Kunze ex Klotzsch) Fée	h	-	>4000	fl
	<i>Campyloneurum asplundii</i> (C. Chr.) Ching	h	-	>4000	fl
	<i>Melpomene peruviana</i> (Desv.) A.R. Sm. & R.C. Moran	h	-	>4000	fl
	<i>Pleopeltis pycnocarpa</i> (C. Chr.) A.R. Sm.	h	-	3935	PG 3047
Pteridaceae	<i>Argyrochosma nivea</i> (Poir.) Windham	h	-	>4000	fl
	<i>Cheilanthes pilosa</i> Goldm.	h	-	>4000	fl
	<i>Cheilanthes pruinata</i> Kaulf.	h	-	>4000	fl
	<i>Cheilanthes scariosa</i> (Sw.) C. Presl	h	-	>4000	fl
	<i>Pellaea ovata</i> (Desv.) Weath.	h	-	>4000	fl
Salviniaceae	<i>Azolla filiculoides</i> Lam.	h	-	>4000	fl
Woodsiaceae	<i>Cystopteris fragilis</i> (L.) Bernh.	h	-	>4000	fl
	<i>Woodsia montevidensis</i> (Spreng.) Hieron.	h	-	>4000	fl
Gymnosperms					
Ephedraceae	<i>Ephedra rupestris</i> Benth.	s	-	>4000	fl
Monocots					
Alstroemeriaceae	<i>Alstroemeria pygmaea</i> Herb.	h	-	3500-4400	MW 2000/68, CV 6840
	<i>Bomarea dulcis</i> (Hook.) Beauverd	h	-	4499	Hill 3819, Raimondi A. 10229
	<i>Bomarea involucrosa</i> (Herb.) Baker	h	-	>4000	fl
Amaryllidaceae	<i>Nothoscordum gramineum</i> (Sims) P. Beauv.	h	-	3800	CV 15181
Asparagaceae	<i>Oziroë acaulis</i> (Baker) Speta	h	-	4000	CV 7146
Cyperaceae	<i>Carex boliviensis</i> Van Heurck & Müll. Arg.	h	-	>4000	fl
	<i>Carex bonplandii</i> Kunth	h	-	>4000	fl
	<i>Carex brachycalama</i> Griseb.	h	-	4499	fl
	<i>Carex pichinchensis</i> Kunth	h	-	>4000	Lechler 2519, PG 3817
	<i>Cyperus andinus</i> Palla ex Kük.	h	-	>4000	fl
	<i>Eleocharis albibracteata</i> Nees & Meyen ex Kunth	h	-	4381	fl [Meyen s.n.]
	<i>Oreobolus venezuelensis</i> Steyererm.	h	-	>4000	PG 3083
	<i>Phylloscirpus acaulis</i> (Phil.) Goetgh. & D.A. Simpson	h	-	>4000	fl [PG 1604]
	<i>Phylloscirpus deserticola</i> (Phil.) Dhooghe & Goetgh.	h	-	>4000	León 5315
	<i>Trichophorum rigidum</i> (Boeckeler) Goetgh., Muasya & D.A. Simpson	h	-	>4000	fl

(...)

family	Species	F	E	Elev.	Voucher
	<i>Zameioscirpus atacamensis</i> (Phil.) Dhooge & Goetgh.	h	-	>4000	f!
	<i>Zameioscirpus muticus</i> Dhooge & Goetgh.	h	-	4285	f! [PG1603]
Hydrocharitaceae	<i>Elodea potamogeton</i> (Bertero) Espinosa	h	-	>4000	f!
Iridaceae	<i>Cardenanthus vargasii</i> R.C. Foster	h	x*	4399	PG 3450
	<i>Olsynium acaule</i> (Klatt) Goldblatt	h	-	4301	PG 3421
	<i>Olsynium junceum</i> (E. Mey. ex C. Presl) Goldblatt	h	-	>4000	f!
	<i>Sisyrinchium brevipes</i> Baker	h	-	>4000	f!
	<i>Sisyrinchium caespitificum</i> Kraenzl.	h	-	>4000	f!
	<i>Sisyrinchium chilense</i> Hook.	h	-	>4000	f!
	<i>Sisyrinchium palustre</i> Diels	h	-	>4000	f!
	<i>Sisyrinchium porphyreum</i> Kraenzl.	h	-	4301	PG 3418
Juncaceae	<i>Distichia muscoides</i> Nees & Meyen	h	-	>4000	f!
	<i>Juncus arcticus</i> Willd.	h	-	>4000	f!
	<i>Juncus bufonius</i> L.	h	-	>4000	f!
	<i>Juncus ebracteatus</i> E. Mey.	h	-	>4000	f!
	<i>Juncus stipulatus</i> Nees & Meyen	h	-	4381	PG 3086
	<i>Luzula ecuadoriensis</i> Balslev	h	-	4335	PG 3002
	<i>Luzula racemosa</i> Desv.	h	-	>4000	f!
	<i>Oxychloe andina</i> Phil.	h	-	4700	PG 3452
	<i>Patosia clandestina</i> (Phil.) Buchenau	h	-	4424	PG 3447
Juncaginaceae	<i>Lilaea scilloides</i> (Poir.) Hauman	h	-	>4000	f!
Orchidaceae	<i>Aa rosei</i> Ames	h	-	4100	CV 6846
	<i>Myrosmodes chiogena</i> (Schltr.) C.A. Vargas	h	-	4897	PG 3057b
	<i>Myrosmodes gymnandra</i> (Rchb. f.) C.A. Vargas	h	-	4458	PG 2996
	<i>Myrosmodes paludosa</i> (Rchb. f.) P. Ortiz	h	-	4285-4897	PG 3022, PG 3057a
Poaceae	<i>Aciachne acicularis</i> Lægaard	h	-	>4000	f!
	<i>Aciachne pulvinata</i> Benth.	h	-	3250-3630	Bennett B. 2004, 2099, Lechler 3238
	<i>Aegopogon cenchroides</i> Humb. & Bonpl. ex Willd.	h	-	>4000	f!
	<i>Agrostis breviculmis</i> Hitchc.	h	-	3790-4600	Anonimo 7158, Bennett B. 2283
	<i>Agrostis toluensis</i> Kunth	h	-	>4000	f!
	<i>Alopecurus hitchcockii</i> Parodi	h	-	>4000	f!
	<i>Anatherostipa hans-meyeri</i> (Pilg.) Peñail.	h	-	3940-4335	AW 940, Bennett B. 2759, Lechler 1978, PG 3017
	<i>Anatherostipa obtusa</i> (Nees & Meyen) Peñail.	h	-	>4000	f! [Monheim 160, 10, OT 5175, 5259, Weddell 4488]
	<i>Avena sativa</i> L.	h	-	>4000	f! [OT s.n.]
	<i>Bromus catharticus</i> Vahl	h	-	>4000	f! [Monheim 19]
	<i>Bromus lanatus</i> Kunth	h	-	>4000	f! [CV 16233]
	<i>Bromus pitensis</i> Kunth	h	-	>4000	f!
	<i>Bromus villosissimus</i> Hitchc.	h	-	>4000	CV 7894
	<i>Calamagrostis antoniana</i> (Griseb.) Hack. ex Dusén	h	-	4600	CV 7184
	<i>Calamagrostis breviaristata</i> (Wedd.) Pilg.	h	-	4897	PG 3067
	<i>Calamagrostis brevifolia</i> (J. Presl) Steud.	h	-	>4000	f!
	<i>Calamagrostis chrysantha</i> (J. Presl) Steud.	h	-	>4000	CV 7167, León 5321
	<i>Calamagrostis eminens</i> (J. Presl) Steud.	h	-	>4000	f!
	<i>Calamagrostis heterophylla</i> (Wedd.) Pilg.	h	-	3790	Bennett B. 2287

(...)

family	Species	F	E	Elev.	Voucher
	<i>Calamagrostis intermedia</i> (J. Presl) Steud.	h	-	3550	Metcalfe R. 30433
	<i>Calamagrostis jamesonii</i> Steud.	h	-	>4000	f! [Pearson 75]
	<i>Calamagrostis macrophylla</i> (Pilg.) Pilg.	h	-	>4000	Metcalfe R. 30433
	<i>Calamagrostis minima</i> (Pilg.) Tovar	h	-	>4000	AW 953
	<i>Calamagrostis nitidula</i> Pilg.	h	-	4400	AW 960, León 5319
	<i>Calamagrostis ovata</i> (J. Presl) Steud.	h	-	>4000	CV 7892, Lechler 2057
	<i>Calamagrostis recta</i> (Kunth) Trin. ex Steud.	h	-	3470	Bennett B. 2264
	<i>Calamagrostis rigescens</i> (J. Presl) Scribn.	h	-	>4000	f! [JS 113]
	<i>Calamagrostis rigida</i> (Kunth) Trin. ex Steud.	h	-	>4000	f!
	<i>Calamagrostis rupestris</i> Trin.	h	-	4400	CV 8131
	<i>Calamagrostis setiflora</i> (Wedd.) Pilg.	h	-	>4000	f!
	<i>Calamagrostis spiciformis</i> Hack.	h	-	3300	Bennett B. 2494
	<i>Calamagrostis tarmensis</i> Pilg.	h	-	4288-3470	Bennett B. 2263, PG 3034, 3018
	<i>Calamagrostis trichophylla</i> Pilg.	h	-	>4000	f! [Stordy s.n., Harlam s.n.]
	<i>Calamagrostis vicunarum</i> (Wedd.) Pilg.	h	-	4381	PG 3085
	<i>Dielsiochloa floribunda</i> (Pilg.) Pilg.	h	-	>4000	AW 1028, León 5320, 5342
	<i>Festuca breviaristata</i> Pilg.	h	-	>4000	f!
	<i>Festuca divergens</i> Tovar	h	x*	4559	PG 2998
	<i>Festuca dolichophylla</i> J. Presl	h	-	>4000	f! [JS 447, OT 5243, 5159]
	<i>Festuca humilior</i> Nees & Meyen	h	-	>4000	CV 7897
	<i>Festuca lasiorrhachis</i> Pilg.	h	-	>4000	Ferreira R. 16741
	<i>Festuca orthophylla</i> Pilg.	h	-	>4000	f! [Pearson 46, OT 5102, AW 4087]
	<i>Festuca rigescens</i> (J. Presl) Kunth	h	-	4897	PG 3049
	<i>Festuca setifolia</i> Steud. ex Griseb.	h	x*	3500-4700	Lechler 1826
	<i>Festuca tenuiculmis</i> Tovar	h	x*	>4000	f!
	<i>Hordeum muticum</i> J. Presl	h	-	>4000	f! [JS 115]
	<i>Jarava ichu</i> Ruiz & Pav.	h	-	>4000	f! [Pearsson 10, Stafford D. 63, Rauh 1490, 642]
	<i>Muhlenbergia angustata</i> (J. Presl) Kunth	h	-	>4000	Boeke JD. 2974
	<i>Muhlenbergia ligularis</i> (Hack.) Hitchc.	h	-	3790	Bennett B. 2284
	<i>Muhlenbergia peruviana</i> (P. Beauv.) Steud.	h	-	4100	Bennett B. 2738
	<i>Nassella brachyphylla</i> (Hitchc.) Barkworth	h	-	>4000	CV 7901
	<i>Nassella inconspicua</i> (J. Presl) Barkworth	h	-	>4000	f! [OT 5187, 5333, Harlam s.n.]
	<i>Nassella pubiflora</i> (Trin. & Rupr.) E. Desv.	h	-	>4000	f! [Lechler 1736, Ellenberg 421]
	<i>Paspalum pygmaeum</i> Hack.	h	-	3630	Bennett B. 2010
	<i>Pennisetum clandestinum</i> Hochst. ex Chiov.	h	-	>4000	f!
	<i>Piptochaetium panicoides</i> (Lam.) E. Desv.	h	-	4285	CV 7163, PG 3020
	<i>Poa aequigluma</i> Tovar	h	-	>4000	f!
	<i>Poa annua</i> L.	h	-	2790	Bennett B. 2289
	<i>Poa apiculata</i> Refulio	h	x	4999	PG 2985
	<i>Poa ayacuchensis</i> Tovar	h	x	4335	PG 3019
	<i>Poa calycina</i> (J. Presl) Kunth	h	-	>4000	CV 7891
	<i>Poa dentigluma</i> Tovar	h	-	>4000	f!
	<i>Poa gilgiana</i> Pilg.	h	-	>4000	CV 7902
	<i>Poa glaberrima</i> Tovar	h	-	4288	PG 3035

(...)

family	Species	F	E	Elev.	Voucher
	<i>Poa gymnantha</i> Pilg.	h	-	>4000	CV 7174
	<i>Poa horridula</i> Pilg.	h	-	>4000	Lechler 2682, Bennett B. 2323
	<i>Poa humillima</i> Pilg.	h	-	>4000	f!
	<i>Poa lepidula</i> (Nees & Meyen) Soreng & L.J. Gillespie	h	-	4600-5300	CV 8136, León 5323
	<i>Poa macusaniensis</i> (E.H.L. Krause) Refulio	h	-	4381	Lechler 1836, PG 3091
	<i>Poa marshallii</i> Tovar	h	-	>4000	f!
	<i>Poa perligulata</i> Pilg.	h	-	>4000	f!
	<i>Poa peruviana</i> Jacq.	h	-	>4000	f!
	<i>Poa pseudoaequigluma</i> Tovar	h	-	>4000	f!
	<i>Poa serpaiana</i> Refulio	h	-	4381	PG 3084
	<i>Poa spicigera</i> Tovar	h	-	>4000	f!
	<i>Polypogon interruptus</i> Kunth	h	-	>4000	f!
	<i>Trisetum spicatum</i> (L.) K. Richt.	h	-	4285	PG 3024
	<i>Vulpia dertonensis</i> (All.) Gola	h	-	>4000	f!
	<i>Vulpia myuros</i> (L.) C.C. Gmel.	h	-	3690	Bennett B. 2660
Potamogetonaceae	<i>Stuckenia punensis</i> (A. Galán) A. Galán	h	-	>4000	f! [Iltis 1441]
	<i>Stuckenia pectinata</i> (L.) Börner	h	-	4360	CV 7132
Eudicots					
Amaranthaceae	<i>Gomphrena meyeniana</i> Walp.	h	-	4360	CV 7121
Apiaceae	<i>Azorella biloba</i> (Schltdl.) Wedd.	h	-	4150-4399	CV 6838, CV 6848, Lechler s.n., PG 3448
	<i>Azorella diapiensoides</i> A. Gray	h	-	>4000	f! [CV 13010, JS 210; 214, Stordy 32]
	<i>Azorella multifida</i> (Ruiz & Pav.) Pers.	h	-	4300	CV 6841
	<i>Bowlesia palmata</i> Ruiz & Pav.	h	-	3935	PG 3043
	<i>Bowlesia tropaeolifolia</i> Gillies & Hook.	h	-	3900	Hoogte L. 2402
	<i>Chaerophyllum andicola</i> (Kunth) K.F. Chung	h	-	4300	CV 1308, Hoogte L. 850
	<i>Daucus montanus</i> Humb. & Bonpl. ex Spreng.	h	-	>4000	f!
	<i>Lilaeopsis macloviana</i> (Gand.) A.W. Hill	h	-	>4000	f! [Aguilar 135]
	<i>Niphogeton scabra</i> (H. Wolff) J.F. Macbr.	h	-	>4000	f!
Apocynaceae	<i>Philibertia lysimachioides</i> (Wedd.) T. Mey.	s	-	>4000	f!
Asteraceae	<i>Achyrocline alata</i> (Kunth) DC.	h	-	>4000	f!
	<i>Achyrocline ramosissima</i> Britton ex Rusby	h	-	>4000	f! [Aguilar 441, FP 13484, JS 93]
	<i>Ageratina glechonophylla</i> (Less.) R.M. King & H. Rob.	s	-	>4000	f!
	<i>Aldama helianthoides</i> (Rich.) E.E. Schill. & Panero	h	-	3935	PG 3040
	<i>Baccharis caespitosa</i> (Ruiz & Pav.) Pers.	s	-	4000-5300	León 5338
	<i>Baccharis genistelloides</i> (Lam.) Pers.	s	-	>4000	Boeke JD. 3105
	<i>Baccharis tola</i> Phil.	a	-	>4000	f!
	<i>Baccharis tricuneata</i> (L. f.) Pers.	a	-	>4000	Boeke JD. 3168
	<i>Bidens andicola</i> Kunth	h	-	>4000	f!
	<i>Chersodoma antennaria</i> (Wedd.) Cabrera	h	-	4522	PG 3432
	<i>Chersodoma jodopappa</i> (Sch. Bip.) Cabrera	s	-	>4000	f! [PG 3311]
	<i>Chuiriraga jussieui</i> J.F. Gmel.	s	-	>4000	Stafford D. 1111, 420
	<i>Conyza artemisioides</i> Meyen & Walp.	h	-	>4000	PG 3072
	<i>Conyza bonariensis</i> (L.) Cronquist	h	-	>4000	f!
	<i>Conyza coronopifolia</i> Kunth	h	-	3935	PG 3038

(...)

family	Species	F	E	Elev.	Voucher
	<i>Conyza deserticola</i> Phil.	h	-	3935	PG 3039
	<i>Cotula mexicana</i> (DC.) Cabrera	h	-	>4000	fl
	<i>Cuatrecasiella isernii</i> (Cuatrec.) H. Rob.	h	-	>4000	fl
	<i>Diplostephium carabayense</i> Wedd.	s	x	>4000	fl
	<i>Erigeron lanceolatus</i> Wedd.	h	-	>4000	fl
	<i>Erigeron rosulatus</i> Wedd.	h	-	>4000	León 5327
	<i>Facelis plumosa</i> (Wedd.) Sch. Bip.	h	-	>4000	fl [FP 13459]
	<i>Galinsoga mandonii</i> Sch.Bip. ex Baker	h	-	3935	PG 3046
	<i>Gamochaeta americana</i> (Mill.) Wedd.	h	-	>4000	Boeke JD. 3053
	<i>Gamochaeta humilis</i> Wedd.	h	-	4400	CV 17630
	<i>Gamochaeta purpurea</i> (L.) Cabrera	h	-	>4000	fl [Aguilar, 418, CV s.n, 16254, Rauh P650]
	<i>Gnaphalium dombeyanum</i> DC.	h	-	>4000	Boeke JD. 3163
	<i>Gnaphalium lacteum</i> Meyen & Walp.	h	-	>4000	fl [FP 13444]
	<i>Gnaphalium polium</i> Wedd.	h	-	>4000	fl [Aguilar, 417, FP 13444, 13470, 13470a]
	<i>Hieracium leptocephalium</i> Benth.	h	-	>4000	fl
	<i>Hieracium neoherrerae</i> Zahn	h	-	4288	PG 3030
	<i>Hypochaeris chillensis</i> (Kunth) Britton	h	-	4288	PG 3029
	<i>Hypochaeris echegarayi</i> Hieron.	h	-	>4000	fl
	<i>Hypochaeris meyeniana</i> (Walp.) Benth. & Hook. f. ex Griseb.	h	-	>4000	fl
	<i>Hypochaeris sessiliflora</i> Kunth	h	-	>4000	fl
	<i>Hypochaeris taraxacoides</i> Ball	h	-	>4000	fl
	<i>Jalcochila ecuadorensis</i> M.O. Dillon & Sagást.	h	-	4516	PG 3102
	<i>Laennecia gnaphalioides</i> (Kunth) Cass.	h	-	>4000	fl
	<i>Leucheria daucifolia</i> (D. Don) Crisci	h	-	>4000	JS J. 474
	<i>Misbrookea strigosissima</i> (A. Gray) V.A. Funk	h	-	>4000	fl
	<i>Mniodes aretioides</i> (Wedd.) Cuatrec.	s	-	4500	Dillon M. 1083
	<i>Mniodes conoidea</i> (Wedd.) S. E. Freire	h	-	4500	Dillon M. 1082
	<i>Mniodes kunthiana</i> (DC.) S. E. Freire	h	-	>4000	León 5326
	<i>Mniodes longifolia</i> (Cuatrec. & Aristeg.) S. E. Freire	h	-	>4000	fl
	<i>Mniodes piptolepis</i> (Wedd.) S. E. Freire	h	-	>4000	fl [Aguilar s.n., 427, Mexia 4262, FP13368, 13437]
	<i>Mniodes schultzii</i> (Wedd.) S. E. Freire	h	-	>4000	fl [Stafford D. 746]
	<i>Mniodes subspicata</i> (Wedd.) S.E. Freire	h	-	>4000	fl [Dillon M., 1070, FP 13369, 13471]
	<i>Noticastrum marginatum</i> (Kunth) Cuatrec.	h	-	>4000	fl
	<i>Novenia acaulis</i> (Benth. & Hook. f. ex B.D. Jacks.) S.E. Freire & F.H. Hellw.	h	-	4300	Martín 2092
	<i>Oriastrum stuebelii</i> (Hieron.) A.M.R. Davies	h	-	>4000	fl
	<i>Oritrophium hieracioides</i> (Wedd.) Cuatrec.	h	-	4285	PG 3026
	<i>Oritrophium limnophilum</i> (Sch. Bip.) Cuatrec.	h	-	>4000	fl
	<i>Paranephelius ovatus</i> A. Gray ex Wedd.	h	-	>4000	fl
	<i>Perezia ciliosa</i> (Phil.) Reiche	h	-	>4000	fl
	<i>Perezia coerulescens</i> Wedd.	h	-		CV 7169, 9626, Dillon M. 1089
	<i>Perezia multiflora</i> (Bonpl.) Less.	h	-	>4000	fl [Zuniga 33, FP 13429]
	<i>Perezia pungens</i> (Bonpl.) Less.	h	-	>4000	Diaz s.n.
	<i>Perezia pygmaea</i> Wedd.	h	-	4381	PG 3090, Monheim 55
	<i>Perezia sublyrata</i> Domke	h	-	>4000	fl

(...)

family	Species	F	E	Elev.	Voucher
	<i>Senecio adenophylloides</i> Sch. Bip.	a	-	4520	PG 3070, 3071
	<i>Senecio adenophyllus</i> Meyen & Walp.	a	-	>4000	f!
	<i>Senecio algens</i> Wedd.	h	-	4335-5024	PG 3010, PG 3825
	<i>Senecio breviscapus</i> DC.	h	-	>4000	f!
	<i>Senecio candollei</i> Wedd.	h	-	4335	PG 3005
	<i>Senecio canescens</i> (Bonpl.) Cuatrec.	h	-	4500	Dillon M. 1080
	<i>Senecio canoi</i> P. Gonzáles, Montesinos & Ed. Navarro	h	x	4520-4742	PG 3429, 3428, 3441
	<i>Senecio chavanilloensis</i> Cuatrec.	h	-	>4000	Boeke JD. 3000
	<i>Senecio collinus</i> DC.	a	x	>4000	f!
	<i>Senecio condimentarius</i> Cabrera	h	-	>4000	f!
	<i>Senecio evacoides</i> Sch. Bip.	h	-	>4000	f!
	<i>Senecio ferreyrae</i> Cabrera	s	x	>4000	f!
	<i>Senecio gamolepis</i> Cabrera	s	-	4857	Linares E. 2733
	<i>Senecio humillimus</i> Sch. Bip.	s	-	4762	PG 3442, 3443
	<i>Senecio hyoseridifolius</i> Wedd.	h	-	>4000	f!
	<i>Senecio jarae</i> Phil.	h	-	4346	Linares E. 2712
	<i>Senecio macrorrhizus</i> Wedd.	h	x	>4000	f!
	<i>Senecio modestus</i> Wedd.	h	-	4499	PG 3814
	<i>Senecio moqueguensis</i> Montesinos	h	x	4496	PG 3422
	<i>Senecio nivalis</i> (Kunth) Cuatrec.	h	-	4520-4897	PG 3822, 3056
	<i>Senecio nutans</i> Sch. Bip.	a	-	>4000	f!
	<i>Senecio pinnatilobatus</i> Sch. Bip.	a	-	4499	PG 3821
	<i>Senecio rhizomatus</i> Rusby	h	-	>4000	f!
	<i>Senecio rudbeckiifolius</i> Meyen & Walp.	a	-	>4000	f!
	<i>Senecio rufescens</i> DC.	a	-	>4000	f!
	<i>Senecio serratifolius</i> (Meyen & Walp.) Cuatrec.	h	-	4897	PG 3053
	<i>Senecio tassaensis</i> Montesinos	h	x	4762	PG 3444, 3445
	<i>Senecio Vulgaris</i> L.	h	-	>4000	f!
	<i>Soliva neglecta</i> Cabrera	h	-	>4000	f!
	<i>Sonchus asper</i> (L.) Hill	h	-	>4000	f!
	<i>Stevia macbridei</i> B.L. Rob.	h	-	3935	PG 3045
	<i>Stevia mandonii</i> Sch. Bip.	h	-	>4000	f!
	<i>Stuckertiella capitata</i> (Wedd.) Beauverd	h	-	>4000	f! [Iltis 1426, JS 504]
	<i>Tagetes multiflora</i> Kunth	h	-	>4000	f!
	<i>Taraxacum fernandezianum</i> Dahlst. ex Skottsbo.	h	-	>4000	f!
	<i>Werneria apiculata</i> Sch. Bip.	h	-	>4000	f!
	<i>Werneria glaberrima</i> Phil.	h	-	>4000	f!
	<i>Werneria heteroloba</i> Wedd.	h	-	4939	PG 2995
	<i>Werneria nubigena</i> Kunth	h	-	>4000	f!
	<i>Werneria obtusiloba</i> S.F. Blake	h	-	4939	PG 2994
	<i>Werneria orbignyana</i> Wedd.	h	-	4335-4499	PG 3004, 3820
	<i>Werneria pectinata</i> Lingelsh.	h	-	>4000	f!
	<i>Werneria pumila</i> Kunth	h	-	>4000	f!
	<i>Werneria pygmaea</i> Gillies ex Hook. & Arn.	h	-	4381	PG 3088, 3089

(...)

family	Species	F	E	Elev.	Voucher
	<i>Werneria solivifolia</i> Sch. Bip.	h	-	4412	PG 3094
	<i>Xenophyllum ciliolatum</i> (A. Gray) V.A. Funk	h	-	4897-5230	PG 3061, 3837, 3104
	<i>Xenophyllum dactylophyllum</i> (Sch. Bip.) V.A. Funk	s	-	5024	PG 3829
	<i>Xenophyllum digitatum</i> (Wedd.) V.A. Funk	h	-	4897-5230	PG 3048, 2978, 3103
	<i>Xenophyllum marcidum</i> (S.F. Blake) V.A. Funk	h	-	5072-5230	PG 3832, 3106
	<i>Xenophyllum pseudodigitatum</i> (Rockh.) V.A. Funk	s	-	5024-5072	PG 3828, 3833
	<i>Xenophyllum staffordiae</i> (Sandwith) V.A. Funk	s	x	4897	PG 3055
Begoniaceae	<i>Begonia veitchii</i> Hook. f.	h	-	>4000	fl
Berberidaceae	<i>Berberis agapatensis</i> Lechl.	a	-	>4000	Lechler 2646
	<i>Berberis carinata</i> Lechl.	a	-	>4000	Lechler 2644
	<i>Berberis rectinervis</i> Rusby	a	-	3850	CV 6853
Boraginaceae	<i>Hackelia revoluta</i> (Ruiz & Pav.) I.M. Johnst.	h	-	4335	PG 3003
	<i>Phacelia secunda</i> J.F. Gmel.	h	-	4000-4150	CV 12534, 7003, Metcalf R. 30708
	<i>Plagiobothrys humilis</i> (Ruiz & Pav.) I.M. Johnst.	h	-	4520	PG 3079
Brassicaceae	<i>Brassica rapa</i> subsp. <i>campestris</i> (L.) Clapham	h	-	>4000	fl
	<i>Brayopsis alpaminae</i> Gilg & Muschl.	h	-	4520	PG 3075
	<i>Brayopsis monimocalyx</i> O.E. Schulz	h	-	4400-4999	CV 17627, PG 2991
	<i>Capsella bursa-pastoris</i> (L.) Medik.	h	-	>4000	fl
	<i>Cardamine bonariensis</i> Pers.	h	-	>4000	fl [Lecher 1811]
	<i>Descurainia athrocarpa</i> (A. Gray) O.E. Schulz	h	-	4330-4897	CV 7008, PG 3051
	<i>Descurainia myriophylla</i> (Willd. ex DC.) R.E. Fr.	h	-	3935	PG 3042
	<i>Draba brackenridgei</i> A. Gray	h	-	4897	PG 3064
	<i>Draba cuzcoensis</i> O.E. Schulz	h	-	4762	PG 3438
	<i>Draba loayzana</i> Al-Shehbaz	h	-	4736	Navarro E. 1092
	<i>Draba punoensis</i> Al-Shehbaz, Ed. Navarro, H. Trinidad & A. Cano	h	x	4658-4762	Navarro E. 1087, PG 3437
	<i>Englerocharis blanca-leoniae</i> A-Shehbaz, P. Gonzáles & A. Cano	h	x	4897	PG 3059
	<i>Englerocharis peruviana</i> Muschl.	h	-	4644	Linares E. 2708
	<i>Lepidium bipinnatifidum</i> Desv.	h	-	4999-4520	PG 2992, 3080
	<i>Lepidium chichicara</i> Desv.	h	-	>4000	fl [Meyen s.n. CV 18541, PG 3307, 3308]
	<i>Lepidium meyenii</i> Walp.	h	-	4300-4762	CV 7134, 7011, PG 3439
	<i>Mancoa hispida</i> Wedd.	h	-	4520	PG 3077
	<i>Rorippa nana</i> (Schltdl.) J.F. Macbr.	h	-	>4000	Lechler 2583
	<i>Weberbaueria insisa</i> A-Shehbaz, P. Gonzáles & A. Cano	h	-	4522-5000	PG 3430, 3107
	<i>Weberbaueria peruviana</i> (DC.) Al-Shehbaz	h	-	>4000	MW 2000/120
	<i>Weberbaueria spathulifolia</i> (A. Gray) O.E. Schulz	h	-	4320-4892	CV 6837, 6837, PG 3060
Cactaceae	<i>Austrocylindropuntia floccosa</i> (Salm-Dyck) F. Ritter	c	-	>4000	fl
	<i>Austrocylindropuntia lagopus</i> (K. Schum.) I. Crook, J. Arnold & M. Lowry	c	-	4360-4400	CV 7136, 22550
	<i>Echinopsis maximiliana</i> Heyder ex A. Dietr.	c	-	>4000	fl
Calceolariaceae	<i>Calceolaria plectranthifolia</i> Walp.	h	-	4288	PG 3028
	<i>Calceolaria scapiflora</i> (Ruiz & Pav.) Benth.	h	-	>4000	fl
Campanulaceae	<i>Hypsela reniformis</i> (Kunth) C. Presl	h	-	>4000	fl [JS 100, Mandon 489]
	<i>Lysipomia acaulis</i> Kunth	h	-	>4000	fl
	<i>Lysipomia glandulifera</i> (Schltdl. ex Wedd.) Schltdl. ex E. Wimm.	h	-	>4000	Lechler 2076
	<i>Lysipomia laciniata</i> A. DC.	h	-	3595-4424	Lechler 1956, Raimondi A. 11597, Sharpe 109, CV 1290, 10724, PG 3446
					(...)

family	Species	F	E	Elev.	Voucher
Caprifoliaceae	<i>Lysipomia sphagnophila</i> Griseb. ex Wedd.	h	-	>4000	Lechler 2641
	<i>Wahlenbergia peruviana</i> A. Gray	h	-	>4000	f!
	<i>Wahlenbergia urcosensis</i> E. Wimm.	h	x	>4000	f!
	<i>Belonanthus spathulatus</i> (Ruiz & Pav.) Schmale	h	-	>4000	f!
	<i>Stangea paulae</i> Graebn.	h	x	5024-084	PG 3826, 2979
	<i>Stangea rhizantha</i> (A. Gray) Killip	h	x	4450-4897	CV 7012, PG 3050
	<i>Stangea wandae</i> Graebn.	h	-	>4000	f! [Stafford D. 737, 1268]
	<i>Valeriana andina</i> Britton	h	-	4335	PG 3015
	<i>Valeriana baltana</i> Graebn.	h	x	>4000	Ferreyra 531
	<i>Valeriana cephalantha</i> Schltld.	h	x	>4000	Lechler 2044
	<i>Valeriana coarctata</i> Ruiz & Pav.	h	-	4644	Linares E. 2690
	<i>Valeriana globularioides</i> Graebn.	h	-	4897	PG 3054
	<i>Valeriana herrerae</i> Killip	h	x	4000	CV 7145
	<i>Valeriana johannae</i> Weberl.	h	x	>4000	Bernardi L. 16774
	<i>Valeriana micropterina</i> Wedd.	h	-	4288-4496	PG 3032, 3001, 3016, 3425, Lechler 1962
	<i>Valeriana niphobia</i> Briq.	h	-	>4000	Hohenacker R. 1962
	<i>Valeriana nivalis</i> Wedd.	h	-	4496	PG 3427
<i>Valeriana pennellii</i> Killip	h	-	>4000	f! [CV 31]	
<i>Valeriana plectritoides</i> Graebn.	h	x	3800	CV 6863, AW 928a	
<i>Valeriana pycnantha</i> A. Gray	h	-	4600	CV 7180	
Caryophyllaceae	<i>Alsine rupestris</i> Muschl.	h	x	>4000	AW 1042
	<i>Arenaria digyna</i> Willd. ex D.F.K. Schltld.	h	-	4999	PG 2990
	<i>Arenaria lanuginosa</i> (Michx.) Rohrb.	h	-	4000	CV 7148
	<i>Cardionema ramosissimum</i> (Weinm.) A. Nelson & J.F. Macbr.	h	-	4000	CV 7154
	<i>Cerastium danguyi</i> J.F. Macbr.	h	-	>4000	f!
	<i>Cerastium glomeratum</i> Thuill.	h	-	>4000	f!
	<i>Cerastium mucronatum</i> Wedd.	h	-	>4000	f!
	<i>Cerastium peruvianum</i> Muschl.	h	-	4400	CV 7036
	<i>Cerastium subspicatum</i> Wedd.	h	-	>4000	Weddell H. 4653
	<i>Drymaria divaricata</i> var. <i>stricta</i> (Rusby) J.A. Duke	h	-	>4000	Lechler 1947
	<i>Paronychia andina</i> A. Gray	h	-	4450	CV 7015
	<i>Paronychia mandoniana</i> Rohrb.	h	-	>4000	f! [Lecher 1760]
	<i>Pycnophyllum aschersonianum</i> Muschl.	h	x	>4000	f!
	<i>Pycnophyllum bryoides</i> (Phil.) Rohrb.	h	-	4400	AW 501b
	<i>Pycnophyllum glomeratum</i> Mattf.	h	-	4400-5024	AW 501a, 952, PG 3081, 3827
	<i>Pycnophyllum lechlerianum</i> Rohrb.	h	x	4999	PG 2988
	<i>Pycnophyllum molle</i> Remy	h	-	4335-4400	PG 3009, AW 946, 950
	<i>Pycnophyllum weberbaueri</i> Muschl.	h	-	4399-5024	PG 3451, 3823
	<i>Silene mandonii</i> (Rohrb.) Bocquet	h	-	>4000	f!
	<i>Silene thysanodes</i> Fenzl	h	-	>4000	f!
	<i>Spergularia fasciculata</i> Phil.	h	-	>4000	f! [Lecher 1772]
	<i>Stellaria cuspidata</i> D.F.K. Schltld.	h	-	3800	AW 925
	<i>Stellaria media</i> (L.) Vill.	h	-	>4000	f!
	<i>Stellaria weddellii</i> Pedersen	h	-	>4000	f!

(...)

family	Species	F	E	Elev.	Voucher
Convolvulaceae	<i>Dichondra microcalyx</i> (Hallier f.) Fabris	h	-	>4000	f!
Crassulaceae	<i>Crassula closiana</i> (Gay) Reiche	h	-	>4000	f!
	<i>Crassula connata</i> (Ruiz & Pav.) A. Berger	h	-	>4000	f! [FP 13043]
Elatinaceae	<i>Elatine triandra</i> Schkuhr	h	-	>4000	Lechler 2687
Escalloniaceae	<i>Escallonia myrtilloides</i> L. f.	a	-	>4000	Boeke JD. 3020
Euphorbiaceae	<i>Euphorbia huanchahana</i> (Klotzsch & Garcke) Boiss.	h	-	>4000	f!
Fabaceae	<i>Adesmia schickendantzii</i> Griseb.	s	-	4192	Galán 3896
	<i>Astragalus arequipensis</i> Vogel	h	-	>4000	JS J. 966
	<i>Astragalus dielsii</i> J.F. Macbr.	h	-	4897	PG 3058
	<i>Astragalus dillinghamii</i> J.F. Macbr.	h	x	4412-4999	PG 3100, 2983
	<i>Astragalus diminutivus</i> (Phil.) Gómez-Sosa	h	-	4200	CV 15179
	<i>Astragalus garbancillo</i> Cav.	h	-	>4000	f! [JS 85]
	<i>Astragalus minimus</i> Vogel	h	-	>4000	f! [Meyen 67]
	<i>Astragalus pilgeri</i> J.F. Macbr.	h	-	4450	CV 7026
	<i>Astragalus uniflorus</i> DC.	h	-	>4000	f!
	<i>Astragalus weddellianus</i> (Kuntze) I.M. Johnst.	h	-	3750-4400	JS J. 549; 424, CV 7026, 7030
	<i>Lupinus ananeanus</i> Ulbr.	h	-	>4000	f!
	<i>Lupinus aridulus</i> C.P. Sm.	h	-	4335	PG 3006
	<i>Lupinus breviscapus</i> Ulbr.	h	-	>4000	f!
	<i>Lupinus eriocladius</i> Ulbr.	h	-	4000	CV 7153
	<i>Lupinus gibertianus</i> C.P. Sm.	h	-	3500-4600	CV 6852, 7170, Lechler 1842
	<i>Lupinus microphyllus</i> Desr.	h	-	>4000	f!
	<i>Lupinus peruvianus</i> Ulbr.	h	-	4335	PG 3008, 3012, Antezana F. 16
	<i>Lupinus pulvinaris</i> Ulbr.	h	-	4500	AW 985
	<i>Lupinus ulbrichianus</i> C.P. Sm.	h	-	>4000	AW 870
	<i>Medicago polymorpha</i> L.	h	-	>4000	f!
	<i>Trifolium amabile</i> Kunth	h	-	>4000	f! [CV 21253]
	<i>Trifolium repens</i> L.	h	-	>4000	f!
	<i>Vicia graminea</i> Sm.	h	-	>4000	f! [CV 12499, 18540, 20915]
Gentianaceae	<i>Gentiana sedifolia</i> Kunth	h	-	>4000	f! [FP 13418; 13426, JS 98]
	<i>Gentianella armerioides</i> (Griseb. ex Gilg) J.S. Pringle	h	-	>4000	Lechler 2000a
	<i>Gentianella bridgesii</i> (Gilg) Fabris ex T.N. Ho & S.W. Liu	h	-	4000	AW 1048
	<i>Gentianella campanuliformis</i> (Reimers) Fabris	h	x	4999	PG 2986
	<i>Gentianella centamalensis</i> (Gilg) Zarucchi	h	x	>4000	f!
	<i>Gentianella dolichopoda</i> (Gilg) J.S. Pringle	h	x*	4381	PG 3092
	<i>Gentianella lobelioides</i> (Gilg) Zarucchi	h	x	4600	AW 955
	<i>Gentianella persquarrosa</i> (Reimers) J.S. Pringle	h	x	4939	PG 2993
	<i>Gentianella potamophila</i> (Gilg) Zarucchi	h	x	>4000	f!
	<i>Gentianella primuloides</i> (Gilg) J.S. Pringle	h	-	>4000	Lechler 2002
	<i>Gentianella punicea</i> (Wedd.) Holub	h	-	4000	Weddell H. s.n.
	<i>Gentianella sandiense</i> (Gilg) J.S. Pringle	h	-	3600	AW 917a
	<i>Gentianella scarlatiflora</i> (Gilg) J.S. Pringle	h	x*	4300	CV 15180
	<i>Gentianella scarlatina</i> (Gilg) Zarucchi	h	-	>4000	AW 1047
	<i>Halenia caespitosa</i> Gilg	h	-	14500 ft	Stafford D. 1117

(...)

family	Species	F	E	Elev.	Voucher	
Geraniaceae	<i>Halenia umbellata</i> (Ruiz & Pav.) Gilg	h	-	3600-4288	MW 2000/123, Hoogte L. 2399, PG 3031	
	<i>Erodium cicutarium</i> (L.) L'Hér. ex Aiton	h	-	>4000	f! [FP 13401]	
	<i>Erodium moschatum</i> (L.) L'Hér. ex Aiton	h	-	>4000	f!	
	<i>Geranium core-core</i> Steud.	h	-	3935	PG 3037, 3041	
	<i>Geranium crassipes</i> Hook. ex A. Gray	h	-	>4000	Lechler 1985	
	<i>Geranium dielsianum</i> R. Knuth	h	-	>4000	f!	
	<i>Geranium fallax</i> Steud.	h	-	>4000	Lechler 1907	
	<i>Geranium sessiliflorum</i> Cav.	h	-	>4000	f! [AW 1003a, FP 13439]	
Grossulariaceae	<i>Geranium weddellii</i> Briq.	h	-	4336	JS J. 537	
	<i>Ribes bolivianum</i> Jancz.	a	-	3850	CV 6864	
Haloragaceae	<i>Ribes brachybotrys</i> (Wedd.) Jancz.	a	-	4800	Hoogte L. 2212	
	<i>Myriophyllum quitense</i> Kunth	h	-	>4000	f! [FP 13425, JS 835]	
Hypericaceae	<i>Hypericum silenoides</i> Juss.	h	-	>4000	f!	
Lamiaceae	<i>Hedeoma mandoniana</i> Wedd.	h	-	>4000	f!	
	<i>Lamium amplexicaule</i> L.	h	-	>4000	f!	
	<i>Lepechinia meyenii</i> (Walp.) Epling	h	-	>4000	f!	
	<i>Stachys pusilla</i> (Wedd.) Briq.	h	-	>4000	f!	
Loasaceae	<i>Caiophora contorta</i> (Desr.) C. Presl	h	-	4330	CV 7009	
	<i>Caiophora horrida</i> (Britton) Urb. & Gilg	h	-	>4000	f! [CV 6830, 13009, 20880]	
	<i>Caiophora rosulata</i> (Wedd.) Urb. & Gilg	h	-	>4000	f!	
Malvaceae	<i>Acaulimalva steinbachii</i> Krapov.	h	-	>4000	f!	
	<i>Nototriche anthemidifolia</i> (J. Rémy) A.W. Hill	h	-	4412	PG 3099	
	<i>Nototriche armeriifolia</i> A.W. Hill	h	x	>4000	f!	
	<i>Nototriche azurella</i> A.W. Hill	h	x*	4600	AW 957	
	<i>Nototriche carabayensis</i> P. González, M. Chanco & Ed. Navarro	h	x	5072	PG 3835	
	<i>Nototriche condensata</i> (Baker f.) A.W. Hill	h	-	>4000	Lechler 1972	
	<i>Nototriche congesta</i> A.W. Hill	h	x*	4500	AW 1018	
	<i>Nototriche cupuliforme</i> Krapov.	h	x	4360	CV 7119	
	<i>Nototriche digitulifolia</i> A.W. Hill	h	x	4897	PG 3052	
	<i>Nototriche erinacea</i> Hill ex Burt	h	x	4897	PG 3062	
	<i>Nototriche flabellata</i> (Wedd.) A.W. Hill	h	-	4400-4897	Lechler 1972, AW 961, PG 3066	
	<i>Nototriche longirostris</i> (Wedd.) A.W. Hill	h	-	4360	CV 7118	
	<i>Nototriche longituba</i> Burt & Hill	h	x	>4000	f! [Howell Williams s.n.]	
	<i>Nototriche obcuneata</i> (Baker f.) A.W. Hill	h	-	4500-5084	AW 983, 1023, PG 3834, 2976	
	<i>Nototriche pediculariifolia</i> (Meyen) A.W. Hill	h	-	4500-4700	AW 1023, CV 7896, 7185	
	<i>Nototriche pellicea</i> A.W. Hill	h	x	4412	PG 3098	
	<i>Nototriche purpurascens</i> A.W. Hill	h	-	4897	PG 3065	
	<i>Nototriche sepaliloba</i> Hochr.	h	x*	4885	Linares E. 2722	
	<i>Nototriche staffordiae</i> Burt & Hill	h	x	>4000	f! [Stafford D. 1263]	
	<i>Nototriche sulphurea</i> A.W. Hill	h	-	4700	AW 219, 963	
	<i>Tarasa urbaniana</i> (Ulbr.) Krapov.	h	-	3600-4520	AW 927, CV 1312, PG 3076	
	Montiaceae	<i>Calandrinia acaulis</i> Kunth	h	-	4300	AW 956, CV 6839
		<i>Montia fontana</i> L.	h	-	>4000	f!
Onagraceae	<i>Epilobium denticulatum</i> Ruiz & Pav.	h	-	4335	AW 357, PG 3014	

(...)

family	Species	F	E	Elev.	Voucher
Orobanchaceae	<i>Oenothera multicaulis</i> Ruiz & Pav.	h	-	>4000	f! [JS 104]
	<i>Neobartsia bartsioides</i> (Hook.) Edwin	h	-	>4000	JS J. 970
	<i>Neobartsia diffusa</i> (Benth.) Uribe-Convers & Tank	h	-	>4000	León 5343, Linares 2742
	<i>Neobartsia elongata</i> (Wedd.) Uribe-Convers & Tank	h	-	4335-4499	PG 3013, 3426, 3818
	<i>Neobartsia fiebrigii</i> (Diels) Uribe-Convers & Tank	h	-	>4000	f!
	<i>Neobartsia inaequalis</i> (Benth.) Uribe-Convers & Tank	h	-	>4000	f! [AW 510, 919]
	<i>Neobartsia pedicularoides</i> (Benth.) Uribe-Convers & Tank	h	-	>4000	f!
	<i>Castilleja pumila</i> (Benth.) Wedd.	h	-	>4000	f! [FP 13428, JS 938, Rauh 47]
Oxalidaceae	<i>Castilleja virgatoides</i> Edwin	h	-	>4000	f!
	<i>Oxalis calachaccensis</i> R. Knuth	h	-	>4000	f! [CV 6825]
	<i>Oxalis cuzcensis</i> R. Knuth	h	-	4520	PG 3073
	<i>Oxalis nubigena</i> Walp.	h	-	>4000	f! [FP 13394, Meyen s.n.]
	<i>Oxalis oreocharis</i> Diels	h	-	4450	CV 7012
	<i>Oxalis pinguiculacea</i> R. Knuth	h	-	3800-4450	CV 6844, CV 7024
Phrymaceae	<i>Oxalis tuberosa</i> Molina	h	-	>4000	AW 295
	<i>Mimulus glabratus</i> Kunth	h	-	>4000	f!
Piperaceae	<i>Peperomia peruviana</i> Dahlst.	h	-	4644	Linares E. 2692
	<i>Peperomia verruculosa</i> Dahlst. ex A.W. Hill	h	-	4450	CV 7013
Plantaginaceae	<i>Callitriche heteropoda</i> Engelm. ex Hegelm.	h	-	>4000	f!
	<i>Callitriche nubigena</i> Fassett	h	-	>4000	f!
	<i>Ourisia biflora</i> Wedd.	h	-	14500 ft	Stafford D. 1112
	<i>Ourisia muscosa</i> Wedd.	h	-	4762	PG 3435
	<i>Plantago australis</i> Lam.	h	-	>4000	f! [CV 21263]
	<i>Plantago lamprophylla</i> Pilg.	h	-	4520	PG 3082
	<i>Plantago linearis</i> Kunth	h	-	4516	PG 3101
	<i>Plantago nubicola</i> (Decne.) Rahn	h	-	4335-4522	Hoogte L. 2997, PG 3011, 3449, 3434, CV 17621, AW 994
	<i>Plantago orbignyana</i> Steinh. ex Decne.	h	-	>4000	f! [Meyen s.n.]
	<i>Plantago rigida</i> Kunth	h	-	4000-4381	AW 1050, Weddell H. s.n., PG 3087
	<i>Plantago sericea</i> Ruiz & Pav.	h	-	>4000	f! [CV 20887, 20923, 21259]
	<i>Plantago tubulosa</i> Decne.	h	-	>4000	f!
	<i>Veronica serpyllifolia</i> L.	h	-	3000	CV 1283, 9637
	Polemoniaceae	<i>Gilia laciniata</i> Ruiz & Pav.	h	-	>4000
<i>Phlox gracilis</i> (Douglas ex Hook.) Greene		h	-	>4000	Hill s.n.
Polygonaceae	<i>Muehlenbeckia volcanica</i> (Benth.) Endl.	h	-	>4000	AW 849
Ranunculaceae	<i>Caltha sagittata</i> Cav.	h	-	>4000	Lechler 1953, Weddell H. s.n., Linares 2680
	<i>Krapfia haemantha</i> (Ulbr.) Tamura	h	-	4335-4897	Raimondi A. s.n., PG 3000, 3068
	<i>Oreithales integrifolia</i> (DC.) Schldl.	h	-	4500	Lechler 2706, CV 22556
	<i>Ranunculus breviscapus</i> DC.	h	-	>4000	f!
	<i>Ranunculus filamentosus</i> Wedd.	h	-	4400-4999	CV 7033, PG 3431, 2984
	<i>Ranunculus flagellifolius</i> Nakai	h	-	>4000	f!
	<i>Ranunculus limoselloides</i> Turcz.	h	-	>4000	f!
	<i>Ranunculus praemorsus</i> Kunth ex DC.	h	-	2500	AW 597, Lechler 2709, Raimondi A. s.n.
	<i>Ranunculus trichophyllus</i> Chaix ex Vill.	h	-	>4000	f!
	<i>Ranunculus uniflorus</i> Phil. ex Reiche	h	-	4381	PG 3093

(...)

family	Species	F	E	Elev.	Voucher
Rosaceae	<i>Aphanes andicola</i> Rothm.	h	-	4520	PG 3078
	<i>Lachemilla barbata</i> (C. Presl) Rothm.	h	-	>4000	f!
	<i>Lachemilla diplophylla</i> (Diels) Rothm.	h	-	>4000	f!
	<i>Lachemilla frigida</i> (Wedd.) Rothm.	h	-	4285	PG 3023
	<i>Lachemilla pinnata</i> (Ruiz & Pav.) Rothm.	h	-	4400	CV 17623
	<i>Lachemilla sandiensis</i> Rothm.	h	x	>4000	AW 945
	<i>Lachemilla vulcanica</i> (Schltdl. & Cham.) Rydb.	h	-	3600	AW 926
	<i>Polylepis besseri</i> Hieron.	ab	-	3800-4100	CV 12528, 7139
	<i>Tetraglochin cristatum</i> (Britton) Rothm.	s	-	>4000	f! [Tupayachi A. 1570, CV 1, 20927, 22545]
Rubiaceae	<i>Galium aparine</i> L.	h	-	3935	PG 3044
	<i>Galium corymbosum</i> Ruiz & Pav.	h	-	>4000	f! [FP 13407]
Saxifragaceae	<i>Saxifraga magellanica</i> Poir.	h	-	3500-4400	CV 6859, 6857, 7172, AW 183
Scrophulariaceae	<i>Limosella subulata</i> E. Ives	h	-	>4000	f!
Solanaceae	<i>Nicotiana pavonii</i> Dunal	h	-	>4000	Metcalf R. 30712
	<i>Nicotiana undulata</i> Ruiz & Pav.	h	-	>4000	f! [CV 9627, JS 902, 632]
	<i>Salpichroa amoena</i> Benoist	s	-	3800-4260	MW 2000/119, Särkinen T. 4056, PG 3036
	<i>Salpichroa glandulosa</i> (Hook.) Miers	s	-	4260	PG 3036
	<i>Salpichroa lehmannii</i> Dammer	s	-	4200-4496	MW 2000/53, Särkinen T. 4039, PG 3424
	<i>Solanum acaule</i> Bitter	h	-	4335-4450	PG 3007, 3095, CV 7023
	<i>Solanum grandidentatum</i> Phil.	h	-	4200-4596	Beltrán H. 6453, MW 2000/54
	<i>Solanum weddellii</i> Phil.	h	-	4200-4200	Särkinen T. 4038
	Urticaceae	<i>Urtica echinata</i> Benth.	h	-	>4000
<i>Urtica flabellata</i> Kunth		h	-	4600-4700	AW 969
<i>Urtica urens</i> L.		h	-	>4000	f!
Verbenaceae	<i>Verbena litoralis</i> Kunth	h	-	>4000	f!
Violaceae	<i>Viola enmae</i> P. Gonzáles	h	x	4300-4400	PG 3420, 3096
	<i>Viola ferreyrae</i> P. Gonzáles	h	x	4300-4400	PG 3419, 3097
	<i>Viola micranthella</i> Wedd.	h	-	4335	PG 2999
	<i>Viola pusillima</i> Wedd.	h	-	>4000	f! [PG 1605]
	<i>Viola pygmaea</i> Juss. ex Poir.	h	-	4300-4360	MW 2000/63, CV 7125