


# Binding of Telomeric DNA G-Quadruplexes by Abietane Diterpene Natural Products

Sarah Ostrowski, Tarryn Franciosa, Ruel E. McKnight

State University of New York at Geneseo, Department of Chemistry, Geneseo NY 14454



## Introduction

G-quadruplexes are non-canonical higher order DNA structures formed from guanine rich sequences, made up of stacked G-tetrads stabilized by Hoogsteen base pairing and  $K^+$  ions. G-quadruplexes are overrepresented in the promoter regions of oncogenes and the 5'UTR of mRNA<sup>1</sup>. As a result, G-Quadruplexes have been implicated as targets for possible anti-cancer therapeutic agents to treat previously "undruggable" targets like the c-myc and ras oncogenes<sup>1</sup>.

The human telomeric repeat,  $[5'G_3(T_2AG_3)_3]$ , is a repeating, single stranded DNA sequence that can form G-quadruplexes. Telomerase, an enzyme expressed in ~90% of all cancers, is responsible for extending telomeric repeats, making cancer cells immortal. It has been shown that stabilization of telomeric G-quadruplexes can inhibit telomerase activity and therefore block the survival of cancer cells<sup>2</sup>.

The compounds used in the study are a group of abietane diterpene natural products from *Hyptis verticillata*, a plant native to the Caribbean and central America that has been used traditionally as an ethnomedicine that has been shown to have a range of therapeutic effects. Some of these effects include anti-microbial, anti-inflammatory, and even anti-cancer activities<sup>3</sup>. Using several biophysical techniques, we have investigated the binding characteristics of these compounds to G-quadruplex DNA as a possible rationale for their observed anti-cancer therapeutic effects.

## Experimental

### Sample Preparation

DNA G-Quadruplexes were formed in solution using a 10 mM phosphate buffer, pH 7, with 100mM  $K^+$  added. Characteristic CD spectra confirmed the presence of quadruplexes in solution.

### Circular dichroism

CD measurements were performed using a Jasco J-815 CD Spectrophotometer. For each run, the abietane solution was added to a solution of G-quadruplex DNA and the spectra were measured from 200-600nm.

### Thermal Denaturation

$T_m$  measurements were performed using a CARY-5000 UV-Vis-NIR Spectrophotometer. Absorption spectra of G-quadruplex-abietane complex were recorded at 295 nm as the temperature was increased from 25.00°C to 95.00°C.

### Thioflavin-T Fluorescence Displacement

Fluorescent studies were performed using a PTI Quantamaster™ 40 UV-Vis Spectrofluorometer from 400-600nm. A Fluorogenic dye that binds specifically to DNA G-quadruplexes, Thioflavin-T (ThT), was used as a fluorescent indicator for the quadruplexes. A solution of ThT bound G-quadruplexes was prepared and the abietane solution was added. The solution was excited at 425 nm and fluorescence intensity was measured.




Figure 1. Left: A picture of *Hyptis verticillata*, the plant from which these compounds are derived<sup>4</sup>. Right: A generalized structure of the abietane diterpenes used.




Figure 2. A simple diagram demonstrating the mechanism of the Thioflavin-T displacement assay, Adapted from Jamroskovic et al.<sup>7</sup>

## Results

### Thioflavin Displacement




Figure 4. Thioflavin-T displacement graphs of compounds 1, 2 and 3. As the abietanes were bound, the fluorescence intensity changed based on whether or no the abietanes were able to displace Thioflavin-T from its binding site on the quadruplexes.

### Circular Dichroism




Figure 5. CD Spectra of three abietanes (compound 1, compound 2, compound 3) after being titrated into G-quadruplex solution. Structures of compounds 2 and 3 are shown.

### Thermal Denaturation




Figure 6. Thermal Denaturation plots of compounds 1, 2, and 3. Each compound was run alongside a DNA only control.




Figure 3. A simplified diagram of DNA G-quadruplexes formed from human telomeric DNA<sup>5</sup>.

Table 1.  $\Delta T_m$  values of each compound determined from the thermal denaturation studies. Apparent Stern-Volmer constants were determined for each compound via the Thioflavin-T displacement studies.

Compound	$\Delta T_m$ (°C)	Stern-Volmer Constant ( $mM^{-1}$ )
1	5.7	0.0075
2	5.4	0.0169
3	-17.86	-
4	8.1	0.0151
5	8.4	0.0118
6	8.46	0.0136
7	-	-
8	2.2	-

## Conclusions

- Displacement studies showed that 5 of the 8 abietane diterpenes interacted with the DNA-ThT complex, quenching the fluorescence.
- CD Spectra indicated that of the 8 compounds tested, compounds 1, 2, and 3 showed the most significant interactions with the telomeric G-quadruplexes.
- The unique spectra of compound 3 indicates a shift from antiparallel quadruplexes to a different form of quadruplex, although it is unclear what that form may be. The positive peaks at 295nm and negative peaks at 240nm shown by the spectra for compounds 1 and 2 indicate an increase in stabilization of the antiparallel form of G-quadruplex<sup>8</sup>.
- Thermal denaturation studies showed that the binding of all the compounds resulted in an increase in thermal stability ( $T_m$ ) at higher temperatures, after an initial structural transition. An early transition/species was most evident for 3.
- The spectra we have recorded for all compounds, specifically compound 3, is very interesting and warrants further investigation.

## References

- Balasubramanian et al. Targeting G-quadruplexes in gene promoters: a novel anticancer strategy? Nat Rev Drug Discov. 2011; 10:261-275
- Sun et al. Inhibition of human telomerase by a G-quadruplex-interactive compound. J. Med. Chem. 1997; 40:2113-2116
- Picking et al. *Hyptis verticillata* Jacq: A review of its traditional uses, phytochemistry, pharmacology and toxicology. Journal of Ethnopharmacology. 2013; 147:16-41
- Mohanty et al. Thioflavin T as an Efficient Inducer and Selective Fluorescent Sensor for the Human Telomeric G-Quadruplex DNA. Journal of the American Chemical Society. 2013; 135:367-376  
<https://regionalconservation.org/ircs/database/plants/PlantPage.asp?TXCODE=Hyptvert>
- Bhasikuttan, Mohanty. Targeting G-quadruplex Structures with extrinsic fluorogenic dyes: promising fluorescence sensors. Chem. Commun. 2015; 51:7581-7597
- Jamroskovic et al. Identification of Compounds that Selectively Stabilize Specific G-Quadruplex Structures by using a Thioflavin T-Displacement Assay as a tool. Chemistry – A European Journal. 2016; 52:18932-18943
- Dai et al. Polymorphism of human telomeric quadruplex structures. Biochimie. 2008;90(8):1172-1183

## Acknowledgments

We acknowledge Professor Roy Porter at the University of the West Indies for the abietanes used in this study, the Geneseo Foundation and the Department of Chemistry for the instruments used to collect our data, Dr. Kazushige Yokoyama at SUNY Geneseo for the Thioflavin-T used in our displacement studies, and Medhi Elmouchtari, a Geneseo alum who began Dr. McKnight's work on G-quadruplexes.