

The Emerging Roles of Human Immunity-Related GTPase M (IRGM) Gene

Harish Changotra

Department of Biotechnology and Bioinformatics,
Jaypee University of Information Technology,
Solan, Himachal Pradesh, India

Article Info

Article history:

Received Nov 25th 2014

Revised Dec 29th, 2014

Accepted Dec 31st 2014

Keyword:

IRGM
autophagy
pseudogene
SNP
Lysosome

ABSTRACT

Immunity-related GTPase family M (IRGM) protein, discovered for the first time in 1990, belongs to IRG family of proteins and is divided into five subfamilies (IRGA, IRGB, IRGC, IRGD and IRGM). These are responsive to interferon- γ and play a pivotal role in immune response to pathogens in mice. Owing to lack of interferon response element in the promoter region, human IRGM does not respond to interferon- γ stimulation, which is why it was earlier thought to be non-functional gene. Moreover, evolutionary history suggests that this gene was non-functional for ~25 million years ago. Bioinformatics has been instrumental in elucidating its evolutionary history as well as functions, especially in its interactions with the proteins of autophagy pathway. Recently, several studies have demonstrated the various functions of IRGM in limiting different pathogens in humans. This review discusses how and various roles played by IRGM unraveled in the recent years.

Copyright © 2015 *International Journal for Computational Biology*,
<http://www.ijcb.in>, All rights reserved.

Corresponding Author:

Harish Chagnotra,
Department of Biotechnology and
Bioinformatics,
Jaypee University of Information
Technology,
Solan, Himachal Pradesh, India
Email: hchangotra@yahoo.com
harish.changotra@juit.ac.in



How to Cite:

First Author name *et. al.* Title of the Manuscript.
IJCB. 201x; Volume 4 (Issue 1): Page 55-61.

1. INTRODUCTION

Immunity-related GTPase family M (IRGM) protein belongs to immunity-related GTPase (IRG) family of proteins, which was described first time in 1990s, and is further divided into five subfamilies namely IRGA, IRGB, IRGC, IRGD and IRGM based on homology of GTP-binding domain. Interferons can induce the IRG family genes in mice and these genes have been shown to play a critical role in resistance to a range of intracellular pathogens that comprises *Mycobacterium tuberculosis*, *Salmonella typhimurium*, *Listeria monocytogenes*, *Toxoplasma gondii*, and *Chlamydia trachomatis* (1-4). The IRG proteins consist of an N-terminal terminal GTP-binding domain, called G-domain, and a highly variable C-terminal region. The IRG family G-domain includes all five GTP-binding motifs (5). IRG genes have homology only to conserved G-domain of other GTPases while other regions of these proteins are variable. N- and C-terminal regions of these have characteristic features that distinguish this family of proteins from other P-loop GTPases. The IRG proteins are divided into two structural subfamilies, namely GMS and GKS. The latter is based on a methionine (M) substitution in place of lysine (K) in the G1 motif (GX4GK/MS) and is a typical characteristic of the GMS type protein (6). The IRG family contains 23 genes in mice and these are found as a tandem copies in the genome.

They have been shown to play role in immune response to pathogens and are one of the strongest pathogen resistance systems in mice.

The IRG family in humans contains two genes; an IRGC and truncated gene IRGM. Human IRGM is homologous to mouse IRGM genes of GMS subfamily and is truncated in humans and codes for N- and C-terminally truncated G-domain.

2. HUMAN IRGM GENE STRUCTURAL ORGANIZATION AND ITS EVOLUTION:

The human *irgm* gene, unlike its mouse ortholog, is not inducible by interferon-gamma (IFN- γ) as promoter region has lost the interferons response element (6, 7). IRGM gene is localized at chromosome 5q33.1 and comprises 5 exons, a long coding first exon and 4 shorter exons that extend more than 50 kb downstream from the first exon. This gene could be expressed in five different 3'-splice isoforms that have different C-terminal ends but their endogenous expression needs to be documented for its all forms. The endogenous IRGM has been shown to localize in mitochondria (8). The mRNA transcripts of five 3'-splicing isoforms (IRGM a - e) extends ~30kb 3' of the first long coding exon. All these different transcripts containing the G-domain of IRGM have been cloned applying 5'- and 3' rapid amplification of cDNA ends (RACE) technology from cultured human cell lines (6, 7). From all known transcripts, IRGMa is the shortest; its termination occurs at the polyadenylation signal sequence at the start of the second intron and made up of 2 exons that encode polypeptide of 181 amino acids. The other transcripts (IRGM b - e) are longer and include two or more downstream exons. Transcripts of this gene have a long 5'- untranslated region (UTR) and consists of promoter region made up by endogenous retroviral nine (ERV9: U3 and U5 region), Alu sequence (AluSc), and a 3'-UTR. The latter encompasses alternatively spliced intronic sequence and exon-intron boundaries and an upstream termination codon in IRGM c - e. Based on the presence of this termination codon, these would be expected to undergo rapid RNA degradation via nonsense-mediated decay. Additionally, there exists a 20.1 kb deletion polymorphism immediately upstream to IRGM gene which has been shown to associate with its altered expression and risk factor for Crohn's disease (9).

Moreover, IRGM has N- and C-terminal regions both with truncation and could not be induced by interferons. This suggested the functions this gene is other than its mouse ortholog. Moreover, because of this reason it was earlier thought be a pseudogene. Bekpen et al (7) have done comprehensive analysis of this gene and have shown that IRG gene cluster was non-functional during primate evolution, approximately 50 million years ago; when the divergence of the anthropoids from the prosimians took place, and the copy number of this gene was reduce to one. Sequence analysis of new and old world monkey species showed that insertion of Alu sequence in the open reading frame (ORF) in the anthropoid common ancestor interrupted this gene making it pseudogene. Further, the function of IRGM was restored, ~20 million years ago, in the ancestors of apes and humans by the insertion of an ERV9 element at 5'- upstream region of IRGM. In addition to this, there are evidences showing the existence of both functional as well as non-functional gene (7, 10). The promoter region corresponds to the ERV9 U3 long terminal repeats which is without interferon response element and could be a possible explanation why this gene is not responsive to the interferons.

3. ROLE OF HUMAN IRGM IN DISEASES:

Earlier, it thought that this gene is non-functional since it was unresponsive to the interferons, unlike other members of its family (6). N- and C-terminal regions of IRGM are truncated which further supports it may have non-functional nature. Moreover, evolutionary history also suggested that this gene was disrupted and non-functional for about 25 million years ago. But, recently scientists have started to unravel the roles played by IRGM and the data suggested critical roles of this gene in controlling infection with the various pathogens in humans (7, 11-14). *Bioinformatics has been instrumental in elucidating its evolutionary history as well as functions especially in its interactions with the proteins of autophagy pathway* (7, 15).

Macroautophagy (referred as autophagy thereafter) pathway which is a conserved eukaryotic catabolic process by which long-lived proteins and damaged organelles are sequestered in the cytoplasm and removed for recycling through lysosomal processes (16). Mechanistically, autophagy starts with the formation of double-membrane vesicles structures called autophagosomes or autophagic vacuoles that engulf cytosolic material, which include damaged organelles or denatured proteins. The autophagosomes so formed fuse with the lysosomes to form single-membrane autolysosomes. Lysosomal enzymes degrade engulfed organelles and regenerate smaller molecules (peptides, amino acids and fatty acids), which are used as a source of energy as well as in anabolic pathways (16-18). Autophagy has been described as a mechanism of both innate and adaptive immunity against intracellular bacteria, viruses, parasites and other diseases (19, 20). Either hosts or

pathogens exploit this process for their benefit and its dysfunction has been implicated in many diseases that include neurodegenerative diseases, muscle diseases, cardiac diseases and infectious diseases (21).

a. IRGM in viral infections:

Currently, sufficient data are available which demonstrate that number of viruses has developed various tactics to modulate autophagy for their benefits. Gre'goire et al (15) has mapped the molecular interactions between the proteins involved in the autophagy pathway and viral proteins. Their data demonstrated the physical interactions between 44 human autophagy-associated proteins and 83 proteins of several RNA viruses belonging to various different families, namely *Bornaviridae*, *Coronaviridae*, *Paramyxoviridae*, *Flaviviridae*, *Orthomyxoviridae*, *Retroviridae* and *Togaviridae*. IRGM was the most commonly targeted protein, interacted with the autophagy pathway proteins, ATG5, ATG10, MAP1CL3C and SH3GLB1 and can interact with 12 viral proteins of Mumps virus (MuV), Hepatitis C virus (HCV), Measles virus (MeV), HIV and Chikungunya virus (ChikV) (15). All these proteins regulate one of the earlier steps of the of autophagosome formation, suggesting that IRGM might play a role in the nucleation or the elongation step of autophagic through its interactions with these proteins. These interactions could be facilitated by viral infection (15), which in turn modulate autophagy for their benefit (22).

Infection with MeV induces autophagosomes and MeV virions production could be reduced by inhibiting autophagy suggesting that it uses this pathway to promote its replication (23). siRNA mediated reduction of IRGM expression resulted in decreased levels of MeV replication. Furthermore, it has been shown that IRGM interacts with MeV-C non-structural protein and induces autophagy which implicate MeV might utilize IRGM mediated modulation of autophagy for its replication (15, 23, 24).

HIV-1 also modulates autophagy but the strategy depends on the type of cells infected. It can infect both CD4+ T cells and macrophages resulting in production of high levels of virions. But, its infection leads to death of CD4+ T cells while its cytopathic effects are resisted by macrophages (25, 26). When uninfected CD4+ T cells were co-cultured with lymphocytes expressing Env protein, it induced autophagy that was through binding of Env to its receptors CD4 and CXCR4, but independent of CD4 or CXCR4 signaling pathways that leads to apoptotic cell death of these T cells (27, 28). Additionally, HIV inhibits autophagy in infected CD+ T cells to enhance its replication (28). Dendritic cells (DCs) infected with HIV shows inhibited autophagy and which is through mTOR signaling pathway (29), while in macrophages shows induced autophagy, which is through interactions of HIV-1-NEF and BECN1 (28, 30). Furthermore, NEF proteins can interact with IRGM and overexpression of NEF in HeLa cells shows higher levels of IRGM-dependent autophagosomes accumulation (15). NEF and IRGM has also been shown to co-localize, therefore, NEF-IRGM might be involved in induction of autophagy.

HCV also induces autophagy which is independent of mTOR and inhibition of autophagy results in reduction of autophagy. It has been shown that HCV non-structural proteins NS3-5B induce autophagy. Recent data demonstrated IRGM interactions with HCV-NS3 and siRNA mediated reduction in IRGM expression impair HCV replication (15, 31-36).

Chikungunya virus infection also induces autophagy and is required for its replication, as inhibition and stimulation of the same, respectively, results in impairment and augmentation of the replication of ChikV (37, 38). IRGM interacts with ChikV-NS2 and E3 proteins which point to role of this protein in viral replication that needs to be tested further (15). *Therefore, several viruses target IRGM and manipulate autophagy for their benefit.*

b. Bacterial diseases:

Singh et al (39) for the first time have shown that IRGM is a functional gene and is involved in the control intracellular pathogen, i.e. *Mycobacterium tuberculosis* (*M. tuberculosis*), through autophagy-mediated pathway. They treated U937 cells, human macrophage cell line, with hIFN- γ (which induces autophagy in macrophages) followed by inhibition of IRGM expression using siRNA, a decrease in LC3 puncta were observed compared to controls. They have also shown rapamycin induced autophagy also inhibited LC3II formation suggesting crucial role of IRGM in proper functioning of this pathway (39). Moreover, it has also been shown that IRGM-dependent autophagy induction is needed for *M. tuberculosis* phagosome maturation in human macrophages and knockdown of IRGM results in augmented survival. Therefore, IRGM plays an important role in limiting mycobacteria in macrophages. Further, genome wide association studies have shown the involvement of IRGM polymorphisms in the susceptibility to *M. tuberculosis*. Itemann et al (40) found that *IRGM* -261TT genotype linked to *M. tuberculosis* infection risk but not *M. africanum* clades in a Ghanese

population while Chen et al (19) studied Chinese population and found -1208A/G promoter polymorphisms associated with susceptibility to tuberculosis. King et al (41) showed single nucleotide polymorphism rs10065172 C/T present in IRGM exon region associated with its decreased expression in lymphoblastoid cell line predispose humans to the increased risk to tuberculosis.

Mycobacterium leprae infection of CD4+ T cells, monocytes and monocyte-derived macrophages from leprosy patients were found to have higher levels of both, protein and mRNA of IRGM. Moreover, levels of IRGM were related to the severity of the disease, implicating its role in defending infection of *M. leprae* (12).

Based on data that suggest requirement of autophagy in the clearance of *Salmonella typhimurium* (42, 43), Huett et al., (44) reasoned that IRGM might be involved in controlling infection with this pathogen and have demonstrated that IRGM plays an essential role in autophagy mediated removal of this pathogen. siRNA mediated knockdown of IRGM reduced the number of bacteria within autophagosomes while its overexpression showed increased autophagic activity.

c. Other diseases:

IRGM has also been shown to play a crucial role in Crohn's disease (CD), which is a chronic inflammatory bowel disease. The mechanisms causing this disease are still not completely understood but involve heightened inflammatory responses and impaired bacterial clearance (45, 46). Recent studies have suggested links between autophagy and Crohn's diseases. Genome-wide association studies (GWAS) have revealed single nucleotide polymorphisms (SNPs) in autophagy genes including IRGM, which are known to influence autophagic processing, associated with susceptibility to Crohn's disease (47-49). Several SNPs (rs13361189, rs10065172, rs4958847, rs1000113, rs931058, rs11747270, rs1000113, rs931058 etc.) have been shown to be associated with susceptibility to Crohn's disease in various populations (50-52). Moreover, a protective exonic synonymous SNP (C313T) binds with the miR-196 and downregulates IRGM protective variant but not risk-associated variant. This binding compromises control of replication of intracellular adherent invasive *Escherichia coli* associated with Crohn's disease, predisposing these individuals to higher risk (53).

4. CONCLUSION:

The role of IRGM in humans has started to emerge. Recently, several studies have shown the involvement of this protein in various diseases that includes viral, bacterial, parasitic and inflammatory diseases. *Bioinformatics has played a pivotal role in deciphering its evolution as well as functions of IRGM.* Further detailed studies are needed to find how IRGM is involved in various diseases and to find if this could be targeted to modulated autophagy to control these diseases.

ACKNOWLEDGMENTS:

Harish Changotra is thankful to Department of Science and Technology, Government of India and Department of Biotechnology, Government of India, respectively, for the grants SB/FT/LS-440/2012 and BT/PR6784/GBD/27/466/2012.

REFERENCES:

1. MacMicking JD, Taylor GA, McKinney JD. Immune control of tuberculosis by IFN-gamma-inducible LRG-47. *Science*. 2003;302(5645):654-9. Epub 2003/10/25.
2. Taylor GA, Collazo CM, Yap GS, Nguyen K, Gregorio TA, Taylor LS, et al. Pathogen-specific loss of host resistance in mice lacking the IFN-gamma-inducible gene IGTP. *Proceedings of the National Academy of Sciences of the United States of America*. 2000;97(2):751-5. Epub 2000/01/19.
3. Taylor GA, Feng CG, Sher A. p47 GTPases: regulators of immunity to intracellular pathogens. *Nature reviews Immunology*. 2004;4(2):100-9. Epub 2004/03/26.
4. Taylor GA. IRG proteins: key mediators of interferon-regulated host resistance to intracellular pathogens. *Cell Microbiol*. 2007;9(5):1099-107. Epub 2007/03/16.
5. Ghosh A, Uthaiyah R, Howard J, Herrmann C, Wolf E. Crystal structure of IIGP1: a paradigm for interferon-inducible p47 resistance GTPases. *Molecular cell*. 2004;15(5):727-39. Epub 2004/09/08.

6. Bekpen C, Hunn JP, Rohde C, Parvanova I, Guethlein L, Dunn DM, et al. The interferon-inducible p47 (IRG) GTPases in vertebrates: loss of the cell autonomous resistance mechanism in the human lineage. *Genome biology*. 2005;6(11):R92. Epub 2005/11/10.
7. Bekpen C, Marques-Bonet T, Alkan C, Antonacci F, Leogrande MB, Ventura M, et al. Death and resurrection of the human IRGM gene. *PLoS Genet*. 2009;5(3):e1000403. Epub 2009/03/07.
8. Singh SB, Ornatowski W, Vergne I, Naylor J, Delgado M, Roberts E, et al. Human IRGM regulates autophagy and cell-autonomous immunity functions through mitochondria. *Nat Cell Biol*. 2010;12(12):1154-65. Epub 2010/11/26.
9. McCarroll SA, Huett A, Kuballa P, Chilewski SD, Landry A, Goyette P, et al. Deletion polymorphism upstream of IRGM associated with altered IRGM expression and Crohn's disease. *Nat Genet*. 2008;40(9):1107-12. Epub 2009/01/24.
10. Zhang ZD, Frankish A, Hunt T, Harrow J, Gerstein M. Identification and analysis of unitary pseudogenes: historic and contemporary gene losses in humans and other primates. *Genome biology*. 2010;11(3):R26. Epub 2010/03/10.
11. Brain O, Cooney R, Simmons A, Jewell D. Functional consequences of mutations in the autophagy genes in the pathogenesis of Crohn's disease. *Inflamm Bowel Dis*. 2012;18(4):778-81. Epub 2011/08/11.
12. Yang D, Chen J, Zhang L, Cha Z, Han S, Shi W, et al. Mycobacterium leprae upregulates IRGM expression in monocytes and monocyte-derived macrophages. *Inflammation*. 2014;37(4):1028-34. Epub 2014/01/29.
13. Kimura T, Watanabe E, Sakamoto T, Takasu O, Ikeda T, Ikeda K, et al. Autophagy-related IRGM polymorphism is associated with mortality of patients with severe sepsis. *PLoS One*. 2014;9(3):e91522. Epub 2014/03/15.
14. Xia F, Li R, Wang C, Yang S, Tian L, Dong H, et al. IRGM1 regulates oxidized LDL uptake by macrophage via actin-dependent receptor internalization during atherosclerosis. *Sci Rep*. 2013;3:1867. Epub 2013/05/22.
15. Gregoire IP, Richetta C, Meyniel-Schicklin L, Borel S, Pradezynski F, Diaz O, et al. IRGM is a common target of RNA viruses that subvert the autophagy network. *PLoS Pathog*. 2011;7(12):e1002422. Epub 2011/12/17.
16. Mizushima N, Levine B, Cuervo AM, Klionsky DJ. Autophagy fights disease through cellular self-digestion. *Nature*. 2008;451(7182):1069-75. Epub 2008/02/29.
17. Kelekar A. Autophagy. *Annals of the New York Academy of Sciences*. 2005;1066:259-71. Epub 2006/03/15.
18. He C, Klionsky DJ. Regulation mechanisms and signaling pathways of autophagy. *Annual review of genetics*. 2009;43:67-93. Epub 2009/08/06.
19. Che N, Li S, Gao T, Zhang Z, Han Y, Zhang X, et al. Identification of a novel IRGM promoter single nucleotide polymorphism associated with tuberculosis. *Clin Chim Acta*. 2010;411(21-22):1645-9. Epub 2010/06/16.
20. Kudchodkar SB, Levine B. Viruses and autophagy. *Reviews in medical virology*. 2009;19(6):359-78. Epub 2009/09/15.
21. Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Cell*. 2008;132(1):27-42. Epub 2008/01/15.
22. Petkova DS, Viret C, Faure M. IRGM in autophagy and viral infections. *Front Immunol*. 2012;3:426. Epub 2013/01/22.
23. Joubert PE, Meiffren G, Gregoire IP, Pontini G, Richetta C, Flacher M, et al. Autophagy induction by the pathogen receptor CD46. *Cell Host Microbe*. 2009;6(4):354-66. Epub 2009/10/20.
24. Meiffren G, Joubert PE, Gregoire IP, Codogno P, Roubourdin-Combe C, Faure M. Pathogen recognition by the cell surface receptor CD46 induces autophagy. *Autophagy*. 2010;6(2):299-300. Epub 2010/01/21.
25. Borel S, Espert L, Biard-Piechaczyk M. Macroautophagy Regulation during HIV-1 Infection of CD4+ T Cells and Macrophages. *Front Immunol*. 2012;3:97. Epub 2012/05/16.
26. Carter CA, Ehrlich LS. Cell biology of HIV-1 infection of macrophages. *Annual review of microbiology*. 2008;62:425-43. Epub 2008/09/13.
27. Espert L, Denizot M, Grimaldi M, Robert-Hebmann V, Gay B, Varbanov M, et al. Autophagy is involved in T cell death after binding of HIV-1 envelope proteins to CXCR4. *The Journal of clinical investigation*. 2006;116(8):2161-72. Epub 2006/08/04.
28. Espert L, Varbanov M, Robert-Hebmann V, Sagnier S, Robbins I, Sanchez F, et al. Differential role of autophagy in CD4 T cells and macrophages during X4 and R5 HIV-1 infection. *PLoS One*. 2009;4(6):e5787. Epub 2009/06/06.

29. Blanchet FP, Moris A, Nikolic DS, Lehmann M, Cardinaud S, Stalder R, et al. Human immunodeficiency virus-1 inhibition of immunoamphisomes in dendritic cells impairs early innate and adaptive immune responses. *Immunity*. 2010;32(5):654-69. Epub 2010/05/11.
30. Kyei GB, Dinkins C, Davis AS, Roberts E, Singh SB, Dong C, et al. Autophagy pathway intersects with HIV-1 biosynthesis and regulates viral yields in macrophages. *The Journal of cell biology*. 2009;186(2):255-68. Epub 2009/07/29.
31. Su WC, Chao TC, Huang YL, Weng SC, Jeng KS, Lai MM. Rab5 and class III phosphoinositide 3-kinase Vps34 are involved in hepatitis C virus NS4B-induced autophagy. *Journal of virology*. 2011;85(20):10561-71. Epub 2011/08/13.
32. Shrivastava S, Bhanja Chowdhury J, Steele R, Ray R, Ray RB. Hepatitis C virus upregulates Beclin1 for induction of autophagy and activates mTOR signaling. *Journal of virology*. 2012;86(16):8705-12. Epub 2012/06/08.
33. Sir D, Chen WL, Choi J, Wakita T, Yen TS, Ou JH. Induction of incomplete autophagic response by hepatitis C virus via the unfolded protein response. *Hepatology*. 2008;48(4):1054-61. Epub 2008/08/09.
34. Mohl BP, Tedbury PR, Griffin S, Harris M. Hepatitis C virus-induced autophagy is independent of the unfolded protein response. *Journal of virology*. 2012;86(19):10724-32. Epub 2012/07/28.
35. Dreux M, Gastaminza P, Wieland SF, Chisari FV. The autophagy machinery is required to initiate hepatitis C virus replication. *Proceedings of the National Academy of Sciences of the United States of America*. 2009;106(33):14046-51. Epub 2009/08/12.
36. Tanida I, Fukasawa M, Ueno T, Kominami E, Wakita T, Hanada K. Knockdown of autophagy-related gene decreases the production of infectious hepatitis C virus particles. *Autophagy*. 2009;5(7):937-45. Epub 2009/07/25.
37. Joubert PE, Werneke S, de la Calle C, Guivel-Benhassine F, Giodini A, Peduto L, et al. Chikungunya-induced cell death is limited by ER and oxidative stress-induced autophagy. *Autophagy*. 2012;8(8):1261-3. Epub 2012/07/28.
38. Krejbich-Trotot P, Gay B, Li-Pat-Yuen G, Hoarau JJ, Jaffar-Bandjee MC, Briant L, et al. Chikungunya triggers an autophagic process which promotes viral replication. *Virology journal*. 2011;8:432. Epub 2011/09/10.
39. Singh SB, Davis AS, Taylor GA, Deretic V. Human IRGM induces autophagy to eliminate intracellular mycobacteria. *Science*. 2006;313(5792):1438-41. Epub 2006/08/05.
40. Intemann CD, Thye T, Niemann S, Browne EN, Amanua Chinbuah M, Enimil A, et al. Autophagy gene variant IRGM -261T contributes to protection from tuberculosis caused by *Mycobacterium tuberculosis* but not by *M. africanum* strains. *PLoS Pathog*. 2009;5(9):e1000577. Epub 2009/09/15.
41. King KY, Lew JD, Ha NP, Lin JS, Ma X, Graviss EA, et al. Polymorphic allele of human IRGM1 is associated with susceptibility to tuberculosis in African Americans. *PLoS One*. 2011;6(1):e16317. Epub 2011/02/02.
42. Rioux JD, Xavier RJ, Taylor KD, Silverberg MS, Goyette P, Huett A, et al. Genome-wide association study identifies new susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis. *Nat Genet*. 2007;39(5):596-604. Epub 2007/04/17.
43. Birmingham CL, Smith AC, Bakowski MA, Yoshimori T, Brumell JH. Autophagy controls *Salmonella* infection in response to damage to the *Salmonella*-containing vacuole. *J Biol Chem*. 2006;281(16):11374-83. Epub 2006/02/24.
44. Huett A, McCarroll SA, Daly MJ, Xavier RJ. On the level: IRGM gene function is all about expression. *Autophagy*. 2009;5(1):96-9. Epub 2008/11/26.
45. Podolsky DK. Inflammatory bowel disease. *The New England journal of medicine*. 2002;347(6):417-29. Epub 2002/08/09.
46. Choi AJ, Ryter SW. Autophagy in inflammatory diseases. *International journal of cell biology*. 2011;2011:732798. Epub 2011/12/23.
47. Massey DC, Parkes M. Genome-wide association scanning highlights two autophagy genes, ATG16L1 and IRGM, as being significantly associated with Crohn's disease. *Autophagy*. 2007;3(6):649-51. Epub 2007/10/09.
48. Barrett JC, Hansoul S, Nicolae DL, Cho JH, Duerr RH, Rioux JD, et al. Genome-wide association defines more than 30 distinct susceptibility loci for Crohn's disease. *Nat Genet*. 2008;40(8):955-62. Epub 2008/07/01.
49. Parkes M, Barrett JC, Prescott NJ, Tremelling M, Anderson CA, Fisher SA, et al. Sequence variants in the autophagy gene IRGM and multiple other replicating loci contribute to Crohn's disease susceptibility. *Nat Genet*. 2007;39(7):830-2. Epub 2007/06/08.
50. Glas J, Seiderer J, Bues S, Stallhofer J, Fries C, Olszak T, et al. IRGM variants and susceptibility to inflammatory bowel disease in the German population. *PLoS One*. 2013;8(1):e54338. Epub 2013/02/01.

51. Jung C, Colombel JF, Lemann M, Beaugerie L, Allez M, Cosnes J, et al. Genotype/phenotype analyses for 53 Crohn's disease associated genetic polymorphisms. *PLoS One*. 2012;7(12):e52223. Epub 2013/01/10.
52. Baskaran K, Pugazhendhi S, Ramakrishna BS. Association of IRGM gene mutations with inflammatory bowel disease in the Indian population. *PLoS One*. 2014;9(9):e106863. Epub 2014/09/06.
53. Brest P, Lapaquette P, Souidi M, Lebrigand K, Cesaro A, Vouret-Craviari V, et al. A synonymous variant in IRGM alters a binding site for miR-196 and causes deregulation of IRGM-dependent xenophagy in Crohn's disease. *Nat Genet*. 2011;43(3):242-5. Epub 2011/02/01.