

A Volume Limited Sample of Cataclysmic Variables from *Gaia* DR2: Space Density and Population Properties

A. F. Pala,^{1,2*} B. T. Gänsicke,² E. Breedt,³ C. Knigge,⁴ J. J. Hermes,⁵
 N. P. Gentile Fusillo,^{1,2} M. A. Hollands,² T. Naylor,⁶ I. Pelisoli,⁷ M. R. Schreiber,⁸
 S. Toonen,⁹ A. Aungwerojwit,¹⁰ E. Cukanovaite,² E. Denny,^{11,12} C. J. Manser,²
 M. L. Pretorius,^{13,14} S. Scaringi,¹⁵ O. Toloza²

¹European Southern Observatory, Karl Schwarzschild Straße 2, Garching, 85748, Germany

²Department of Physics, University of Warwick, Coventry, CV4 7AL, UK

³Institute of Astronomy, University of Cambridge, Cambridge, CB3 0HA, UK

⁴School of Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, UK

⁵Department of Astronomy, Boston University, 725 Commonwealth Ave., Boston, MA 02215, USA

⁶School of Physics, University of Exeter, Stocker Road, Exeter EX4 4QL, UK

⁷Institut für Physik und Astronomie, Universitätsstandort Golm, Karl-Liebknecht-Str 24/25, D-14476 Potsdam, Germany

⁸Instituto de Física y Astronomía, Millennium Nucleus for Planet Formation (NPF), Universidad de Valparaíso, 2360102 Valparaíso, Chile

⁹Institute for Gravitational Wave Astronomy, School of Physics and Astronomy, West office 237, Birmingham B15 2TT, Edgbaston, UK

¹⁰Department of Physics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand

¹¹University of North Carolina, Department of Physics and Astronomy, Chapel Hill, NC - 27599-3255, USA

¹²Gemini Observatory, Southern Operations Center, Casilla 603, La Serena, Chile

¹³Department of Astronomy, University of Cape Town, Private Bag X3, Rondebosch 7701, South Africa

¹⁴South African Astronomical Observatory, PO Box 9, Observatory 7935, Cape Town, South Africa

¹⁵Department of Physics and Astronomy, Texas Tech University, Lubbock, TX 79409-1051, USA

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We present the first volume-limited sample of cataclysmic variables (CVs), selected using the accurate parallaxes provided by the second data release (DR2) of the ESA *Gaia* space mission. The sample is composed of 42 CVs within 150 pc, including two new systems discovered using the *Gaia* data, and is (77 ± 10) per cent complete. We use this sample to study the intrinsic properties of the Galactic CV population. In particular, the CV space density we derive, $\rho = (4.8^{+0.6}_{-0.8}) \times 10^{-6} \text{ pc}^{-3}$, is lower than predicted by most binary population synthesis studies. We also find a low fraction of period bounce CVs, seven per cent, and an average white dwarf mass of $\langle M_{\text{WD}} \rangle = (0.83 \pm 0.17) M_{\odot}$. Both findings confirm previous results, ruling out the presence of observational biases affecting these measurements, as has been suggested in the past. The observed fraction of period bounce CVs falls well below theoretical predictions, by at least a factor of five, and remains one of the open problems in the current understanding of CV evolution. Conversely, the average white dwarf mass supports the presence of additional mechanisms of angular momentum loss that have been accounted for in the latest evolutionary models. The fraction of magnetic CVs in the 150 pc sample is remarkably high at 36 per cent. This is in striking contrast with the absence of magnetic white dwarfs in the detached population of CV progenitors, and underlines that the evolution of magnetic systems has to be included in the next generation of population models.

Key words: Hertzsprung-Russell and colour-magnitude diagrams – cataclysmic variables – stars: statistics – stars:evolution

1 INTRODUCTION

Cataclysmic variables (CVs) are compact interacting binaries containing a white dwarf accreting from a Roche-lobe

* E-mail: apala@eso.org

filling donor star (see Warner 1995, for a comprehensive review). In most systems, these companions are low-mass, late-type stars. If the white dwarf is not magnetised ($B \lesssim 1$ MG), the mass lost from the donor forms an accretion disc around the white dwarf. In the presence of stronger magnetic fields, the disc is either truncated at the magnetospheric radius of the white dwarf ($1 \text{ MG} \lesssim B \lesssim 10 \text{ MG}$) or is fully suppressed ($B \gtrsim 10 \text{ MG}$). In these magnetic CVs, known as intermediate polars (IPs) and polars, respectively, the accretion flow follows the field lines and accretes onto the white dwarf at its magnetic poles.

The evolution of CVs is, as for all types of interacting binaries, dictated by orbital angular momentum losses (AMLs) and by the internal structure of the companion star (Rappaport et al. 1983; Paczynski & Sienkiewicz 1983; Spruit & Ritter 1983; Knigge et al. 2011). In fact, the time scale at which the secondary star loses mass is comparable to its thermal time scale, resulting in a donor which is slightly out of thermal equilibrium and hotter and bloated compared to an isolated main sequence star of the same mass. This deviation from thermal equilibrium is thought to be the cause of the major features observed in the CV orbital period distribution: the “period gap” and the “period minimum”, as further detailed below.

As angular momentum is removed from the system, the orbital separation decreases and, consequently CVs evolve from long to short orbital periods (Rappaport et al. 1983; Paczynski & Sienkiewicz 1983; Spruit & Ritter 1983). At long orbital periods ($P_{\text{orb}} \gtrsim 3 \text{ h}$) CV evolution is driven by magnetic wind braking (MB) and gravitational wave radiation (GWR). The ongoing mass transfer monotonously erodes the secondary star which, at $P_{\text{orb}} \approx 3 \text{ h}$, becomes fully convective. In the standard framework of CV evolution, it is assumed that a re-configuration of the magnetic fields on the donor results in a greatly reduced efficiency of MB from that point onwards, and the secondary star detaches from its Roche lobe. In the period range $2 \text{ h} \lesssim P_{\text{orb}} \lesssim 3 \text{ h}$, the so-called period gap, the system evolves as a detached binary whilst still losing angular momentum through GWR. Observational support for the disrupted MB hypothesis is provided by the observed properties of the post-common envelope binaries in the period range $2 \text{ h} \lesssim P_{\text{orb}} \lesssim 3 \text{ h}$ (Schreiber et al. 2010; Zorotovic et al. 2016). At $P_{\text{orb}} \approx 2 \text{ h}$, the orbital separation is such that the donor fills its Roche lobe again and the accretion process resumes.

Below the period gap, CVs keep evolving towards shorter orbital periods until they reach the period minimum, $P_{\text{orb}} \approx 80 \text{ min}$. At this stage, the time-scale on which the secondary star loses mass becomes much shorter compared to its thermal time-scale. The donor is driven out of thermal equilibrium and stops shrinking in response to the mass loss. Consequently the system starts evolving back towards longer orbital periods, becoming a “period bouncer”.

With MB being much more efficient than GWR in removing angular momentum from the system, CVs above the period gap are predicted to have mass accretion rate orders of magnitudes higher ($\dot{M} \sim 10^{-9} - 10^{-8} M_{\odot} \text{ yr}^{-1}$, Spruit & Ritter 1983) than those of the CVs below the period gap ($\dot{M} \sim 5 \times 10^{-11} M_{\odot} \text{ yr}^{-1}$, Patterson 1984). While this is roughly in agreement with the accretion rates estimated from observations (Townsend & Gänsicke 2009; Pala et al. 2017), the theoretical framework outlined above fails to re-

produce a number of observational properties of the Galactic population of CVs: (i) the predicted fractions of CVs above and below the period gap (≈ 1 per cent and ≈ 99 per cent, respectively, de Kool 1992; Kolb 1993; Howell et al. 2001) are in clear disagreement with the observations (e.g. ≈ 23 per cent and ≈ 77 per cent, Gänsicke et al. 2009, though the observed samples are typically magnitude-limited, and hence biased towards more luminous CVs); (ii) period bouncers are expected to be the main component ($\approx 40 - 70$ per cent) of the present-day Galactic CV population but only a small number of such systems has been identified (Patterson et al. 2005; Unda-Sanzana et al. 2008; Littlefair et al. 2006; Patterson 2011; Kato et al. 2015, 2016; McAllister et al. 2017; Neustroev et al. 2017; Pala et al. 2018); (iii) there are clues of the presence of additional AML mechanisms that are not accounted for by the standard model of CV evolution (Patterson 1998; Knigge et al. 2011; Schreiber et al. 2016; Pala et al. 2017; Zorotovic & Schreiber 2017; Belloni et al. 2018; Liu & Li 2019), although the physical origin of this enhanced AML is still unclear.

A key parameter that provides stringent constraints on the models of CV formation and evolution is their space density, ρ_0 . Binary population synthesis studies carried out by de Kool (1992) and Politano (1996) suggested CV space densities of $\approx 2 \times 10^{-5} - 2.0 \times 10^{-4} \text{ pc}^{-3}$. More recent works by Goliash & Nelson (2015), which also accounts for the presence of CVs containing nuclear evolved donors, and by Belloni et al. (2018) provide an estimate of $\approx (1.0 \pm 0.5) \times 10^{-5} \text{ pc}^{-3}$ and $\lesssim 2 \times 10^{-5} \text{ pc}^{-3}$, respectively, comparable to the earlier results.

These predicted values are systematically larger than the ones derived from observations. For example, Thomas & Beuermann (1998) used the *ROSAT* All Sky Survey to infer $\rho_0 \approx 6.1 \times 10^{-7} \text{ pc}^{-3}$ for polars. Later studies based on the *ROSAT* Bright Survey (RBS) and the *ROSAT* North Ecliptic Pole (NEP) suggested $\rho_0 = 4_{-2}^{+6} \times 10^{-6} \text{ pc}^{-3}$ (Pretorius et al. 2007a; Pretorius & Knigge 2012). Most recently, Hernández Santisteban et al. (2018) estimated an upper limit on the space density of period bounce CVs from a search for eclipsing systems in Stripe82 from the Sloan Digital Sky Survey (SDSS, York et al. 2000), finding $\rho_0 \lesssim 2 \times 10^{-5} \text{ pc}^{-3}$.

In the past, an accurate measurement of the CV space density has been challenged by the lack of accurate distances. In April 2018, the European Space Agency (ESA) *Gaia* space mission has delivered parallaxes for more than one billion stars in its second data release DR2 (Gaia Collaboration et al. 2016, 2018), providing the first opportunity to construct a volume-limited sample of CVs and to derive their intrinsic properties. Schwöpe (2018) carried out the first application of the *Gaia* data in this context, and, using the distances from *Gaia* for a X-ray selected sample of CVs from RBS, NEP and *Swift*/BAT survey, derived the space densities of IPs, $\rho_0 < 1.3 \times 10^{-7} \text{ pc}^{-3}$, and non-magnetic CVs, $\rho_0 < 5.1 \times 10^{-6} \text{ pc}^{-3}$. These results are based on the assumptions that the X-ray selected sample is complete and representative of the intrinsic population. However, as discussed by Pretorius & Knigge (2012), it is possible that a large fraction of faint CVs may not have been detected in the RBS and NEP surveys, and that the space density derived from the corresponding X-ray selected CV sample could be easily underestimated by a factor ≈ 2 .




Figure 1. Hertzsprung-Russell diagram of the *Gaia* sources with reliable astrometry (Equation 4) located within 150 pc (grey), showing the position of the 150 pc CVs (pink dots, Section 3) and of the different types of contaminants (Section 2) that have been mistakenly identified as CVs or CV candidates within the literature (Appendices A, B and C). For comparison, SDSS CVs with accurate and clean SDSS photometry (as defined in Section 4) are shown by the black crosses, however the majority of these systems are at distances $d > 150$ pc. Systems discarded for having high astrometric excess noise (AEN) in *Gaia* are shown in yellow.

Here we present a study of the first volume-limited sample of CVs within 150 pc, selected by combining the *Gaia* DR2 parallaxes and the available information from a large number of spectroscopic and photometric surveys.

The sample contains a total of 42 objects and provides the first direct insight into the intrinsic properties of the Galactic population of CVs.

2 SAMPLE SELECTION

The *Gaia* space mission provides the first opportunity to construct a volume-limited sample of CVs that allows us to infer the intrinsic properties of the Galactic CV population.

The first step towards this goal is the choice of an optimal volume, which is sufficiently large to be representative of the entire CV population and robust against small number statistics, and not subject to distance (magnitude) related observational biases.

The volume enclosed within 150 pc represents a good compromise between the two requirements: assuming the typical space density derived from binary population synthesis studies, $\lesssim 2 \times 10^{-5} \text{ pc}^{-3}$ (Belloni et al. 2018), it is expected to contain ≈ 140 CVs, and at $d = 150$ pc even the systems with the lowest accretion rates¹ should be bright

enough to have accurate astrometric solutions (typical uncertainties on the parallaxes are ≈ 0.2 mas for $G \lesssim 19$ mag, see Figure 7 in Gaia Collaboration et al. 2018). Restricting $d \leq 150$ pc also reduces the uncertainties in the derived space densities related to the unknown age and scale height of the CV population (see Section 3).

The next step is to compile a list of all CVs and CV candidates that could plausibly be within $d \leq 150$ pc. CVs are mainly discovered thanks to their outbursting properties. In fact, CV accretion discs undergo thermal instabilities called dwarf nova outbursts (Osaki 1974; Meyer & Meyer-Hofmeister 1984; Hameury et al. 1998), during which CV systems brighten up to 2–9 mag and these outbursts can last for days up to weeks (Warner 1995; Maza & Gonzalez 1983). Many surveys search the sky nightly for transient events, such as the Catalina Real-time Transient Survey (CRTS, Drake et al. 2009), the All-Sky Automated Survey and the All-Sky Automated Survey for Supernovae (ASAS and ASAS-SN, Pojmanski 1997; Shappee et al. 2014; Kochanek et al. 2017), the Mobile Astronomical System of Telescope Robots (MASTER, Lipunov et al. 2010), the Palomar Transient Factory (PTF, Law et al. 2009) and the Intermediate PTF (iPTF, Kulkarni 2013), the Asteroid Terrestrial-impact Last Alert System (ATLAS; Tonry et al. 2018), the

at a distance of 187 pc ($\varpi = 5.3 \pm 0.3$ mas) and is as bright as $G = 18.9$ mag (Pala et al. 2018).

¹ e.g. QZLib, one of the few period bouncers known, is located




Figure 2. Identification spectra of the two new CVs we discovered within 150 pc. *Gaia* J051903.99+630340.4 (top) presents the typical spectral appearance of an SU UMa CV as it is also confirmed by its orbital period, $P_{\text{orb}} \simeq 126$ min (Appendix C), which locates it below the period gap. Dwarf nova outbursts of this system have been detected by ASAS-SN in February 2012, March 2016 and September 2018, during which the system brightened, on average, by $\simeq 5.9$ mag. However, in the ASAS-SN observations, the CV is blended with a bright ($V = 10.1$ mag) nearby companion (within $\simeq 7''$, see Section 7). Consequently, during each dwarf nova outburst, only a variation of $\simeq 0.6$ mag is observed in the overall brightness recorded by ASAS-SN of the two stars blended together, thus preventing an earlier detection of *Gaia* J051903.99+630340.4 as a CV. Conversely, *Gaia* J154008.28-392917.6 (bottom) is a low accretion rate system of the WZ Sge type, likely located at the period minimum (note its similarity with e.g. EZ Lyn in Figure 8), which has never been observed in outburst so far.

Zwicky Transient Facility (ZTF; [Graham et al. 2019](#)) and the Gaia Photometric Science Alerts ([Hodgkin et al. 2013](#)), which have been very successful in the identification of thousands of CVs in outbursts (e.g. [Breidt et al. 2014](#)).

Alternatively, CVs can be identified thanks to their blue colours (see e.g. [Gänsicke 2005](#)) and their X-ray emission (the latter favouring magnetic systems since the X-ray emission arises mainly from small, hot region near the magnetic poles of polars and IPs). Finally, CVs are one of by-products of SDSS, with over 300 new CVs discovered in the last decade via the detection of strong emission disc lines in their spectra ([Szkody et al. 2002, 2003, 2004, 2005, 2006, 2007, 2009, 2011](#)). Whilst the spectroscopic method is less affected by selection biases, it is the most expensive in term of telescope time.

New systems are continuously discovered and an up-to-date catalogue of these discoveries is missing. We therefore searched the catalogues of the aforementioned transient surveys and the International Variable Star Index (VSX, [Watson et al. 2006](#), compiled by the American Association of Variable Star Observers, AAVSO) to collect all information regarding newly identified CVs and CV candidates. Combining these findings with the Ritter & Kolb catalogue ([Ritter & Kolb 2003](#)), a collection of the observational properties of

all CVs with an orbital period determination (1429 systems), we built a list of $\simeq 8000$ systems.

We cross-matched this list with the *Gaia* DR2 catalogue. In order to take into account proper motions and the low precision ($\simeq 10''$) of the coordinates reported in some of the catalogues, we first performed a cross-match with a $30''$ search radius, resulting in $\simeq 364\,000$ objects. Using the *Gaia* proper motions and radial velocities (whenever available), we calculated the corresponding coordinates at epoch 2000. We then performed a second cross-match between our list of CVs and this catalogue, selecting the closest source within $10''$ radius, which resulted in $\simeq 6400$ objects. Since our focus is on the properties of the 150 pc CV sample, we applied a cut in parallax:

$$\varpi + 3\sigma_{\varpi} \geq 6.66 \text{ mas} \quad (1)$$

This selection returned 166 systems which, within their parallax uncertainties, are located within 150 pc (Figure 1).

More than half of these 166 systems have been identified as CV candidates in transient surveys because they have shown one or more brightenings that resembled a dwarf nova outburst, but have no photometric or spectroscopic follow-up confirming their CV nature. Therefore, for each system, we inspected the literature, the CRTS, AAVSO and ASAS-SN light curve archives for their outburst history, and SDSS




Figure 3. PanSTARRS *g*-band (left) and *r*-band (right) images of OGLE-BLG-DN-0040. The position of OGLE-BLG-DN-0040 is highlighted with a blue circle, while the crosses show the *Gaia* detections, one of which (cyan cross) is located only $\approx 1.4''$ away from OGLE-BLG-DN-0040. However, OGLE-BLG-DN-0040 is fainter than the *Gaia* detection limit in quiescence so does not have an entry in the *Gaia* DR2 archive. This is actually a spurious association, as can be clearly seen in the *r*-band image.

for serendipitous spectroscopy. In this way, we identified 28 objects that are mistakenly classified as CVs or CV candidates in the literature (Table D1), many of them single low-mass stars which show flaring phenomena that can be mistaken for dwarf nova outbursts. More details are provided in the Appendices A, B & C.

Finally, we searched for *Gaia* sources in the white dwarf locus of the HR diagram (defined by the colour cuts from Hollands et al. 2018, see their equations 2, 3 and 4) that showed large anomalous uncertainties in their *G* magnitudes, weighted over the number of observations in the given band. Such large values can be interpreted as intrinsic stellar variability arising e.g. from pulsations, debris transits, magnetism or on-going accretion (Hermes et al., in preparation). Among the different candidates, two systems stood out because of their large intrinsic variability and for being overluminous to the canonical white dwarf cooling tracks. Spectroscopic follow-up later confirmed these as new CVs located within 150 pc: *Gaia* J051903.99+630340.4 and *Gaia* J154008.28–392917.6 (Figure 2 and Appendix D, which we named with the identifier *Gaia* JHHMSS.SS±DDMMSS.S, defined with respect to their coordinates at equinox 2000 and epoch 2000, as computed using the *Gaia* ICRS coordinates and proper motions).

2.1 Astrometric accuracy

Along with positions, magnitudes and proper motions, i.e. the kinematic parameters used to obtain the astrometric solution for each source, *Gaia* DR2 provides a series of additional parameters that allow to evaluate the accuracy of this solution. In particular, the `astrometric_excess_noise`

(AEN) represents the errors introduced in the astrometric modelling (see Lindegren et al. 2012) and, ideally, should be zero. A possible selection to remove sources with poor astrometric solution is to impose `astrometric_excess_noise` < 1 (Lindegren et al. 2018). However, we noticed that Z Cha, for which the *Gaia* parallax ($\varpi = 8.7 \pm 0.1$ mas) implies a distance of 116 ± 2 pc, in good agreement with the distance estimated by Beuermann (2006), 112 ± 8 pc, has `astrometric_excess_noise` = 1.08. Therefore, in order to build a sample that is the most complete as possible we applied a more generous cut, considering the parallaxes of those sources for which `astrometric_excess_noise` > 2 as unreliable.

2.2 Spurious *Gaia* detections

Ten CV candidates are located in crowded regions and the cross-match between our sample and the *Gaia* catalogue returned a spurious detection (Figure 3). These sources are actually fainter than the *Gaia* detection limit and do not have an entry in the *Gaia* DR2 archive. Our list also contained EU Cnc for which *Gaia* DR2 provides a parallax of $\varpi = 1.8 \pm 2.5$ mas and therefore satisfies the condition in Equation 1. However, this CV is located in an open cluster for which the distance has been determined as $d = 785 \pm 50$ pc (Belloni et al. 1993) and hence we discarded this system.

3 THE 150 PC CV SAMPLE

Gaia DR2 provided parallaxes for about 1.3 billion sources. Converting these measurements into distances is not always trivial as the mere inversion of the parallax can introduce

Table 1. 150 pc CV sample from *Gaia* DR2, sorted by increasing distance. The CV types are as follows: UG, dwarf nova of U Gem sub-type; UGSU, dwarf nova of SU UMa sub-type; UGWZ, dwarf nova of WZ Sge sub-type; IP, intermediate-polar; AM, polar; NL, novalike; AM CVn, AM CVn star. SDSS *ugriz* photometry of the systems highlighted with a star is used to assess the completeness of the *Gaia* 150 pc sample (see Section 4.2).

System	<i>Gaia</i> DR2 ID	ϖ (mas)	σ_ϖ (mas)	P_{orb} (min)	Type	G (mag)	G_{BP} (mag)	G_{RP} (mag)	Distance (pc)
WZ Sge	1809844934461976832	22.16	0.04	81.63	UGWZ	15.21	15.21	15.06	45.13 ± 0.08
VW Hyi	4653893040002306432	18.53	0.02	106.95	UGSU	13.84	13.94	13.45	53.96 ± 0.06
EX Hya	6185040879503491584	17.56	0.04	98.26	IP	13.21	13.23	12.88	56.9 ± 0.1
GP Com	3938156295111047680	13.73	0.06	46.57	AM CVn	15.95	15.89	15.91	72.8 ± 0.3
V455 And	1920126431748251776	13.24	0.06	81.08	IP	16.06	16.13	15.71	75.5 ± 0.3
GD 552	2208124536065383424	12.35	0.05	102.73	UGWZ	16.46	16.46	16.18	81.0 ± 0.3
ASASSN-14dx*	2488974302977323008	12.34	0.04	82.81	UGWZ	14.96	14.92	14.69	81.0 ± 0.3
AM Her	2123837555230207744	11.40	0.02	185.65	AM	13.58	13.86	12.85	87.8 ± 0.1
IX Vel	5515820034889610112	11.04	0.03	279.25	NL	9.32	9.34	9.27	90.6 ± 0.2
OY Car	5242787486412627072	11.01	0.03	90.89	UGSU	15.62	15.64	15.21	90.8 ± 0.2
AE Aqr	4226332451596335616	10.97	0.06	592.78	IP	10.95	11.47	10.26	91.2 ± 0.5
U Gem	674214551557961984	10.71	0.03	254.74	UG	13.91	14.38	13.11	93.4 ± 0.3
V396 Hya	3503987633230546688	10.69	0.15	65.10	AM CVn	17.66	17.70	17.45	94 ± 1
BW Scl	2307289214897332480	10.60	0.10	78.23	UGWZ	16.26	16.26	16.10	94.4 ± 0.9
V627 Peg	1800384942558699008	10.03	0.07	78.51	UGWZ	15.67	15.65	15.27	99.7 ± 0.7
AR UMa	783921244796958208	9.87	0.12	115.92	AM	16.26	16.35	15.78	101 ± 1
1RXS J105010.3–140431	3750072904055666176	9.14	0.11	88.56	UGWZ	17.17	17.21	17.08	109 ± 1
TCP J21040470+4631129	2163612727665972096	9.13	0.12	77.07 ^(a)	UGWZ	17.77	17.87	17.29	109 ± 2
V2051 Oph	4111991385628196224	8.90	0.07	89.90	UGSU	15.37	15.46	14.87	112.4 ± 0.9
V834 Cen	6096905573613586944	8.90	0.21	101.52	AM	16.66	16.82	16.07	113 ± 3
GW Lib	6226943645600487552	8.87	0.08	76.78	UGWZ	16.49	16.49	16.32	113 ± 1
ST LMi	3996419759863758592	8.83	0.08	113.89	AM	16.13	–	–	113 ± 1
SS Cyg	1972957892448494592	8.72	0.05	396.19	UG	11.69	12.11	10.95	114.6 ± 0.6
V884 Her	4503256687122329088	8.69	0.02	113.01	AM	13.49	13.57	13.18	115.1 ± 0.3
Z Cha	5210507882302442368	8.66	0.12	107.28	UGSU	15.85	15.94	15.19	116 ± 2
<i>Gaia</i> J051903.99+630340.4	285957277597658240	8.59	0.04	126.	UGSU	15.17	15.30	14.77	116.4 ± 0.5
V2301 Oph	4476137370261520000	8.24	0.08	112.97	AM	16.75	16.94	15.86	121 ± 1
V893 Sco	6039131391540808832	8.06	0.05	109.38	UGSU	14.65	14.76	14.25	124.1 ± 0.8
QZ Vir*	3800596876396315648	7.81	0.07	84.70	UGSU	16.06	16.12	15.76	128 ± 1
V1040 Cen	5343601913741261312	7.80	0.03	87.11	UGSU	14.04	14.11	13.69	128.1 ± 0.4
SDSS J125044.42+154957.3	3934459045528378368	7.79	0.18	86.3	AM	18.22	18.30	17.95	129 ± 3
V379 Tel	6658737220627065984	7.65	0.07	101.03	AM	16.19	16.69	15.45	131 ± 1
BL Hyi	4697621824327141248	7.65	0.07	113.64	AM	17.25	17.45	16.70	131 ± 1
MR Ser*	1203639265875666304	7.59	0.05	113.47	AM	16.23	16.47	15.64	131.8 ± 0.8
V3885 Sgr	6688624794231054976	7.54	0.08	298.31	NL	10.25	10.28	10.16	133 ± 1
<i>Gaia</i> J154008.28–392917.6	6008982469163902464	7.49	0.11	–	UGWZ	17.36	17.39	17.23	134 ± 2
VV Pup	5719598950133755392	7.30	0.05	100.44	AM	15.93	16.04	15.48	137.0 ± 0.9
VY Aqr	6896767366186700416	7.24	0.14	90.85	UGSU	16.86	16.96	16.44	138 ± 3
IP Peg	2824150286583562496	7.08	0.05	227.82	UG	14.71	15.27	13.88	141.2 ± 1.0
HT Cas	426306363477869696	7.07	0.06	106.05	UGSU	16.35	16.48	15.81	141 ± 1
SDSS J102905.21+485515.2	834947865750806272	6.96	0.24	91.33	UGWZ	18.16	17.94	17.84	144 ± 5
EZ Lyn*	935056333580267392	6.87	0.15	84.97	UGWZ	17.81	17.84	17.71	146 ± 3
V379 Vir*	3699606286708406912	6.7	0.2	88.4	AM	18.01	18.02	17.93	150 ± 4
V355 UMa	1558322303741820928	6.66	0.09	82.52	UGWZ	17.38	17.35	17.27	150 ± 2

Notes. (a): spectroscopic period (ATel #13009).

some biases in the distance estimate, especially when the fractional error on the parallax is larger than 20 per cent (see e.g. Bailer-Jones 2015; Luri et al. 2018). This is the case for many systems in our sample, in particular for those that are further away and have poor parallax measurements. Therefore, to estimate the distances, we used a statistical approach in which, following the prescription by Bailer-Jones (2015), we defined a likelihood and an exponentially decreasing volume density prior. The latter contains the Galactic CV scale height, h , which is a function of the system age: older CVs (i.e. period bouncers and short period systems that have not evolved through the period minimum yet, $P_{\text{orb}} \lesssim 2$ h) have

larger scale heights, $h \simeq 260$ – 450 pc, while younger CVs (i.e. long period systems, $P_{\text{orb}} \gtrsim 2$ h) have smaller scale heights, $h \simeq 120$ pc (Pretorius et al. 2007b). For the systems with a measured orbital period we defined the relative scale heights following Pretorius et al. (2007b):

$$h = \begin{cases} 120 \text{ pc} & \text{for long period systems } (P_{\text{orb}} \gtrsim 2 \text{ h}) \\ 260 \text{ pc} & \text{for short period systems } (P_{\text{orb}} \lesssim 2 \text{ h}) \\ 450 \text{ pc} & \text{for period bouncer CVs} \end{cases} \quad (2)$$

Our sample contains three period bouncer candidates, GD 552 (Unda-Sanzana et al. 2008), SDSS J102905.21+485515.2 (Thorstensen et al. 2016)




Figure 4. The spatial distribution of CVs in the 150 pc sample in equatorial (top) and Galactic (bottom) coordinates shows no evident correlation with the different sub-types, suggesting that the 150 pc CV sample is not affected by obvious selection biases.

and 1RXS J105010.3–140431 (Patterson 2011; Pala et al. 2017), for which we assumed a scale height of $h = 450$ pc. An orbital period determination is not available for *Gaia* J154008.28–392917.6. Nonetheless, its spectral appearances suggest that this is a low accretion rate systems of the WZ Sge sub-type. These CVs are found below the period gap ($P_{\text{orb}} \lesssim 2$ h) and thus, for *Gaia* J154008.28–392917.6, we assumed a scale height of $h = 260$ pc.

We determined the distance as the median and the corresponding uncertainties as the 16th and the 84th percentiles of the posterior distribution of each system. From the posterior distribution, we also determined the probability for each object to be located within 150 pc. According to these distances, we discarded 55 systems (Tables D1) for which $d > 150$ pc.

The final 150 pc sample (Table 1) consists of 42 CVs and two AM CVn systems² (GP Com and V396 Hya). Although there are many similarities between AM CVn stars and CVs, the formation channels and evolutionary histories of these two classes of systems are different and we do not include the two AM CVn in the following discussion (see Ramsay et al. 2018 for a detailed study of these stars using the *Gaia* DR2 data).

Figure 4 show the spatial distribution of this sample and it is colour-coded according to the CV type. The systems appear to be uniformly distributed on the sky, with no evident correlation between their location and their sub-types. In order to evaluate whether the position of the 150 pc CVs on the sky is consistent with a uniform distribution

from a statistical point of view, we generated 1000 samples, each consisting of 10^5 objects uniformly distributed in a volume enclosed within 150 pc, assuming an exponentially decreasing space density (Equation 5) with $h = 280$ pc, i.e. the average value of the scale heights of different CV sub-types (Equation 2). For each sample, we randomly selected 42 objects and applied a two sample Kolmogorov-Smirnov (KS) test between the pairwise cumulative distributions of these objects and that of the CVs in the 150 pc sample. We found that the average KS statistics is $D \approx 0.03$, lower than the 95 per cent critical level, $D_{\text{critic}} \approx 0.29$, with a p -value ≈ 0.7 . This implies that the null hypothesis, i.e. uniform distribution on the sky, cannot be rejected thus suggesting that 150 pc sample is not affected by obvious selection biases.

4 COMPLETENESS

A key property of any sample used to determine a space density is its *completeness*, in combination with a good understanding of potential observational selection effects. We use two independent tests to assess the completeness of the 150 pc CV sample.

4.1 Previously known CVs with distances

A simple assessment of the completeness of the 150 pc CV sample is to establish the fraction of known CVs with *reliable* pre-*Gaia* distance measurements of $d \leq 150$ pc recovered by *Gaia*. Except parallaxes, all distance measurements are indirect, and we briefly review the main methods used for CVs to qualify what we consider as reliable distances³.

- Accurate trigonometric parallaxes from both space- (Duerbeck 1999; Harrison et al. 2004) and ground-based observations (Thorstensen 2003; Thorstensen et al. 2008).
- Modelling the ultraviolet flux of the white dwarf. This method requires a clean detection of the white dwarf which, owing to the strong contamination from the accretion disc in the optical, is best detected in the ultraviolet. By performing a fit to the spectroscopic data, the distance to the system can be measured by assuming a mass-radius relationship, since the scaling factor between the best-fit model is related to the ratio between the distance and the stellar radius (e.g. Gänsicke et al. 1999).
- Modelling the red/infrared photometric contribution of the donor. The emitting area of the secondary is constrained by the Roche geometry, i.e. by the orbital period and the mass ratio. When the secondary is detected in the near-infrared, the distance to the system can be estimated using the “Bailey method” (Bailey 1981) from its K magnitude in combination with the mass-radius relationship for CV secondary stars (e.g. Knigge et al. 2011).

³ Ak et al. (2007) proposed a period-luminosity-colour relation, i.e. an alternative empirical calibrations in which the absolute magnitude of the system is determined from its period and its 2MASS (Two Micron All Sky Survey, Skrutskie et al. 2006) colours. However, its application to short-period dwarf novae appeared to be problematic (Patterson 2011) and we excluded the distances derived with this method from the following analysis.

² An AM CVn system consists of a white dwarf accreting from another white dwarf or a partially degenerate helium star, often via an accretion disc. Their orbital periods are shorter than that of CVs and their optical spectra do not contain any hydrogen.

- d) Analysis of the eclipse light curves. Similarly to the modelling the SED, this method provides both the effective temperature and the radius of the white dwarf, allowing to estimate the distance to the system (Littlefair et al. 2008; McAllister et al. 2019).
- e) $M_V - P_{\text{orb}}$ relationship. The radius of the accretion disc is a relatively fixed fraction of that of the white dwarf Roche lobe ($R_{\text{disc}} \simeq 0.7R_{\text{Roche-lobe}}$, Sulkanen et al. 1981), and scales with the mass ratio and the orbital separation, i.e. with P_{orb} . This, combined with the fact that disc outbursts occur when the accretion disc effective temperature rises to $\simeq 8000$ K allows the use of dwarf nova outbursts as standard candles. This method was first introduced by Warner (1987) and later refined by Patterson (2011) using a sample of 46 CVs with good distance estimates. One of the main sources of uncertainty is the often unknown inclination of the system.

We identified 49 CVs with published pre-*Gaia* distance measurements of $d \leq 150$ pc (Table D3). For systems with multiple distance estimates, we considered the most reliable measurement following the order of the list above. *Gaia* re-identified all these 49 CVs, confirming that 28 of them are located within 150 pc. The remaining 21 CVs are found to be located further than their pre-*Gaia* distances, the majority of which were estimated by modelling the red/infrared photometric contribution of the donor. This method systematically underestimates the distance to the systems (Figure 5) owing to an unaccounted contamination by light from the accretion disc which results in an overestimate of the brightness of the donor. In contrast, the distances estimated from the modelling of the SED in the ultraviolet are more accurate since the white dwarf is the dominant source of emission in this wavelength range.

Since *Gaia* re-identified all the previously known CVs located within 150 pc, we conclude that the completeness of our sample is not limited by the ability of *Gaia* in obtaining astrometry of CVs but rather by the efficiency of the methods available to identify CVs in the first place. As discussed in Section 2, the discovery of CVs has been strongly biased towards highly variable systems, such as dwarf novae with a high duty-cycle, and disfavoured the discovery of systems with both low mass transfer rates (WZ Sge dwarf novae, which have outburst recurrence times of the order of decades), and high mass transfer rates (nova-like CVs in which the disc remains in a hot state).

Discovering > 300 new CVs, SDSS has demonstrated that the spectroscopic identification of CVs is largely independent of CV sub-type. In fact, follow-up of the SDSS CVs led to the unambiguous confirmation of a pile-up of intrinsically faint CVs near the period minimum (Gänsicke et al. 2009). The SDSS CVs represent the currently least biased sample of CVs, and we make use of the *ugriz* photometry to (i) estimate the completeness of our *Gaia* CV sample and (ii) search for CVs within 150 pc that have so far been missed.

4.2 Completeness of the 150 pc CV sample

Because the majority of known CVs fall within the colour space of quasars, we can make use of the highly efficient spectroscopic follow-up of SDSS quasar candidates to assess the completeness of the 150 pc CV sample. The uniform spa-




Figure 5. Comparison between pre-*Gaia* distances estimated using different methods and distances based on the *Gaia* parallaxes. The most accurate pre-*Gaia* distances have been determined from ground and space based parallax measurements (pink). Whereas modelling ultraviolet spectra of white dwarf dominated CVs (blue) also provides reliable distance measurements, estimates based on modelling the red/infrared photometric contribution of the donor stars (orange) systematically overestimate the distance. Although subject to large uncertainties, the distance estimates from the $M_V - P_{\text{orb}}$ relationship are consistent with the *Gaia* determinations for most of the systems in the sample.

tial distribution of the 150 pc CVs (Section 3) suggests that the properties of the local CV population are not evidently correlated with their location on the sky and that CVs are identified with equal success at low and high Galactic latitudes. Moreover, the typical CV scale height ($h = 280$ pc) is much larger than the radius ($h = 150$ pc) of the volume we are considering and consequently the correlation of the distribution of the 150 pc CVs with the Galactic latitude is negligible. For this reason, although the SDSS footprint covers only one third of the sky and avoids the Galactic plane, the properties of the SDSS CVs can be safely extended to the whole sky and hence, to the 150 pc *Gaia* CV sample.

4.2.1 Definition of the spatial and colour footprint followed-up by the SDSS quasar search

With the aim to study extragalactic objects, SDSS acquired photometric observations for 7 500 square degree of sky and subsequently performed spectroscopic follow-up for a subset of the photometric objects. SDSS selected the spectroscopic quasar targets according to their colours (Richards et al. 2002). Within the vicinity of a quasar in the four dimensional ($u-g; g-r; r-i; i-z$) colour space, all photometric objects have the same probability to be spectroscopically observed by the Legacy (York et al. 2000) and BOSS (Baryon Oscillation Spectroscopic Survey (Dawson et al. 2013)).

Consequently, the probability for an object to be observed spectroscopically depends on its location on the sky (i.e. whether it is located within the Legacy/BOSS footprint) and on its colour similarity to quasars. To de-

Table 2. Summary of the cross-match between *Gaia* and SDSS and the relative sub-samples used to estimate the completeness in Section 4.

Description	Number of objects
Total number of source in <i>Gaia</i> DR2 with $\varpi + 3\sigma_{\varpi} \geq 6.66$ mas	2 100 094
Number of source with reliable astrometry (Equation 4)	910 187
of which in SDSS	303 723
of which in SDSS DR14	299 501
of which in SDSS DR7	4222
of which within the Legacy and BOSS footprint	155 099
of which in the quasar colour space	42 823
of which with similar colour to CVs	5300
of which with a spectrum	2811

fine this parameter space we used the CasJobs⁴ service to query the PlateX table imposing `programName == "boss" || programName == "legacy"`. This returned the coordinates of the centres of the 4235 spectroscopic plates observed during the Legacy and BOSS surveys. Each spectroscopic plate has a field of view of 1.49° in radius, therefore objects located within $\leq 1.49^\circ$ from the plate centre that have (i) fiber magnitudes, i.e the flux contained within the aperture of a spectroscopic fiber, fainter than 15 mag in *g* or *r*, and 14.5 mag in *i* and (ii) $i < 19.1$ mag, are all potential spectroscopic targets. These magnitude limits are imposed by the fixed 15 min exposure time of the SDSS observing strategy: objects brighter than the first set of limits saturate the SDSS detectors and objects fainter than $i = 19.1$ are so numerous that they were not systematically followed-up spectroscopically. We used the quasar catalogue from the SDSS DR7 (Schneider et al. 2010) to calculate the colour distance, i.e. the nearest neighbour distance, between each pair of quasar. We found that ≈ 99 per cent of them have another quasar within ≈ 0.29 mag. We can hence assume that all objects found within 0.29 mag from a quasar are located within the colour space in which SDSS selected its spectroscopic targets and had therefore a chance to be observed spectroscopically.

CVs occupy a sub-region of the colour space in which quasars are found. This sub-region can be defined in an analogous fashion as done above for the case of quasars. We calculated the colours ($u - g; g - r; r - i; i - z$) of all known CVs with SDSS photometry. To avoid contamination from objects with poor photometry or objects mistakenly classified as CVs, we only considered SDSS sources that could be matched to CVs that were confirmed either by spectroscopy, or by a published orbital period measurement, and that have accurate (photometric errors < 5 per cent) and clean SDSS photometry, i.e.:

$$\text{clean} = 1 \ \& \ \text{mode} = 1 \ \& \ \text{type} = 6. \quad (3)$$

These flags ensure selection of stars (`type=6`) rather than galaxies and avoids blended photometry or multiple detections of the same source. We also required that the selected objects have reliable *Gaia* parallax and colours (Equation 4). The final sample contains high-quality photometry of 418 CVs, the majority of which are located further than 150 pc. In order to build a reference CV sample that is as representative as possible of the overall SDSS CV population, we computed the apparent magnitudes that SDSS CVs with




Figure 6. Frequency distribution (top) and cumulative distribution (bottom) of the colour distance in the four dimensional space ($u - g; g - r; r - i; i - z$) for the 130 SDSS CVs, showing that 95 per cent of them have another CV within ≈ 0.27 mag (vertical blue line).

$d > 150$ pc would have if they were located at $d = 150$ pc. This allows us to remove those systems that, if they were closer, would have not been observed by SDSS owing to the bright limit in the target selection. Moreover, we only considered those CVs that fulfil the conditions listed above (location on the sky and colour similarity with quasars), which reduced the reference sample to 130 CVs (black crosses in Figure 1).

We then defined a four-dimensional “sphere” in colour space, centred on each of these 130 CVs, and calculated the colour distance between each pair. Figure 6 shows that the distribution peaks at a colour distance of ≈ 0.06 mag, however, to enclose 95 per cent of the sample, we defined a colour radius of ≈ 0.27 mag. All objects located within ≤ 0.27 mag from any of this 130 reference CVs are hence potential CV candidates. From their spectral completeness (i.e. the ratio between the objects with a spectrum and the total number of objects observed photometrically) combined with the fraction of spectroscopically observed objects drawn from that

⁴ <https://skyserver.sdss.org/casjobs/>




Figure 7. Two-dimensional projection of the four-dimensional ($u-g;g-r;r-i;i-z$) colour space onto ($u-g;g-r$) of the sources within 150 pc (black) that are found within the quasar colour space (grey). The four-dimensional colour space spanned by CVs has been defined by 130 reference CVs (green dots, Section 4.2.1), and the projection of that colour space onto ($u-g;g-r$) is shown by the light orange cloud. CV candidates contained within this four-dimensional colour space are shown in pink. Among them, four confirmed CVs (dark blue) are known. Given the spectral completeness of ≈ 53 per cent, \approx two more CVs are expected to be identified, implying an overall completeness of ≈ 71 per cent for the 150 pc CV sample.

colour space that actually are CVs, one can then estimate how many CVs are expected to be found among the candidates. Comparing the expected number of CVs with that of known CVs within the reference spatial and colour footprint then provides the completeness of the sample. In the following Section, we apply this method to the SDSS CVs in order to derive the completeness of the 150 pc *Gaia* CV sample.

4.2.2 Application to the 150 pc sample

We performed a cross-match between *Gaia* and SDSS⁵, querying the *Gaia* archive for all objects that, including their parallax uncertainties, are located within 150 pc (Equation 1), which returned $\approx 2\,100\,000$ objects. Among them, many systems have inaccurate astrometry and $G_{BP} - G_{RP}$ colours due to faintness, blended double stars or other astrometric effects. Following the prescription from Lindgren et al. (2018, their appendix C), we selected only those sources with the most reliable astrometry by applying the following cuts,

$$\begin{aligned} \text{astrometric_excess_noise} &< 1 \\ \text{phot_bp_rp_excess_factor} &> 1 + 0.015 \times \text{bp_rp}^2 \\ \text{phot_bp_rp_excess_factor} &< 1.3 + 0.06 \times \text{bp_rp}^2 \end{aligned} \quad (4)$$

⁵ While the *Gaia* consortium already provides the cross-match with SDSS (`gaiadr2.sdssdr9.best_neighbour`), we found that $\approx 82\,000$ objects common to SDSS and *Gaia* are missing from that table, as well as ≈ 1000 associations of spurious *Gaia* data with SDSS sources.

which leaves $\approx 910\,000$ objects with reliable *Gaia* colours and astrometry (grey dots in Figure 1).

We then queried the SDSS DR14 archive using the CasJobs website and retrieved the SDSS coordinates, photometry, and Modified Julian Date of the observations for the closest objects within $30''$ to each *Gaia* entry. Using the *Gaia* proper motions, we calculated the coordinates of the *Gaia* sources at the epoch of the SDSS observations, and then performed a two arc-second radius cross-match with the SDSS objects, thus obtaining the best epoch-matched association between each *Gaia* entry and the SDSS DR14 sample for a total of $\approx 300\,000$ objects.

Data releases later than DR7 provide more accurate photometry based on an improved background subtraction. As a side effect of this re-reduction of the photometry, some sources nearby bright stars that were present in DR7 are no longer included in the later data releases. To recover these lost sources, we applied the procedure outlined above also to the SDSS DR7 catalogue, retrieving photometry for an additional ≈ 4000 sources that are no longer included in the subsequent data releases.

The final cross-match between the *Gaia* source within 150 pc and SDSS contains 303 723 objects (Table 2).

Among these 303 723 objects, 5300 (2811 of which have a spectrum) fall within the spatial and colour footprint of our SDSS CV reference sample defined above and are hence CV candidates (Figure 7). Four of these are known CVs (V379 Vir, EZ Lyn, MR Ser, ASASSN-14dx), three of which (V379 Vir, EZ Lyn and MR Ser) have a spectrum (Figure 8).

The spectra of these 2811 objects were then visually inspected, but no additional CV was identified. This is not surprising given the extensive search for CVs in the SDSS data that have been carried out in the past by P. Szkody and collaborators (Szkody et al. 2002, 2003, 2004, 2005, 2006, 2007, 2009, 2011). The spectral completeness of the CV candidate sample results $2811/5300 \approx 53$ per cent. Considering that there are three CVs with a spectrum, we can hence estimate that a total of \approx six CVs were expected to be identified. Since only four CVs have been detected, we can conclude that the sample is ≈ 71 per cent complete (Figure 9). This estimate of the completeness is subject to small number statistics, and we derive a more robust value making use of the much larger sample of CVs with CRTS light curves.

4.2.3 Additional constraints from CRTS light curves

Additional constraints on the completeness can be derived from the long-term light curves of the 2489 CV candidates identified in Section 4.2.1 for which no SDSS spectroscopy is available, which provide the possibility to discover new CVs via the detection of their intrinsic variability. CRTS light curves are the most suitable for this task since, spanning more than 10 years, they are sensitive to the low mass transfer rate systems with disc outburst recurrence times of years (e.g. Breed et al. 2014). Moreover, covering $\approx 30\,000 \text{ deg}^2$ of the sky between $-75^\circ < \delta < 65^\circ$ and $|b| \gtrsim 15^\circ$ (Drake et al. 2009), the CRTS footprint largely overlaps with that of SDSS, allowing us to combine the spectral completeness constraints derived in the previous section with the presence of possible variability.

Whereas the CRTS observing strategy aimed at obtain-




Figure 8. SDSS spectra of the CVs in the 150 pc SDSS sample. The blue and the orange dashed lines highlight the position of the Balmer series and of some of the most common He I lines, respectively. EZ Lyn shows the typical spectral appearance of the low-accreting WZ Sge type CVs, with the white dwarf signature clearly recognisable in the broad Balmer lines absorption features. Magnetic CVs undergo phases of high mass transfer (high states) alternated with phases in which accretion almost ceases (low states). Among the magnetic systems in the 150 pc sample observed by SDSS, V379 Vir have been caught in a low state, while MR Ser has been observed in a high state.

ing data with ≈ 2 week cadence, seasonal visibility and adverse weather condition can result in quite a different number of observations per field and therefore the CRTS data for a given object span a wide range in quality. Therefore it is necessary to define a criterion that identifies intrinsic variability opposed to possible scatter due to poor weather conditions, cosmic rays contamination and possible calibration issues. To this purpose, we used a sample of 1807 hydrogen-atmosphere white dwarfs from [Gentile Fusillo et al. \(2019\)](#), flagged as “DA” in their table 4) which are located within 150 pc, have been observed by CRTS, but are not variable (a small number of white dwarfs exhibit variability due to pulsations or rotations, though with amplitudes that are much lower than the variability seen among CVs). Since these systems are not intrinsic variable, in an ideal data set, all data points of a light curve should be consistent with the median value within their uncertainties. Consequently the related reduced chi-squared χ^2_ν should converge to ≈ 1 as the number of degrees of freedom, $\nu = N - 1$ (where N is the number of observations), increases, whereas the standard deviation of the normalised residuals should be $|\sigma_{\text{RES}}| < 2$. We queried the CRTS database⁶ for the light curves of the white dwarf test sample. We considered only observations with $V \geq 13$ mag to avoid saturated exposures and computed the χ^2_ν associ-

ated with each light curve. We found that χ^2_ν shows a much larger scatter than expected (Figure 10), reflecting the presence of low-quality photometric observations which mimic the presence of intrinsic variability in the systems. Using those white dwarfs that appear intrinsically variable owing to the low quality of their CRTS observations (Figure 10, orange and pink points), we defined the following two cuts that remove systems with poor CRTS data:

$$\chi^2_\nu > \frac{300}{\nu} + 3.5 \quad \text{or} \quad |\sigma_{\text{RES}}| > 2 \quad (5)$$

Three per cent of the objects satisfies the condition above and this fraction represents the intrinsic systematic uncertainty associate with this selection method.

To verify the quality of our criteria, we applied the same cuts to a control sample of 456 CVs from [Drake et al. \(2014\)](#) for which CRTS light curves having at least three data points⁷ with $V \geq 13$ mag are available. Our method outlined above is able to identify as variable 93 per cent of these CVs. The 7 per cent of systems that we are unable

⁷ Ideally, three data points represent a lower limit for the number of observations required to identify a CV candidate since two observations during quiescence allow to define a quiescent reference level with respect of which any significantly bright (even single) outlier can be indicative of the occurrence of a disc outburst.

⁶ <http://nesssi.cacr.caltech.edu/DataRelease/>




Figure 9. Flowchart of the process used to estimate the completeness of the 150 pc sample. The starting point is the cross-match between *Gaia* and SDSS (a), which returns 303 723 sources. A subsample of these, 155 099 (b), falls within the footprint that the SDSS Legacy and BOSS surveys observed spectroscopically. Among them, 42 823 have colours similar to quasars (c) and hence have all the same probability to have a spectrum in SDSS. Thanks to their colour similarity with the 130 SDSS CVs (d) that occupy a sub-region of the quasar colour space (see Section 4.2.1), 5300 CV candidates are identified, 2811 of which have a spectrum (e). This implies a spectral completeness of 53 per cent (f). This sample contains four known CVs, three of which have a spectrum (g). We can hence estimate that ≈ 6 CVs should be identified in total (h). Four CVs have already been found and we can conclude that the CV sample is ≈ 71 per cent complete (i).

to recover are those showing low variability or having a low number of observations ($N \lesssim 10$).

We then repeated the same exercise for the 2489 CV candidates without a spectrum (Table 3 and Figure 11, left branch), since these are the objects for which the lack of additional information prevents us from establishing their true nature. As before, we queried the CRTS database for unsaturated exposures and we considered only light curves with three or more data points, which were available for 1983 objects.

Applying Equation 5, we find that 3.5 per cent of them are variable. This fraction needs to be corrected for the fact that our method is unable to recover seven per cent of the 456 CVs from the control sample from Drake et al. (2014), thus resulting in a total fraction of 10.5 per cent (i.e. 208) variable objects, which are possible CV candidates. These need to be complemented with the remaining 2489-1983=506 sources for which no information (neither a SDSS spectrum nor a CRTS light curve) is available and that could




Figure 10. Sample of 1807 hydrogen-atmosphere white dwarfs used to define the criteria for identifying variable systems. Systems above (orange points) or below the threshold (light blue) but with $|\sigma_{\text{RES}}| > 2$ (pink dots) are identified as variable.

also hide a not yet identified CV. The total 714 systems, representing 13 per cent of the 5300 candidates identified in the previous section, are the sources for which we are unable to unambiguously establish whether they are CVs and set an upper limit for the completeness of our sample to 87 per cent.

A lower limit on the completeness can be derived imposing more stringent constraints on the CRTS light curve. Ideally, 200 points are in principle sufficient to identify short term variability (not necessarily related to disc instabilities, such as eclipses) in low mass accreting short period systems (Parsons et al. 2013). However, in order to obtain a conservative lower limit on the completeness, we decided to impose a more stringent constraint, assuming that a minimum of 300 data points are necessary to identify variable systems. In this case, our control sample is reduced to 71 CVs from Drake et al. (2014) for which CRTS light curves with at least 300 unsaturated exposures are available and, with our selection criteria defined in Equation 5, we are able to recover as variable 97 per cent of them.

Among the 2489 CV candidates without a spectrum, 839 have a CRTS light curves matching the new requirement of 300 data points (Table 3 and Figure 11, right branch) and one per cent of them results variable according to Equation 5. Accounting for the fact that, in this case, our method is unable to recover three per cent of CVs, results in a total fraction of four per cent variable objects, i.e. 34. Adding these to the 2489-839=1650 objects for which we have no information about (no SDSS spectra nor CRTS light curves) results in a total of 1684 CV candidates. These correspond to 32 per cent of the initial 5300 candidates, implying a lower limit of 68 per cent in the completeness of our sample.

Considering the upper and lower limits derived above, we conclude that our sample is (77 ± 10) per cent complete, where the error is estimated as the sum of the maximum uncertainties in our method (seven per cent and three per cent respectively), representing the efficiency in recovering CVs as a function of the number of observations available.




Figure 11. Flowchart of the process used to estimate the completeness of the 150 pc sample using CRTS. The starting point is the 5300 CV candidates identified in Section 4.2.2. 2489 of them do not have a spectrum and thus lack of additional information required to establishing their true nature. Using their CRTS light curves we establish the number of objects for which we are unable to unambiguously establish whether they are CVs, thus obtaining an upper and lower limit for the completeness of the 150 pc CV sample.

Table 3. Summary of the different sub-samples used to estimate the completeness from CRTS light curves (Section 4.2.3).

Description	Number of objects
Source with colour similarity to CVs	5300
of which without a spectrum	2489
of which with a CRTS light curve with more than three unsaturated data points	1983
of which variable	208
of which without a CRTS light curve with more than three unsaturated data points	506
of which with a CRTS light curve with more than 300 unsaturated data points	839
of which variable	34
of which without a CRTS light curve with more than 300 unsaturated data points	1650

4.3 Summary on the completeness

Using two independent methods, i.e. the count of objects with a spectrum (Section 4.2.2) and the count of objects found to be variable (Section 4.2.3), we obtained two values for the completeness, ≈ 71 per cent and 77 ± 10 per cent respectively, that are in good agreement among each other, thus supporting the validity of our result.

Because of the larger sample size of CVs used for the compute of the completeness carried out in Section 4.2.3, we adopt (77 ± 10) per cent as the final value. As stated above, the CVs and CV candidates used in this analysis have only been selected according to their position on the sky and colours and hence the completeness estimate can be safely extended to the whole *Gaia* 150 pc CV sample. Given the (77 ± 10) per cent completeness, we can estimate that ≈ 12 CVs are still to be discovered within 150 pc.

The methods we employed to derive the completeness are based on the SDSS quasar target selection, and are biased against red, donor-dominated CVs. Moreover, such systems, if located within 150 pc are too bright to be selected as spectroscopic targets by SDSS owing to the saturation limit of the SDSS photometry. Nonetheless, it is reasonable to assume that all SU UMa and U Gem CVs within 150 pc have already been identified since their brightening during their relatively frequent outbursts would have unlikely been missed by the many time-domain surveys and amateur observers that regularly scan the sky for transient events (see e.g. Breedt et al. 2014 for a case study of CRTS). An example is U Gem, the prototype for its class, is known since 1855 (Pogson 1857) and has been observed photometrically by SDSS, but not spectroscopically owing to its brightness ($g = 14.6$ mag).

Instead, we cannot exclude that some WZSge systems, nova-like CVs and polars within 150 pc remain to be identified: ASASSN-14dx, a WZSge star located at $d = 81.0 \pm 0.3$ pc with a quiescent magnitude of $V \approx 16.2$ mag (Thorstensen et al. 2016), and TCP J21040470+4631129, another WZSge-type CV located at $d = 109 \pm 2$ pc with a quiescent magnitude of $V \approx 17.7$ mag (Atel #12936), have only been discovered in 2014 and 2019, respectively, following a dwarf nova outburst. We can conclude that the missing systems are most likely those showing low or no variability, such as (i) low mass transfer rate systems with outburst recurrence times of decades, (ii) high mass transfer rate systems (i.e. nova-likes) with stable, hot accretion discs and (iii) strongly magnetic disc-less systems that do not experience disc outburst, which have been in a low-state during the *ROSAT* All Sky X-ray survey, and other serendipitous X-ray observations.

As a final remark, it is important noticing that the colour footprint in which we searched for CV candidates (Figure 7) includes the region in which these low variability systems are expected to be located and therefore the methods we used to derive the completeness are optimally accounting for those objects that most likely are still missing in the 150 pc CV sample.

5 SPACE DENSITY

To determine the space density of CVs, we assumed a Galactic model following the prescription of Pretorius et al.

Table 4. CV space densities for different scale heights. Both the values and the number of systems (N_{CV}) derived from the analysis of the *Gaia* data and the corresponding ones corrected for the completeness of the sample are reported.

h (pc)	d_{lim} (pc)	150	
	Completeness N_{CV}	77% 42	100% 54
		ρ_0 [10^{-6} pc $^{-3}$]	
100		$4.8^{+0.8}_{-1.0}$	$6.3^{+0.9}_{-1.1}$
280		$3.7^{+0.6}_{-0.8}$	$4.8^{+0.6}_{-0.8}$
500		$3.4^{+0.5}_{-0.6}$	$4.3^{+0.5}_{-0.7}$

(2007a) and approximated the Galaxy as an axisymmetric disc, with no halo, no bulge, no spiral structure and no thick disc. We assumed the following density profile:

$$\rho = \rho_0 \exp\left(-\frac{|z|}{h}\right) \quad (6)$$

where z is the distance above the Galactic plane, ρ_0 the space density in the mid-plane, and h the scale height of the CV population. We ignored any possible dependencies from the radial distance from the Galactic centre since they are negligible within the volume we are considering.

ρ_0 is then calculated as the ratio between the number of CVs (N_{CV}) found in the effective volume (V_{eff}) for a given Galactic model (i.e. for a given scale height) and the effective volume itself. One of the unknowns in the determination of the space density is the scale height of the CV population. As discussed in Section 3, h depends on the age of the population, and with the small number of CVs within 150 pc, it is not possible to independently measure h . We therefore assumed $h = 100, 280$ and 500 pc and determined ρ_0 using a Monte Carlo approach to evaluate the effective volume enclosed within a given distance (middle panel of Figure 12). For our distance limit of 150 pc, we found a variation of ≈ 30 per cent of ρ_0 for the two extreme cases of h ($\approx 4.8 \times 10^{-6}$ pc $^{-3}$ for $h = 100$ pc, and $\approx 3.4 \times 10^{-6}$ pc $^{-3}$ for $h = 500$ pc respectively, see Table 4). However, the 150 pc CV sample is dominated by old CVs and hence the higher values we assumed for h are more likely to be more representative of the properties of the observed sample.

In order to investigate for the presence of possible biases as a function of distance, we divided the 150 pc volume in four bins of equal volume (bottom panel of Figure 12). Within the uncertainties ρ_0 remains constant in the four bins, although its value decreases in the outer bin, reflecting the presence of possible detection biases. However, the cumulative distributions of the different CV sub-types (Figure 13) show all the same trend suggesting that, whether present, these detection biases affect all CV sub-classes in the same way.

Considering the volume enclosed within 150 pc, and accounting for the Poisson uncertainties, a conservative measurement of the space density of known CVs results $\rho_0 = (3.7^{+0.6}_{-0.8}) \times 10^{-6}$ pc $^{-3}$ for $h = 280$ pc (solid line in the middle panel Figure 12). The related 3σ uncertainties (dotted lines in the middle panel Figure 12) also include the uncertainty due to the unknown scale height of the Galactic CV population. Accounting for the possibility that the *Gaia* CV sample




Figure 12. *Top:* cumulative distribution of the 150 pc CV sample as a function of the distance (black line) in comparison with the prediction by the Galactic model assuming different scale heights (coloured lines as indicated). *Middle:* ρ_0 as a function of distance for different scale heights. The black solid line corresponds to our conservative value of the CV space density $\rho_0 = (3.7_{-0.8}^{+0.6}) \times 10^{-6} \text{ pc}^{-3}$ for $h = 280 \text{ pc}$. The dashed and dotted lines represent the relative 1σ and 3σ uncertainties, respectively. *Bottom:* CV space density in four bins of equal volumes. The error bars report the Poissonian uncertainties.

is only 77 per cent complete, would imply a space density of $\rho_0 = (4.8_{-0.8}^{+0.6}) \times 10^{-6} \text{ pc}^{-3}$ for $h = 280 \text{ pc}$, which would still be consistent with the conservative value derived above without introducing any correction for incompleteness.

Prior to our *Gaia*-based analysis, the most reliable space density had been estimated using an X-ray selected sample of CVs, $\rho_0 = 4_{-2}^{+6} \times 10^{-6} \text{ pc}^{-3}$ (Pretorius & Knigge 2012). Schwobe (2018) recently re-visited this sample, making use of the *Gaia* distances, and derived $\rho_0 < 5.1 \times 10^{-6} \text{ pc}^{-3}$. Both studies assumed that this magnitude-limited sample of X-ray selected CVs is complete and representative of the intrinsic population, leaving systematic uncertainties

of a factor two in the space density measurement (Pretorius & Knigge 2012).

Our measurement, $\rho_0 = (4.8_{-0.8}^{+0.6}) \times 10^{-6} \text{ pc}^{-3}$, is in good agreement with these previous estimates. However, using a volume-limited sample and the accurate astrometry of *Gaia*, we were able to reduce the uncertainty on the CV density by an order of magnitude.

6 COMPARISON WITH THE MODELS OF CV EVOLUTION

6.1 The intrinsic population

The volume-limited sample of CVs obtained from *Gaia* provides, for the first time, a direct insight into the intrinsic properties of the Galactic population of CVs and allows a direct comparison between the observations and theoretical predictions.

Most models of CV evolution predict that ≈ 99 per cent of the present day CVs should be found below the period gap, with a large fraction of them (≈ 40 – 70 per cent) having already evolved through the period minimum (Kolb 1993; Howell et al. 2001; Goliašch & Nelson 2015). The results from *Gaia* show instead a different picture, with (83 ± 6) per cent of the CVs in the 150 pc sample below the period gap and (17 ± 6) per cent above (Figure 14). More importantly, the *Gaia* 150 pc CV sample contains only three plausible candidate period bouncers, GD 552 (Unda-Sanzana et al. 2008), SDSS J102905.21+485515.2 (Thorstensen et al. 2016) and 1RXS J105010.3–140431 (Patterson 2011; Pala et al. 2017), with WZ Sge, V455 And and EZ Lyn being three additional weaker candidates (they all have brown-dwarf companions, but are right at the period minimum). Thus seven to at most 14 per cent of the 150 pc CVs below the gap are period bouncers, a much smaller fraction than predicted by the population models. This discrepancy could reflect the selection biases discussed above, i.e. it could be possible that a number of period bouncers have not yet been identified because of their low quiescent variability and long outburst recurrence time. Assuming that the *Gaia* 150 pc CV sample is 77 per cent complete (Section 4), ≈ 12 CVs are still to be identified within 150 pc. In the most favourable case in which all these are period bouncers would result into a fraction of (31 ± 7) per cent, bringing the observations into a marginal agreement with the prediction by Goliašch & Nelson (2015), ≈ 40 per cent, although the fraction of CVs above, (13 ± 5) per cent, and below the gap, (87 ± 5) per cent, would still be quite different from the theoretical predictions.

This observed disagreement can be potentially resolved by accounting for the presence of additional AML mechanisms besides the mere MB and GWR. Schreiber et al. (2016) proposed a model in which an empirical consequential AML (eCAML), i.e. a mechanism of AML arising from the mass transfer process itself, is generated by the friction between the secondary star and the white dwarf ejecta during nova explosions. The eCAML could lead to dynamically unstable mass transfer in CVs hosting low-mass white dwarfs, which would not survive as semi-detached binaries but would merge into single objects (see also Nelemans et al. 2016). As a consequence, the fraction of CVs above and below the period gap would become, respectively, ≈ 85 per cent and ≈ 15 per cent, and would be in good agreement with what derived from the analysis of the *Gaia* 150 pc CV sample, (83 ± 6) per cent and (17 ± 6) per cent, even when accounting for incompleteness, (87 ± 5) per cent and (13 ± 5) per cent. The space density predicted by the eCAML model, $\rho_0 \lesssim 2 \times 10^{-5} \text{ pc}^{-3}$ (Belloni et al. 2018), is about a factor 4 higher than the space density derived from the study of the *Gaia* 150 pc sample, $\rho_0 = (4.8^{+0.6}_{-0.8}) \times 10^{-6} \text{ pc}^{-3}$. This difference likely reflects the general uncertainties on the parameters (such as initial mass ratio distribution, initial separa-




Figure 13. Cumulative distribution of the 150 pc CVs as a function of the distance for the different subtypes. The different distributions shows all similar trends, suggesting the absence of clear selection effects in the 150 pc sample.




Figure 14. Period distribution for the CVs in the 150 pc sample. The majority of the systems are found below the period and, accounting also for the two WZ Sge-type stars without an orbital period determination, make up for ≈ 83 per cent of the observed systems.

Table 5. Literature effective temperatures and masses for the CV white dwarfs in the 150 pc sample, ordered according to their orbital periods.

System	P_{orb} (min)	T_{eff} (K)	$\langle \dot{M} \rangle$ ($10^{-11} M_{\odot}/\text{yr}^{-1}$)	M_{WD} (M_{\odot})	Comment	Reference
GW Lib	76.78	$16995 \pm 812 \downarrow$	13 ± 3	0.84 ± 0.02	Pulsating, brightenings	17, 21
BW Scl	78.23	15480 ± 900	10 ± 7	–		16
V627 Peg	78.51	16292 ± 753	13 ± 9	–		17
V455 And	81.08	11799 ± 750	4 ± 2	–	Eclipsing, pulsating	8
WZ Sge	81.63	14900 ± 250	7.4 ± 1.3	0.85 ± 0.04		1, 2
V355 UMa	82.52	12924 ± 1250	5 ± 4	–	Pulsating	33
EZ Lyn	84.97	13595 ± 99	6 ± 4	–	Pulsating, brightenings	27
1RXS J105010.3–140431	88.56	11622 ± 277	3 ± 2	–	Period bouncer candidate	17
V2051 Oph	89.90	–	–	0.78 ± 0.06	Eclipsing	19, 28
VY Aqr	90.85	15148 ± 100	10 ± 6	–		23
OY Car	90.89	15000 ± 2000	8 ± 5	0.84 ± 0.04	Eclipsing	11, 12
EX Hya	98.26	–	–	0.790 ± 0.026	Magnetic	5,6,7,32
VV Pup	100.44	12251 ± 600	4 ± 3	–	Magnetic	20
V834 Cen	101.52	14927 ± 900	10 ± 6	–	Magnetic	20
GD 552	102.73	11118 ± 400	2.9 ± 1.9	–	Period bouncer candidate	9
HT Cas	106.05	14000 ± 1000	22 ± 8	0.61 ± 0.04	Eclipsing	25, 26
VW Hyi	106.95	20000 ± 1000	50 ± 34	$0.71^{+0.18}_{-0.26}$		3, 4
Z Cha	107.28	15700 ± 550	10 ± 3	0.84 ± 0.09	Eclipsing	29, 30
MR Ser	113.47	14816 ± 900	9 ± 6	–	Magnetic	20
BL Hyi	113.64	13818 ± 900	7 ± 5	–	Magnetic	20
ST LMi	113.89	11005 ± 500	2.8 ± 1.8	–	Magnetic	20
AM Her	185.65	19800 ± 700	33 ± 17	$0.78^{+0.12}_{-0.17}$	Magnetic	10
IP Peg	227.82	–	–	1.16 ± 0.02	Eclipsing	24
U Gem	254.74	30000 ± 1000	31 ± 6	1.2 ± 0.05		14, 15
IX Vel	279.25	–	–	0.8 ± 0.2		31
SS Cyg	396.19	–	–	0.81 ± 0.19		22
AE Aqr	592.78	–	–	0.63 ± 0.05	Magnetic, evolved donor	13

References. (1) Sion et al. (1995), (2) Steeghs et al. (2007), (3) Gänsicke & Beuermann (1996), (4) Smith et al. (2006), (5) Eisenbart et al. (2002), (6) Belle et al. (2003), (7) Beuermann & Reinsch (2008), (8) Araujo-Betancor et al. (2005a), (9) Unda-Sanzana et al. (2008), (10) Gänsicke et al. (2006), (11) Horne et al. (1994), (12) Littlefair et al. (2008), (13) Echevarría et al. (2008), (14) Long et al. (2006), (15) Echevarría et al. (2007), (16) Gänsicke et al. (2005), (17) Pala et al. (2017), (19) Baptista et al. (1998), (20) Araujo-Betancor et al. (2005a), (21) van Spaandonk et al. (2010), (22) Bitner et al. (2007), (23) Sion et al. (2003), (24) Copperwheat et al. (2010), (25) Feline et al. (2005), (26) Horne et al. (1991), (27) Szkody et al. (2013), (28) Saito & Baptista (2006), (29) Robinson et al. (1995), (30) Wade & Horne (1988), (31) Neustroev et al. (2011), (32) Suleimanov et al. (2019), (33) Szkody et al. (2010)

Notes. For BW Scl, V455 And, V355 UMa, EZ Lyn, VY Aqr, VV Pup, V834 Cen, GD 552, MR Ser, BL Hyi and ST LMi, the white dwarf T_{eff} reported in the literature has been determined via spectroscopic analyses assuming $\log g = 8$. However, this assumption does not reflect the average mass of CV white dwarfs $M_{\text{WD}} \approx 0.8 M_{\odot}$, i.e. $\log g = 8.35$. For these systems, the T_{eff} reported in this table have been re-computed for $\log g = 8.35$ using equation 1 from Pala et al. (2017).

tion distribution, initial binary fraction, common-envelope and magnetic braking efficiency) employed in binary population synthesis studies. Moreover, the space density derived by Belloni et al. (2018) has been derived from a CV population consisting of 80 per cent period bouncers, while the fraction of such systems is only five per cent in the observed 150 pc sample. Considering only systems that have not evolved through the period minimum yet, the space density predicted by the eCAML model, $\rho_0 \approx 4.5^{+4.5}_{-2.3} \times 10^{-6} \text{ pc}^{-3}$ (Belloni et al. 2019), perfectly agrees with the observation. This implies that either the large fraction of period bounce CVs still has to be identified (although this is not likely the case, see Section 4.2.2), or that the current models fail to properly describe the post period minimum evolution of CVs. Alternatively, it is possible that the time scales required for a CV to evolve to the period minimum are much longer than current models (including eCAML) suggest.

Finally, we note in passing that the 150 pc sample contains some of the most peculiar CVs known:

- AR UMa is the polar with the highest magnetic field,

$\langle B \rangle = 230 \text{ MG}$ (Schmidt et al. 1996). This suggests that strong magnetic fields are probably not as rare as thought (Ferrario et al. 2015) and underlines the urgency to better understand the origin of the magnetic CVs.

- AE Aqr is a post thermal time scale mass transfer system (see Section 6.4). It is also an IP, with the fastest spinning white dwarf, $P_{\text{spin}} = 33 \text{ s}$ (Patterson et al. 1980) among all CVs. This rapid spin results in a propeller mechanism which prevents the mass lost from the donor to reach the white dwarf surface, instead being expelled in the surrounding space.

- EZ Lyn is characterised by quasi-periodic brightening events superimposed to a sinusoidal photometric modulations (Zharikov et al. 2008, 2013) that, in the case of its twin system SDSS J123813.73-033933.0, have been interpreted as the interplay between spiral arms and small amplitude thermal instabilities in the accretion disc (Pala et al. 2019). Similar behaviour, although less periodic, has been observed also in GW Lib, where it could be associated with fluctuation in the mass accretion rate (Toloza et al. 2016).

- V445 And, “The CV that has it all”, displays a graz-




Figure 15. *Left:* effective temperature for magnetic (squares) and non-magnetic (circles) CV white dwarfs from Townsley & Gänsicke (2009) and Pala et al. (2017) (grey). Magnetic and non-magnetic systems within 150 pc from Table 5 are shown in blue and red, respectively. The period gap is highlighted by the grey band. The solid lines represent the evolutionary tracks from Pala et al. (2017) for a typical CV ($M_{\text{WD}} = 0.8 M_{\odot}$, with an initial secondary mass of $M_2 = 0.65 M_{\odot}$ and an initial $P_{\text{orb}} = 12 \text{ h}$) in which AML is driven by both MB and GWR above the period gap while, below the period gap, only GWR (black) or GWR plus a residual MB (red) drive the evolution of the system. *Right:* white dwarf mass distribution for the CVs in the 150 pc sample as reported from the literature (Table 5).

ing eclipse, permanent superhumps, non-radial white dwarf pulsations and a spectroscopic period much longer ($P_{\text{spec}} \approx 3.5 \text{ h}$) than the orbital one ($P_{\text{orb}} \approx 81.08 \text{ min}$, Araujo-Betancor et al. 2005a), the origin of which is still not understood.

The prevalence of several of such peculiar systems suggests that the Galactic population of CVs is intrinsically very variegated and that these systems do not represent exceptional cases. The unexpected behaviour of systems like EZ Lyn and V455 And suggests that the physics describing the accretion process is still far from being completely understood. Moreover, the existence of systems such as ARUMa and AE Aqr is not accounted for by the model of CV evolution and provides another clue of the incompleteness of the current theories describing the evolution of close interacting binaries.

6.2 Mass accretion rates and white dwarf masses

Mass accretion results in compressional heating of the white dwarfs in CVs (Sion 1995; Townsley & Bildsten 2004) and, therefore, the white dwarf effective temperature (T_{eff}) provides a direct measurement of its secular mean accretion rate ($\langle \dot{M} \rangle$, Townsley & Bildsten 2003). The different efficiencies of MB and GWR in removing angular momentum from the binary orbit (Section 1) cause long period CVs to have $\langle \dot{M} \rangle$ about one order of magnitude higher compared to those of short period CVs. Consequently, long period systems should host hotter white dwarfs compared to short period systems and hence T_{eff} measurements provide a direct insight into

the evolutionary stage of the systems (Townsley & Gänsicke 2009; Pala et al. 2017).

Among the 42 CVs found within 150 pc, 21 have a published T_{eff} that can be considered reliable⁸ and we can hence estimate their $\langle \dot{M} \rangle$ using equation 2 from Townsley & Gänsicke (2009). Table 5 lists the corresponding T_{eff} values as compiled from Townsley & Gänsicke (2009) and Pala et al. (2017), and includes four additional systems EZ Lyn, GD 552, Z Cha and V355 UMa. For the systems for which the T_{eff} from the literature has been determined via spectroscopic analyses assuming $\log g = 8$, we re-computed the corresponding T_{eff} for $\log g = 8.35$ using equation 1 from Pala et al. (2017), in order to reflect the average observed mass of CV white dwarfs $M_{\text{WD}} \approx 0.8 M_{\odot}$, i.e. $\log g = 8.35$. The errors on $\langle \dot{M} \rangle$ have been derived accounting for the uncertainties on both T_{eff} and the white dwarf mass (M_{WD}). For EZ Lyn, VY Aqr and V355 UMa we adopt ten per cent of their T_{eff} values as uncertainty, as no errors on T_{eff} were published. For the systems without a mass measurement, we assumed $M_{\text{WD}} = 0.83 \pm 0.23 M_{\odot}$, corresponding to the average mass of CV white dwarfs (Zorotovic et al. 2011).

⁸ Following the prescription by Townsley & Gänsicke (2009), a T_{eff} measurement is considered reliable when it has been derived (i) from the analysis of an ultraviolet spectrum in which the white dwarf signature has been unambiguously identified from the detection of a broad Ly α absorption profile and, possibly, sharp absorption metal lines, or (ii) from the analysis of the eclipse light curve in which both the white dwarf ingress and egress have been clearly detected.

The left panel of Figure 15 shows the effective temperature as a function of the orbital period. For comparison, also the values from Townsley & Gänsicke (2009) and Pala et al. (2017) are reported, together with the evolutionary tracks for a typical CV ($M_{\text{WD}} = 0.8 M_{\odot}$, with an initial secondary mass of $M_2 = 0.65 M_{\odot}$ and an initial $P_{\text{orb}} = 12$ h, Pala et al. 2017). With only two 150 pc CVs having a T_{eff} at $P_{\text{orb}} \gtrsim 180$ min, the comparison with these tracks is only meaningful at short orbital periods. The T_{eff} distribution of the intrinsic population shows some fundamental discrepancies between the theory and the observations that have been already highlighted by Pala et al. (2017) from the study of a large sample of CVs observed with the *Hubble Space Telescope*. In particular, the systems with the period minimum are characterised by a large scatter in T_{eff} as the white dwarf temperatures due to compressional heating are very sensitive to the white dwarf mass (Townsley & Gänsicke 2009). The recent work by Belloni et al. (2019) has shown that the mass distribution of CV white dwarfs is indeed the main reason behind this spread. Moreover, the systems below the period gap host hotter white dwarfs than suggested by the standard models (black track). This is also true in the case of the period bouncer track, which appears to be steeper than the theoretical predictions. As already discussed by Pala et al. (2017), these findings suggest that additional AML mechanisms are present (red track) besides pure GR in this period range. These additional AML mechanisms could also imply that period bounce CVs evolve faster than predicted by the models potentially explaining the lack of such evolved systems in the intrinsic population.

As shown in Figure 15 (left), the effective temperatures of systems with similar orbital periods but different distances (coloured against grey points) are fully consistent among each other, allowing us to rule out the presence of observational biases as possible explanation for the discrepancies between theory and observations. Our results, instead, highlights the need for a revision of the current models of compact binary evolution.

Finally, the masses of 14 CV white dwarfs in the 150 pc sample are available in the literature (Table 5, right panel of Figure 15). The average mass results $\langle M_{\text{WD}} \rangle = 0.83 \pm 0.17$, in perfect agreement with the measurement from Zorotovic et al. (2011), $\langle M_{\text{WD}} \rangle = 0.83 \pm 0.23 M_{\odot}$. As also shown by Zorotovic et al. (2011), this result confirms that the higher mass of CV white dwarfs compared to that of single white dwarfs and their detached progenitors, $\langle M_{\text{WD}} \rangle \simeq 0.6 M_{\odot}$ (Koester et al. 1979; Liebert et al. 2005b; Kepler et al. 2007), cannot be related to an observational bias.

The average mass of CV white dwarfs cannot be explained (i) invoking different parent populations for the present day CVs and the present day pre-CVs (e.g. Zorotovic et al. 2011) or (ii) assuming mass growth during nova cycles or through thermal time-scale mass transfer (Wijnen et al. 2015). Instead, the eCAML proposed by Schreiber et al. (2016) could mitigate the discrepancy between the theory and the observations. The eCAML leads to merger of the two stellar components in systems hosting low mass white dwarfs, which would then disappear from the CV population thus naturally explaining the observed high average mass of CV white dwarfs. Nonetheless, the exact mechanism behind the additional CAML and the reason for its dependence on white dwarf mass are unclear.




Figure 16. CV subtype contribution to the 150 pc sample. More than one third of the observed CVs host a magnetic white dwarf, in clear contrast with the total absence of magnetic white dwarfs in the parent population of post common envelope binaries (Liebert et al. 2005a, 2015).

6.3 Magnetic systems

Pretorius et al. (2013) derived the space density of magnetic CVs as $\rho_{\text{mCV}} = 1.3^{+0.6}_{-0.4} \times 10^{-6} \text{ pc}^{-3}$. Comparing this value with the space density of non-magnetic CVs $\rho_0 = 4^{+6}_{-2} \times 10^{-6} \text{ pc}^{-3}$ from Pretorius & Knigge (2012), implies that about one third of CVs are magnetic.

From the analysis of the *Gaia* parallaxes, we found that a large fraction, (36 ± 7) per cent, of the CVs identified within 150 pc contain a magnetic white dwarf (Figure 16). This corresponds to a space density⁹ of $\rho_{\text{mCV}} = 1.3^{+0.3}_{-0.4} \times 10^{-6} \text{ pc}^{-3}$ (Table 6), consistent with the result by Pretorius et al. (2013).

Magnetic CVs are thought to follow a different evolutionary path compared to non-magnetic system. Owing to the coupling with the magnetic field of the white dwarf, MB is reduced or even completely suppressed in the system with the strongest magnetic fields (Li et al. 1994). Consequently, IPs and polars evolve slower than non-magnetic CVs, thus explaining the high fraction of observed magnetic CVs (Araujo-Betancor et al. 2005b).

However, this high incidence of magnetism in CVs is not reflected in the fraction of magnetic white dwarfs observed in their parent population, i.e. the post common envelope binaries (PCEBs), with no confirmed magnetic white dwarf detected in any of these detached binaries (Liebert et al. 2005a, 2015). Different scenarios have been proposed to explain the observed fraction of magnetic CVs, such as Ap and Bp progenitors which preserve a fossil magnetic field while becoming white dwarfs (Angel et al. 1981) or interaction during the common envelope phase (Tout et al. 2008). How-

⁹ Note that the values reported in Table 6 cannot be corrected for incompleteness since this would require the knowledge of the contribution of each subtype to the population of CVs that still has to be discovered.

Table 6. Space densities for different CV sub-types computed considering the volume enclosed within $d_{\text{lim}} = 150$ pc assuming a scale height $h = 280$ pc.

Subtype	N_{CV}	$N_{\text{CV}}/N_{\text{tot}}$	ρ_0 [10^{-6} pc $^{-3}$]
WZ Sge	12	(28 ± 7) %	1.0 ± 0.3
SU UMa	10	(24 ± 7) %	0.8 $^{+0.2}_{-0.4}$
U Gem	3	(7 ± 4) %	0.2 $^{+0.2}_{-0.1}$
NL	2	(5 ± 3) %	1.5 $^{+0.2}_{-0.3}$
Magnetic	15	(36 ± 7) %	1.3 $^{+0.3}_{-0.4}$

ever, they have been unable to explain the lack of magnetic white dwarfs in the population of PCEB and, consequently, the origin of magnetic CVs remains unclear.

6.4 Systems with an evolved donor

The stability of the mass transfer process requires a mass ratio $q = M_2/M_{\text{WD}} \lesssim 1$ (where M_2 is the mass of the secondary star, Frank et al. 2002). However, it has been shown that CVs hosting massive donors ($M_2 \approx 1.5 M_{\odot}$) could survive a phase of thermal-time-scale mass transfer during which the accreted material burns steadily onto the white dwarf (Schenker et al. 2002). During the thermal-time-scale mass transfer, the donor is stripped of its envelope and the surviving system is a normal CV in which the white dwarf accretes from the remnant core of its companion, rich of CNO processed material. These CVs are predicted to make up for ≈ 30 per cent of the Galactic CV population (Schenker et al. 2002).

These systems can be easily recognised in the ultraviolet from their enhanced NV/CIV line flux ratios compared to those of CVs that have formed through the standard channel and the observed fraction of CVs with an evolved donor has been found to be ≈ 15 per cent (Gänsicke et al. 2003). In the 150 pc CV sample, we identified only two of such CVs: AE Aqr (Jameson et al. 1980) and V2301 Oph, (Schmidt & Stockman 2001), corresponding to a fraction of (5 ± 3) per cent. The higher fraction found by Gänsicke et al. (2003) can be explained by an observational bias, these systems host more massive (and hence larger) and brighter donors compared to those of normal CVs and therefore are easily detected even at large distances. Instead, the theoretically predicted fraction by Schenker et al. (2002), 30 per cent, is much higher compared to the observed fraction of CV with evolved donors in the 150 pc sample.

CVs with evolved donors represent an evolutionary link between CVs and the more compact AM CVn stars. The formation of the latter is still poorly understood but three different pathways have been proposed (Nelemans 2005). In the first scenario, the systems are formed from a double white dwarf binary in which the less massive star is brought to contact with its Roche lobe by the orbital shrinkage due to GWR. Alternatively, the progenitors could be a binary containing a white dwarf plus a non-degenerate, helium core burning star. Finally, AM CVn could descend from CVs with an evolved donor. Hosting massive secondary stars, these systems are able to evolve to P_{orb} much shorter than the period minimum. Owing to the mass transfer process, the




Figure 17. Hertzsprung-Russell diagram of the *Gaia* sources within 150 pc (grey) showing, as a reference, the position occupied by the Sun (Casagrande & Vandenberg 2018) and by V379 Tel (pink), *Gaia* J154008.28-392917.6 (blue) and *Gaia* J051903.99+630340.4 (orange) and their common proper motion companions (black), which result being a M-dwarf, a white dwarf (WD) and a G-type star, respectively.

secondary is progressively eroded, till only its Helium-rich core is left and an AM CVn star is born. The fraction of CVs with an evolved donor that we derived could provide an upper limit on the number of AM CVn that are expected to form through this channel, yielding a valuable observational test for the contribution of the different formation channels to the overall population of these compact systems.

7 TERTIARY COMPANIONS

The formation and the evolution of a binary can be influenced by the presence of a nearby third body orbiting the system. For example, the third body can give rise to Lidov-Kozai cycles that can induce periodical variation in the eccentricity and the inclination of the inner binary with respect to the orbital plane of the external body (see Naoz 2016 for a review). This gravitational interaction can hence affect the mass transfer process and potentially even lead to mergers (e.g. Toonen et al. 2016). In order to fully constrain the formation and evolution of CVs, it is important to assess the multiplicity fractions of these systems.

In the past, the most successful method to identify wide orbit companion was the identification of period modulations in the long term light curves of the system and several studies have suggested that some CVs could be part of triple systems. Few examples are: VY Scl (Martínez-Pais et al. 2000), DP Leo (Beuermann et al. 2011), FS Aur, (Chavez et al. 2012) and LX Ser (Li et al. 2017), which have been suggested to host circumbinary sub-stellar objects or giant planets. However, the observed modulations could also be explained by the magnetic activity of the donor star and, owing to the lack of a direct detection of the third bodies, it is difficult to disentangle the two scenarios.

The accurate parallaxes and proper motions delivered

Table 7. 150 pc CVs with a common proper motion companion.

System	Gaia DR2 ID	P_{orb}	Type	ϖ	μ_{RA}	μ_{Dec}	Δ	Angular separation	D	$-E_{\text{bind}}$	
		(min)		(mas)	(mas yr $^{-1}$)	(mas yr $^{-1}$)		(pc)	(AU)	(erg)	
V379 Tel	6658737220627065984 6658737388128701184	101.03	AM M2V	7.65(7) 7.52(5)	-60.8(1) -60.64(8)	-17.75(9) -18.44(6)	2.9	30.8''	2.3 ± 1.5	$(5 \pm 3) \times 10^5$	$(1.5 \pm 1.0) \times 10^{40}$
Gaia J154008.28-392917.6	6008982469163902464 6002961479076981632	-	UGWZ WD	7.49(11) 7.6(8)	64.8(2) 52(2)	1.6(2) 4(1)	2.4	1.1°	3 ± 13	$(6 \pm 26) \times 10^5$	$(2 \pm 8) \times 10^{40}$
Gaia J051903.99+630340.4 TYC 4084-172-1	285957277597658240 285957277597658368	126:	UGSU G3V	8.59(4) 8.43(3)	-13.07(5) -11.19(3)	-45.30(5) -44.38(4)	11	6.8''	2.2 ± 0.7	$(4.6 \pm 1.4) \times 10^5$	$(3.7 \pm 1.1) \times 10^{40}$

Notes. $-E_{\text{bind}}$ has been computed assuming $M_3 = 0.4 M_{\odot}$, $M_3 = 0.6 M_{\odot}$ and $M_3 = 1 M_{\odot}$ for V379 Tel, Gaia J154008.28-392917.6 and Gaia J051903.99+630340.4, respectively.

by the Gaia space mission in its DR2 offer the first opportunity to carry out a systematic search for third components on wide orbits to the 150 pc CVs by searching for common proper motion companions.

In order to identify these objects, we performed a cone search of 3 pc radius around each of the 150 pc CVs, and selected those objects for which

$$\Delta = \sqrt{\left(\frac{\Delta\mu_{\text{RA}}}{\sigma_{\mu_{\text{RA}}}}\right)^2 + \left(\frac{\Delta\mu_{\text{Dec}}}{\sigma_{\mu_{\text{Dec}}}}\right)^2} < 3 \quad (7)$$

where $\Delta\mu$ are the differences in the proper motion components and σ_{μ} are the quadrature sum of the corresponding 3σ uncertainties.

We find that two CVs, V379 Tel and Gaia J154008.28-392917.6, satisfy this condition. Using the Gaia coordinates and parallaxes, we computed the separation between the inner binary and the third body which resulted $D \simeq 2.5$ pc (Table 7). Although it did not satisfy the previous condition ($\Delta \simeq 11$), we cannot ignore that Gaia J051903.99+630340.4 has a nearby companion (TYC 4084-172-1) located at a similar distance ($\simeq 2.2$ pc) and therefore we also include this object in the following discussion.

Given the relatively large orbital separations, we computed the gravitational binding energy:

$$-E_{\text{bind}} = G \frac{(M_{\text{WD}} + M_2)M_3}{D} \quad (8)$$

of each triplet assuming that the inner binary is a typical short period CV (as the three systems listed in Table 7 are all located below the period gap), with a white dwarf of mass $M_{\text{WD}} = 0.83 M_{\odot}$ and a donor of mass $M_2 = 0.14 M_{\odot}$, typical of a CV below the period gap (Knigge et al. 2011). We assumed that the third body is on a circular orbit around the inner binary and we estimated their possible masses accordingly to their position on the Hertzsprung-Russell diagram (Figure 17). The third companions result being a M-dwarf of spectral type \simeq M2V (as we estimated using as a reference the table¹⁰ from Pecaute & Mamajek 2013), a white dwarf (WD) and a G-type star for V379 Tel, Gaia J154008.28-392917.6 and Gaia J051903.99+630340.4, respectively. We therefore assumed:

- $M_3 = 0.4 M_{\odot}$, corresponding to the mass of a M2V star

(Pecaute & Mamajek 2013), in the case of the tertiary companion of V379 Tel;

- $M_3 = 0.6 M_{\odot}$, corresponding to the average mass of single white dwarfs (Liebert et al. 2005b), in the case of the tertiary companion of Gaia J154008.28-392917.6;
- $M_3 = 1 M_{\odot}$ in the case of TYC 4084-172-1, given its spectral type G3V and surface gravity of $\log(g) = 4.2(2)$ (Frasca et al. 2018, see also Figure A4).

The binding energy we derived are listed in Table 7. The related uncertainties have been computed accounting for those on the Gaia coordinates and parallaxes of each pair of objects, and are dominated by the latter. At this stage, their large values combined with the unknown uncertainties related to the theoretical limit for stellar binding energies, $-E_{\text{bind}} \simeq 1 \times 10^{41}$ (Burgasser et al. 2007), make it impossible to unambiguously establish whether these CVs are part of hierarchical systems and additional observational efforts are required in order to finally assess their hierarchical structure.

8 CONCLUSIONS

Making use of the accurate astrometry delivered by the ESA Gaia space mission in its DR2, we carry out the first detailed study of the volume-limited sample of 42 CVs located within 150 pc. Combining the Gaia data with the photometric and spectroscopic observations from SDSS and CRTS, we estimate the sample to be (77 ± 10) per cent complete. This is mainly dictated by the efficiency of the discovery methods employed in detecting new CVs, which are biased towards systems accreting at intermediate mass rates, that can be easily detected in the X-rays or thanks to their dwarf nova outbursts.

Assuming $h = 280$ pc as a typical scale height for the Galactic CV population, we estimate the CV space density, which results $\rho_0 = (4.8^{+0.6}_{-0.8}) \times 10^{-6} \text{ pc}^{-3}$. Thanks to the exquisite Gaia data, we reduce the uncertainty on ρ_0 by a factor of ten compared to the pre-Gaia estimates. The uncertainties we derive take well into account for the estimated completeness of the sample and for possible different values of the scale height in the range 100 – 500 pc. Nonetheless, given that the 150 pc CV sample is dominated by short period systems representing the old component of the Galactic CV population, it is reasonable to assume that the larger values of h are likely the most realistic.

The advent of the Gaia space mission provides also the

¹⁰ http://www.pas.rochester.edu/~emamajek/EEM_dwarf_UBVIJHK_colors_Teff.txt

unique opportunity to study the intrinsic properties of the Galactic CV population and to constrain the models describing the formation and evolution of these systems. We find that the observed space density is significantly lower than predicted by the current available models of CV evolution. Moreover, the fractions of CVs above (17 per cent) and below (83 per cent) the period gap are in clear disagreement with the theoretical predictions (1 per cent and 99 per cent, respectively). Both discrepancies can be resolved by the recently proposed eCAML model in which CVs hosting low-mass white dwarfs merge owing to frictional AML arising from nova explosions. Consequently, the Galactic CV population would be composed by a lower absolute number of systems that would naturally imply a lower space density. Moreover, the fractions of CVs predicted by the eCAML model results 15 per cent and 85 per cent above and below the period gap respectively, in better agreement with the observed value. The disappearance of the CVs hosting low-mass white dwarfs would also be consistent with the average masses of the CVs white dwarfs in the 150 pc sample, $\langle M_{\text{WD}} \rangle = 0.83 \pm 0.17 M_{\odot}$, being higher than the masses of their detached progenitors (i.e. PCEB, $\langle M_{\text{WD}} \rangle \simeq 0.6 M_{\odot}$, Zorotovic et al. 2011). The need to include additional mechanisms of AML in the models is also supported by the observed effective temperatures and, consequently, mass accretion rates of the 150 pc CVs, which, below the period gap, are found to be accreting at higher rates than theoretically predicted. However, the observed fraction of period bounce CVs, 7-14 per cent is much lower than current models (including eCAML) predicts, 40 – 80 per cent and it is possible that the time scales required for a CV to evolve to the period minimum are much longer than theoretically predicted.

Studying the composition of the 150 pc CV sample, we identify a large fraction of magnetic CVs, 36 per cent. This finding is particularly intriguing given that no confirmed magnetic white dwarf is known among the PCEBs. All the models proposed to explain the observed fraction of magnetic CVs predict also the existence of magnetic white dwarfs in PCEBs and, consequently, this high incidence of magnetism among CV white dwarfs remains unclear.

We also show that the fraction of CVs hosting nuclear evolved donors is $\simeq 5$ per cent, lower than the pre-*Gaia* observational estimate, $\simeq 15$ per cent. Most likely, this difference arises from an observational bias since these systems are brighter than normal CVs and are easily detected even at large distances. Moreover, the observed fraction of CVs with evolved donors is significantly lower than predicted by the theory (30 per cent).

Finally, we find that three CVs have a common proper motion companion. However, the lack of accurate system parameters does not allow to draw definite conclusion on whether they form hierarchical triple systems and additional observations are required to finally establish whether these common proper motion companion pairs are gravitationally bound.

ACKNOWLEDGEMENTS

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and

Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

The research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP/2007–2013) / ERC Grant Agreement n. 320964 (WDTracer).

Based on observations made with ESO Telescopes at Paranal Observatory under programme ID 0101.C-0646(A).

Based on observations obtained under programme ID SO2018B-015 at the Southern Astrophysical Research (SOAR) telescope, which is a joint project of the Ministério da Ciência, Tecnologia, Inovações e Comunicações (MCTIC) do Brasil, the U.S. National Optical Astronomy Observatory (NOAO), the University of North Carolina at Chapel Hill (UNC), and Michigan State University (MSU).

The research leading to these results has received funding from the European Research Council under the European Union’s Horizon 2020 research and innovation programme n. 677706 (WD3D).

The work presented in this article made large use of TOPCAT and STILTS Table/VOTable Processing Software (Taylor 2005).

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy Office of Science. The SDSS-III web site is <http://www.sdss3.org/>.

SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Partici-

pation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

A.A. acknowledges generous supports from Faculty of Science and Naresuan University M.R.S. thanks for support from Fondecyt (grant 1181404). O.T. was supported by a Leverhulme Trust Research Project Grant. B.T.G. and O.T. were supported by the UK STFC grant ST/P000495.

REFERENCES

- Ak T., Bilir S., Ak S., Retter A., 2007, *New Astron.*, **12**, 446
- Angel J. R. P., Borra E. F., Landstreet J. D., 1981, *ApJS*, **45**, 457
- Araujo-Betancor S., et al., 2005a, *A&A*, **430**, 629
- Araujo-Betancor S., Gänsicke B. T., Long K. S., Beuermann K., de Martino D., Sion E. M., Szkody P., 2005b, *ApJ*, **622**, 589
- Bailer-Jones C. A. L., 2015, *PASP*, **127**, 994
- Bailey J., 1981, *MNRAS*, **197**, 31
- Banzatti A., Pascucci I., Edwards S., Fang M., Gorti U., Flock M., 2019, *ApJ*, **870**, 76
- Baptista R., Catalan M. S., Horne K., Zilli D., 1998, *MNRAS*, **300**, 233
- Belle K. E., Howell S. B., Sion E. M., Long K. S., Szkody P., 2003, *ApJ*, **587**, 373
- Belloni T., Verbunt F., Schmitt J. H. M. M., 1993, *A&A*, **269**, 175
- Belloni D., Schreiber M. R., Zorotovic M., Izkiewicz K., Hurley J. R., Giersz M., Lagos F., 2018, *MNRAS*, **478**, 5626
- Belloni D., Schreiber M. R., Pala A. F., Gänsicke B. T., Zorotovic M., Rodrigues C. V., 2019, arXiv e-prints, p. [arXiv:1910.06333](https://arxiv.org/abs/1910.06333)
- Bernardini F., de Martino D., Falanga M., Mukai K., Matt G., Bonnet-Bidaud J.-M., Masetti N., Mouchet M., 2012, *A&A*, **542**, A22
- Beuermann K., 2006, *A&A*, **460**, 783
- Beuermann K., Reinsch K., 2008, *A&A*, **480**, 199
- Beuermann K., Schwöpe A., Weissieker H., Motch C., 1985, *Space Sci. Rev.*, **40**, 135
- Beuermann K., Harrison T. E., McArthur B. E., Benedict G. F., Gänsicke B. T., 2003, *A&A*, **412**, 821
- Beuermann K., et al., 2011, *A&A*, **526**, A53
- Bitner M. A., Robinson E. L., Behr B. B., 2007, *ApJ*, **662**, 564
- Breedt E., et al., 2014, *MNRAS*, **443**, 3174
- Burgasser A. J., Reid I. N., Siegler N., Close L., Allen P., Lowrance P., Gizis J., 2007, *Protostars and Planets V*, pp 427–441
- Casagrande L., VandenBerg D. A., 2018, *MNRAS*, **479**, L102
- Chavez C. E., Tovmassian G., Aguilar L. A., Zharikov S., Henden A. A., 2012, *A&A*, **538**, A122
- Clemens J. C., Crain J. A., Anderson R., 2004, in Moorwood A. F. M., Iye M., eds, *Proc. SPIE Vol. 5492, Ground-based Instrumentation for Astronomy*. pp 331–340, doi:10.1117/12.550069
- Copperwheat C. M., Marsh T. R., Dhillon V. S., Littlefair S. P., Hickman R., Gänsicke B. T., Southworth J., 2010, *MNRAS*, **402**, 1824
- Dawson K. S., et al., 2013, *AJ*, **145**, 10
- Denisenko D., 2018, *The Astronomer's Telegram*, **11917**
- Denisenko D. V., Larin I., 2018, preprint, ([arXiv:1807.04574](https://arxiv.org/abs/1807.04574))
- Downes R. A., Webbink R. F., Shara M. M., Ritter H., Kolb U., Duerbeck H. W., 2001, *Publications of the Astronomical Society of the Pacific*, **113**, 764
- Drake A. J., et al., 2009, *ApJ*, **696**, 870
- Drake A. J., et al., 2014, *MNRAS*, **441**, 1186
- Duerbeck H. W., 1999, *Information Bulletin on Variable Stars*, **4731**
- Echevarría J., de la Fuente E., Costero R., 2007, *AJ*, **134**, 262
- Echevarría J., Smith R. C., Costero R., Zharikov S., Michel R., 2008, *MNRAS*, **387**, 1563
- Eisenbart S., Beuermann K., Reinsch K., Gänsicke B. T., 2002, *A&A*, **382**, 984
- Fang M., et al., 2018, *ApJ*, **868**, 28
- Feline W. J., Dhillon V. S., Marsh T. R., Watson C. A., Littlefair S. P., 2005, *MNRAS*, **364**, 1158
- Ferrario L., de Martino D., Gänsicke B. T., 2015, *Space Sci. Rev.*, **191**, 111
- Frank J., King A., Raine D. J., 2002, *Accretion Power in Astrophysics: Third Edition*
- Frasca A., Guillout P., Klutsch A., Ferrero R. F., Marilli E., Biazzo K., Gandolfi D., Montes D., 2018, *A&A*, **612**, A96
- Gaia Collaboration et al., 2016, *A&A*, **595**, A1
- Gaia Collaboration et al., 2018, *A&A*, **616**, A1
- Gänsicke B. T., 2005, in Hameury J.-M., Lasota J.-P., eds, *Astronomical Society of the Pacific Conference Series Vol. 330, The Astrophysics of Cataclysmic Variables and Related Objects*. p. 3 ([arXiv:astro-ph/0410412](https://arxiv.org/abs/astro-ph/0410412))
- Gänsicke B. T., Beuermann K., 1996, *A&A*, **309**, L47
- Gänsicke B. T., Koester D., 1999, *A&A*, **346**, 151
- Gänsicke B. T., Sion E. M., Beuermann K., Fabian D., Cheng F. H., Krautter J., 1999, *A&A*, **347**, 178
- Gänsicke B. T., et al., 2003, *ApJ*, **594**, 443
- Gänsicke B. T., Szkody P., Howell S. B., Sion E. M., 2005, *ApJ*, **629**, 451
- Gänsicke B. T., Long K. S., Barstow M. A., Hubeny I., 2006, *ApJ*, **639**, 1039
- Gänsicke B. T., et al., 2009, *MNRAS*, **397**, 2170
- Gentile Fusillo N. P., et al., 2019, *MNRAS*, **482**, 4570
- Goliasch J., Nelson L., 2015, *ApJ*, **809**, 80
- Graham M. J., et al., 2019, *Publications of the Astronomical Society of the Pacific*, **131**, 078001
- Hamann F., Persson S. E., 1992, *ApJS*, **82**, 247
- Hameury J.-M., Menou K., Dubus G., Lasota J.-P., Hure J.-M., 1998, *MNRAS*, **298**, 1048
- Harrison T. E., Johnson J. J., McArthur B. E., Benedict G. F., Szkody P., Howell S. B., Gelino D. M., 2004, *AJ*, **127**, 460
- Hernández Santisteban J. V., Knigge C., Pretorius M. L., Sullivan M., Warner B., 2018, *MNRAS*, **473**, 3241
- Hodgkin S. T., Wyrzykowski L., Blagorodnova N., Kozlov S., 2013, *Philosophical Transactions of the Royal Society of London Series A*, **371**, 20120239
- Hollands M. A., Tremblay P. E., Gänsicke B. T., Gentile-Fusillo N. P., Toonen S., 2018, *MNRAS*, **480**, 3942
- Horne K., Wood J. H., Stiening R. F., 1991, *ApJ*, **378**, 271
- Horne K., Marsh T. R., Cheng F. H., Hubeny I., Lanz T., 1994, *ApJ*, **426**, 294
- Howell S. B., Nelson L. A., Rappaport S., 2001, *ApJ*, **550**, 897
- Jameson R. F., King A. R., Sherrington M. R., 1980, *MNRAS*, **191**, 559
- Kato T., et al., 2015, *PASJ*, **67**, 105
- Kato T., et al., 2016, *PASJ*, **68**, 65
- Kepler S. O., Kleinman S. J., Nitta A., Koester D., Castanheira B. G., Giovannini O., Costa A. F. M., Althaus L., 2007, *MNRAS*, **375**, 1315
- Knigge C., Baraffe I., Patterson J., 2011, *ApJS*, **194**, 28
- Kochanek C. S., et al., 2017, *PASP*, **129**, 104502
- Koester D., Schulz H., Weidemann V., 1979, *A&A*, **76**, 262
- Kolb U., 1993, *A&A*, **271**, 149

- Kulkarni S. R., 2013, *The Astronomer's Telegram*, **4807**
- Larin I., Denisenko D., Pogrebisskiy S., 2018, *The Astronomer's Telegram*, **11401**
- Law N. M., et al., 2009, *PASP*, **121**, 1395
- Lépine S., Bergeron P., Lanning H. H., 2011, *AJ*, **141**, 96
- Li J. K., Wu K. W., Wickramasinghe D. T., 1994, *MNRAS*, **268**, 61
- Li K., et al., 2017, *PASJ*, **69**, 28
- Liebert J., et al., 2005a, *AJ*, **129**, 2376
- Liebert J., Bergeron P., Holberg J. B., 2005b, *ApJS*, **156**, 47
- Liebert J., Ferrario L., Wickramasinghe D. T., Smith P. S., 2015, *ApJ*, **804**, 93
- Lindgren L., Lammers U., Hobbs D., O'Mullane W., Bastian U., Hernández J., 2012, *A&A*, **538**, A78
- Lindgren L., et al., 2018, *A&A*, **616**, A2
- Linnell A. P., Godon P., Hubeny I., Sion E. M., Szkody P., 2007, *ApJ*, **662**, 1204
- Linnell A. P., Godon P., Hubeny I., Sion E. M., Szkody P., Barrett P. E., 2009, *ApJ*, **703**, 1839
- Lipunov V., et al., 2010, *Advances in Astronomy*, **2010**, 349171
- Littlefair S. P., Dhillon V. S., Marsh T. R., Gänsicke B. T., Southworth J., Watson C. A., 2006, *Science*, **314**, 1578
- Littlefair S. P., Dhillon V. S., Marsh T. R., Gänsicke B. T., Southworth J., Baraffe I., Watson C. A., Copperwheat C., 2008, *MNRAS*, **388**, 1582
- Liu W.-M., Li X.-D., 2019, *ApJ*, **870**, 22
- Long K. S., Brammer G., Froning C. S., 2006, *ApJ*, **648**, 541
- Luri X., et al., 2018, preprint, ([arXiv:1804.09376](https://arxiv.org/abs/1804.09376))
- Martínez-Pais I. G., Martín-Hernández N. L., Casares J., Rodríguez-Gil P., 2000, *ApJ*, **538**, 315
- Maza J., Gonzalez L. E., 1983, *IAU Circ.*, **3854**
- McAllister M. J., et al., 2017, *MNRAS*, **467**, 1024
- McAllister M., et al., 2019, *MNRAS*, **486**, 5535
- McArthur B. E., et al., 2001, *ApJ*, **560**, 907
- Meyer F., Meyer-Hofmeister E., 1984, *A&A*, **132**, 143
- Miller-Jones J. C. A., Sivakoff G. R., Knigge C., Körding E. G., Templeton M., Waagen E. O., 2013, *Science*, **340**, 950
- Naos S., 2016, *ARA&A*, **54**, 441
- Natta A., Testi L., Alcalá J. M., Rigliaco E., Covino E., Stelzer B., D'Elia V., 2014, *A&A*, **569**, A5
- Nelemans G., 2005, in Hameury J.-M., Lasota J.-P., eds, *Astronomical Society of the Pacific Conference Series Vol. 330, The Astrophysics of Cataclysmic Variables and Related Objects*. p. 27 ([arXiv:astro-ph/0409676](https://arxiv.org/abs/astro-ph/0409676))
- Nelemans G., Siess L., Repetto S., Toonen S., Phinney E. S., 2016, *ApJ*, **817**, 69
- Neustroev V. V., Suleimanov V. F., Borisov N. V., Belyakov K. V., Shearer A., 2011, *MNRAS*, **410**, 963
- Neustroev V. V., et al., 2017, *MNRAS*, **467**, 597
- Nisini B., Antonucci S., Alcalá J. M., Giannini T., Manara C. F., Natta A., Fedele D., Biazzo K., 2018, *A&A*, **609**, A87
- Osaki Y., 1974, *PASJ*, **26**, 429
- Paczynski B., Sienkiewicz R., 1983, *The Astrophysical Journal*, **268**, 825
- Pala A. F., et al., 2017, *MNRAS*, **466**, 2855
- Pala A. F., Schmidtbreick L., Tappert C., Gänsicke B. T., Mehner A., 2018, *MNRAS*, **481**, 2523
- Pala A. F., et al., 2019, *MNRAS*, **483**, 1080
- Parsons S. G., et al., 2013, *MNRAS*, **429**, 256
- Patterson J., 1984, *ApJS*, **54**, 443
- Patterson J., 1998, *PASP*, **110**, 1132
- Patterson J., 2011, *MNRAS*, **411**, 2695
- Patterson J., Branch D., Chincarini G., Robinson E. L., 1980, *ApJ*, **240**, L133
- Patterson J., Thorstensen J. R., Kemp J., 2005, *PASP*, **117**, 427
- Pecaut M. J., Mamajek E. E., 2013, *ApJS*, **208**, 9
- Pogson N., 1857, *MNRAS*, **17**, 200
- Pojmanski G., 1997, *Acta Astron.*, **47**, 467
- Politano M., 1996, *ApJ*, **465**, 338
- Pretorius M. L., Knigge C., 2012, *MNRAS*, **419**, 1442
- Pretorius M. L., Knigge C., Kolb U., 2007a, *MNRAS*, **374**, 1495
- Pretorius M. L., Knigge C., O'Donoghue D., Henry J. P., Gioia I. M., Mullis C. R., 2007b, *MNRAS*, **382**, 1279
- Pretorius M. L., Knigge C., Schwöpe A. D., 2013, *MNRAS*, **432**, 570
- Ramsay G., Schreiber M. R., Gänsicke B. T., Wheatley P. J., 2017, *A&A*, **604**, A107
- Ramsay G., et al., 2018, *A&A*, **620**, A141
- Rappaport S., Verbunt F., Joss P. C., 1983, *ApJ*, **275**, 713
- Richards G. T., et al., 2002, *AJ*, **123**, 2945
- Ritter H., Kolb U., 2003, *A&A*, **404**, 301
- Rixon G., et al., 2014, *The Astronomer's Telegram*, **6593**
- Robinson E. L., et al., 1995, *ApJ*, **443**, 295
- Saito R. K., Baptista R., 2006, *AJ*, **131**, 2185
- Schenker K., King A. R., Kolb U., Wynn G. A., Zhang Z., 2002, *MNRAS*, **337**, 1105
- Schmidt G. D., Stockman H. S., 2001, *ApJ*, **548**, 410
- Schmidt G. D., Szkody P., Smith P. S., Silber A., Tovmassian G., Hoard D. W., Gänsicke B. T., de Martino D., 1996, *ApJ*, **473**, 483
- Schneider D. P., et al., 2010, *AJ*, **139**, 2360
- Schreiber M. R., et al., 2010, *A&A*, **513**, L7
- Schreiber M. R., Zorotovic M., Wijnen T. P. G., 2016, *MNRAS*, **455**, L16
- Schwöpe A. D., 2018, *A&A*, **619**, A62
- Schwöpe A. D., Schwarz R., Greiner J., 1999, *A&A*, **348**, 861
- Shappee B. J., et al., 2014, *ApJ*, **788**, 48
- Sherrington M. R., Jameson R. F., Bailey J., Giles A. B., 1982, *MNRAS*, **200**, 861
- Sion E. M., 1995, *ApJ*, **438**, 876
- Sion E. M., Cheng F. H., Long K. S., Szkody P., Gilliland R. L., Huang M., Hubeny I., 1995, *ApJ*, **439**, 957
- Sion E. M., Szkody P., Cheng F., Gänsicke B. T., Howell S. B., 2003, *ApJ*, **583**, 907
- Skrutskie M. F., et al., 2006, *AJ*, **131**, 1163
- Smith A. J., Haswell C. A., Hynes R. I., 2006, *MNRAS*, **369**, 1537
- Soderblom D. R., Hillenbrand L. A., Jeffries R. D., Mamajek E. E., Naylor T., 2014, *Protostars and Planets VI*, pp 219–241
- Sproats L. N., Howell S. B., Mason K. O., 1996, *MNRAS*, **282**, 1211
- Spruit H. C., Ritter H., 1983, *A&A*, **124**, 267
- Steehgs D., Howell S. B., Knigge C., Gänsicke B. T., Sion E. M., Welsh W. F., 2007, *ApJ*, **667**, 442
- Suleimanov V. F., Doroshenko V., Werner K., 2019, *MNRAS*, **482**, 3622
- Sulkanen M. E., Brasure L. W., Patterson J., 1981, *ApJ*, **244**, 579
- Szkody P., Silber A., 1996, *AJ*, **112**, 289
- Szkody P., et al., 2002, *AJ*, **123**, 430
- Szkody P., et al., 2003, *AJ*, **126**, 1499
- Szkody P., et al., 2004, *AJ*, **128**, 1882
- Szkody P., et al., 2005, *AJ*, **129**, 2386
- Szkody P., et al., 2006, *AJ*, **131**, 973
- Szkody P., et al., 2007, *AJ*, **134**, 185
- Szkody P., et al., 2009, *AJ*, **137**, 4011
- Szkody P., et al., 2010, *ApJ*, **710**, 64
- Szkody P., et al., 2011, *AJ*, **142**, 181
- Szkody P., Mukadam A. S., Sion E. M., Gänsicke B. T., Henden A., Townsley D., 2013, *AJ*, **145**, 121
- Taylor M. B., 2005, *TOPCAT & STIL: Starlink Table/VOTable Processing Software*. p. 29
- Thomas H.-C., Beuermann K., 1998, in Breitschwerdt D., Freyberg M. J., Truemper J., eds, *Lecture Notes in Physics*, Berlin Springer Verlag Vol. 506, IAU Colloq. 166: The Local Bubble and Beyond. pp 247–250, doi:10.1007/BFb0104728
- Thorstensen J. R., 2003, *AJ*, **126**, 3017

- Thorstensen J. R., Lépine S., Shara M., 2008, *AJ*, **136**, 2107
- Thorstensen J. R., Schwarz R., Schwobe A. D., Staude A., Vogel J., Krumpe M., Kohnert J., Nebot Gómez-Morán A., 2009, *PASP*, **121**, 465
- Thorstensen J. R., Alper E. H., Weil K. E., 2016, *AJ*, **152**, 226
- Toloza O., et al., 2016, *MNRAS*, **459**, 3929
- Tonry J. L., et al., 2018, *ApJ*, **867**, 105
- Toonen S., Hamers A., Portegies Zwart S., 2016, *Computational Astrophysics and Cosmology*, **3**, 6
- Tout C. A., Wickramasinghe D. T., Liebert J., Ferrario L., Pringle J. E., 2008, *MNRAS*, **387**, 897
- Townsley D. M., Bildsten L., 2003, *The Astrophysical Journal*, **596**, L227
- Townsley D. M., Bildsten L., 2004, *ApJ*, **600**, 390
- Townsley D. M., Gänsicke B. T., 2009, *The Astrophysical Journal*, **693**, 1007
- Unda-Sanzana E., et al., 2008, *MNRAS*, **388**, 889
- Wade R. A., Horne K., 1988, *ApJ*, **324**, 411
- Warner B., 1987, *MNRAS*, **227**, 23
- Warner B., 1995, Cambridge Astrophysics Series, **28**
- Watson C. L., Henden A. A., Price A., 2006, Society for Astronomical Sciences Annual Symposium, **25**, 47
- Wijnen T. P. G., Zorotovic M., Schreiber M. R., 2015, *A&A*, **577**, A143
- York D. G., et al., 2000, *AJ*, **120**, 1579
- Zharikov S. V., et al., 2008, *A&A*, **486**, 505
- Zharikov S., Tovmassian G., Aviles A., Michel R., Gonzalez-Buitrago D., García-Díaz M. T., 2013, *A&A*, **549**, A77
- Zorotovic M., Schreiber M. R., 2017, *MNRAS*, **466**, L63
- Zorotovic M., Schreiber M. R., Gänsicke B. T., 2011, *A&A*, **536**, A42
- Zorotovic M., et al., 2016, *MNRAS*, **457**, 3867
- de Kool M., 1992, *A&A*, **261**, 188
- van Spaandonk L., Steeghs D., Marsh T. R., Parsons S. G., 2010, *ApJ*, **715**, L109

APPENDIX A: FLARING RED DWARFS

12 objects in our sample are located on the main sequence or very close to it (empty and yellow diamonds in Figure 1). These objects are often classified as CV candidates because they have shown some transient phenomena that has been interpreted as a likely dwarf nova outburst. However, these systems are much redder than the typical CVs and it is therefore more likely that they are actually flaring red dwarfs. This is the case, for example, of MASTER OT J143453.02+023616.1 and MASTER OT J120525.84+621743.3: their SDSS spectra confirm their red dwarf nature (top and middle panel of Figure A1). The remaining 10 systems have similar colors and we conclude that they are also flaring red dwarfs.

APPENDIX B: YOUNG STELLAR OBJECTS

Three CV candidates, SSS J155929.1-223618 (Watson et al. 2006), Larin 2 (Larin et al. 2018; Denisenko & Larin 2018) and SSS J035055.8-204817 show particularly red colours but have also a *Galex* detection. Larin et al. (2018) suggested that Larin 2 could be a long period magnetic CVs with a large infrared contribution from the secondary. Later on Denisenko (2018) argued that wind-driven accretion is going on in this system, where the wind circularisation radius is smaller than the Roche lobe radius.




Figure A1. Sample systems reported as CV candidates in the literature. The SDSS spectra reveal that they are actually flaring red dwarfs (top and middle panels) and a quasar (bottom panel).

We acquired VLT/X-shooter and WHT/ISIS spectroscopic for these systems, as well as one SOAR spectrum for SSS J035055.8-204817, (Figure A2), where we detected the presence of the lithium absorption line at 6707 Å. Lithium is expected to be depleted once the core is hot enough for it to be burnt, and since low-mass stars at this stage are fully convective, its presence in the stellar photosphere suggests youth (Soderblom et al. 2014). Moreover, the forbidden lines of [O I] (5577 Å and 6300 Å) and [S II] (6730 Å) are commonly observed in young stellar objects (YSOs, Fang et al. 2018) but not in CVs. Finally, the metal emission lines are much narrower than the hydrogen lines and these different line widths suggest that the first arise from a hot corona close to the stellar surface while the seconds originate far-out in the magnetosphere (Hamann & Persson 1992). We therefore concluded that both systems are likely YSOs (brown diamonds in Figure 1).

Recently, Denisenko & Larin (2018) identified another CV candidate within 150 pc, DDE 158, owing to its colours and variability properties similar to those of Larin 2. Similarly to SSS J155929.1-223618, Larin 2 and SSS J035055.8-204817, the SOAR spectrum of DDE 158 presented in the vsnet-alert #23884 is characterised by metal emission lines narrower than the hydrogen ones, suggesting that they do not originate in a typical CV accretion disc. The non-detection of the forbidden [O I] (5577 Å) line does not allow to rule out a YSO (has suggested in the vsnet-alert #23884) since several YSOs in which this feature has not been detected (even in medium resolution spectra, see e.g. Natta et al. 2014; Nisini et al. 2018; Banzatti et al. 2019) are known. Finally, the red portion of the spectrum resembles the emission of a M-dwarf star. If DDE 158 was a CV, a secondary star of such spectral type would imply the presence of a white




Figure A2. VLT/X-shooter (top and middle panels), SOAR (grey, bottom panel) and WHT/ISIS (black, bottom panel) spectra of three YSOs that have been mistakenly classified as CVs (Watson et al. 2006; Larin et al. 2018). In all the three systems, the metal emission lines are much narrower than the hydrogen lines, suggesting that the first arise from a hot corona close to the stellar surface while the seconds originate far-out in the magnetosphere. The panels on the left show a zoom in the regions in which Li I and/or the forbidden lines of [O I] (6300 Å) and [S II] (6730 Å) are detected, which are all characteristic of YSOs.

dwarf as hot as $\gtrsim 15000$ K. Given the distance as derived from the *Gaia* parallax (≈ 103 pc), such white dwarf would contribute significantly to the overall system emission, causing the spectrum to be much bluer than actually observed. We can thus conclude that also DDE 158 is likely another YSO and therefore we do not include it in our sample.

APPENDIX C: OTHER NON-CV SYSTEMS

Three systems listed in the literature as CVs, SDSS J121929.46+471522.8 (Szkody et al. 2006), NSV 15401 (Downes et al. 2001) and Gaia14abg (Rixon et al. 2014), are actually single white dwarfs.

The SDSS spectrum of SDSS J121929.46+471522.8 shows a blue continuum and a weak emission line at $\lambda = 6539.4$ Å but no absorption and/or emission lines typical of a CV (Figure A3). This is likely a cold ($T_{\text{eff}} \approx 8000$ K) white dwarf whose atmosphere is dominated by helium, which explain the absence of absorption features. The origin of the emission is however unclear. Its wavelength seems to suggest that it arises from a O II transition. However, this is quite unlikely because (i) the formation of O II lines require much higher temperatures than the one we estimated for SDSS1219 and (ii) many other stronger O II lines should be detected in the spectrum. Contamination from a nearby fibre in the SDSS plate, centred onto a source showing strong emission in this wavelength is, more likely, the origin of the observed emission in SDSS1219.

The spectrum of NSV 15401 (aka Lan 159) presented by Lépine et al. (2011, see their figure 2) clearly show that this is a single white dwarf while Gaia14abg was among the first objects identified by the *Gaia* alerts and was affected by a cross-matching problem.

We also discarded five detached binaries (Ret1 REF, BPM 18764 REF), one pre-polar (WXL Mi), and MLS 101009:010045+010019, whose SDSS spectrum shows that this system is actually a quasar (bottom panel of Figure A1).

APPENDIX D: NEW CVS FROM *Gaia*

We observed *Gaia* J051903.99+630340.4 with the Intermediate-dispersion Spectrograph and Imaging System (ISIS) spectrograph mounted on the William Herschel Telescope in La Palma (Spain). We used a 1.2'' slit combined with the R600B and R600R gratings, centred at 4540 Å and 6561 Å and providing a nominal spectral resolution of $R \approx 2000$ and $R \approx 4000$, respectively. *Gaia* J051903.99+630340.4 shows a spectrum typical of a SUUMa star, dominated by a blue continuum and strong double-peaked Balmer and He I emission lines (top panel of Figure 2). Although this class of CVs is characterised by relatively short outburst recurrence times (of the order of months up to one year), it is possible that the dwarf nova outbursts of *Gaia* J051903.99+630340.4 have been missed given the presence of the much brighter nearby companion




Figure A3. Photometric SED (top) and SDSS spectrum (middle) of SDSS1219 along with a black body of $T_{\text{eff}} = 8000$ K. On the bottom panel is shown the spectrum of a galaxy observed by a nearby fiber on the same SDSS plate. This galaxy saturated the SDSS detector in the wavelength region (grey band, also highlighted in the middle panel) in which the emission lines is observed in SDSS1219 and it most likely the origin of this anomalous feature.




Figure A4. WHT/ISIS spectrum of TYC4084-172-1, a G3V star (Frasca et al. 2018), located 2.2 pc away from *Gaia* J051903.99+630340.4

(TYC 4084-172-1, $G = 9.6$ mag, Section 7). With the ISIS spectrograph, we also obtained phase resolved spectroscopic observations. From a fit to the position of the $H\alpha$ emission, we estimated a $P_{\text{orb}} \simeq 126$ min, consistent with the SU UMa classification.

We performed a spectroscopic follow-up of *Gaia* J154008.28–392917.6 using the Goodman spectrograph (Clemens et al. 2004) mounted on the Southern Astrophysical Research (SOAR) telescope in Cerro Pachón (Chile). We used a $1''$ slit and the red camera to acquired eight spectra of 300 s exposure each, using a 930 line/mm grating covering the wavelength range 3650 Å–5200 Å. The average spectrum is shown in the bottom panel of Figure 2. *Gaia* J154008.28–392917.6 resembles a typical low accreting system, with the signature of both the white dwarf (pressure broadened Balmer absorption lines) and the accretion disc (double-peaked Balmer emission lines) clearly detected in its spectrum. Given its spectral similarities with other CVs at the period minimum (see e.g. EZ Lyn in Figure 8), we classified *Gaia* J154008.28–392917.6 as WZ Sge-type CV likely located close (or even having already evolved through) the period minimum.

Table D1: CVs with unreliable *Gaia* parallaxes (Section 2.1 & 2.2) and objects within 150 pc that have been mistakenly identified as CVs in the literature. The flags are as follows: FRD, likely flaring red dwarf; SWD, single white dwarf; DB, detached binary; YSO, young stellar object; Q, quasar; AEN, astrometric excess noise greater than 2; SGD, spurious *Gaia* detection.

System	α	δ	ϖ (mas)	σ_{ϖ} (mas)	Astrometric excess noise	Flag
MASTER OT J015119.13–643046.6	01:51:19.39	–64:30:45.18	25.1	0.3	1.15	FRD
MASTER OT J031121.54–601851.0	03:11:21.43	–60:18:50.24	19.07	0.13	0.96	AEN
OGLE-BLG-DN-0128	17:47:29.66	–34:42:44.55	18	1	2.12	AEN
N SMC 2012	00:32:55.06	–74:20:19.7	14.60	0.09	0.50	SGD
SDSS J121929.46+471522.8	12:19:29.32	+47:15:22.89	14.3	0.1	0.46	SWD
CSS 131106:032129+180827	03:21:28.62	+18:08:27.05	12.8	0.5	1.64	FRD
MASTER OT J020836.79–104018.8	02:08:36.74	–10:40:18.79	12.2	0.4	1.58	FRD
MASTER OT J194753.58–475722.9	19:47:53.60	–47:57:23.14	12	2	10.18	AEN
MASTER OT J072448.87+533952.1	07:24:48.86	+53:39:51.49	10.6	0.1	0.42	FRD
Gaia16bvf	19:25:17.71	+08:39:20.40	10	2	6.69	AEN
V1454 Cyg	19:53:38.50	+35:21:45.62	10.21	1.79	6.8	AEN
WX LMi	10:26:27.52	+38:45:02.01	10.1	0.1	0.22	DB
Gaia14abg	17:30:47.93	+50:00:16.65	9.5	0.1	0.0	SWD
Ret1	03:34:34.43	–64:00:57.88	9.35	0.02	0.14	DB
BPM 18764	08:02:00.41	–53:27:49.36	9.2	0.1	0.3	DB
OGLE-BLG-DN-0040	17:34:23.99	–23:32:44.43	9	1	2.92	SGD
ASASSN-14ib	04:22:12.22	–03:25:13.47	8.98	3.25	2.89	AEN
MASTER OT J143453.02+023616.1	14:34:53.08	+02:36:16.15	8.4	0.6	0.0	FRD
NSV 15401	01:55:10.12	+69:42:40.14	8.32	0.08	0.13	SWD
SBS 1316+577A	13:18:00.68	+57:28:04.00	8.23	0.05	0.36	FRD
SSS J035055.8–204817	03:50:56.02	–20:48:15.95	7.98	0.06	0.0	YSO
Larin 2	12:48:50.77	–41:26:54.65	7.95	0.13	0.32	YSO
Gaia17cuc	10:26:20.55	–44:18:49.68	7.8	0.1	0.39	FRD
ASASSN-17eo	20:02:14.34	+31:36:34.66	8	1	3.17	AEN
SSS J162131.9–230140	16:21:31.92	–23:01:40.73	7.3	0.1	0.59	FRD
SSS J155929.1–223618	15:59:29.20	–22:36:17.82	7.1	0.1	0.34	YSO
MASTER OT J100950.32+471815.8	10:09:50.23	+47:18:16.76	7.0	0.3	0.68	FRD
SSS J155147.2–211323	15:51:47.08	–21:13:23.86	6.89	0.13	0.61	FRD
OGLE-GD-ECL-02234	10:45:49.79	–61:29:57.04	6.79	0.08	0.53	FRD
ASASSN-15ep	08:21:06.24	–72:20:12.09	6.2	0.1	0.71	FRD
OGLE-BLG-DN-0428	18:00:11.42	–29:41:38.40	6	1	2.78	SGD
MASTER OT J012916.47+321859.0	01:29:16.48	+32:18:58.84	6	1	2.46	FRD, AEN
NSV 18024	08:44:35.16	–37:58:02.84	6.1	0.9	7.71	AEN
MACHO 401.48296.2600	17:58:32.38	–27:52:44.12	6	1	2.47	AEN
Gaia17brd	20:15:08.25	+20:40:31.19	6	4	3.4	AEN
MASTER OT J140957.49+290922.7	14:09:57.47	+29:09:22.79	5.8	0.6	0.0	FRD
V1419 Aql	19:13:06.79	+01:34:23.24	6	2	5.41	AEN
MASTER OT J120525.84+621743.3	12:05:25.88	+62:17:43.04	5.7	0.5	2.23	FRD, AEN
MASTER OT J020404.19+741804.6	02:04:03.60	+74:18:02.45	5.6	1.3	4.56	AEN
Gaia17cva	19:45:37.72	+28:05:32.88	6	2	1.46	SGD
OGLE-BLG-DN-0266	17:54:53.98	–21:22:40.19	5	2	3.86	AEN
Gaia17aoi	13:24:44.33	–14:23:35.65	5	1	2.45	AEN
OGLE-BLG-DN-0824	18:10:04.90	–29:05:23.58	5	1	3.61	SGD, AEN
CSS 111021:220328+141059	22:03:28.11	+14:11:00.49	4	1	0.0	SGD
OGLE-BLG-DN-0011	17:17:26.01	–28:33:23.79	4	1	4.47	SGD, AEN
MASTER OT J203824.15+174242.3	20:38:24.10	+17:42:43.15	4	2	2.98	AEN
OGLE-BLG-DN-0087	17:43:07.76	–34:19:28.10	3	1	3.2	SGD, AEN
OGLE-BLG-DN-0156	17:49:49.23	–21:22:13.57	3	2	5.32	SGD, AEN
CSS 150422:172535+231215	17:25:34.90	+23:12:14.33	3	3	5.14	AEN
MASTER OT J051042.59+513540.0	05:10:42.60	+51:35:39.82	3	1	3.14	AEN
CSS 081201:213947+170658	21:39:47.16	+17:06:56.53	3	2	1.65	SGD
Gaia16bfi	16:37:08.62	–67:36:56.46	5.1	1.9	4.70	AEN
ASASSN-13cv	22:10:25.35	+30:46:10.06	4.5	0.8	1.19	FRD
OGLE-BLG-DN-0584	18:03:46.14	–27:15:33.70	4.2	0.8	1.74	FRD

Table D1 – continued from previous page

System	α	δ	ϖ (mas)	σ_{ϖ} (mas)	Astrometric excess noise	Flag
ASASSN-17gc	19:51:36.94	-00:59:04.05	4	2	0.71	SGD
ASASSN-16cd	19:06:38.16	+33:09:03.17	3.4	1.5	5.5	SGD, AEN
ASASSN-17nm	09:46:09.11	-57:14:20.41	3.1	1.2	0.0	SGD
Gaia17aok	21:52:55.72	59:18:18.71	3.1	1.2	0.0	SGD
CSS 120313:131043-042600	13:10:42.68	-04:26:00.73	3	2	2.2	AEN
OGLE-BLG-DN-0216	17:52:24.02	-32:16:51.54	2.2	1.5	2.7	SGD, AEN
EL Aql	18:56:01.87	-03:19:18.80	2	3	2.68	AEN
MASTER OT J182201.93+324906.7	18:22:01.80	+32:49:00.57	2.0	2.0	4.85	SGD, AEN
MLS 101009:010045+010019	01:00:44.71	+01:00:18.48	2	2	1.53	Q
EU Cnc	08:51:27.17	+11:46:56.94	2	2	0.0	SGD
OGLE-BLG-DN-0181	17:51:01.18	-29:14:38.30	2	2	2.82	AEN
Gaia17afs	17:35:17.86	+01:32:48.79	1.1	1.9	3.0	SGD, AEN
OGLE-BLG-DN-0054	17:38:24.00	-21:54:26.87	2	2	2.13	AEN
DO Vul	19:52:10.72	+19:34:42.14	0.8	2	4.78	AEN
Gaia17bqf	19:59:19.93	+16:24:40.31	-0.2	2	3.07	AEN
V1722 Aql	19:14:09.74	+15:16:38.25	-1	3	3.07	SGD, AEN

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.

Table D2. CV and CV candidates with parallaxes $\varpi + 3\sigma_{\varpi} \geq 6.66$ mas that are located further than 150 pc.

System	α	δ	ϖ (mas)	σ_{ϖ} (mas)	Distance (pc)	$P(d < 150 \text{ pc})$ (%)
V1108 Her	18:39:26.14	+26:04:09.96	6.6	0.1	152 ± 3	0.24
EK TrA	15:14:00.10	-65:05:36.65	6.58	0.04	152 ± 1	0.02
BZ UMa	08:53:44.22	+57:48:40.35	6.56	0.06	153 ± 1	0.04
EI Psc	23:29:54.17	+06:28:12.11	6.55	0.07	153 ± 2	0.04
MASTER OT J050806.84+712352.0	05:08:06.78	+71:23:51.69	6.5	2.8	487 ⁺⁴⁸⁹ ₋₂₅₅	0.04
FL Psc	00:25:11.04	+12:17:11.81	6.5	0.1	154 ± 3	0.11
V348 Pav	19:56:48.05	-60:34:30.00	6.48	0.08	154 ± 2	0.006
ASASSN-16jg	14:45:35.72	-39:20:26.73	6.4	0.3	158 ⁺⁹ ₋₈	0.17
1RXS J083842.1-282723	08:38:43.33	-28:27:00.95	6.4	0.1	157 ± 3	0.004
SDSS J150551.58+065948.7	15:05:51.61	+06:59:48.49	6.3	0.4	162 ⁺¹³ ₋₁₁	0.14
EF Eri	03:14:13.41	-22:35:43.77	6.3	0.3	161 ± 7	0.05
MASTER OT J220559.40-341434.9	22:05:59.47	-34:14:34.15	6.2	0.3	161 ± 7	0.04
CSS 081221:050716+125314	05:07:16.24	+12:53:14.16	5.4	2.3	485 ⁺⁴⁶⁷ ₋₂₃₇	0.02
ASASSN-18bh	01:09:52.87	+47:57:11.11	5.2	1.7	361 ⁺³⁵⁴ ₋₁₄₆	0.02
ASASSN-14ip	20:50:23.43	-48:37:13.71	5.0	1.1	253 ⁺¹¹⁵ ₋₆₀	0.01
V498 Hya	08:45:55.06	+03:39:29.28	4.9	2.0	469 ⁺⁴⁴⁶ ₋₂₁₇	0.01
ASASSN-15td	12:15:13.70	-01:46:41.56	4.9	2.3	527 ⁺⁴⁷⁸ ₋₂₅₅	0.01
OGLE-BLG-DN-0183	17:51:05.70	-28:03:37.84	4.8	1.8	440 ⁺⁴¹⁶ ₋₁₉₃	0.01
ASASSN-16ee	08:35:42.43	-31:21:47.92	4.5	1.3	335 ⁺²⁵⁴ ₋₁₁₀	0.006
CSS 170417:080539+354055	08:05:38.98	+35:40:54.94	4.3	1.7	461 ⁺⁴¹³ ₋₁₉₆	0.006
ASASSN-13bd	23:59:58.00	-12:54:32.68	4.3	2.2	575 ⁺⁴⁸⁵ ₋₂₇₁	0.007
SDSS J125641.29-015852.0	12:56:41.29	-01:58:51.74	4.0	1.1	350 ⁺²⁰² ₋₉₆	0.0007
CSS 080927:212522-102627	21:25:21.76	-10:26:28.18	4.0	1.6	471 ⁺⁴⁰⁴ ₋₁₉₄	0.003
MASTER OT J122126.39-311248.3	12:21:26.40	-31:12:48.49	3.8	1.0	363 ⁺²¹⁷ ₋₁₀₄	0.0002
CSS 111103:074400+415504	07:44:00.47	+41:55:03.56	3.7	2.4	636 ⁺⁴⁹⁹ ₋₂₉₅	0.004
IK Leo	10:21:46.45	+23:49:25.91	3.6	1.4	499 ⁺⁴⁰² ₋₁₉₈	0.0008
ASASSN-15rj	02:59:38.35	+44:57:04.77	3.6	1.6	529 ⁺⁴²⁹ ₋₂₂₀	0.001
MASTER OT J070740.72+702630.0	07:07:40.55	+70:26:30.30	3.5	1.1	416 ⁺²⁸⁷ ₋₁₃₅	0.0002
ASASSN-15ef	16:49:40.59	-17:50:09.72	3.5	1.6	549 ⁺⁴⁴⁴ ₋₂₃₄	0.001
CSS 110124:032934+182530	03:29:33.92	+18:25:29.57	3.5	1.1	406 ⁺²⁶⁵ ₋₁₂₆	0.0001
ASASSN-17jf	20:29:17.10	-43:40:19.18	3.5	1.1	426 ⁺²⁹⁷ ₋₁₄₀	0.0001
ASASSN-17mw	02:49:07.33	+48:51:01.16	3.5	1.1	408 ⁺²⁶⁵ ₋₁₂₇	0.00009
Gaia16apf	00:34:33.39	+54:28:42.04	3.5	1.3	488 ⁺³⁸¹ ₋₁₈₆	0.0004
ASASSN-15gm	19:37:13.59	-22:57:06.31	3.5	1.6	544 ⁺⁴³⁵ ₋₂₂₆	0.0009
ASASSN-16jb	17:50:44.96	-25:58:37.45	3.42	1.2	456 ⁺³³⁴ ₋₁₆₀	0.0002
ASASSN-16do	06:34:12.71	-32:59:49.49	3.1	1.3	525 ⁺³⁹² ₋₁₉₈	0.0001
ASASSN-13ck	00:11:33.73	+04:51:22.43	3.1	1.6	590 ⁺⁴⁵² ₋₂₄₇	0.0005
ASASSN-15px	23:08:57.87	-65:59:32.49	3.1	1.3	547 ⁺⁴¹³ ₋₂₁₄	0.0002
CSS 110406:152159+261223	15:21:58.84	+26:12:23.30	2.7	2.1	689 ⁺⁴⁹⁹ ₋₃₀₃	0.001
CSS 101108:022436+372021	02:24:36.44	+37:20:21.40	2.7	1.8	671 ⁺⁴⁸⁵ ₋₂₈₇	0.0005
KK Cnc	08:07:14.25	+11:38:12.32	2.5	1.6	670 ⁺⁴⁷⁶ ₋₂₇₈	0.0002
ASASSN-16jk	15:40:24.84	+23:07:50.86	2.5	1.4	655 ⁺⁴⁵⁷ ₋₂₆₀	0.00004
MASTER OT J152701.21-314433.6	15:27:01.25	-31:44:35.30	2.5	1.5	667 ⁺⁴⁶⁸ ₋₂₇₀	0.00008
ASASSN-15aw	01:57:46.15	+51:10:23.88	2.4	1.7	695 ⁺⁴⁸⁷ ₋₂₉₂	0.0002
CSS 100108:081031+002429	08:10:30.61	+00:24:28.32	2.4	2.2	716 ⁺⁵⁰⁴ ₋₃₁₂	0.0008
ASASSN-15nf	20:12:42.71	+15:44:44.92	2.4	2.4	724 ⁺⁵⁰⁹ ₋₃₁₉	0.001
CSS 110430:091710+314309	09:17:09.87	+31:43:07.59	2.3	1.5	686 ⁺⁴⁷² ₋₃₇₆	0.00005
ASASSN-17bi	02:16:05.42	+68:39:03.61	2.3	1.5	694 ⁺⁴⁷⁷ ₋₂₈₂	0.00007
CSS 090928:030141+241541	03:01:40.52	+24:15:41.35	2.3	2.5	732 ⁺⁵¹¹ ₋₃₂₃	0.002
ASASSN-14kk	01:32:02.78	-10:43:57.72	2.3	1.5	689 ⁺⁴⁷¹ ₋₂₇₆	0.00004
MASTER OT J211855.08+280314.9	21:18:55.10	+28:03:15.39	2.2	3.1	744 ⁺⁵²⁰ ₋₃₃₅	0.002
SSS 110125:103550-424610	10:35:49.64	-42:46:10.14	2.1	2.2	741 ⁺⁵⁰⁸ ₋₃₂₀	0.0006
CSS 150822:232026+221833	23:20:26.23	+22:18:34.05	0.3	3.0	814 ⁺⁵²³ ₋₃₄₆	0.0007
CSS 090926:230711+294010	23:07:11.34	+29:40:11.33	0.2	2.2	857 ⁺⁵¹⁹ ₋₃₄₄	0.00007

Table D3. CVs with pre-*Gaia* distance estimates $d \leq 150$ pc.

System	<i>Gaia</i> DR2 ID	P_{orb} (min)	Distance (pc)	Method	Reference	<i>Gaia</i>					Distance (pc)
						G (mag)	G_{BP} (mag)	G_{RP} (mag)	ϖ (mas)	σ_{ϖ} (mas)	
WZ Sge	1809844934461976832	81.63	43.30 ^{+1.60} _{-1.50}	a	1	15.21	15.21	15.06	22.16	0.04	45.13±0.08
AY Lyr	2096934223687181696	105.55	52	c	2	17.93	17.96	17.50	2.22	0.13	452 ⁺³² ₋₂₂
XZ Eri	5097770801875122432	88.7	66	c	2	19.25	19.28	19.00	3.0	0.3	331 ⁺⁴⁴ ₋₂₄
VW Hyi	4653893040002306432	106.95	64 ⁺²⁰ ₋₁₇	e	3	13.84	13.94	14.45	18.53	0.02	53.96±0.06
EX Hya	6185040879503491584	98.26	64.5±1.2	a	4	13.21	13.23	12.88	17.56	0.04	56.95±0.13
GD 552	2208124536065383424	102.73	74±4	e	5	16.46	16.46	16.18	12.35	0.05	81.0±0.3
QZ Vir	3800596876396315648	84.70	76	c	2	16.06	16.12	15.76	7.81	0.07	128 ± 1
AM Her	2123837555230207744	185.65	79 ⁺⁸ ₋₆	a	1	13.58	13.86	12.85	11.40	0.02	87.76±0.14
1RXS J105010.3-140431	3750072904055666176	88.56	80±20	e	5	17.17	17.21	17.08	9.14	0.11	109±1
AR UMa	783921244796958208	115.92	86 ⁺¹⁰ ₋₈	a	6	16.26	16.35	15.78	9.87	0.12	101±1
V455 And	1920126431748251776	81.08	90±15	b	7	16.06	16.13	15.71	13.24	0.06	75.5±0.3
ASASSN-14ag	3071240270519385856	86.85	90	e	8	16.18	16.17	15.70	5.63	0.09	178±3
IX Vel*	5515820034889610112	279.25	96±1	a	9	9.32	9.34	9.27	11.04	0.03	90.6±0.2
VY Aqr	6896767366186700416	90.85	97 ⁺¹⁵ ₋₁₂	a	1	16.86	16.96	16.44	7.24	0.14	138±3
U Gem	674214551557961984	254.74	97±7	c	10	13.91	14.38	13.11	10.71	0.03	93.3±0.3
IGR J18308-1232	4153024090088033280	–	100	c	11	17.65	18.07	16.89	0.49	0.15	1595 ⁺⁵¹⁴ ₋₁₉₃
V426 Oph	4471872295941149056	410.83	100	e	12	12.37	12.65	11.77	5.20	0.04	192.5±1.5
WW Cet	2427474150870397056	253.15	100	c	2	13.55	13.65	12.96	4.59	0.05	218±2
OY Car	5242787486412627072	90.89	100	c	13	15.62	15.64	15.21	11.01	0.03	90.8±0.2
V2051 Oph	4111991385628196224	89.90	102 ±16	e	5	15.37	15.46	14.87	8.90	0.07	112.4±0.9
AE Aqr*	4226332451596335616	592.78	102 ⁺⁴² ₋₂₃	a	1	10.95	11.47	10.26	10.97	0.06	91.2±0.5
GW Lib	6226943645600487552	76.78	104 ⁺³⁰ ₋₂₀	a	1	16.49	16.49	16.32	8.87	0.08	113±1
DI UMa	1013298268207936128	78.59	107	c	2	17.75	16.86	16.82	1.46	0.08	685 ⁺⁴³ ₋₃₁
V3885 Sgr*	6688624794231054976	298.31	110±30	a	14	10.25	10.28	10.16	7.54	0.08	133±1
HU Aqr	6911950900211768704	125.02	111	c	2	16.47	16.66	15.88	5.20	0.06	192±2
Z Cha	5210507882302442368	107.28	112±8	c	10	15.85	15.94	15.19	8.66	0.12	115±1
AH Eri	3176908972944418816	344.30	113	c	2	17.46	17.85	16.75	0.82	0.09	1129 ⁺¹³⁴ ₋₈₀
EF Eri	5099482805904892288	81.02	113 ⁺¹⁹ ₋₁₆	a	1	18.21	18.17	18.11	6.3	0.3	161 ± 7
BK Lyn	702296666944246784	107.97	114	c	2	14.52	14.48	14.45	1.98	0.07	505 ⁺²⁰ ₋₁₆
SS Cyg*	1972957892448494592	396.19	114±2	a	15	11.69	12.11	10.95	8.72	0.05	114.6±0.6
ST LMi	3996419759863758592	113.89	115 ⁺²¹ ₋₂₂	b	16	16.13	–	–	8.83	0.08	113±1
DH Aql	4200218019655998720	–	116	c	2	17.97	18.28	17.43	3.6	0.2	282 ⁺¹⁸ ₋₁₃
IQ Eri	5078976609103251456	–	116 ⁺¹¹⁶ ₋₅₈	e	3	17.18	17.11	16.85	5.19	0.17	193 ⁺⁷ ₋₆
IP Peg	2824150286583562496	227.82	124	e	12	14.71	15.27	13.88	7.08	0.05	141±1
VV Pup	5719598950133755392	100.44	124 ⁺¹⁷ ₋₁₄	a	6	15.93	16.04	15.48	7.30	0.05	137±0.9
MR Ser	1203639265875666304	113.47	126 ⁺¹⁴ ₋₁₂	a	6	16.23	16.47	15.64	7.59	0.05	131.8±0.9
TV Col	2901783160488793728	329.18	128±1	a	17	13.93	14.03	13.61	1.95	0.02	512±4
IGR J17195-4100	5959894875620104064	240.34	130	c	11	15.33	15.55	14.75	1.53	0.04	653 ⁺³⁰ ₋₂₇
EQ Cet	5041907811522399488	92.82	130	c	18	17.34	17.41	16.78	3.51	0.11	285 ⁺¹⁰ ₋₈
BL Hyi	4697621824327141248	113.64	130	c	19	17.25	17.45	16.70	7.65	0.07	131 ± 1
BW Scl	2307289214897332480	78.23	131±18	b	20	16.26	16.26	16.10	10.60	0.10	94.4±0.9
HT Cas	426306363477869696	106.05	131 ⁺²² ₋₁₇	a	8	16.35	16.48	15.80	7.07	0.06	141 ± 1
RX And	374510294830244992	302.24	135	e	12	13.17	13.38	12.50	5.03	0.05	199±2
V406 Vir	3681313024562519552	80.52	140±35	e	5	17.72	17.71	17.55	5.91	0.16	169±5
V834 Cen	6096905573613586944	101.52	144 ⁺¹⁸ ₋₂₃	b	16	16.66	16.82	16.07	8.9	0.2	113±3
V405 Peg	2838503311371673472	255.81	149	a	21	15.28	16.05	14.29	5.78	0.06	173±2
V2301 Oph	4476137370261520000	112.97	150	b	22	16.75	16.94	15.86	8.24	0.08	121±1
SX LMi	733329416268149376	96.72	150	c	2	16.69	16.77	16.30	3.08	0.12	325 ⁺¹⁴ ₋₁₁
CU Vel	5524430207364715520	113.04	150±50	b	23	16.71	16.93	16.15	6.29	0.06	159±2

Notes. Systems highlighted with a star have a parallax measurement from *Gaia* DR1 (Ramsay et al. 2017) but we do not report them in this table since they do not represent a pre-*Gaia* distance determination. The method abbreviations recall the list in Section 4. The distances reported in the last column have been determined from the *Gaia* parallaxes as described in Section 3.

References: (1) Thorstensen (2003), (2) Sproats et al. (1996), (3) Pretorius & Knigge (2012), (4) Beuermann et al. (2003), (5) Patterson (2011), (6) Thorstensen et al. (2008), (7) Araujo-Betancor et al. (2005a), (8) Thorstensen et al. (2016), (9) Linnell et al. (2007), (10) Beuermann (2006), (11) Bernardini et al. (2012), (12) Warner (1987), (13) Sherrington et al. (1982), (14) Linnell et al. (2009), (15) Miller-Jones et al. (2013), (16) Araujo-Betancor et al. (2005b), (17) McArthur et al. (2001), (18) Schwobe et al. (1999), (19) Beuermann et al. (1985), (20) Gänsicke et al. (2005), (21) Thorstensen et al. (2009), (22) Szkody & Silber (1996), (23) Gänsicke & Koester (1999).