



University of Dundee

Transition to quantitative faecal immunochemical testing from guaiac faecal occult blood testing in a fully rolled-out population-based national bowel screening programme

Clark, Gavin R. C.; Strachan, Judith; Carey, Francis; Godfrey, Thomas G.; Irvine, Audrey; McPherson, Alisson

Published in:
Gut

DOI:
[10.1136/gutjnl-2019-320297](https://doi.org/10.1136/gutjnl-2019-320297)

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Clark, G. R. C., Strachan, J., Carey, F., Godfrey, T. G., Irvine, A., McPherson, A., Brand, J., Anderson, A. S., Fraser, C., & Steele, R. J. C. (2020). Transition to quantitative faecal immunochemical testing from guaiac faecal occult blood testing in a fully rolled-out population-based national bowel screening programme. *Gut*. <https://doi.org/10.1136/gutjnl-2019-320297>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Supplementary Tables

Supplementary Table 1. Uptake by age (years), sex, socio-economic deprivation

(Scottish Index of Multiple Deprivation: SIMD) and screening history

	FOBT (95% CI)	FIT (95% CI)
Uptake		
Total	56.4% (56.3%-56.5%)	63.9% (63.8%-64%)
Age group (years)		
50-54	47.3% (47.1%-47.5%)	57.5% (57.4%-57.7%)
55-59	53.8% (53.5%-54.0%)	60.3% (60.1%-60.5%)
60-64	59.7% (59.5%-59.9%)	66.1% (65.9%-66.3%)
65-69	66.2% (65.9%-66.4%)	71.3% (71.1%-71.6%)
70-74	63% (62.7%-63.2%)	69.6% (69.3%-69.8%)
Sex		
Male	53.2% (53.1%-53.4%)	61.6% (61.4%-61.7%)
Female	59.6% (59.4%-59.7%)	66.1% (66%-66.3%)
SIMD		
1 - most deprived	43.3% (43.1%-43.6%)	51.7% (51.4%-51.9%)
2	51.4% (51.2%-51.7%)	59.3% (59.1%-59.5%)
3	57.7% (57.5%-58%)	65.2% (65.0%-65.4%)
4	62.4% (62.2%-62.6%)	69.3% (69.1%-69.5%)
5 - least deprived	66.1% (65.8%-66.3%)	72.5% (72.3%-72.7%)
Screening history		
First round	45.8% (45.5%-46.1%)	59.1% (58.8%-59.5%)
Participated in previous round	87.5% (87.4%-87.6%)	92.3% (92.3%-92.4%)
Did not participate in previous round	38.6% (38.2%-39%)	52.9% (52.5%-53.2%)
Never participated	8.8% (8.7%-9%)	18.9% (18.8%-19.1%)

Supplementary Table 2. Positivity of Faecal Occult Blood Test (FOBT) and Faecal Immunochemical Test (FIT) by age (years), sex and socio-economic deprivation (Scottish Index of Multiple Deprivation: SIMD)

	FOBT (95% CI)	FIT (95% CI)
Positivity		
Total	2.2% (2.1%-2.2%)	3.1% (3%-3.1%)
Age group (years)		
50-54	1.6% (1.6%-1.7%)	2.3% (2.2%-2.4%)
55-59	1.8% (1.7%-1.9%)	2.7% (2.6%-2.8%)
60-64	1.8% (1.7%-1.9%)	3.1% (3%-3.2%)
65-69	2.9% (2.8%-3%)	3.7% (3.5%-3.8%)
70-74	3% (2.9%-3.1%)	4% (3.9%-4.2%)
Sex		
Male	2.6% (2.5%-2.7%)	3.6% (3.5%-3.7%)
Female	1.8% (1.7%-1.8%)	2.6% (2.5%-2.6%)
SIMD		
1 - most deprived	3.2% (3.1%-3.4%)	4.3% (4.1%-4.4%)
2	2.6% (2.5%-2.7%)	3.5% (3.4%-3.6%)
3	2.1% (2.0%-2.2%)	3.0% (3.0%-3.1%)
4	1.8% (1.7%-1.9%)	2.7% (2.6%-2.7%)
5 - least deprived	1.5% (1.5%-1.6%)	2.4% (2.3%-2.5%)

Supplementary Table 3. Colorectal cancer (CRC) positive predictive value (PPV) by age (years), sex and socio-economic deprivation (Scottish Index of Multiple Deprivation: SIMD)

	FOBT (95% CI)	FIT (95% CI)
CRC PPV		
Total	6.4% (5.9%-6.9%)	5.2% (4.8%-5.5%)
Age group (years)		
50-54	4.4% (3.5%-5.6%)	3.1% (2.5%-3.8%)
55-59	5.2% (4.2%-6.5%)	3.9% (3.2%-4.8%)
60-64	6.4% (5.3%-7.8%)	5% (4.3%-5.8%)
65-69	6.7% (5.7%-7.9%)	6.4% (5.5%-7.4%)
70-74	8.6% (7.4%-10.1%)	7.0% (6.2%-8.0%)
Sex		
Male	6.8% (6.1%-7.5%)	5.3% (4.8%-5.8%)
Female	5.9% (5.2%-6.7%)	5.0% (4.4%-5.5%)
SIMD		
1 - most deprived	5.3% (4.3%-6.4%)	3.8% (3.1%-4.6%)
2	5.5% (4.6%-6.7%)	4.7% (4%-5.6%)
3	6.0% (5.0%-7.3%)	5.3% (4.6%-6.2%)
4	6.5% (5.4%-7.8%)	6.0% (5.2%-6.9%)
5 - least deprived	9.1% (7.7%-10.7%)	6.0% (5.2%-7.0%)

Supplementary Table 4. Higher-risk (HR) adenoma positive predictive value (PPV)

by age (years), sex and socio-economic deprivation (Scottish Index of Multiple

Deprivation: SIMD)

	FOBT (95% CI)	FIT (95% CI)
HR adenoma PPV		
Total	19.3% (18.5%-20.2%)	24.3% (23.6%-25.2%)
Age group (years)		
50-54	16.3% (14.5%-18.3%)	20.0% (18.5%-21.6%)
55-59	18.1% (16.2%-20.1%)	22.6% (20.9%-24.3%)
60-64	18.5% (16.6%-20.5%)	24.8% (23.4%-26.3%)
65-69	20.7% (19%-22.5%)	27.5% (25.8%-29.2%)
70-74	22.3% (20.3%-24.3%)	26.2% (24.6%-27.8%)
Sex		
Male	23.1% (21.9%-24.3%)	28.8% (27.8%-29.8%)
Female	14.6% (13.5%-15.8%)	18.6% (17.6%-19.6%)
SIMD		
1 - most deprived	19.7% (17.8%-21.6%)	23.5% (21.9%-25.1%)
2	16.7% (15.1%-18.5%)	24.7% (23.2%-26.4%)
3	19.3% (17.5%-21.3%)	24% (22.5%-25.6%)
4	20.2% (18.4%-22.2%)	25.1% (23.5%-26.7%)
5 - least deprived	21.3% (19.2%-23.5%)	24.2% (22.6%-25.9%)