

Northumbria Research Link

Citation: El, Gennady, Gammal, A. and Kamchatnov, A.M. (2007) Generation of oblique dark solitons in supersonic flow of Bose-Einstein condensate past an obstacle. Nuclear Physics A, 790 (1-4). 771c-775c. ISSN 0375-9474

Published by: Elsevier

URL: <https://doi.org/10.1016/j.nuclphysa.2007.03.024>
<<https://doi.org/10.1016/j.nuclphysa.2007.03.024>>

This version was downloaded from Northumbria Research Link: <http://nrl.northumbria.ac.uk/42788/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



UniversityLibrary



Northumbria
University
NEWCASTLE

Generation of oblique dark solitons in supersonic flow of Bose-Einstein condensate past an obstacle

G.A. El^a, A. Gammal^b, and A.M. Kamchatnov^c

^aDepartment of Mathematical Sciences, Loughborough University, Loughborough LE11 3TU, UK

^bInstituto de Física, Universidade de São Paulo, 05315-970, C.P.66318 São Paulo, Brazil

^cInstitute of Spectroscopy, Russian Academy of Sciences, Troitsk, Moscow Region, 142190, Russia

1. Introduction

Nonlinear and dispersive properties of Bose-Einstein condensate (BEC) provide a possibility of formation of various nonlinear structures such as vortices and bright and dark solitons (see, e.g., [1]). Yet another type of nonlinear wave patterns has been observed in a series of experiments on the BEC flow past macroscopic obstacles [2]. In [3] these structures have been associated with spatial dispersive shock waves. Spatial dispersive shock waves represent dispersive analogs of the well-known viscous spatial shocks (oblique jumps of compression) occurring in supersonic flows of compressible fluids past obstacles. In a viscous fluid, the shock can be represented as a narrow region within which strong dissipation processes take place and the thermodynamic parameters of the flow undergo sharp change. On the contrary, if viscosity is negligibly small compared with dispersion effects, the shock discontinuity resolves into an expanding in space oscillatory structure which transforms gradually, as the distance from the obstacle increases, into a “fan” of stationary solitons. If the obstacle is small enough, then such a “fan” reduces to a single spatial dark soliton [4]. Here we shall present the theory of these new structures in BEC.

2. Generation of spatial dark solitons in an expanding BEC

In the mean-field approximation, dynamics of BEC is described very well by the Gross-Pitaevskii (GP) equation [1]

$$i\frac{\partial\psi}{\partial t} = -\frac{1}{2}\Delta\psi + V(\mathbf{r})\psi + |\psi|^2\psi, \quad (1)$$

which is written in standard non-dimensional units (see, e.g., [3]). Here $\psi(\mathbf{r})$ denotes the condensate order parameter and $V(\mathbf{r})$ is the potential of the external forces acting on the condensate, as, e.g., the confining potential of the trap and/or the potential of an obstacle inside the BEC. In [2], the flow arising due to free expansion of a BEC after its release from the trap was actually two-dimensional, $\mathbf{r} = (x, y)$, and radially symmetrical. To illustrate




Figure 1. Generation of a pair of spatial dark solitons by an impenetrable disc of radius $r = 1$ placed at $(0,0)$ in a BEC radially expanding from the center at $(-25,0)$. The density plot is shown at $t = 12$. Initial radius of the condensate $R = 25$.

the process of generation of waves by the obstacle, we have studied the time-dependent numerical solution of the GP equation (1) where $V(\mathbf{r})$ corresponds to interaction of the obstacle with the condensate. In our simulations, the flow was modeled by a free expansion of a BEC after its release from strongly confining trap, and the obstacle was modeled by an impenetrable disk with radius $r = 1$. The results of these simulations are shown in Fig. 1 where a pair of spatial dark solitons is clearly seen behind the obstacle. It is clear that radial symmetry is not essential for a phenomenon of generation of such solitons. In fact, in real experiments the flow near the obstacle can be approximated by a uniform flow with constant velocity M where in our notation M is the Mach number. The conditions for good enough accuracy of such an approximation can be found as follows.

It is convenient to temporarily return to dimensional units. To be definite, we shall consider here free expansion of a BEC in a pancake geometry, that is the condensate is supposed to be confined in axial direction by a strong potential, so that its axial size is given by $a_{\parallel} = \sqrt{\hbar/m\omega_{\parallel}}$, ω_{\parallel} being the frequency of the harmonic potential in the axial direction. Let the transverse frequency ω_{\perp} be small enough so that the Thomas-Fermi approximation is applied to the distribution of BEC in radial (transverse) direction. Then, before switching off the transverse potential the stationary radial density distribution is given by

$$n(r) = C (1 - r^2/l^2), \quad C = m\omega_{\perp}^2 l^2 / 2g_{2D}, \quad (2)$$

where $l = \left((16/\sqrt{2\pi}) N a_{\parallel}^3 a_s \omega_{\parallel}^2 / \omega_{\perp}^2 \right)^{1/4}$ is the radius of BEC before its release from the trap, and $g_{2D} = g/\sqrt{2\pi}a_{\parallel}$ is the effective nonlinearity constant in the pancake geometry.

An explicit time-dependent solution of the cylindrical GP equation for the condensate freely expanding in radial direction from the initial equilibrium state (2) is given by (see,

e.g., [5])

$$n(r, t) = \frac{C}{1 + \omega_{\perp}^2 t^2} \left[1 - \frac{r^2}{l^2(1 + \omega_{\perp}^2 t^2)} \right], \quad u(r, t) = \frac{\omega_{\perp}^2 tr}{1 + \omega_{\perp}^2 t^2}. \quad (3)$$

Now, let us consider the regions of BEC not too far from the axis, $r \ll l\omega t$, for $t \gg \omega_{\perp}^{-1}$. Then in these approximations we obtain

$$n(r, t) \cong C/\omega^2 t^2, \quad u(r, t) \cong r/t, \quad (4)$$

that is the density is practically uniform but varying with time. In the same approximation the local healing length ξ and the local sound velocity c_s are given by $\xi = \hbar t/ml$, $c_s = l/\sqrt{2}t$. Hence, the local Mach number $M = u/c_s \cong \sqrt{2}r/l$ asymptotically does not depend on t . Thus, we get a supersonic flow past obstacle if we place the obstacle at the distance $d > l/\sqrt{2}$ from the axis.

Now, the flow can be considered as uniform if the size of the obstacle (scaled here as the healing length ξ for a chosen moment of observation) satisfies the condition $\xi/d \ll 1$. For $d \sim l$ this gives

$$t \ll ml^2/\hbar. \quad (5)$$

This condition does not contradict to the condition $t \gg 1/\omega_{\perp}$ if $Na_s/a_{\parallel} \gg 1$ which is nothing but the standard condition of applicability of the Thomas-Fermi approximation supposed in the beginning of our discussion.

At last, the characteristic time of establishing the soliton distribution is about $t \sim \xi/c_s \sim m\xi^2/\hbar$ and also satisfies the condition (5), so that the flow can be considered as quasi-stationary.

Thus, we have shown for this example of BEC dynamics that the conditions of homogeneity, uniformity and quasi-stationarity of the flow can be satisfied in experiments with BEC. Thus, we can simplify our treatment of oblique dark solitons by supposing that the flow is uniform and homogeneous, $\mathbf{u} = (M, 0)$ at $|x| \rightarrow \infty$, where M denotes the ratio of the asymptotic velocity of the flow to the sound velocity corresponding to the asymptotic value of the density $n = n_0 = 1$. Let at the initial moment $t = 0$ there be no disturbance in the condensate so that it is described by the plane wave function

$$\psi(x, y)|_{t=0} = \exp(iMx) \quad (6)$$

corresponding to a uniform condensate flow. To be specific, let us take $M = 5$. The density plot at $t = 20$ is shown in Fig. 2. It can be clearly seen how a pair of oblique solitons is formed behind the obstacle. As our simulations show, their length grows with time and, except in the vicinity of the end points, the density distributions does not demonstrate any vorticity.

3. Oblique soliton: analytic solution

To obtain an analytic description of spatial dark solitons in a BEC, we notice that far enough from the obstacle its potential can be neglected, so we transform Eq. (1) to the




Figure 2. Generation of a pair of oblique dark solitons after “switching on” a uniform supersonic flow ($M = 5$) past a disk-shaped impenetrable obstacle of radius $r = 1$ located at $(0, 0)$. The direction of the flow is from left to right. Density plot is shown for $t = 20$. The dark structures correspond to oblique dark solitons, which in turn generate the vortex streets near the end points.




Figure 3. Cross sections of the density distributions for $x = 20$ (dashed line), $x = 60$ (solid line) and $y > 0$ obtained from numerical solution of the GP equation (1) ($M = 5$, $t = 20$). These are compared with profiles (13) with slope $a = 10$ shown as functions of y at the same values of x ($x = 20$ corresponds to “crosses” and $x = 60$ to “circles”).

so-called “hydrodynamic form” by substitution

$$\psi(\mathbf{r}) = \sqrt{n(\mathbf{r})} \exp\left(i \int^{\mathbf{r}} \mathbf{u}(\mathbf{r}') d\mathbf{r}'\right) \exp(-i\mu t), \quad (7)$$

Substituting (7) into (1) and separating real and imaginary parts we obtain a system of equations for the density $n(x, y)$ and the velocity field $\mathbf{u} = (u(x, y), v(x, y))$,

$$\begin{aligned} (nu)_x + (nv)_y &= 0, \\ uu_x + vu_y + n_x + \left(\frac{n_x^2 + n_y^2}{8n^2} - \frac{n_{xx} + n_{yy}}{4n}\right)_x &= 0, \\ uv_x + vv_y + n_y + \left(\frac{n_x^2 + n_y^2}{8n^2} - \frac{n_{xx} + n_{yy}}{4n}\right)_y &= 0, \end{aligned} \quad (8)$$

where we have omitted tildes for convenience. Our aim now is to find the solution of this system under the conditions that the BEC flow is uniform at infinity

$$n = 1, \quad u = M, \quad v = 0 \quad \text{at} \quad |x| \rightarrow \infty, \quad (9)$$

where M denotes the ratio of the asymptotic velocity of the flow to the sound velocity, i.e. the Mach number.

Let us seek a solution of the form $n = n(\theta)$, $u = u(\theta)$, $v = v(\theta)$, where $\theta = x - ay$ and a denotes the slope of a line with a given phase (e.g., location of a wave crest) with the

y -axis. Substitution of this *ansatz* into the system (8) followed by a simple integration, yields expressions for the velocity components in terms of the density

$$u = \frac{M(1 + a^2n)}{(1 + a^2)n}, \quad v = -\frac{aM(1 - n)}{(1 + a^2)n}, \quad (10)$$

where the integration constants were chosen according to condition (9). Besides that, we get the ordinary differential equation for the density

$$\frac{1}{4}(1 + a^2)(n_\theta^2 - 2nn_{\theta\theta}) + 2n^3 - (2 + p)n^2 + p = 0, \quad (11)$$

where

$$p = M^2/(1 + a^2). \quad (12)$$

Equation (11) with the condition (9) for the density at infinity can be easily integrated (see [4] for details) to give the desired solution in the form of a dark soliton

$$n(\theta) = 1 - \frac{1 - p}{\cosh^2[\sqrt{1 - p}\theta/\sqrt{1 + a^2}]}. \quad (13)$$

The velocity components can then be found by substitution of this solution into Eqs. (10). Thus, formula (13) gives the exact dark spatial soliton solution of the GP equation. We call it ‘‘oblique’’ because it is always inclined with respect to the direction of the supersonic flow. To compare the analytic solution (13) with the results of numerical simulations, we have calculated the parameter p from Eq. (12) using the value of the slope a inferred from the numerical solution. A comparison of the theoretical profile of the oblique dark soliton given by Eq. (13) with the corresponding part of the density profile in the full numerical solution is shown in Fig. 3. The excellent agreement between these two profiles confirms that the line patterns in Fig. 2 are indeed oblique dark solitons generated by the obstacle.

The developed here theory explains qualitatively the patterns observed in the experiment [2] where the large amplitude nonlinear waves generated by the BEC flow past an obstacle are clearly seen.

AMK thanks EPSRC and RFBR (grant 05-02-17351) and AG thanks FAPESP and CNPq for financial support.

REFERENCES

1. L.P. Pitaevskii and S. Stringari, Bose-Einstein Condensation, Cambridge University Press, Cambridge, 2003.
2. E.A. Cornell, talk at the Conference on Nonlinear Waves, Integrable Systems and their Applications (Colorado Springs, June 2005); available at <http://jilawww.colorado.edu/bec/papers.html>.
3. G.A. El and A.M. Kamchatnov, Phys. Lett A 350 (2006) 192 ; erratum: Phys. Lett. A 352 (2006) 554 .
4. G.A. El, A. Gammal and A.M. Kamchatnov, Oblique dark solitons in supersonic flow of a Bose-Einstein condensate, Phys. Rev. Lett. (2006) (to appear); also e-print nlin.PS/0604044.
5. A.M. Kamchatnov, JETP 94 (2004) 908.