



Contents lists available at SciVerse ScienceDirect

Biochimica et Biophysica Acta

journal homepage: www.elsevier.com/locate/bbadis

Implications for oxidative stress and astrocytes following 26S proteasomal depletion in mouse forebrain neurones[•]

Jamal Elkhazaz^a, Aslihan Ugun-Klusek^b, Tim Constantin-Teodosiu^a, Karen Lawler^a, R John Mayer^a, Ellen Billett^b, James Lowe^c, Lynn Bedford^a

^a School of Biomedical Sciences, University of Nottingham, Nottingham, UK

^b School of Science and Technology, Nottingham Trent University, Nottingham, UK

^c Division of Histopathology, School of Molecular Medical Sciences, University of Nottingham, UK

article info

Article history:
Received 25 March 2013
Received in revised form 25 June 2013
Accepted 1 July 2013
Available online xxxx

Keywords:
Neurodegeneration
Ubiquitin proteasome system
26S proteasome
Oxidative stress
Peroxiredoxin 6
Astrocytes

abstract

Neurodegenerative diseases are characterized by progressive degeneration of selective neurones in the nervous system, but the underlying mechanisms involved in neuroprotection and neurodegeneration remain unclear. Dysfunction of the ubiquitin proteasome system is one of the proposed hypotheses for the cause and progression of neuronal loss. We have performed quantitative two-dimensional fluorescence difference in-gel electrophoresis combined with peptide mass fingerprinting to reveal proteome changes associated with neurodegeneration following 26S proteasomal depletion in mouse forebrain neurones. Differentially expressed proteins were validated by Western blotting, biochemical assays and immunohistochemistry. Of significance was increased expression of the antioxidant enzyme peroxiredoxin 6 (PRDX6) in astrocytes, associated with oxidative stress. Interestingly, PRDX6 is a bifunctional enzyme with antioxidant peroxidase and phospholipase A₂ (PLA₂) activities. The PLA₂ activity of PRDX6 was also increased following 26S proteasomal depletion and may be involved in neuroprotective or neurodegenerative mechanisms. This is the first in vivo report of oxidative stress caused directly by neuronal proteasome dysfunction in the mammalian brain. The results contribute to understanding neuronal–glial interactions in disease pathogenesis, provide an in vivo link between prominent disease hypotheses and importantly, are of relevance to a heterogeneous spectrum of neurodegenerative diseases.

© 2013 The Authors. Published by Elsevier B.V. All rights reserved.

1. Introduction

Neurodegenerative diseases are characterized by the progressive degeneration of selective neurones of the nervous system. Abnormal protein aggregation, impaired protein degradation, mitochondrial dysfunction and oxidative stress are key hypotheses for cause and progression of major human neurodegenerative diseases, including Alzheimer's disease (AD) and Parkinson's disease (PD) [1–3]. Attention has also turned to the contribution of glial cells to neurodegeneration [4], but the underlying mechanisms involved in neuroprotection and neurodegeneration in the nervous system remain unclear.

The ubiquitin proteasome system (UPS) is the major intracellular pathway for regulated degradation of unwanted proteins and central to normal cellular homeostasis [5]. A sequence of enzymes covalently attach polyubiquitin chains to unwanted proteins as a signal for

degradation by the 26S proteasome. Studies in human brain and disease models have implicated dysfunction of the UPS in the pathological changes that lead to neurodegeneration [6–9]. Ubiquitin-positive protein inclusions are a common feature of human neurodegenerative diseases [6]. Also, in PD and dementia with Lewy bodies (DLB) patients, altered proteasome activity and subunit expression has been reported [7]. In a significant study, we showed that genetic depletion of 26S proteasomes in mouse brain neurones caused neurodegeneration and the formation of protein inclusions resembling human pale bodies, the precursor of Lewy bodies, providing a compelling link between UPS-mediated protein degradation and neurodegeneration [10].

Proteomic studies are of considerable interest to identify much-needed novel pathogenic mechanisms connected to neurodegenerative disease. There are inherent difficulties with a study of a mixture of cell-types in the brain, but it is essential to investigate in vivo models to identify the importance of neuronal–glial cellular interactions during disease development that are not revealed in studies of cell lines composed of a single clonal cell-type. We have employed a quantitative two-dimensional fluorescence difference in-gel electrophoresis (2D-DIGE) proteomic approach to reveal proteome changes associated with cortical neurodegeneration following 26S proteasomal depletion in our unique mouse model.

[•] This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Corresponding author at: D37b School of Biomedical Sciences, University of Nottingham, Queens Medical Centre, Nottingham, NG2 7UH, UK. Tel.: +44 1158230512.

E-mail address: lynn.bedford@nottingham.ac.uk (L. Bedford).

Protein changes identified in our 2D-DIGE study were validated by alternative approaches, namely 1D and 2D Western blotting as well as biochemical and immunohistochemical investigations to further understand their significance. The results show new information linking UPS dysfunction to oxidative stress in the brain in vivo and the importance of understanding neuronal–glial interactions during disease progression.

2. Materials and methods

2.1. 26S proteasomal depletion mouse model

Neurone-specific 26S proteasome-depleted micewerecreatedusing Cre/loxP conditional gene targeting as described in detail previously [10]. For forebrain, including cortex, neurone-specific inactivation of Psmc1, Cre recombinase was expressed under the control of the calcium calmodulin-dependent protein kinase II α promoter (Psmc1^{fl/fl}; CaMKII α -Cre) [10]. CaMKII α is expressed in post-mitotic neurones from approximately post-natal week 2 [11,12]. Appropriate litter-mate mice were used as controls.

All procedures were carried out under personal and project licenses granted by the UK Home Office in accordance with the Animals (Scientific Procedures) Act 1986 and with ethical approval from the University of Nottingham Ethical Review Committee.

2.2. 2D fluorescence difference in-gel electrophoresis (2D-DIGE)

Mouse cortex was homogenized in lysis buffer containing 30 mM Tris–HCl pH 8.8, 8 M Urea and 4% (w/v) CHAPS, followed by centrifugation at 20,000 g for 5 min at 4 °C and collection of supernatant. Protein estimation used the Bio-Rad (Bradford) protein assay kit. CyDye labeling was performed according to the manufacturer's instructions (GE Healthcare) and incorporating a dye swap. 15 μ g of each sample was labeled with Cy3 and Cy5, and a pooled sample was labeled with Cy2 containing equal amounts of all samples as an internal standard. 10 mM lysine was used to stop labeling. First dimension isoelectric focusing (IEF) used a Bio-Rad Protein IEF Cell and 7 cm 3–10 non-linear pH gradient IPG strips (Bio-Rad). Strips were passively rehydrated for 1 h, actively rehydrated for 13 h and 40 min at 50 V followed by IEF (250 V for 20 min linear, 4000 V for 2 h linear, 4000 V for 10,000 V/h rapid). The strips were incubated in 2% (w/v) dithiothreitol in equilibration buffer [50 mM Tris–HCl pH 8.8, 6 M Urea, 2% (w/v) sodium dodecyl sulfate (SDS) and 20% (v/v) glycerol] and then 2.5% (w/v) iodoacetamide in equilibration buffer for 15 min each. Strips were placed on top of 12–18% gradient SDS-PAGE resolving gels for the second dimension. A Fujifilm FLA-5100 scanner was used to scan each gel at the corresponding wavelengths to the CyDyes. Images were analyzed using SameSpots software (Progenesis) with a 1.2-fold change set as the cut off value and ANOVA (P b 0.05).

2.3. Mass spectrometry analysis

For identification of protein spots, gels were either silver (GE Healthcare) or Coomassie blue (Cheshire Sciences) stained using a mass spectrometry-compatible protocol. Spots were excised from the gel manually and washed three times with 50 μ L of acetonitrile (ACN)/25 mM NH₄HCO₃ (2:1) for 15 min each followed by 50 μ L of 25 mM NH₄HCO₃ for 10 min. Gel pieces were then air dried for 15 min and rehydrated in 5 μ L of 12.5 ng/ μ L sequencing grade trypsin (Promega) on ice for 20 min. 5 μ L of 25 mM NH₄HCO₃ was added to each tube and incubated at 37 °C for 4 h. Tryptic digests were collected, dried in a vacuum concentrator (Eppendorf) and 2 μ L of 50% (v/v) ACN/0.1% (v/v) trifluoroacetic acid (TFA) was added to each tube. Finally, 0.5 μ L of sample was transferred to the MALDI plate followed by 0.5 μ L of 10 mg (w/v) α -cyano-4-hydroxycinnamic acid matrix (LaserBio Laboratories) in 50% (v/v) ACN/0.1% (v/v)

TFA. Peptide mass fingerprints were generated using a MALDI-TOF mass spectrometer (Bruker Daltonics Ultraflex III MALDI-TOF). Proteins were identified using the Mascot search engine (<http://www.matrixscience.com>); stating “Mus musculus” species, carbamidomethyl fixed and oxidized methionine as variable modifications and 100 ppm peptide tolerance. Positive identity was given by scores over 56 (comparing Swiss-Prot database) and their molecular mass and pI were compared to the position of the spot on the 2D gel.

2.4. Western blot analysis

Mouse cortex was homogenized as described in 2D-DIGE. 50–100 μ g was mixed with 2 \times reducing sample buffer [150 mM Tris–HCl pH 6.8, 8 M Urea, 10% (v/v) SDS, 20% (v/v) glycerol, 10% (v/v) mercaptoethanol, 3% (w/v) dithiothreitol, 0.1% (w/v) bromophenol blue]. Proteins were separated using 12% SDS-PAGE and transferred to nitrocellulose membrane. Blocking was for 1 h in 5% (w/v) Marvel in Tris-buffered saline containing 0.1% (v/v) Tween 20. Incubation in primary and corresponding horseradish peroxidase-conjugated secondary antibodies (Sigma) was overnight at 4 °C and for 1 h at room temperature respectively in blocking solution. The primary antibodies used were: 1:1000 vimentin (GeneTex), peroxiredoxin 6 (GeneTex), glucose-regulated protein (Cell Signaling), protein disulphide isomerase (GeneTex), CCAAT-enhancer-binding protein homologous protein (Cell Signaling) and glial fibrillary acidic protein (Sigma); 1:500 fumerate hydratase (GeneTex); 1:250 stathmin (GeneTex); 1:200 X-box binding protein (Santa Cruz). Proteins were detected using enhanced chemiluminescence (Pierce). The band intensity was calculated with Quantity One 1-D analysis Software and/or Aida. For 2D Western blot analysis, samples were separated as described in 2D-DIGE and following the second dimension processed to Western blotting as described here.

2.5. Reactive oxygen species assay

Levels of reactive oxygen species were examined using the 2,7-dichlorofluorescein diacetate — cellular reactive oxygen species detection assay kit (abcam) according to the manufacturer's instructions. Detection used fluorescent spectroscopy with excitation and emission of 485 nm and 520 nm respectively.

2.6. Lipid peroxidation

Malondialdehyde (MDA) concentration was determined as an indicator of lipid peroxidation products based on the method of Erdelmeier et al. [13]. Mouse cortex was homogenized in 5 mM butylated hydroxytoluene in 20 mM phosphate buffer pH 7.4, followed by centrifugation at 3000 g for 10 min at 4 °C. Protein estimation used the Bio-Rad (Bradford) protein assay kit. 300 μ L (9.5 μ g/ μ L) was hydrolyzed using HCl pH 1–2 and incubated at 60 °C for 80 min. 60 μ L of sample was mixed with 195 μ L of 10.3 mM N-methyl-2-phenylindol in 3:1 (v/v) acetonitrile:methanol and then 45 μ L of concentrated HCl, incubated at 45 °C for 60 min and centrifuged at 15,000 g for 10 min to clarify. Absorbance was measured spectrophotometrically at 586 nm. Concentration of malondialdehyde (μ M/mg protein) was calculated using 1,1,3,3-tetramethoxypropane as a standard.

2.7. Phospholipase A₂ assay

Phospholipase A₂ (PLA₂) activity (U/ml/mg protein) was determined using EnzChek Phospholipase A2 Assay kit (Invitrogen) according to the manufacturer's instructions. Detection used fluorescent spectroscopy with excitation and emission of 485 nm and 520 nm respectively. MJ33 inhibitor was used at 3 mol% based on previous reports [14,15].

2.8. Protein oxidation

Protein carbonyl content (nM/mg protein) was determined utilizing the 2,4-dinitrophenylhydrazine (DNPH) reaction following Cayman's Protein Carbonyl Colorimetric Assay Kit according to the manufacturer's instructions. Absorbance was measured spectrophotometrically at 375 nm.

2.9. Immunohistochemistry

Mice were perfusion-fixed with 0.9% saline followed by 4% paraformaldehyde in phosphate buffered saline pH 7.4. The brains were then processed to paraffin with chloroform as the clearing agent. Immunohistochemistry was performed as directed in Vector Laboratories M.O.M Immunodetection [GFAP (Sigma)] or Vectastain Elite Rabbit IgG [PRDX6 (GeneTex) and MAP2 (abcam)] ABC kits using 0.01 M citrate buffer containing 0.05% Tween-20 pH 6 for antigen retrieval and appropriate fluorescently-conjugated secondary antibodies.

2.10. Statistical analysis

Results are expressed as mean average \pm SEM. Statistical differences were analyzed by ANOVA and Student's t-test with significance set as indicated.

3. Results

3.1. 26S proteasomal depletion mouse model of neurodegeneration

Generation of neurone-specific 26S proteasome-depleted mice has been described in detail previously [10]. To summarize, the Cre/loxP system spatially restricts inactivation of an essential subunit of the 19S regulatory particle of the 26S proteasome, ATPase Psmc1. PSMC1 is necessary for the assembly and activity of the 26S proteasome [10]. For forebrain neurone-specific inactivation of Psmc1, including cortex, floxed Psmc1 mice were crossed with mice expressing Cre recombinase under the control of the calcium calmodulin-dependent protein kinase II α promoter (Psmc1^{fl/fl}; CaMKII α -Cre). CaMKII α is expressed in post-mitotic neurones from approximately post-natal week 2. We previously showed that 26S proteasomal depletion in mouse cortical brain neurones caused neurodegeneration and the formation of intraneuronal inclusion bodies accompanied by reactive gliosis at 6 weeks of age [10]. The study here investigates proteomic changes accompanying neurodegeneration in the mouse cortex.

3.2. Differentially-expressed proteins in 26S proteasome-depleted cortex

Cortices from individual 6 week-old 26S proteasome-depleted and control (n = 4) mice were compared using 2D-DIGE proteomic analysis and Progenesis SameSpots to identify differentially-expressed proteins. Fig. 1 shows a representative 2D gel image. The expression level of 24 spots showed statistically significant changes between 26S proteasome-depleted and control animals (1.2-fold, ANOVA p b 0.05). Supplementary Table 1 lists the 19 proteins that were identified by peptide mass fingerprinting. The 2D-DIGE results and protein identifications were validated by Western blot analysis of selected differentially-expressed proteins based on antibody availability. We confirmed by 1D Western blotting that expression of glial fibrillary acidic protein (GFAP; Fig. 1, spot 2), vimentin (VIME; Fig. 1, spot 1) and peroxiredoxin 6 (PRDX6; Fig. 1, spot 10) was significantly increased while mitochondrial fumarate hydratase (FUMH; Fig. 1, spot 4) and stathmin (STMN1; Fig. 1, spot 11) were significantly decreased in 26S proteasome-depleted vs. control cortex consistent with the 2D-DIGE analysis (Fig. 2). GFAP and VIME are associated with the intermediate filament system in astrocytes and their up-regulation is a hallmark of astrocyte activation and the resulting




Fig. 1. Representative 2D gel image of mouse brain cortex homogenate labeled with Cy5 dye and differentially-expressed protein spots between 26S proteasome-depleted and control mouse cortices highlighted (spots 1–24). Numbered spots showed 1.2-fold change with statistical significance (ANOVA p b 0.05). Spots 1–16 were identified by peptide mass fingerprinting (Supplementary Table 1).

reactive gliosis [16]. This confirms our previously reported reactive astroglia by GFAP immunostaining of cortical brain sections following neuronal 26S proteasomal depletion [10]. PRDX6 has a well-known role as an antioxidant enzyme and its up-regulation in the cortex following 26S proteasomal depletion is suggestive of oxidative stress [17–22]. FUMH is a key enzyme of the tricarboxylic acid (TCA) cycle and STMN1 has an important function in microtubule dynamics [23].

2D gel electrophoresis can separate isoforms of the same protein, whereas 1D Western blotting provides a single band of total protein. Fig. 3 shows differential expression of GFAP isoforms in 6 week-old mice using 2D Western blot analysis. Two predominant isoforms of GFAP were detected in the control and 26S proteasome-depleted mouse cortices (Fig. 3; spots 1 and 2). Four additional GFAP isoforms were detected in the 26S proteasome-depleted cortex (Fig. 3; spots 3–6). Interestingly, GFAP was also identified in 2D-DIGE spot 9 (Fig. 1 and Supplementary Table 1) that may correspond to spot 5 or 6 in Fig. 3. We cannot exclude that the novel isoforms of GFAP detected in the 26S proteasome-depleted cortex may be present in the control cortex, but below the level of detection by this approach. GFAP isoforms may be associated with astrocyte subtypes that have specific functions and neuropathological conditions in the brain [24,25]. Alternatively, spots 5 and 6 may be GFAP protein breakdown products [26].

3.3. Neuronal 26S proteasomal depletion causes oxidative stress

Since oxidative stress is a pivotal factor in neuronal death in neurodegenerative diseases, we further investigated the antioxidant enzyme PRDX6 and oxidative stress in the mouse cortex following 26S proteasomal depletion.

To investigate the levels of reactive oxygen species (ROS) in 26S proteasome-depleted and control mouse cortices we used 2',7'-dichlorofluorescein diacetate fluorogenic dye. This is the most widely used assay for measuring oxidative stress [27]. The levels of ROS were significantly increased in 26S proteasome-depleted cortices at 2 and half weeks of age (t-test p b 0.05; Fig. 4A). Because CaMKII α is expressed in cortical neurones from approximately postnatal week 2, the data indicate that the ensuing loss of PSMC1 and 26S proteasome activity causes oxidative stress. There was no significant difference in ROS levels between 26S proteasome-depleted and control mouse cortices at 3 weeks-old (Fig. 4B). At 4 and 6 weeks




Fig. 2. Validation and quantitation of identified protein changes following 26S proteasomal depletion in mouse cortex. 1D Western blot analysis of total cortical homogenates from 6 week-old control and 26S proteasome-depleted (*Psmc1^{fl/fl}; CaMKII α -Cre*) mice for GFAP, VIME, PRDX6, FUMH and STMN1. A representative β -actin loading control is shown; this was performed for each Western blot. (B–F) Densitometry used QuantityOne software. Values were normalized to β -actin and represented as % vs. controls. Error bars represent SEM. n = 3, *p < 0.05, **p < 0.01 (Student's t-test).

290 of age there was a significant decrease in the levels of ROS in 26S
 291 proteasome-depleted cortices compared to controls (t-test p < 0.01;
 292 Fig. 4C and D). Linear regression analysis showed a significant correla-
 293 tion between age and the levels of ROS in 26S proteasome-depleted
 294 cortex (p < 0.05; Fig. 4E). There was also a significant correlation
 295 between age and the levels of PRDX6 protein expression in 26S
 296 proteasome-depleted cortex between 2 and 6 weeks-old (p < 0.01;
 297 Fig. 4E). Importantly, there was an inverse relationship between the
 298 levels of PRDX6 protein expression and ROS in 26S proteasome-

depleted mouse cortex with increasing age, indicative of an antioxidant
 response of PRDX6 (Fig. 4E).

3.4. Neuronal 26S proteasomal depletion causes increased lipid peroxidation

High polyunsaturated fatty acid content makes the brain particu-
 larly susceptible to oxidative stress-associated lipid damage. Also,
 lipid peroxidation is known to be an autocatalytic process, amplifying
 the destructive effects of the initial free radical [28,29]. Quantitation




Fig. 3. Protein isoform expression of GFAP revealed by 2D Western blotting of control and 26S proteasome-depleted (*Psmc1^{fl/fl}; CaMKII α -Cre*) mouse cortices. Arrows indicate six GFAP isoforms.




Fig. 4. Increased PRDX6 protein expression is associated with decreased reactive oxygen species (ROS). Levels of ROS (A–D) in control and 26S proteasome-depleted (*Psmc1^{fl/fl}; CaMKII α -Cre*) cortices. Data represented as mean \pm SEM. n = 6, *p < 0.05, **p < 0.01 (Student's t-test). (E) Inverse relationship between the levels of PRDX6 protein expression and ROS in 26S proteasome-depleted mouse cortex with increasing age. Data represented as mean \pm SEM.

of malondialdehyde (MDA), a toxic secondary product of membrane lipid peroxidation, in cortical tissue homogenates between 4 and 6 weeks of age identified significantly increased levels of MDA in 5 and 6 week-old 26S proteasome-depleted mice compared to controls (t-test p < 0.01; Fig. 5A–C), indicating that lipid oxidation is increased following neuronal 26S proteasomal depletion.

Protein carbonyls are hallmarks of the oxidative status of proteins. Therefore, to further investigate oxidative stress, we evaluated carbonyl content spectrophotometrically using a reaction with 2,4-dinitrophenylhydrazine in mouse cortical tissue homogenates.

No significant difference in the levels of protein carbonyls was observed between 26S proteasome-depleted and control mouse cortices at 6 weeks-old (Supplementary Fig. 1).

3.5. Increased phospholipase A₂ activity in 26S proteasome-depleted cortex

Interestingly, PRDX6 is a bifunctional enzyme with peroxidase and phospholipase A₂ (PLA₂) activities [30]. The PLA₂ activity of PRDX6 has not been studied as widely as the peroxidase-associated antioxidant properties. Quantitation of PLA₂ activity in 26S proteasome-depleted and control cortical homogenates between 4 and 6 weeks of age showed significantly increased activity in the 6 week-old 26S proteasome-depleted mouse cortex (t-test p < 0.01; Fig. 5D–F). The chemical inhibitor MJ33 that has previously been shown to have some (although not total) specificity for PRDX6 PLA₂ activity [14,15] significantly decreased PLA₂ activity in 26S proteasome-depleted cortex, suggesting that some of the PLA₂ activity was associated with PRDX6 (Fig. 5F).

3.6. Astrocytic localization of PRDX6

To investigate the cellular localization of PRDX6 we performed double immunofluorescent labeling of brain sections with PRDX6 and GFAP or 200 kD neurofilament heavy chain (NF-H) for astrocytes and neurones respectively (Fig. 6 and Supplementary Fig. 2). PRDX6 immunolabeled cells with the characteristic morphology of astrocytes in control and 26S proteasome-depleted cortices from 6 week-old mice (Fig. 6). Double-labeling with GFAP confirmed the localization of PRDX6 in astrocytes in the 26S proteasome-depleted mice (Fig. 6; right-hand panel). GFAP is the most widely used marker for immunohistochemical identification of astrocytes and labels reactive astrocytes that are responding to central nervous system (CNS) damage, but it is recognized that not all non-reactive astrocytes in the healthy CNS are identified by GFAP [31]. Therefore, PRDX6 expression in the control mouse brain is in non-reactive astrocytes that are not immunohistochemically labeled by GFAP (Fig. 6; left-hand panel). Importantly, we noted a much higher diffuse PRDX6 staining in the 26S proteasome-depleted cortical brain sections compared to the control (Fig. 6; compare i and ii), suggesting PRDX6 may be secreted by activated astrocytes in response to the neuronal changes. The expression of PRDX6 did not co-localize with NF-H in mouse cortical neurones (Supplementary Fig. 2).

4. Discussion

Proteomic studies of human post-mortem brain and disease models are of considerable interest to understand pathogenic mechanisms connected to neurodegenerative disease. This study has identified and validated several differentially-expressed proteins accompanying neurodegeneration in the mouse cortex following neuronal 26S proteasomal depletion. Among these, the antioxidant enzyme PRDX6 was significantly increased. Since oxidative stress is a pivotal factor in human neurodegenerative diseases [32,33], supported by animal and cellular models [33–35], we further investigated PRDX6 and oxidative stress in the 26S proteasome-depleted mouse cortex.

Here we have shown a significant inverse relationship between the levels of PRDX6 protein expression and ROS following 26S proteasomal depletion in the mouse cortical neurones, indicative of oxidative stress and an antioxidant response of PRDX6. Lipid peroxidation was also significantly increased in the cortex of 26S proteasome-depleted mice. Similar to the other PRDXs and glutathione peroxidase family, PRDX6 can reduce hydrogen peroxide and short chain hydroperoxides, but PRDX6 can also directly bind and reduce phospholipid hydroperoxides [36,37]. This characteristic plays an important role in its antioxidant defense [19,20]. Studies in cell and mouse models demonstrate that




Fig. 5. Neuronal 26S proteasomal depletion causes increased lipid peroxidation and PRDX6 PLA₂ activity. Quantitation of MDA (A–C) and PLA₂ activity (D–F) in control and 26S proteasome-depleted (*Psmc1^{fl/fl}; CaMKIIα-Cre*) cortices at 4, 5 and 6 weeks of age. (F) PLA₂ activity is significantly decreased by chemical inhibitor MJ33 at 6 weeks of age. Data presented as mean ± SEM. n = 4, **p < 0.01 (Student's t-test).

375 decreased expression or overexpression of PRDX6 results in increased
376 sensitivity or resistance to oxidant stress respectively [17–22].

377 PRDX6 is a bifunctional enzyme with peroxidase and PLA₂ activities
378 [30]. Increased PLA₂ activity in 26S proteasome-depleted cortex corre-
379 lates with increased PRDX6 expression. Given previous studies have
380 shown that the PLA₂ activity of PRDX6 is sensitive to MJ33 [14,15], we
381 propose that the decreased PLA₂ activity in the presence of MJ33 is
382 partly attributable to PRDX6. However, we recognize that MJ33 is not
383 totally specific for PRDX6 and that other phospholipases that have
384 not been investigated in this study presumably explain the MJ33-
385 insensitive PLA₂ activity [38].

386 The PLA₂ activity of PRDX6 has been associated with several cellular
387 functions. PLA₂ enzyme activity liberates both a free fatty acid and
388 lysophosphatidylcholine from phosphatidylcholine substrates and has
389 been implicated in oxidative stress-induced apoptosis and inflamma-
390 tion [39–42]. Importantly, a recent study in pulmonary microvascular
391 endothelial cells suggested that the PLA₂ activity of PRDX6 may also
392 play a role in antioxidant protection provided by PRDX6 [43].

393 The cellular distribution of PRDX6 in our mouse model is similar to
394 previous studies in mouse and human brain showing expression of
395 PRDX6 mainly in astrocytes [44–49]. A study in mouse brain neural
396 cell types showed differential expression patterns of the six mammalian
397 isoforms of the PRDX family and only PRDX6 was found in astrocytes,
398 which may be indicative of a specific role in their function [46]. Impor-
399 tantly, an increase in PRDX6 and the number and staining intensity of
400 PRDX6-positive astrocytes has been described in human brain regions
401 affected in AD, PD and DLB, as well as other neurodegenerative disease
402 mouse models [44,45,47,49,50]. Since oxidative stress is regarded as a
403 fundamental process in the events that lead to neurodegeneration, the
404 antioxidant function of PRDX6 may play an important neuroprotective
405 response of the astrocyte [32,51,52]. Further support for PRDX6 in this

context was shown in parkin-deficient mice, where PRDX6 was
406 downregulated [53].

407 We also noted a much higher diffuse PRDX6 staining in the 26S
408 proteasome-depleted cortical brain sections compared to the control,
409 suggesting PRDX6 may be secreted by activated astrocytes. This is
410 supported by previous studies that have suggested that this enzyme
411 may be a secreted protein [44,54]. Evidence suggests PRDX6 is
412 present at very low levels in neurones and an early study in PD and
413 DLB disease brains demonstrated the presence of PRDX6 in Lewy
414 bodies [44,46–49]. However, PRDX6 expression was not detectable
415 in neurones or inclusion bodies in our mouse model.

416 This is the first in vivo report of oxidative stress caused directly by
417 neuronal 26S proteasome dysfunction in the mammalian brain. Our
418 findings are supported by cellular studies using chemical proteasome
419 inhibitors [55–57]. Various antioxidant defenses have also been dem-
420 onstrated in response to proteasome inhibitor oxidative stress [57],
421 but PRDX6 has not been described previously and most likely because
422 studies were not focused on the brain. Since the high polyunsaturated
423 fatty acid content makes the brain particularly susceptible to lipid
424 peroxidation, the unique ability of PRDX6 to reduce phospholipid
425 hydroperoxides may play an important role in antioxidant protection
426 in the brain.

427 Increased astrocytic PRDX6 expression was associated with de-
428 creased levels of ROS, and together with the presence of oxidative
429 stress, supports an antioxidant neuroprotective role of astrogliosis
430 in response to neurodegeneration caused by 26S proteasome deple-
431 tion in mouse brain neurones. However, the astrocytic network has
432 a wide range of activities that can be both beneficial and detrimental
433 such as energy metabolism and the release of inflammatory molec-
434 ules respectively [31,58]. We emphasize that the PLA₂ activity of
435 PRDX6 may also be involved in the production of further mediators
436




Fig. 6. Astrocytic localization of PRDX6 in the mouse cortex. Double immunofluorescent labeling of cortical brain sections from control and 26S proteasome-depleted (*Psmc1^{fl/fl}; CaMKIIα-Cre*) 6 week-old mice with GFAP (green) and PRDX6 (red). DAPI (blue) was used as a fluorescent nuclear counterstain. Enlarged views of the boxed areas are shown (zoom). Note much higher diffuse PRDX6 staining in the 26S proteasome-depleted cortical brain sections in addition to the more focused staining in astrocytes (compare i and ii). Scale bar, 50 μ m.

and may reflect mitochondrial dysfunction. We recently reported that 26S proteasomal depletion in mouse brain neurones leads to the formation of inclusions composed predominantly of morphologically abnormal mitochondria with disrupted or disintegrated cristae [60]. A spectrum of mitochondrial pathologies that may be associated with oxidative stress has been described in human neurodegenerative diseases and associated disease models, including perturbed respiratory chain function, mitochondrial dynamics and clearance [32,33,61]. Therefore, we suggest that mitochondrial dysfunction may be important in the mechanism of oxidative stress and neurodegeneration following 26S proteasome depletion.

Proteasome inhibition is known to induce endoplasmic reticulum (ER) stress and activation of the unfolded protein response (UPR) signaling pathways [62–64]. ER stress is associated with the production of ROS from the ER as well as mitochondria and evidence of ER stress has been shown in various human neurodegenerative diseases, such as AD and PD [65–68]. Investigation of key mammalian ER stress-induced proteins; the chaperone glucose-regulated protein 78 (GRP78), the transcription factor X-box binding protein-1 (XBP1), protein disulphide isomerase (PDI) and the cell death mediator CCAAT-enhancer-bindingprotein homologous protein (CHOP), showed that neuronal 26S proteasomal depletion does not cause activation of the UPR (Supplementary Fig. 3) [69,70]. Taken together, ER stress is not an important source of ROS in this model.

Although we found evidence for increased oxidation of lipids indicative of oxidative stress in the cortex following neuronal 26S proteasome depletion, protein oxidation was not increased. Proteasome function is known to be important for the degradation of oxidatively modified proteins [71–75]. Therefore, we may have expected to find increased protein oxidation following 26S proteasome depletion in the mouse cortex due to increased oxidative stress and/or decreased removal of oxidatively modified proteins. However, in the heterogeneous population of cellular proteasome complexes, the 26S proteasome has a relatively minor role in the removal of oxidatively damaged proteins compared to the 20S proteasome [72,73,76–78]. We previously showed that inactivation of *Psmc1* specifically disrupts 26S proteasome function; assembly and activity of the 20S core proteolytic proteasome was not affected [10]. Therefore, 20S proteasome function in *Psmc1*; *CaMKIIα-Cre* neurones may be sufficient to protect cells from protein oxidative modification. Alternatively, it is possible that the level of protein oxidation was not sufficient for detection in a mixed cell population of targeted (*CaMKIIα*) and non-targeted neurones and glia.

Different quantitative proteomic approaches will favor different subpopulations of proteins. 2D-DIGE fluorescence-based detection provides high sensitivity that is linear over several orders of magnitude [79,80]. The significant advantage of this technology is the ability to multiplex using different fluorescent cyanine dyes, providing greater accuracy of quantitation over conventional 2D gel approaches [79]. However, the percentage of lysine residues in proteins may affect labeling efficiency and current in-gel digestion and mass spectrometers limit identification of lesser abundant proteins detected by 2D-DIGE [79]. Together with the well-known limitations of 2D gel electrophoresis, i.e. hydrophobic proteins, dynamic range and quantitative distribution issues, our study may miss some important molecular players involved in the orchestration of cellular events following neuronal 26S proteasomal depletion [79–81].

In conclusion, we reveal that oxidative stress may contribute to the cellular events leading to neurodegeneration following UPS dysfunction, providing a novel intersection between two prominent hypotheses in disease pathogenesis. Increased astrocytic expression of PRDX6 also reveals innovative information regarding the role of neuronal–glial interactions and astroglialosis in neurodegeneration.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.bbadis.2013.07.002>.

related to cellular signaling functions that may be protective or deleterious in the progressive neuronal loss.

The intercellular signaling molecule(s) and mechanism(s) that modulate reactive astrogliosis in response to neurodegeneration following 26S proteasomal depletion in neurones will require further study. Diverse molecules have been suggested that can be released by all central nervous system cell-types, including ROS [58,59]. Interestingly, we report here that mitochondrial FUMH, a key enzyme of the TCA cycle, is decreased in the 26S proteasome-depleted cortex

512 Acknowledgements

513 This work was supported by Parkinson's UK.

514 References

- 515 [1] A.H. Schapira, P. Jenner, Etiology and pathogenesis of Parkinson's disease, *Mov.*
516 *Disord.* 26 (2011) 1049–1055.
- 517 [2] K. Leuner, W.E. Muller, A.S. Reichert, From mitochondrial dysfunction to amyloid
518 beta formation: novel insights into the pathogenesis of Alzheimer's disease, *Mol.*
519 *Neurobiol.* 46 (2012) 186–193.
- 520 [3] T. Morawe, C. Hiebel, A. Kern, C. Behl, Protein homeostasis, aging and Alzheimer's
521 disease, *Mol. Neurobiol.* 46 (2012) 41–54.
- 522 [4] E. Croisier, M.B. Graeber, Glial degeneration and reactive gliosis in alpha-
523 synucleinopathies: the emerging concept of primary gliodegeneration, *Acta*
524 *Neuropathol.* 112 (2006) 517–530.
- 525 [5] A.L. Goldberg, Protein degradation and protection against misfolded or damaged
526 proteins, *Nature* 426 (2003) 895–899.
- 527 [6] J. Lowe, A. Blanchard, K. Morrell, G. Lennox, L. Reynolds, M. Billett, M. Landon,
528 R.J. Mayer, Ubiquitin is a common factor in intermediate filament inclusion
529 bodies of diverse type in man, including those of Parkinson's disease, Pick's
530 disease, and Alzheimer's disease, as well as Rosenthal fibres in cerebellar
531 astrocytomas, cytoplasmic bodies in muscle, and Mallory bodies in alcoholic
532 liver disease, *J. Pathol.* 155 (1988) 9–15.
- 533 [7] D. Ebrahimi-Fakhari, L. Wahlster, P.J. McLean, Protein degradation pathways in
534 Parkinson's disease: curse or blessing, *Acta Neuropathol.* 124 (2012) 153–172.
- 535 [8] F.J. Dennissen, N. Kholod, F.W. van Leeuwen, The ubiquitin proteasome system in
536 neurodegenerative diseases: culprit, accomplice or victim? *Prog. Neurobiol.* 96
537 (2012) 190–207.
- 538 [9] M. Orre, W. Kamphuis, S. Dooves, L. Kooijman, E.T. Chan, C.J. Kirk, V. Dimayuga
539 Smith, S. Koot, C. Mamber, A.H. Jansen, H. Ova, E.M. Hol, Reactive glia show
540 increased immunoproteasome activity in Alzheimer's disease, *Brain* 136 (2013)
541 1415–1431.
- 542 [10] L. Bedford, D. Hay, A. Devoy, S. Paine, D.G. Powe, R. Seth, T. Gray, I. Topham,
543 K. Fone, N. Rezvani, M. Mee, T. Soane, R. Layfield, P.W. Sheppard, T. Ebendal,
544 D. Usoskin, J. Lowe, R.J. Mayer, Depletion of 26S proteasomes in mouse brain
545 neurons causes neurodegeneration and Lewy-like inclusions resembling human
546 pale bodies, *J. Neurosci.* 28 (2008) 8189–8198.
- 547 [11] M. Mayford, M.E. Bach, Y.Y. Huang, L. Wang, R.D. Hawkins, E.R. Kandel, Control of
548 memory formation through regulated expression of a CaMKII transgene, *Science*
549 274 (1996) 1678–1683.
- 550 [12] J.Z. Tsien, D.F. Chen, D. Gerber, C. Tom, E.H. Mercer, D.J. Anderson, M. Mayford,
551 E.R. Kandel, S. Tonegawa, Subregion- and cell type-restricted gene knockout in
552 mouse brain, *Cell* 87 (1996) 1317–1326.
- 553 [13] I. Erdelmeier, D. Gerard-Monnier, J.C. Yadan, J. Chaudiere, Reactions of N-methyl-
554 2-phenylindole with malondialdehyde and 4-hydroxyalkenals. Mechanistic aspects
555 of the colorimetric assay of lipid peroxidation, *Chem. Res. Toxicol.* 11 (1998)
556 1184–1194.
- 557 [14] A.B. Fisher, C. Dodia, A. Chander, M. Jain, A competitive inhibitor of phospholipase A2
558 decreases surfactant phosphatidylcholine degradation by the rat lung, *Biochem. J.*
559 288 (Pt 2) (1992) 407–411.
- 560 [15] A.B. Fisher, C. Dodia, Role of phospholipase A2 enzymes in degradation of
561 dipalmitoylphosphatidylcholine by granular pneumocytes, *J. Lipid Res.* 37 (1996)
562 1057–1064.
- 563 [16] M. Pekny, M. Nilsson, Astrocyte activation and reactive gliosis, *Glia* 50 (2005)
564 427–434.
- 565 [17] Y. Manevich, T. Sweitzer, J.H. Pak, S.I. Feinstein, V. Muzykantov, A.B. Fisher, I-Cys
566 peroxiredoxin overexpression protects cells against phospholipid peroxidation-
567 mediated membrane damage, *PNAS* 99 (2002) 11599–11604.
- 568 [18] X. Wang, S.A. Phelan, K. Forsman-Semb, E.F. Taylor, C. Petros, A. Brown, C.P. Lerner, B.
569 Paigen, Mice with targeted mutation of peroxiredoxin 6 develop normally but are
570 susceptible to oxidative stress, *J. Biol. Chem.* 278 (2003) 25179–25190.
- 571 [19] Y. Wang, S.I. Feinstein, Y. Manevich, Y.S. Ho, A.B. Fisher, Lung injury and mortality
572 with hyperoxia are increased in peroxiredoxin 6 gene-targeted mice, *Free Radic.*
573 *Biol. Med.* 37 (2004) 1736–1743.
- 574 [20] Y. Wang, S.I. Feinstein, Y. Manevich, Y.S. Ho, A.B. Fisher, Peroxiredoxin 6
575 gene-targeted mice show increased lung injury with paraquat-induced oxidative
576 stress, *Antioxid. Redox Signal.* 8 (2006) 229–237.
- 577 [21] Y. Wang, S.A. Phelan, Y. Manevich, S.I. Feinstein, A.B. Fisher, Transgenic mice
578 overexpressing peroxiredoxin 6 show increased resistance to lung injury in
579 hyperoxia, *Am. J. Respir. Cell Mol. Biol.* 34 (2006) 481–486.
- 580 [22] Y. Wang, S.I. Feinstein, A.B. Fisher, Peroxiredoxin 6 as an antioxidant enzyme:
581 protection of lung alveolar epithelial type II cells from H₂O₂-induced oxidative
582 stress, *J. Cell. Biochem.* 104 (2008) 1274–1285.
- 583 [23] L. Jourdain, P. Curmi, A. Sobel, D. Pantaloni, M.F. Carrier, Stathmin: a tubulin-
584 sequestering protein which forms a ternary T2S complex with two tubulin molecules,
585 *Biochemistry* 36 (1997) 10817–10821.
- 586 [24] J. Middeldorp, E.M. Hol, GFAP in health and disease, *Prog. Neurobiol.* 93 (2011)
587 421–443.
- 588 [25] W. Kamphuis, C. Mamber, M. Moeton, L. Kooijman, J.A. Sluijs, A.H. Jansen, M. Verwee,
589 L.R. de Groot, V.D. Smith, S. Rangarajan, J.J. Rodriguez, M. Orre, E.M. Hol, GFAP
590 isoforms in adult mouse brain with a focus on neurogenic astrocytes and reactive
591 astrogliosis in mouse models of Alzheimer disease, *PLoS One* 7 (2012) e42823.
- [26] J.S. Zoltewicz, D. Scharf, B. Yang, A. Chawla, K.J. Newsom, L. Fang, Characterization
of antibodies that detect human GFAP after traumatic brain injury, *Biomark.*
Insights 7 (2012) 71–79.
- [27] B. Kalyanaraman, V. Darley-Usmar, K.J. Davies, P.A. Dennerly, H.J. Forman, M.B. Grisham,
G.E. Mann, K. Moore, L.J. Roberts II, H. Ischiropoulos, Measuring reactive oxygen and
nitrogen species with fluorescent probes: challenges and limitations, *Free Radic. Biol.*
Med. 52 (2012) 1–6.
- [28] B.A. Freeman, J.D. Crapo, Biology of disease: free radicals and tissue injury, *Lab.*
Invest. 47 (1982) 412–426.
- [29] T.F. Slater, Free-radical mechanisms in tissue injury, *Biochem. J.* 222 (1984) 1–15.
- [30] A.B. Fisher, Peroxiredoxin 6: a bifunctional enzyme with glutathione peroxidase
and phospholipase A(2) activities, *Antioxid. Redox Signal.* 15 (2011) 831–844.
- [31] M.V. Sofroniew, H.V. Vinters, Astrocytes: biology and pathology, *Acta Neuropathol.*
119 (2010) 7–35.
- [32] S. Gandhi, A.Y. Abramov, Mechanism of oxidative stress in neurodegeneration, *606*
Oxid. Med. Cell. Longev. (2012), (428010).
- [33] M. Varcin, E. Bentea, Y. Michotte, S. Sarre, Oxidative stress in genetic mouse
models of Parkinson's disease, *Oxid. Med. Cell. Longev.* (2012), (624925).
- [34] J.K. Andersen, Oxidative stress in neurodegeneration: cause or consequence? *Nat.*
Med. 10 (2004) S18–25, (Suppl.).
- [35] D. Pratico, Evidence of oxidative stress in Alzheimer's disease brain and antioxidant
therapy: lights and shadows, *Ann. N. Y. Acad. Sci.* 1147 (2008) 70–78.
- [36] A.B. Fisher, C. Dodia, Y. Manevich, J.W. Chen, S.I. Feinstein, Phospholipid hydro-
peroxides are substrates for non-selenium glutathione peroxidase, *J. Biol. Chem.*
274 (1999) 21326–21334.
- [37] Y. Manevich, T. Shuvaeva, C. Dodia, A. Kazi, S.I. Feinstein, A.B. Fisher, Binding of
peroxiredoxin 6 to substrate determines differential phospholipid hydroperoxide
peroxidase and phospholipase A(2) activities, *Arch. Biochem. Biophys.* 485 (2009)
139–149.
- [38] M.K. Jain, B.Z. Yu, J. Rogers, G.N. Ranadive, O.G. Berg, Interfacial catalysis by phospho-
lipase A2: dissociation constants for calcium, substrate, products, and competitive
inhibitors, *Biochemistry* 30 (1991) 7306–7317.
- [39] P.J. Leavey, C. Gonzalez-Aller, G. Thurman, M. Kleinberg, L. Rinckel, D.W.
Ambruso, S. Freeman, F.A. Kuypers, D.R. Ambruso, A 29-kDa protein associated
with p67phox expresses both peroxiredoxin and phospholipase A2 activity
and enhances superoxide anion production by a cell-free system of NADPH
oxidase activity, *J. Biol. Chem.* 277 (2002) 45181–45187.
- [40] S. Chatterjee, S.I. Feinstein, C. Dodia, E. Sorokina, Y.C. Lien, S. Nguyen, K. Debolt,
D. Speicher, A.B. Fisher, Peroxiredoxin 6 phosphorylation and subsequent
phospholipase A2 activity are required for agonist-mediated activation of
NADPH oxidase in mouse pulmonary microvascular endothelium and alveolar
macrophages, *J. Biol. Chem.* 286 (2011) 11696–11706.
- [41] S.Y. Kim, E. Chun, K.Y. Lee, Phospholipase A(2) of peroxiredoxin 6 has a critical
role in tumor necrosis factor-induced apoptosis, *Cell Death Differ.* 18 (2011)
1573–1583.
- [42] D.R. Ambruso, M.A. Ellison, G.W. Thurman, T.L. Leto, Peroxiredoxin 6 translocates
to the plasma membrane during neutrophil activation and is required for optimal
NADPH oxidase activity, *Biochim. Biophys. Acta* 1823 (2012) 306–315.
- [43] Y.C. Lien, S.I. Feinstein, C. Dodia, A.B. Fisher, The roles of peroxidase and phospholipase
A2 activities of peroxiredoxin 6 in protecting pulmonary microvascular endothelial
cells against peroxidative stress, *Antioxid. Redox Signal.* 16 (2012) 440–451.
- [44] J.H. Power, J.M. Shannon, P.C. Blumbers, W.P. Gai, Nonselenium glutathione
peroxidase in human brain: elevated levels in Parkinson's disease and dementia
with Lewy bodies, *Am. J. Pathol.* 161 (2002) 885–894.
- [45] C.W. Strey, D. Spellman, A. Stieber, J.O. Gonatas, X. Wang, J.D. Lambris, N.K. Gonatas,
Dysregulation of stathmin, microtubule-destabilizing protein, and up-regulation of
Hsp25, Hsp27, and the antioxidant peroxiredoxin 6 in a mouse model of familial
amyotrophic lateral sclerosis, *Am. J. Pathol.* 165 (2004) 1701–1718.
- [46] M.H. Jin, Y.H. Lee, J.M. Kim, H.N. Sun, E.Y. Moon, M.H. Shong, S.U. Kim, S.H. Lee,
T.H. Lee, D.Y. Yu, D.S. Lee, Characterization of neural cell types expressing
peroxiredoxins in mouse brain, *Neurosci. Lett.* 381 (2005) 252–257.
- [47] J.H. Power, S. Asad, T.K. Chataway, F. Chegini, J. Manavis, J.A. Temlett, P.H. Jensen, P.C.
Blumbers, W.P. Gai, Peroxiredoxin 6 in human brain: molecular forms, cellular
distribution and association with Alzheimer's disease pathology, *Acta Neuropathol.*
115 (2008) 611–622.
- [48] J. Goemaere, B. Knoop, Peroxiredoxin distribution in the mouse brain with emphasis
on neuronal populations affected in neurodegenerative disorders, *J. Comp. Neurol.*
520 (2011) 258–280.
- [49] K. Yata, S. Oikawa, R. Sasaki, A. Shindo, R. Yang, M. Murata, K. Kanamaru, H. Tomimoto,
Astrocytic neuroprotection through induction of cytoprotective molecules; a
proteomic analysis of mutant P301S tau-transgenic mouse, *Brain Res.* 1410
(2011) 12–23.
- [50] K. Krapfenbauer, E. Engidawork, N. Cairns, M. Fountoulakis, G. Lubec, Aberrant
expression of peroxiredoxin subtypes in neurodegenerative disorders, *Brain*
Res. 967 (2003) 152–160.
- [51] M.F. Beal, Oxidatively modified proteins in aging and disease, *Free Radic. Biol.*
Med. 32 (2002) 797–803.
- [52] C. Pimentel, L. Batista-Nascimento, C. Rodrigues-Pousada, R.A. Menezes, Oxidative
stress in Alzheimer's and Parkinson's diseases: insights from the yeast *Saccharomyces*
cerevisiae, *Oxid. Med. Cell. Longev.* 2012 (2012) 132146.
- [53] J.J. Palacino, D. Sagi, M.S. Goldberg, S. Krauss, C. Motz, M. Wacker, J. Klose, J. Shen,
Mitochondrial dysfunction and oxidative damage in parkin-deficient mice, *J. Biol.*
Chem. 279 (2004) 18614–18622.
- [54] X.Z. Zhang, Z.F. Xiao, C. Li, Z.Q. Xiao, F. Yang, D.J. Li, M.Y. Li, F. Li, Z.C. Chen,
Triosephosphate isomerase and peroxiredoxin 6, two novel serum markers for
human lung squamous cell carcinoma, *Cancer Sci.* 100 (2009) 2396–2401.

- 678 [55] M. Demasi, K.J. Davies, Proteasome inhibitors induce intracellular protein aggregation
679 and cell death by an oxygen-dependent mechanism, *FEBS Lett.* 542 (2003) 89–94.
- 680 [56] Y.H. Ling, L. Liebes, Y. Zou, R. Perez-Soler, Reactive oxygen species generation and
681 mitochondrial dysfunction in the apoptotic response to Bortezomib, a novel
682 proteasome inhibitor, in human H460 non-small cell lung cancer cells, *J. Biol.*
683 *Chem.* 278 (2003) 33714–33723.
- 684 [57] J. Chandra, Oxidative stress by targeted agents promotes cytotoxicity in hematologic
685 malignancies, *Antioxid. Redox Signal.* 11 (2009) 1123–1137.
- 686 [58] M.V. Sofroniew, Molecular dissection of reactive astrogliosis and glial scar formation,
687 *Trends Neurosci.* 32 (2009) 638–647.
- 688 [59] F. Antunes, E. Cadenas, Estimation of H₂O₂ gradients across biomembranes, *FEBS*
689 *Letts.* 475 (2000) 121–126.
- 690 [60] S.M. Paine, G. Anderson, K. Bedford, K. Lawler, R.J. Mayer, J. Lowe, L. Bedford, Pale
691 body-like inclusion formation and neurodegeneration following depletion of 26S
692 proteasomes in mouse brain neurones are independent of alpha-synuclein, *PLoS*
693 *One* 8 (2013) e54711.
- 694 [61] A. Federico, E. Cardaioli, P. Da Pozzo, P. Formichi, G.N. Gallus, E. Radi, Mitochondria,
695 oxidative stress and neurodegeneration, *J. Neurol. Sci.* 322 (2012) 254–262.
- 696 [62] A. Fribley, Q. Zeng, C.Y. Wang, Proteasome inhibitor PS-341 induces apoptosis
697 through induction of endoplasmic reticulum stress-reactive oxygen species in
698 head and neck squamous cell carcinoma cells, *Mol. Cell. Biol.* 24 (2004) 9695–9704.
- 699 [63] M.S. Choy, M.J. Chen, J. Manikandan, Z.F. Peng, A.M. Jenner, A.J. Melendez, N.S. Cheung,
700 Up-regulation of endoplasmic reticulum stress-related genes during the early phase of
701 treatment of cultured cortical neurons by the proteasomal inhibitor lactacystin, *J. Cell.*
702 *Physiol.* 226 (2011) 494–510.
- 703 [64] R. Xiong, D. Siegel, D. Ross, The activation sequence of cellular protein handling
704 systems after proteasomal inhibition in dopaminergic cells, *Chem. Biol. Interact.*
705 204 (2013) 116–124.
- 706 [65] E. Ferreira, I. Baldeiras, I.L. Ferreira, R.O. Costa, A.C. Rego, C.F. Pereira, C.R. Oliveira,
707 Mitochondrial- and endoplasmic reticulum-associated oxidative stress in Alzheimer's
708 disease: from pathogenesis to biomarkers, *Int. J. Cell Biol.* (2012), (735206).
- 709 [66] J.F. Abisambra, U.K. Jinwal, L.J. Blair, J.C. O'Leary III, Q. Li, S. Brady, L. Wang, C.E. Guidi,
710 B. Zhang, B.A. Nordhues, M. Cockman, A. Suntharalingham, P. Li, Y. Jin, C.A. Atkins,
711 C.A. Dickey, Tau accumulation activates the unfolded protein response by impairing
712 endoplasmic reticulum-associated degradation, *J. Neurosci.* 33 (2013) 9498–9507.
- 713 [67] B. Bhandary, A. Marahatta, H.R. Kim, H.J. Chae, An involvement of oxidative stress
714 in endoplasmic reticulum stress and its associated diseases, *Int. J. Mol. Sci.* 14
715 (2013) 434–456.
- 716 [68] T. Omura, M. Kaneko, Y. Okuma, K. Matsubara, Y. Nomura, Endoplasmic reticulum
717 stress and Parkinson's disease: the role of HRD1 in averting apoptosis in neurodegen-
718 erative disease, *Oxid. Med. Cell. Longev.* (2013), (239854).
- 719 [69] J.D. Malhotra, H. Miao, K. Zhang, A. Wolfson, S. Pennathur, S.W. Pipe, R.J. Kaufman,
720 Antioxidants reduce endoplasmic reticulum stress and improve protein secretion,
721 *PNAS* 105 (2008) 18525–18530.
- 722 [70] A. Samali, U. Fitzgerald, S. Deegan, S. Gupta, Methods for monitoring endoplasmic
723 reticulum stress and the unfolded protein response, *Int. J. Cell Biol.* (2010), (830307).
- 724 [71] A.M. Pickering, K.J. Davies, Degradation of damaged proteins: the main function
725 of the 20S proteasome, *Prog. Mol. Biol. Transl. Sci.* 109 (2012) 227–248.
- 726 [72] A.M. Pickering, A.L. Koop, C.Y. Teoh, G. Ermak, T. Grune, K.J. Davies, The
727 immunoproteasome, the 20S proteasome and the PA28alpha proteasome
728 regulator are oxidative-stress-adaptive proteolytic complexes, *Biochem. J.* 432
729 (2010) 585–594.
- 730 [73] T. Grune, B. Catalgol, A. Licht, G. Ermak, A.M. Pickering, J.K. Ngo, K.J. Davies, HSP70
731 mediates dissociation and reassociation of the 26S proteasome during adaptation
732 to oxidative stress, *Free Radic. Biol. Med.* 51 (2011) 1355–1364.
- 733 [74] Q. Ding, E. Dimayuga, J.N. Keller, Proteasome regulation of oxidative stress in aging
734 and age-related diseases of the CNS, *Antioxid. Redox Signal.* 8 (2006) 163–172.
- 735 [75] X. Wang, J. Yen, P. Kaiser, L. Huang, Regulation of the 26S proteasome complex
736 during oxidative stress, *Sci. Signal.* 3 (2010) ra88.
- 737 [76] R. Shringarpure, T. Grune, J. Mehlhase, K.J. Davies, Ubiquitin conjugation is not
738 required for the degradation of oxidized proteins by proteasome, *J. Biol. Chem.*
739 278 (2003) 311–318.
- 740 [77] M. Kastle, S. Reeg, A. Rogowska-Wrzesinska, T. Grune, Chaperones, but not oxidized
741 proteins, are ubiquitinated after oxidative stress, *Free Radic. Biol. Med.* 53 (2012)
742 1468–1477.
- 743 [78] S.A. Hussong, R.J. Kappahn, S.L. Phillips, M. Maldonado, D.A. Ferrington,
744 Immunoproteasome deficiency alters retinal proteasome's response to stress,
745 *J. Neurochem.* 113 (2010) 1481–1490.
- 746 [79] N.S. Tannu, S.E. Hemby, Two-dimensional fluorescence difference gel electrophore-
747 sis for comparative proteomics profiling, *Nat. Protoc.* 1 (2006) 1732–1742.
- 748 [80] T. Rabilloud, M. Chevallet, S. Luche, C. Lelong, Two-dimensional gel
749 electrophoresis in proteomics: past, present and future, *J. Proteomics* 73
750 (2010) 2064–2077.
- 751 [81] S. Beranova-Giorgianni, Proteome analysis by two-dimensional gel electrophore-
752 sis and mass spectrometry: strengths and limitations, *TrAC Trends Anal. Chem.* 22
753 (2003) 273–281.
- 754