



University of Dundee

From mouse to man

Lipworth, Brian; Anderson, W. J.; Short, P. M.

Published in:
British Journal of Pharmacology

DOI:
[10.1111/bph.13335](https://doi.org/10.1111/bph.13335)

Publication date:
2016

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Lipworth, B. J., Anderson, W. J., & Short, P. M. (2016). From mouse to man: predicting biased effects of beta-blockers in asthma. *British Journal of Pharmacology*, 173(1), 248-249. DOI: 10.1111/bph.13335

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



From mouse to man: Predicting biased effects of beta-blockers in asthma

Journal:	<i>British Journal of Pharmacology</i>
Manuscript ID:	Draft
Manuscript Type:	Letter
Date Submitted by the Author:	n/a
Complete List of Authors:	Lipworth, Brian; University of Dundee, Scottish Centre for Respiratory Research Anderson, William; University of Dundee, Scottish Centre for Respiratory Research Short, Philip; University of Dundee, Scottish Centre for Respiratory Research
Major area of pharmacology:	Respiratory pharmacology
Cross-cutting area:	Drug discovery/target validation
Additional area(s):	Adrenoceptors

SCHOLARONE™
Manuscripts

"This is the peer reviewed version of the following article: Lipworth, B. J., Anderson, W. J., and Short, P. M. (2016) From mouse to man: predicting biased effects of beta-blockers in asthma. *British Journal of Pharmacology*, 173: 248–249., which has been published in final form at doi: 10.1111/bph.13335.. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving."

From mouse to man: predicting biased effects of beta-blockers in asthma

B J Lipworth, W J Anderson, P M Short

Scottish Centre for Respiratory Research
Ninewells Hospital and Medical School
Dundee, DD1 9SY, UK

Correspondence: B J Lipworth, Scottish Centre for Respiratory Research, University of Dundee, Ninewells Hospital & Medical School, Dundee, DD1 9SY b.j.lipworth@dundee.ac.uk, Tel: 01382 383188, Fax: 0182 383259

Conflict of interest statement: BJL has received previous grant support from the Chief Scientist Office, Scotland to evaluate effects of propranolol in patients with persistent treated asthma. BJL has also received unrestricted grant support from Chiesi, Meda, Almirall and Teva to evaluate small airways in persistent asthma and COPD; as well as multi-centre pharmaceutical support from Astra Zeneca, Teva, Janssen and Roche. In addition BJL has received personal payment for consultancy and advisory boards with the following pharmaceutical companies: Astra Zeneca, Chiesi, Teva, Boehringer Ingelheim and Meda. BJL has also received personal payment for giving speaker talks with Chiesi, Teva, Meda and Mitsubishi Tanabe as well as support to attend educational meetings from Chiesi, Boehringer Ingelheim and Teva. WJA has no conflict of interest and PMS has no conflict of interest.

We read with interest the elegant data from (Thanawala *et al.*, 2015) in ovalbumin sensitized epinephrine deficient or wild type mice which suggested differential effects of biased signaling with propranolol and nadolol. We were particularly intrigued by the observation in epinephrine deficient mice whereby there was a disconnect between propranolol and nadolol in restoring the asthma phenotype compared to controls. It is tempting to simplistically extrapolate these data in mice to what might happen in human subjects with asthma in order to explain the negative effects of propranolol on airway hyper-responsiveness (AHR), reported in two separate placebo controlled double blind trials in patients receiving inhaled corticosteroids (ICS), which were powered to detect a one doubling dilution difference in the provocative concentration of methacholine (n=18) or histamine (n=16) to produce a 20% fall in forced expiratory volume in 1s (i.e. the PC20 FEV1 threshold) (Anderson *et al.*, 2014; Short *et al.*, 2013). In this regard, in patients with persistent asthma, the PC threshold for FEV1 is closely related to PC threshold for airway resistance (Short *et al.*, 2015).

Upon close inspection of the data for methacholine AHR (Thanawala *et al.*, 2015), the provocative concentration to induce a 100% increase (PC100 threshold) in airway resistance was unaltered in wild mice (n=6) treated with propranolol in contrast to an increase with nadolol (n=7). The blunting of methacholine AHR with nadolol which was statistically significant ($P < 0.05$) amounting to approximately a 0.6 doubling dilution shift compared to vehicle treated mice (n=10). Such an effect in mice with nadolol on AHR would be considered clinically irrelevant in human patients as it is less than the minimal important difference of one doubling dilution shift in PC threshold. It is therefore difficult to extrapolate the magnitude of this effect with nadolol on methacholine AHR in mice to what has previously been reported in two unblinded studies with nadolol in human asthmatic subjects which amounted to an approximate two doubling dilution shift in methacholine PC20, albeit in mild intermittent asthmatics who were not taking ICS (Hanania *et al.*, 2010; Hanania *et al.*, 2008).

It is however unclear how the relative mg dose per body weight of propranolol in mice (80-140mg/L in water) equates to that in humans (80mg slow release tablet /day). Moreover if propranolol at usual therapeutic doses of 80mg/day does indeed confer arrestin independent partial agonist activity at the ERK1/2 activation pathway in humans then one might expect to see an increase in TH2 mediated inflammatory biomarkers. For example in persistent asthmatics there was no worsening in eosinophils, eosinophilic cationic protein or exhaled breath nitric oxide when oral propranolol 80mg/day was added to a low dose of ICS, while a higher dose of ICS in conjunction with oral placebo produced further suppression of the same TH2 biomarkers (Anderson *et al.*, 2014).

Moreover asthma control and disease specific quality of life were also unaltered by propranolol (Anderson *et al.*, 2014; Short *et al.*, 2013).

In order to properly confirm the putative beneficial effects of biased inhibitory signaling in mice this will require a placebo controlled trial to demonstrate clinically relevant improvements in methacholine PC20, inflammatory markers and asthma control with nadolol on top of existing ICS therapy in persistent asthma. The placebo controlled clinical trial (clinicaltrials.gov)

NCT01804218) evaluating effects of nadolol in ICS naïve mild intermittent asthmatics will unfortunately not answer this clinically important question.

References

Anderson WJ, Short PM, Williamson PA, Manoharan A, Lipworth BJ (2014). The inverse agonist propranolol confers no corticosteroid-sparing activity in mild-to-moderate persistent asthma. *Clin Sci (Lond)* **127**(11): 635-643.

Hanania NA, Mannava B, Franklin AE, Lipworth BJ, Williamson PA, Garner WJ, *et al.* (2010). Response to salbutamol in patients with mild asthma treated with nadolol. *Eur Respir J* **36**(4): 963-965.

Hanania NA, Singh S, El-Wali R, Flashner M, Franklin AE, Garner WJ, *et al.* (2008). The safety and effects of the beta-blocker, nadolol, in mild asthma: an open-label pilot study. *Pulm Pharmacol Ther* **21**(1): 134-141.

Short PM, Anderson WJ, Manoharan A, Lipworth BJ (2015). Usefulness of impulse oscillometry for the assessment of airway hyperresponsiveness in mild-to-moderate adult asthma. *Ann Allergy Asthma Immunol* **115**(1): 17-20.

Short PM, Williamson PA, Anderson WJ, Lipworth BJ (2013). Randomized placebo-controlled trial to evaluate chronic dosing effects of propranolol in asthma. *Am J Respir Crit Care Med* **187**(12): 1308-1314.

Thanawala VJ, Valdez DJ, Joshi R, Forkuo GS, Parra S, Knoll BJ, *et al.* (2015). Beta-blockers have differential effects on the murine asthma phenotype. *Br J Pharmacol*.