



## **Providing mental healthcare to immigrants: current challenges and new strategies.**

Giacco, D; Matanov, A; Priebe, S

For additional information about this publication click this link.

<http://qmro.qmul.ac.uk/xmlui/handle/123456789/9533>

Information about this research object was correct at the time of download; we occasionally make corrections to records, please therefore check the published record when citing. For more information contact [scholarlycommunications@qmul.ac.uk](mailto:scholarlycommunications@qmul.ac.uk)



# Providing mental healthcare to immigrants: current challenges and new strategies

*Domenico Giacco, Aleksandra Matanov, and Stefan Priebe*

## Purpose of review

The article reviews recent evidence on improving access to mental healthcare for immigrants and best practice of care provision.

## Recent findings

Language barriers, different beliefs and explanatory models of illness, confidentiality concerns, stigma, reluctance to seek psychological help outside families, and social deprivation may prevent immigrants from accessing mental healthcare. Pathways are influenced by families, primary care practitioners, voluntary organizations, and social services. Interpreting services are often not available, and data documentation on immigrants' use of services is inconsistent. Nonmedical specific services for immigrants can be effective in outreach activities. Cultural training of staff can improve clinicians' attitudes and patients' satisfaction with care. Integrative approaches between primary and mental healthcare, psychoeducational programs, and technological innovations have been developed to improve access to care.

## Summary

Immigrants can face significant barriers in accessing mental healthcare. Strategies to overcome these barriers are as follows: increased coordination and communication between voluntary organizations, social services and mental health services; training of staff on cross-cultural issues; integration of mental healthcare with primary care; psychoeducational initiatives focused on families and broader social groups; and technology-based interventions.

## Keywords

barriers, immigrants, mental healthcare, pathways to care

## INTRODUCTION

Immigrants, that is, people who were born outside the country of residence [1], show higher rates of mental disorders, such as psychosis, mood disorders, anxiety disorders, and particularly posttraumatic stress disorder, compared with indigenous populations [2,3].

Various factors can influence their higher psychiatric morbidity, that is, premigration, migration, and postmigration stressors, long-term adjustment difficulties in another country, and poor living conditions [4].

In the last 50 years, immigration has almost doubled, with currently 191 million immigrants worldwide. Targeted mental healthcare strategies for these populations may be needed [5]. Understanding barriers to care for immigrants and identifying their pathways may help to develop strategies to meet the cultural, religious, and linguistic needs of different immigrant groups [5,6].

## WHAT IS ALREADY KNOWN?

Immigrants face substantial obstacles in accessing healthcare services. Previous studies identified language and literacy issues, cultural differences in explanatory models of illness, restricted legal entitlements, social deprivation, and traumatic experiences as the main barriers to accessing and receiving healthcare [1,7,8]. Immigrants often have specific needs and treatment preferences [9,10]. Good practice recommendations have emphasized the need for organizational flexibility, provision of

Unit for Social and Community Psychiatry, Barts<sup>1</sup> and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK

Correspondence to Professor Stefan Priebe, Unit for Social and Community Psychiatry, Newham Centre for Mental Health, London E13 8SP, UK. Tel: +44 20 75404210; fax: +44 20 75402976; e-mail: s.priebe@qmul.ac.uk

**Curr Opin Psychiatry** 2014, 27:000–000

DOI:10.1097/YCO.0000000000000065

## Provision of services to people with mental illness

### KEY POINTS

- Access to mental healthcare in immigrants may be hindered by language barriers; different beliefs and explanatory models of mental illness; confidentiality concerns; influence of stigma and reluctance to seek help outside immediate social networks; and social deprivation.
- Immigrants often follow nonmedical pathways to care through self-referral, social services, voluntary not-for-profit or religious organizations.
- Interpreting services and cultural competence training for staff are rarely available in mainstream health services, and information on service use by immigrants is scarce.
- Collaboration and sharing of information between mental health services and nonmedical services (voluntary organizations and social services), integration of mental healthcare with physical care, and psychoeducational family programs may increase help-seeking and engagement with services in immigrant populations.
- Technology-based interventions may help support translation of information, reach underserved populations, and deliver culturally tailored psychosocial programs.

good professional interpreting services, raising cultural awareness of staff, delivery of psychoeducational programs, collaboration with families, communities and social services, building positive and stable relationships with patients, and clear guidelines on care entitlements of migrants [1,9,11,12]. Sound research evidence on how mental healthcare should be delivered to benefit immigrants is still scarce.

### PURPOSE OF THE REVIEW

In this article, we review the evidence published between June 2012 and November 2013 on the barriers encountered by immigrants when accessing mental healthcare; their pathways to care; the current state-of-care provision for immigrant groups; and newly developed interventions to improve access to and quality of care provided to immigrants. The included studies were carried out in European countries, the United States, Canada, and Australia, and encompass both economic and political immigrants.

### BARRIERS TO MENTAL HEALTHCARE FOR IMMIGRANTS

Barriers to mental healthcare fall into five categories: language barriers, different beliefs and

explanatory models of mental illness, confidentiality concerns, influence of stigma and reluctance to seek help outside immediate social networks, and social deprivation. These barriers are similar to those in general healthcare, which have been well documented [1,7]. However, the specific nature and social significance of mental disorders can lead to additional problems in accessing care.

### Language barriers

Good communication between clinicians and patients is particularly important in mental healthcare in which diagnostic procedures are based on verbal communication rather than on physical and objective examinations. Mental health practitioners qualitatively assessed in 16 major European cities in the EUGATE study reported language difficulties as a major hindrance to assessing symptoms, establishing a diagnosis, and developing a relationship with immigrant patients [13<sup>1</sup>]. Language problems were also the most prominent barriers to accessing treatment among Latin-American immigrants in the USA, in both community samples and patient groups with depression and PTSD [14<sup>1</sup>,15<sup>1</sup>,16<sup>1</sup>]. Patients reported difficulties in expressing their symptoms and needs to English-speaking clinicians and in understanding their instructions [15<sup>1</sup>,16<sup>1</sup>,17,18<sup>1</sup>]. A preference for bilingual clinicians was a major concern for trauma-exposed Latin-American women [15<sup>1</sup>]. Insufficient language skills may also limit access to specific treatments such as psychotherapy. General practitioners in Denmark were sometimes reluctant to refer immigrants to such treatments because of the lack of bilingual therapists and skilled interpreters [19]. Forty-three percent of psychotherapists surveyed in Hamburg ( $N=485$ ) refused treatment to immigrant patients because of language problems [20<sup>1</sup>]. Patients' difficulties in expressing themselves in a psychotherapeutic setting may be frustrating for both patients and clinicians, with negative effects on therapeutic alliances, while the presence of interpreters introduces further challenges in establishing trust and a therapeutic rapport [15<sup>1</sup>,21<sup>1</sup>].

### Explanatory models of mental illness and expectations of care

Mental health practitioners surveyed across Europe reported that divergent explanatory models of mental illness, such as beliefs in supernatural causes and preferences for physiological interpretations were commonly encountered in non-western immigrant patients [13<sup>1</sup>]. They were perceived as a potential barrier to the diagnostic procedure, making it difficult to assess psychosocial functioning and

AQ1

AQ2

distinguish between culturally accepted beliefs and actual symptoms [13<sup>11</sup>,18<sup>11</sup>,20<sup>11</sup>]. Patients' views and values may clash with those of clinicians, resulting in distrust and difficulties in establishing therapeutic alliances. Subsequently, engagement with both psychopharmacological and psychotherapeutic treatment may be poor [13<sup>11</sup>,21<sup>11</sup>,22]. Adherence to antipsychotic medication was poorer among immigrant patients in Spain than indigenous patients (19 vs. 40%), with particularly low rates among sub-Saharan Africans [23<sup>11</sup>]. However, Danish general practitioners reported that immigrant patients sometimes demanded medication when clinicians themselves believed psychological treatment was more suitable [19]. Different views and expectations may also be linked to a lack of a perceived need for mental healthcare and poor knowledge of services, as found in studies on Latin, Korean, and African-American immigrants in the USA [15<sup>11</sup>,24–26].

### **Stigma and reluctance to seek help outside immediate social networks**

Stigma and negative views of mental disorder can influence treatment seeking in Latin-American immigrant patients with depression and PTSD [15<sup>11</sup>,16<sup>11</sup>], and may explain their concerns about receiving medication. Mental health practitioners providing care for Asian-Americans also reported stigma as an important obstacle to these patients [17]. The significance of families and their role as a source of psychological support for Latin-American and Asian immigrants has been highlighted by both patients and professionals in the USA and Australia [16<sup>11</sup>,17,27]. However, the presence of this support may also lead to reluctance to seek help outside the family. Immigrant patients may prefer self-reliance and have concerns about the family's reputation [16<sup>11</sup>,17,27].

### **Social deprivation**

Specific problems arise when immigrants live in poor socioeconomic conditions. Mental health professionals in European urban settings emphasized the risk of marginalization for immigrant patients [13<sup>11</sup>]. They suggested engaging with social services and communities to increase the social integration. The self-reported need for interpreters in GP consultations among Danish immigrants was linked to modest income levels, poor health, and unemployment [28<sup>11</sup>]. Economic factors such as costs, lack of health insurance, and unemployment were the main barriers to utilizing mental healthcare among Latin-American immigrants in the USA, particularly

in low-income groups [14<sup>11</sup>,15<sup>11</sup>,18<sup>11</sup>]. Homelessness was the most significant risk factor for dropping out of substance abuse treatment in the same population [29<sup>11</sup>].

### **PATHWAYS TO MENTAL HEALTHCARE**

Immigrants in both USA and Europe often follow nonmedical pathways to care through self-referral, social services, voluntary not-for-profit or religious organizations [14<sup>11</sup>,16<sup>11</sup>,30<sup>11</sup>]. Thirty-nine percent of immigrant patients in Italian community mental health centers were referred through such routes [30<sup>11</sup>]. The community outreach activities of these agencies seem to be key in promoting access to care, particularly for marginalized groups of immigrants [30<sup>11</sup>,31<sup>11</sup>]. For example, language assistance programs increased access of Spanish-speaking people to mental health services in California, but only when implemented by community-based organizations [32<sup>11</sup>]. Emergency departments were a primary source of care for low-income Latin-American groups and treatment was offered to patients who screened positive for depression in this setting [18<sup>11</sup>]. Those who dropped out of the subsequent intervention were concerned about receiving medication, but valued counseling delivered by a social worker, particularly over the phone.

### **CURRENT STATE OF MENTAL HEALTHCARE PROVISION**

Information on immigrants' use of mental healthcare and on provision of relevant staff and services is scarce. Several studies published in the period considered in this review have explored this issue.

#### **Service use registers**

The EUGATE study assessed emergency, primary care, and mental healthcare services in 16 European cities [33<sup>11</sup>] and found that 48% of all services kept a database of information on patients' service use. However, immigration status was recorded by only 25% of mental health services, which is still higher than that in other investigated services (19 and 10%).

#### **Provision of interpreting services and bilingual staff**

The EUGATE study also explored the provision of interpreting services in different settings [33<sup>11</sup>]. Many surveyed organizations (42%) did not provide any language assistance; 53% never provided any face-to-face interpreting, and 59% never had any

## Provision of services to people with mental illness

telephone interpreting. Face-to-face interpreting was most often provided by mental health services (35%), followed by emergency (23%) and primary care services (20%). In California, the presence of language assistance programs was associated with improved access to mental healthcare for Spanish-speaking patients [30<sup>■</sup>]. The provision of interpreters increased completion rates in substance abuse treatment for patients from Central and South America [29<sup>■</sup>].

AQ3 Seventy-six percent of mental health services assessed in the EUGATE study reported having members of staff with migrant background [33<sup>■</sup>], and practitioners frequently regarded this as an effective way to improve the attitude toward immigrants in the service. A survey of public outpatient psychotherapists in Hamburg found that only 14% of them were immigrants, whereas 27% of the population in the area was of such background [20<sup>■</sup>]. Almost all bilingual professionals were from the EU countries, although the largest immigrant groups were from outside Europe. Community-based mental health providers in California reported a significantly higher number of Spanish-speaking bilingual staff in comparison to Medi-Cal county providers (180 vs. 17); however, this was not associated with improved treatment access [32<sup>■</sup>].

### Cultural competence

AQ4 Cultural competence training for mental health staff was viewed as important for improving the knowledge, attitudes, and skills of mental health professionals [13<sup>■</sup>]. Ninety-one percent of psychotherapists in Hamburg considered culturally sensitive attitudes necessary for conducting psychotherapy with immigrants, although cultural beliefs and rituals informed the therapeutic approach of just 29% [20<sup>■</sup>]. Seventy-two percent of therapists agreed that advanced training in cross-cultural competence would be helpful, whereas 9% had already completed some form of such training.

Bilingual and bicultural Latin-American therapists in the USA reported using culturally preferred communication styles, language, and self-disclosure to engage with patients of the same immigrant background [34<sup>■</sup>].

### Asylum seekers, refugees, and undocumented migrants

Providing care to asylum seekers, refugees, and undocumented migrants poses specific challenges. Restricted legal entitlements to care and lack of required documents may prevent these patients from accessing healthcare in both Europe and the

USA [15<sup>■</sup>,18<sup>■</sup>,30<sup>■</sup>]. Addressing confidentiality concerns is particularly important as patients may fear being reported to the authorities in the host country or their countries of origin [15<sup>■</sup>,21<sup>■</sup>,35<sup>■</sup>]. Concerns about the ethnic and political background of interpreters can inhibit information sharing and the development of trust [13<sup>■</sup>]. An uncertain immigration status and a fear of repatriation undermine the motivation for psychotherapeutic treatment [21<sup>■</sup>,36<sup>■</sup>], although some refugee patients appear to benefit despite precarious circumstances [36<sup>■</sup>].

Traumatic experiences are frequent in these populations and disclosing them may in itself be a challenge to a therapeutic relationship [36<sup>■</sup>]. Refugee interpreters may feel overwhelmed by the emotional impact of interpreting, particularly for patients with shared histories, and require professional guidance and training in self-care [37<sup>■</sup>]. Developing trust and good therapeutic alliance among trauma-affected patients, psychotherapists, and interpreters was seen as an essential curative factor by all three groups [21<sup>■</sup>].

AQ5 A study exploring mental healthcare provision for refugees, asylum seekers, and undocumented migrants in eight European capitals [31<sup>■</sup>] identified differences in service provision between generic services and those with a specific focus on these groups, frequently found in not-for-profit and voluntary sectors. The latter tend to offer more outreach programs and social support, whereas medical interventions were mostly provided by generic services. The activities of specific and generic services overlapped substantially, highlighting the need for good collaboration and information sharing.

## MENTAL HEALTH INTERVENTIONS FOR IMMIGRANTS

A number of studies in the present review assessed mental health interventions designed to meet the needs of immigrants, with particular emphasis on integrating mental and physical healthcare, developing community and family-oriented approaches, and using new technologies.

### Integrating mental and physical care

General practitioners and primary care services can promote immigrants' access to and engagement with mental healthcare, as they care for a large number of such patients and are able to build trusting relationships [14<sup>■</sup>,16<sup>■</sup>]. A survey of Latin-American immigrant patients in the USA who received mental health treatment at community healthcare clinics, found that 77% originally made an appointment because of physical ailments [26]. Trauma-exposed



Latin-American women expressed a preference for treatment, and in particular individual psychotherapy, delivered by mental health specialists in primary care settings [15<sup>■</sup>].

Approaches to integrate mental health and primary care services have shown positive results in immigrant populations. They successfully reduced depressive symptoms in Asian-Americans [38,39<sup>■</sup>] in both general community health centers and culturally sensitive clinics [39<sup>■</sup>].

### **Community-focused and family-focused psychoeducational approaches**

Providing information on physical and mental health in community settings and utilizing family support are being increasingly used in psychoeducational approaches for immigrants.

Workshops on mental health and physical diseases were effective in increasing knowledge and utilization of services amongst African immigrants in the USA [25].

Culturally tailored approaches for promoting a healthy lifestyle for pregnant and early postpartum Latin-American women in the USA also reduced the risk of depression [40]. Parenting interventions combined with practical and emotional support by befrienders were experienced as psychologically beneficial by asylum-seeking mothers in the UK [41].

A 12-week multifamily group intervention with bilingual facilitators delivered to women with perinatal depression showed good feasibility (13 out of 16 families attended more than 90% of sessions), and improved the psychosocial functioning and support in the families [42<sup>■</sup>].

Family-oriented, culturally tailored psychoeducation intervention for Chinese-American patients with schizophrenia improved symptom levels and quality of life, knowledge of the illness, and social support for the caregivers [43<sup>■</sup>]. Flexibility in delivering the intervention to accommodate the specific needs of families was found to be particularly helpful.

### **Technology-based interventions**

Technology-based interventions have been developed to overcome language barriers, ensure a detailed and valid mental health assessment of immigrants, reach underserved populations, and provide tailored psychosocial approaches.

An iPad-assisted psychosocial risk-assessment was tested on a population of Afghani refugees in Canada [35<sup>■</sup>]. The assessment tool increased the reported intention to visit a psychosocial counselor (72% of experimental vs. 46% of usual care group).

Using Internet videoconferencing (Webcam intervention) for online consultations between psychiatrists and immigrants with depression was found to be more effective than standard primary care in reducing depressive symptoms and improving quality of life [44<sup>■</sup>].

A culturally tailored, problem-solving, web-based intervention was tested in Turkish immigrants with depression [45<sup>■</sup>]. The study had high attrition rates and failed to show a significant effect on depressive symptoms. However, a completers-only analysis suggested better recovery from depression in the experimental group.

### **CONCLUSION**

Research on mental healthcare for immigrants is still inconsistent and has methodological shortcomings. Yet, a number of common barriers preventing access to mental healthcare have been identified in very different immigrant populations. Immigrants, particularly those with poorer socioeconomic status, may share similar pathways to mental healthcare that are different from those of indigenous populations and often involve non-medical agencies. Recent evidence suggests that several strategies should be implemented and tested to improve access to mental healthcare in immigrants:

- (1) Sharing of information between mental health services and existing networks of voluntary organizations and social services, which are in a better position to carry out outreach activities, develop trusting relationships, and direct patients to mental health services.
- (2) Improving the collaboration between 'migrant-specific' (voluntary organizations, charities) and generic mental health services, with regular communication and protocols to avoid overlap of activities.
- (3) Training of mental health professionals on cultural beliefs and explanatory models of mental disorders may improve attitudes to immigrant patients, and help reassure patients about their confidentiality concerns.
- (4) Integrating mental healthcare with physical healthcare can help to engage immigrants, in particular, but not exclusively, when significant physical health and mental health needs are present at the same time.
- (5) Psychoeducational family programs may increase knowledge about mental health problems. Considering the family-centered culture of many immigrant groups, such programs can influence help-seeking.

## Provision of services to people with mental illness

- (6) Technology-based interventions can support a translation of information, enable same-language clinicians to access underserved populations, and support a cost-effective implementation of culturally tailored psychosocial programs.

### Acknowledgements

No funding was received for the preparation of this article.

### Conflicts of interest

The authors declare no conflicts of interest.

### REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

AQ6

1. Priebe S, Sandhu S, Dias S, *et al.* Good practice in healthcare for migrants: views and experiences of care professionals in 16 European countries. *BMC Public Health* 2011; 11:187.
2. Bhugra D, Jones P. Migration and mental illness. *Adv Psychiatr Treat* 2001; 7:216–222.
3. Orozco R, Borges G, Medina-Mora ME, *et al.* A cross-national study on prevalence of mental disorders, service use, and adequacy of treatment among Mexican and Mexican American Populations. *Am J Public Health* 2013; 103:1610–1618.
4. Lindencrona F, Ekblad S, Hauff E. Mental health of recently resettled refugees from the Middle East in Sweden: the impact of preresettlement trauma, resettlement stress and capacity to handle stress. *Soc Psychiatry Psychiatr Epidemiol* 2008; 43:121–131.
5. International Organisation for Migration. International migration, health and human rights. Geneva: International Organisation for Migration; 2013.
6. Heinz A, Kluge U. Mental health in different groups of migrants and ethnic minority within Europe and beyond: regional and cross-national challenges and approaches in research, practice and training. *Eur Psychiatry* 2012; 27:S1–S4.

AQ7

7. Mladovsky P. Migrant health in the EU. *Eurohealth* 2007; 13:9–1.
8. Agudelo Suarez AA, Gil Gonzales D, Vives Cases C. A metasynthesis of qualitative studies regarding opinions and perceptions about barriers and determinants of health services' accessibility in economic migrants. *BMC Health Serv Res* 2012; 12:461.
9. Priebe S, Matanov A, Schor R, *et al.* Good practice in mental healthcare for socially marginalised groups in Europe: a qualitative study of expert views in 14 countries. *BMC Public Health* 2012; 12:248.
10. Cabassa LJ, Lester R, Zayas LH. It is like being in a labyrinth: 'Hispanic immigrants' perceptions of depression and attitudes towards treatments. *J Immigr Minor Health* 2007; 9:1–16.
11. Cambridge J, Singh SP, Johnson M. The need for measurable standards in mental health interpreting: a neglected area. *Psychiatr Bull* 2012; 36:121–124.
12. Bäärnhielm S, Mösko M. Cross-cultural training in mental healthcare: challenges and experiences from Sweden and Germany. *Eur Psychiatry* 2012; 27:S70–S76.
13. Sandhu S, Bjere NV, Dauvrin M, *et al.* Experiences with treating immigrants: a qualitative study in mental health services across 16 European countries. *Soc Psychiatry Psychiatr Epidemiol* 2013; 48:105–116.

This study qualitatively assessed experiences of 48 mental health professionals working in 16 European countries. The interviews highlighted specific challenges to treating immigrants in mental health services, including complications with diagnosis, difficulty in developing trust, and increased risk of marginalization. This is the first time that common challenges in delivering mental healthcare to immigrants were identified across such a wide range of countries.

14. Bridges AJ, Andrews AR, Deen TL. Mental health needs and service utilization by Hispanic immigrants residing in mid-southern United States. *J Transcult Nurs* 2012; 23:359–368.

This study investigated mental health needs and service utilization patterns in a community sample of Hispanic immigrants in the USA. The findings highlighted the importance of economic and language barriers as well as the greater likelihood of participants with mental disorder accessing medical but not psychiatric services.

15. Kaltman S, Hurtado de Mendoza A, Gonzales FA, Serrano A. Preferences for trauma-related mental health services among Latina immigrants from Central America, South America, and Mexico. *Psychol Trauma* 2013. doi: 10.1037/a0031539; Advance online publication March 25.

AQ8

This study explored mental health treatment preferences in trauma-exposed Latin-American immigrant women. The results highlighted a preference for mental health treatment delivered in primary care settings and by mental health professionals, as well as for individual psychotherapy over medication.

16. Hansen MC, Cabassa LJ. Pathways to depression care: help-seeking experiences of low-income Latinos with diabetes and depression. *J Immigrant Minority Health* 2012; 14:1097–1106.

This study investigated pathways to depression care in low-income Latin-American immigrant patients with comorbid diabetes and depression. The importance of family and primary care physicians initiating treatment was highlighted by the findings. Significant barriers to care in this patient group included tendency toward self-reliance, stigma, and language difficulties.

17. Sung J, Mayo N, Lasley C, Ko MJ. Characteristics of collaborative care in increasing access to mental health service in the Asian community. *Fam Syst Health* 2013; 31:307–318.

18. Wells A, Lagomassino IT, Palinkas LA, *et al.* Barriers to depression treatment among low-income, Latino emergency department patients. *Commun Ment Health J* 2013; 49:412–418.

This study assessed perceptions on barriers and facilitators to depression treatment among low-income Latin-American patients in emergency departments in the USA. The study highlighted concerns about medication, transport and employment issues, and lack of immigrant documentation among patients who discontinued treatment, as well as the preference for proactive telephone counseling provided by a social worker.

19. Jensen NK, Norredam M, Priebe S, Krasnik A. How do general practitioners experience providing care to refugees with mental health problems? A qualitative study from Denmark. *BMC Fam Pract* 2013; 14:17.

20. Mosko MO, Gil-Martinez F, Shultz H. Cross-cultural opening in German outpatient mental healthcare service: an exploratory study of structural and procedural aspects. *Clin Psychol Psychother* 2013; 20:434–446.

This study was a comprehensive survey of 485 public outpatient psychotherapists in Hamburg investigating their migration background, cross-cultural encounters, and cross-cultural sensitivity. The challenges in providing treatment to immigrant patients because of language barriers and different cultural values were frequently encountered. Culturally sensitive attitudes were seen as important for working with immigrant patients and the need for advanced cross-cultural sensitivity training was emphasized.

21. Mirdal JM, Rydal E, Essendrop Sondey M. Traumatized refugees, their therapists, and their interpreters: three perspectives on psychological treatment. *Psychol Psychother Theor Res Pract* 2012; 85:436–455.

This study qualitatively assessed how traumatized refugees, their therapists, and interpreters in Denmark perceive curative and hindering factors in psychological therapy, taking into account both successful and unsuccessful encounters. The data were analyzed for each triad and across three groups of participants. The findings emphasized the need for developing trust and good therapeutic alliance between therapists, patients, and interpreters, as well as providing psychoeducation and practical support.

22. Guregard S, Seikkula J. Establishing therapeutic dialogue with refugee families. *Contemp Fam Ther* 2013. doi 10.1007/s10591-013-9263-5; Advance online publication.

23. Forcada I, Pera V, Cruz I, *et al.* Comparison of immigrant and native-born population adherence to antipsychotic treatment in a Spanish health region. *Commun Ment Health J* 2013; 49:199–205.

This study investigated adherence to antipsychotic pharmacological treatment of native and immigrant patients in Spain. At 1-year follow-up, adherence among immigrants was lower than in native patients, particularly in groups such as sub-Saharan Africans.

24. Park SY, Cho S, Park Y, *et al.* Factors associated with mental health service utilization among Korean American immigrants. *Commun Ment Health J* 2013; 49:765–773.

25. Piwowarczyk L, Bishop H, Saia K, *et al.* Pilot evaluation of a health promotion program for African immigrant and refugee women: the UJAMBO program. *J Immigrant Minority Health* 2013; 15:219–223.

26. Ruiz E, Aguirre RTP, Mitschke DB. What leads non-U.S.-Born Latinos to access mental healthcare? *Soc Work Healthcare* 2013; 52:1–19.

27. Poon WC, Joubert L, Harvey C. Experiences of Chinese migrants caring for family members with schizophrenia in Australia. *Soc Work Healthcare* 2013; 52:2–3; 144-165.

28. Harpelund L, Smith Nielsen S, Krasnik A. Self-perceived need for interpreter among immigrants in Denmark. *Scand J Public Health* 2012; 40:457.

This study investigated self-perceived need for an interpreter in GP consultations amongst large non-western immigrant groups in Denmark. The authors concluded that proposed measures for patients paying fees for using interpreters are likely to affect immigrants with modest income, poor health, and who are outside the labor force.

- 29.** Guerrero EG, Campos M, Urada D, Yang JC. Do cultural and linguistic competence matter in Latinos' completion of mandated substance abuse treatment? *Subst Abuse Treat Prev Policy* 2012; 7:34.

This study analyzed client and program data from publicly funded mandatory substance abuse treatment programs in California to investigate whether culturally and linguistically responsive practice improved outcomes for Latin-American patients. The findings established that the availability of Spanish language translators was associated with increased odds of completing the treatment.

- 30.** Tarricone I, Stivanello E, Ferrari S, *et al.* Migrant pathways to community mental health centres in Italy. *Int J Soc Psychiatry* 2012; 58:505–511.

This study compared migrant pathways to four community mental health centers in different provinces in Italy and identified predictors of pathways to care. The findings demonstrated frequent use of nonmedical pathways to care among immigrants, particularly those without a resident's permit.

- 31.** Priebe S, Matanov A, Barros H, *et al.* Mental health-care provision for marginalized groups across Europe: findings from the PROMO study. *Eur J Public Health* 2013; 23:97–103.

This study investigated the provision of mental healthcare for socially marginalized groups, including asylum seekers, refugees, and undocumented migrants in highly deprived areas of eight European capitals. The study highlighted similarities and differences in care provision between generic and group-specific services. The latter services are often found in voluntary and not-for-profit sectors and tend to offer more outreach programs and social care. The authors emphasized the need for more coordination between these two groups of services.

- 32.** Snowden LR, McClellan SR. Spanish-language community-based mental health treatment programs, policy-required language-assistance programming, and mental health treatment access among Spanish-speaking clients. *Am J Public Health* 2013; 103:1628–1633.

This study investigated the effect of implementing language assistance programming on improving accessibility of mental healthcare under Medi-Cal for Spanish-speaking persons in California. The results suggest that the presence of language assistance programs moderately improved access to mental healthcare, but only when implemented by community-based organizations.

- 33.** Kluge U, Bogic M, Devillé W, *et al.* Health services and the treatment of immigrants: data on service use, interpreting services and immigrant staff members in services across Europe. *Eur Psychiatry* 2012; 27:S56–S62.

This study surveyed 240 primary, emergency, and mental health services across 16 EU countries. Availability of data on service use by immigrants, the provision of interpreting services, and immigrant staff numbers were recorded. The findings highlighted scarcity of data on immigrants, patients' service use, and a high number of services, which did not provide any interpreting and did not employ any members of staff with immigrant background. Mental health services were more likely to provide face-to-face interpreting and employ immigrant staff.

- 34.** Gallardo ME. Context and culture: the initial clinical interview with the Latina/o client. *J Contemp Psychother* 2013; 43:43–52.

This study investigated bilingual and bicultural Latin-American psychotherapists and the ways they engaged with immigrant patients. They reported using culturally accepted styles of communication, language, and self-disclosure to build rapport with patients from the same background.

- 35.** Ahmad F, Shakya J, Li J, *et al.* A pilot with computer-assisted psychosocial risk-assessment for refugees. *BMC Med Inform Decis Mak* 2012; 12:7.

This study was a pilot RCT, using a computer-assisted tool to assess Afghani refugees in community mental health centers in Canada. The findings suggest that the tool offers a promising model to integrate medical and social care to address health and settlement needs of refugees.

- 36.** Vincent F, Jenkins H, Larkin M, Clohessy S. Asylum-seekers' experiences of trauma-focused cognitive behaviour therapy for posttraumatic stress disorder: a qualitative study. *Behav Cogn Psychother* 2013; 41:579–593.

This study explored asylum seekers' experiences of trauma-focused cognitive behavioral treatment for PTSD. The findings highlighted that fear of repatriation may impede engagement in therapy, but that some refugee patients still find therapy beneficial.

- 37.** Green H, Sperling D, Carsswel K. Too close to home? Experiences of Kurdish refugee interpreters working in UK mental health services. *JMH*; 21:227–235.

This study qualitatively assessed experiences of refugee interpreters working in UK mental health services. The findings highlighted the challenges interpreters experienced in clinical encounters, and emphasized the need for supervision and training in mental health and self-care.

- 38.** Kwong K, Chung H, Cheal K, *et al.* Depression care management for Chinese Americans in primary care: a feasibility pilot study. *Communit Ment Health J* 2013; 49:157–165.

- 39.** Ratzliff ADH, Ni K, Chan YF, *et al.* A collaborative care approach to depression treatment for Asian Americans. *Psychiatr Serv* 2013; 64:487–490.

This study examined the effectiveness of collaborative care for depression provided to Asian immigrants in the USA in either culturally sensitive or general community health centers. Matched non-Asian population receiving collaborative care in the same general health centers was also assessed. All three groups had similar treatment process and depression outcomes, demonstrating effectiveness of collaborative care for immigrant populations across different settings.

- 40.** Keiffer EC, Caldwell CH, Welmerink DB, *et al.* Effect of the healthy MOMs lifestyle intervention on reducing depressive symptoms among pregnant Latinas. *Am J Commun Psychol* 2013; 51:76–89.

- 41.** O'Shaughnessy R, Nelki J, Chiumento A, *et al.* Sweet mother: evaluation of a pilot mental health service for asylum-seeking mothers and babies. *J Pub Mental Health* 2012; 11:214–228.

- 42.** Valdez CR, Padilla B, Mcardell Moore S, Magana S. Feasibility, acceptability, and preliminary outcomes of the Fortalezas Familiares intervention for Latino families facing maternal depression. *Fam Proc* 2013; 52:394–410.

This study investigated the feasibility and acceptability of community-based, multifamily group intervention for Latin-American families facing maternal depression. The intervention resulted in improved psychological functioning, and increased family and marital support.

- 43.** Kung WW, Tseng YF, Wang Y, *et al.* Pilot study of ethnically sensitive family psychoeducation for Chinese American patients with schizophrenia. *Soc Work Mental Health* 2013; 10:384–408.

This pilot study tested the feasibility and acceptability of a culturally sensitive, family psychoeducation protocol for Chinese-American adult schizophrenic patients and their caregivers. The protocol was well received, leading to improvements in both patient and carer outcomes. The flexibility of implementation was found to be of great importance.

- 44.** Moreno FA, Chong J, Dumbaule J, *et al.* Use of standard Webcam and Internet equipment for telepsychiatry treatment of depression among underserved Hispanics. *Psychiatr Serv* 2012; 63:1213–1217.

This study investigated the effectiveness of depression treatment for Hispanic patients delivered by a psychiatrist via Internet videoconferencing compared to treatment as usual by a primary care provider. Results suggest that telepsychiatry delivered through the Internet using domestic Webcams is effective and acceptable.

- 45.** Unlu Ince B, Cuijpers P, van Hof E, *et al.* Internet-based, culturally sensitive, problem-solving therapy for Turkish migrants with depression: randomized controlled trial. *J Med Internet Res* 2013; 15:e227. doi: 10.2196/jmir.2853; Advanced Online Publication.

This study evaluated the effectiveness of culturally sensitive, self-help, problem-solving intervention delivered via the Internet to Turkish immigrants with depression. There was no significant effect on depressive symptoms compared to a waiting list control groups, but a completers-only analysis indicated possible effectiveness of the intervention when assessed in a larger sample.

AQ10

AQ11

AQ9



Dear Author,

During the preparation of your manuscript for typesetting, some queries have arisen. These are listed below. Please check your typeset proof carefully and mark any corrections in the margin as neatly as possible or compile them as a separate list. This form should then be returned with your marked proof/list of corrections to the Production Editor.

## QUERIES: to be answered by AUTHOR/EDITOR

QUERY NO.	QUERY DETAILS	RESPONSE
<AQ1>	Please provide the expanded form of EUGATE.	
<AQ2>	Please provide the expanded forms of PTSD, GP, RCT.	
<AQ3>	Please check whether the intended meaning of the following sentence is retained after making the edits. 'A survey of public outpatient psychotherapists in Hamburg found that only 14% of them were immigrants, whereas 27% of the population in the area was of such background'.	
<AQ4>	Please check whether the intended meaning of the following sentence is retained after making the edits. 'Seventy-two percent of therapists agreed that advanced training in cross-cultural competence would be helpful, whereas 9% had already completed some form of such training'.	
<AQ5>	Please check whether the intended meaning of the following sentence is retained after making the edits. 'The latter tend to offer more outreach programs and social support, whereas medical interventions were mostly provided by generic services'.	
<AQ6>	Please check the author names in Refs.[1,3,9,13,18,23,24,25,30,31,33,38,40,41,43,45].	
<AQ7>	The page range appearing in Ref.[7] is invalid. Please check.	
<AQ8>	Please update Refs. [15,22], if possible, by providing complete publication details such as volume, year of publication, and page range.	
<AQ9>	The intended meaning of the following sentence is not clear. Please check. 'The findings highlighted scarcity of data on immigrants, patients' service use,	

and a high number of services, which did not provide any interpreting and did not employ any members of staff with immigrant background'.

<AQ10> Please provide the year of publication in Ref.[37].

<AQ11> Please check the page range in Ref.[44].