

Kent Academic Repository

Full text document (pdf)

Citation for published version

Wisman, Arnaud and Shrira, Ilan (2015) The Smell of Death: Evidence that Putrescine Elicits Threat Management Mechanisms: The Smell of Death. *Frontiers in Psychology* . pp. 1-26. ISSN 1664-1078.

DOI

<http://doi.org/10.3389/fpsyg.2015.01274>

Link to record in KAR

<http://kar.kent.ac.uk/50211/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

The Smell of Death: Evidence that Putrescine Elicits Threat Management Mechanisms

Arnaud Wisman and Ilan Shrira

Journal Name: Frontiers in Psychology
ISSN: 1664-1078
Article type: Original Research Article
Received on: 27 May 2015
Accepted on: 10 Aug 2015
Provisional PDF published on: 10 Aug 2015
Frontiers website link: www.frontiersin.org
Citation: Wisman A and Shrira I(2015) The Smell of Death: Evidence that Putrescine Elicits Threat Management Mechanisms. *Front. Psychol.* 6:1274. doi:10.3389/fpsyg.2015.01274
Copyright statement: © 2015 Wisman and Shrira. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](http://creativecommons.org/licenses/by/2.0/). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after rigorous peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

The Smell of Death:

Evidence that Putrescine Elicits Threat Management Mechanisms

Arnaud Wisman^{1*}, Ilan Shrira²

¹School of Psychology, University of Kent, Canterbury, Kent, UK

²Department of Behavioral Sciences, Arkansas Tech University, Russellville, AR, USA

*Correspondence:

Dr. Arnaud Wisman
School of Psychology
University of Kent
Canterbury
Kent CT2 7NP
United Kingdom
a.wisman@kent.ac.uk

Keywords: olfaction, putrescine, threat, threat management, chemosensory cue

Abstract

The ability to detect and respond to chemosensory threat cues in the environment plays a vital role in survival across species. However, little is known about which chemical compounds can act as olfactory threat signals in humans. We hypothesized that brief exposure to putrescine, a chemical compound produced by the breakdown of fatty acids in the decaying tissue of dead bodies, can function as a chemosensory warning signal, activating threat management responses (e.g., heightened alertness, fight-or-flight responses). This hypothesis was tested by gauging people's responses to conscious and non-conscious exposure to putrescine. In Experiment 1, putrescine increased vigilance, as measured by a reaction time task. In Experiments 2 and 3, brief exposure to putrescine (vs. ammonia and a scentless control condition) prompted participants to walk away faster from the exposure site. Experiment 3 also showed that putrescine elicited implicit cognitions related to escape and threat. Experiment 4 found that exposure to putrescine, presented here below the threshold of conscious awareness, increased hostility toward an out-group member. Together, the results are the first to indicate that humans can process putrescine as a warning signal that mobilizes protective responses to deal with relevant threats. The implications of these results are briefly discussed.

1. Introduction

When animals die they release an unpleasant smell. A pungent component of this scent is emitted by putrescine, a volatile diamine that results from the breakdown of fatty acids in the putrefying

1 tissue of dead bodies (Hussain et al., 2013). Interestingly, animal research shows that putrescine
2 can function as a powerful chemosensory signal that prompts the perceiver to leave or avoid the
3 area (Prounis & Shields, 2013; Yao et al., 2009). The aim of the present research is to show that
4 humans respond in a similar way to putrescine, and more generally, that exposure to putrescine
5 triggers threat management behaviors (Blanchard et al., 2001; Neuberg et al., 2011).

6
7 A growing body of research suggests that humans can identify threats via chemosignals
8 (Ackerl et al., 2002; Chen & Haviland-Jones, 2000; de Groot et al., 2012; Mujica-Parodi et al.,
9 2009; Prehn et al., 2006; Zhou & Chen, 2009). For instance, when people are exposed to sweat
10 taken from donors during a fearful experience, perceivers show a heightened startle reflex (Pause
11 et al., 2009; Prehn et al., 2006) and interpret ambiguous facial expressions as fearful (Zhou &
12 Chen, 2009). This transmission of threat-arousing chemosignals is assumed to serve an adaptive
13 function by orienting us to impending dangers. Indeed, the ability to detect and process
14 chemosensory threat cues is vital for the survival of a wide range of species (Stevenson, 2010).
15 However, thus far there is little evidence that humans can, like other organisms, detect olfactory
16 threat cues in the environment through means other than the chemosignals (e.g., body sweat) of
17 conspecifics.

18
19 The decay of tissue and its resulting scent can function as a “necromone” cue that signals
20 an animal’s death to conspecifics. Alarm and avoidance behaviors (necrophobic behaviors) in
21 response to these scents are widespread in the animal kingdom and thought to have evolved at
22 least 420 million years ago (Yao et al., 2009). In fact, recent research shows that necrophobic
23 behavior may have innate underpinnings through the activation of trace amine-associated
24 receptors (TAARs), a group of specialized scent receptors in the olfactory epithelium (Horowitz
25 et al., 2014; Hussain et al., 2013; Li & Liberles, 2015). TAARs are known to detect specific
26 chemicals that evoke behavioral responses, without the need for prior exposure to the scents. For
27 example, in model vertebrates, certain TAARs respond to diamines (e.g., putrescine) by
28 producing avoidant behaviors that likely serve to defend against immediate dangers (Yoon et al.,
29 2015). Thus, it is feasible that we have a chemosensory sensitivity to diamines like putrescine (Li
30 & Liberles, 2015), given that their detection can aid survival (Stevenson, 2010).

31
32 A further advantage of examining putrescine as a threat stimulus is that we know what it
33 is. Despite the impressive amount of indirect support for human chemosignals amassed in recent
34 years, their chemical properties have yet to be identified (Wyatt, 2009). Focusing on a known
35 compound, putrescine, enables us to directly test whether it plays a causal role in human threat
36 responses. In a similar vein, although several studies have shown that chemosensory cues can
37 elicit greater readiness for behavior (Bradley et al., 2001; Prehn et al., 2005), thus far there is
38 little direct evidence that a specific chemical substance can cause overt behavioral changes in
39 humans (Wysocki & Preti, 2004). Since exposure to putrescine elicits specific behaviors in
40 animals (e.g., escape, avoidance), we can examine whether putrescine produces similar behaviors
41 in humans. In sum, putrescine appears to be well-suited to test as a specific chemical compound
42 that can act as a threat signal in humans.

43
44 Chemosensory cues can convey danger in at least two fitness-relevant domains: microbial
45 and predator threats (Stevenson, 2010). First, olfactory information is often central to identifying
46 the presence of pathogens. For example, pathogens can alter the scent of those who become

1 infected, which can be detected by conspecifics (Arakawa et al., 2010; Olsson et al., 2014; Tybur
2 et al., 2011). Similarly, the release of putrescine in decaying tissue co-occurs with the arrival of
3 bacteria, a motivation for others to eschew physical contact with the dead body. A number of
4 species exhibit necrophobic behaviors, and after detecting the scent emanating from dead bodies,
5 usually respond by leaving or avoiding the area (Prounis & Shields, 2013). Second, putrescine
6 released by decaying bodies can signal the risk of predation (Boissy et al., 1986). Since a large
7 proportion of deaths in the wild are the result of predator attacks, putrescine would be a useful
8 alarm cue to stay away (Misslin, 2003).

9
10 In humans, responses to specific scents can develop through learned associations between
11 odors and personal experiences (Degel et al., 2001; Stevenson et al., 1998). For example, based
12 on the cultural expression that when “something smells fishy” it is viewed suspiciously, exposure
13 to fish-like odors arouses suspicion toward others and reduces cooperation, an orientation that is
14 assumed to result from conditioned reactions to this scent (Lee & Schwarz, 2012). Since
15 putrescine can emanate from various sources (Yeoman et al., 2013), people may learn to
16 associate the smell of putrescine with threats, and it is plausible that occasional exposure to
17 putrescine, whenever it occurs, could lead to conditioned threat responses (Stevenson, 2010).
18 However, we render it unlikely that modern humans have strong conscious meaningful
19 associations with the scent of putrescine. Moreover, conscious scent evaluations are often
20 inaccurate, context dependent, and colored by other sensory modalities (Sela & Sobel, 2010). In
21 view of this, it is important to note that responses to aversive chemosensory cues do not require
22 prior learning or conscious evaluation (Dielenberg et al., 2001; Li et al., 2013; Miller & Maner,
23 2010). Indeed, scents can alter our perception, cognition, behavior, and physiology (e.g., heart
24 rate, skin conductance) even when there is no conscious scent detection (Krusemark & Li, 2012;
25 Li et al., 2007; Pause et al., 2009; Sela & Sobel, 2010), and even after olfactory adaptation has
26 set in (de Groot et al., 2012; Smeets & Dijksterhuis, 2014). Thus, neither prior associations with
27 olfactory signals, nor conscious processing, are necessary conditions for people to process them
28 as threatening (Pause, 2012; Williams et al., 2006; Köster, 2002; Sela & Sobel, 2010; Smeets &
29 Dijksterhuis, 2014).

30
31 At the most basic level, threat detection increases vigilance and sharpens our reactions to
32 events in the environment (Williams et al., 2006). For instance, detection of a predator’s scent
33 will interrupt foraging and increase behaviors (e.g., scanning the environment) that facilitate
34 predator detection (Woody & Szechtman, 2011). Once the threat management system is
35 engaged, it produces readiness for fight-or-flight behaviors (Blanchard et al., 1986; Cannon,
36 1927; Gray & McNaughton, 2003; Mobbs et al., 2009). Flight responses seek to escape the
37 situation, whereas fight responses—whether physical or verbal aggression—are typically only
38 used when escape is not possible. In contrast to popular belief that the dominant response to
39 threats is to fight, flight is actually far more common (Misslin, 2003), presumably because nature
40 selects more strongly for strategies that minimize risk. In one study, for example, when people
41 were confronted by a threatening out-group member, they responded with aggressive readiness
42 (fight), but only when there was little possibility of escaping; when given the option, though,
43 participants chose to distance themselves (flight) from the other person (Cesario et al., 2010).

44 45 **2. Overview and Hypotheses**

46

1 Coming full circle, we propose that putrescine can serve as a (non-conscious) signal that initiates
2 threat management responses. Specifically, we hypothesize that brief exposure to putrescine
3 increases vigilance, followed by the readiness to either escape (flight), or engage in aggressive
4 readiness (fight) when escape is not possible. Experiment 1 assessed whether putrescine (vs.
5 ammonia and a neutral scent) increased vigilance as measured by faster responses in a simple
6 reaction time task. Experiments 2 and 3 assessed whether brief exposure to putrescine (vs.
7 ammonia and neutral scent) caused participants to walk away faster from the exposure site after
8 completing the experiment (outdoors). Experiment 3 also tested whether putrescine evoked
9 cognitions related to escape and threat. Finally, Experiment 4 examined whether non-conscious
10 exposure to putrescine increased aggressive readiness (e.g., defensiveness toward an out-group
11 member). All four experiments adhered to the Declaration of Helsinki guidelines, and gained the
12 prior approval by the University Research Ethics Committee. Written consent was obtained from
13 all participants involved in these experiments, and all were fully debriefed.

14

15 **3. Experiment 1: The effect of putrescine on vigilance**

16

17 In Experiment 1, we tested whether brief exposure to putrescine increased vigilance. To measure
18 vigilance, we employed a task closely modeled after the shortened version of the Psychomotor
19 Vigilance Task (PVT; Dinges & Powell, 1985) that assessed participants' reaction times to a red
20 dot that was presented at random intervals on a computer screen.

21

22 In addition, Experiment 1 was designed to determine whether ammonia served as an
23 appropriate aversive control condition. Our pilot testing revealed that ammonia, unlike other
24 aversive scents we had examined (i.e., skatole¹ and indole), was rated similarly to putrescine on
25 repugnance, familiarity, and intensity. Moreover, previous research has used ammonia (NH₃;
26 ammonium hydroxide) as an aversive scent prime (Rieser et al., 1976; Wise et al., 2005).
27 Furthermore, ammonia can increase trigeminal nerve activation associated with vigilance and
28 sensory rejection, via activation of the sympathetic nervous system (Hummel & Kobal, 1992;
29 Sekizawa & Tsubone, 1994). However, some research suggests that unpleasant ambient odors
30 can also decrease reaction times on simple tasks like the current PVT (Millot et al., 2002). In
31 view of this, we made no specific prediction about whether ammonia, like putrescine, would
32 enhance vigilance relative to our scentless control condition.

33

34 **3.1. Method**

35

36 **3.1.1. Participants and Procedure**

37

38 A sample of sixty participants (43 females; $M_{\text{age}} = 21.20$, $SD = 3.20$) completed the study in
39 return for a financial incentive of 3 pounds (approximately \$5).

40

¹ In line with previous research (Wheatley & Haidt, 2005), we pilot-tested the so-called “fart spray” along with skatole, indole, and ammonia, for suitability as an aversive control condition. These ratings are presented in Table 1. As can be seen, ammonia and fart spray were rated similarly to putrescine on all three dimensions of repugnance, familiarity, and intensity, whereas indole and skatole diverged from putrescine on at least one dimension. A disadvantage of fart spray, however, is that we could not ascertain its precise chemical compounds—its manufacturers were reluctant to disclose this information.

1 Participants were randomly assigned to one of three conditions: putrescine ($C_4H_{12}N_2$;
2 Sigma-Aldrich), ammonia (5%; NH_3 ; Sigma-Aldrich), or water. One hour before the start of the
3 experiment, cotton wool pads were blotted with 2 ml of one of the three compounds, and stored
4 separately in small (100 ml) sealable amber jars. Participants were run in our lab individually,
5 and seated in different cubicles to avoid carryover effects of scents. The refreshment rate in each
6 cubicle was 4 to 5 air changes (cycles) per hour. Furthermore, participants were booked at least
7 30 minutes apart in order to ventilate the rooms—by opening the lab room’s window—between
8 sessions. When preparing materials for the experiment, one of the researchers marked the bottom
9 of each jar with a number code, so that the experimenters were unaware of the meaning of these
10 codes. This basic procedure was repeated in our subsequent experiments to keep the
11 experimenters blind to the conditions.

12 Participants were seated in front of a standard PC (equipped with Authorware 7.1
13 software) with a 17-inch screen. They were given instructions (on-screen) to open the jar, sniff
14 the scent inside for 10 seconds, and close the jar. After that, they rated the scent on its intensity
15 (“This scent is intense”; 1 = strongly disagree and 9 = strongly agree), repugnance (“This scent
16 is repugnant”; 1 = strongly disagree and 9 = strongly agree), and familiarity (“This scent is
17 familiar”; 1 = strongly disagree and 9 = strongly agree). Repugnance was included as evaluative
18 rating (alongside the standard measures of intensity and familiarity) because repugnance (or
19 disgust) is often a central component of aversive scents. Participants were then introduced to the
20 adapted PVT, which lasted about five minutes (see Loh et al., 2004). The task instructed them to
21 click on a red dot as quickly as possible whenever they saw the dot on the screen. Ten dots (each
22 measuring 1 cm) were shown at different locations on the screen, and the time between
23 appearances was randomized at variable intervals (2-45 sec). As soon participants clicked on the
24 red dot with the mouse, a screen appeared for five seconds with the message: “prepare for next
25 trial”. Participants received two practice trials first, to get them familiar with the main task of ten
26 trials. Finally, after completing the PVT and filling out a standard demographic questionnaire,
27 they were fully debriefed and thanked for their participation.

28 29 **3.2. Results and Discussion**

30 31 **3.2.1. Hedonic Evaluations**

32
33 We began by testing our prediction, based on our pilot testing, that putrescine and ammonia
34 would not differ from each other on repugnance, familiarity and intensity. As predicted, separate
35 one-way between-subjects ANOVAs revealed that there was no significant difference between
36 ammonia and putrescine on repugnance, $F(1, 38) = 0.38$, $p = .54$, $\eta^2 = .01$, familiarity, $F(1, 38) =$
37 0.26 , $p = .26$, $\eta^2 = .03$, or intensity, $F(1, 38) = 0.14$, $p = .71$, $\eta^2 = .004$ (see Table 2, for
38 descriptive statistics). Moreover, the analyses reported below were not altered when entering all
39 hedonic evaluations as covariates.

40 41 **3.2.2. Reaction Times**

42
43 We examined our main prediction that putrescine, relative to the neutral control condition
44 (water), would elicit faster reaction times. In line with previous PVT research, we applied
45 reciprocal transformation to the raw data (i.e., $1/RT$). This type of transformation is standard
46 within the PVT paradigm, as it reduces the impact of extreme scores and brings them into an

1 acceptable range (Dinges et al., 1987; Dorrian et al., 2004). A one-way between-subjects
 2 ANOVA revealed a difference between the scent conditions, $F(2, 57) = 4.32, p = .018, \eta^2 = .13$.
 3 Post hoc comparisons, with the raw means reported here, showed that putrescine produced faster
 4 reaction times ($M = 1.04, SD = .10$) than the neutral scent ($M = 1.24, SD = .35; p = .013$), but not
 5 compared to ammonia ($M = 1.12, SD = .20; p = .28$). No difference was found between the
 6 neutral and ammonia conditions ($p = .14$).

7
 8 In sum, only putrescine caused participants to react more quickly compared to the neutral
 9 condition, supporting our hypothesis that putrescine increases vigilance. At the same time,
 10 ammonia did not increase vigilance relative to the scentless control condition. Importantly, the
 11 findings show that, consistent with our pilot study, ammonia and putrescine are evaluated similar
 12 on repugnance, familiarity, and intensity, and were similar in the degree of vigilance they
 13 elicited. Consequently, together with previous research (Rieser et al., 1976; Wise et al., 2005),
 14 Experiment 1 indicated that ammonia would serve as an appropriate aversive control condition.
 15 Experiments 2 and 3 investigated our hypothesis that putrescine activates the motivation to
 16 escape the situation (flight).

17 18 **4. Experiment 2: The effect of putrescine on escape behavior**

19
 20 Similar to Experiment 1, Experiment 2 first asked participants to rate a scent prime (putrescine
 21 vs. ammonia vs. neutral) on three dimensions: intensity, familiarity, and repugnance, then we
 22 observed whether it influenced the tendency to escape the situation. To avoid the biases
 23 associated with some operationalizations of flight in prior research (e.g., self-reported intentions,
 24 Gilbert & Gilbert, 2003), we employed an overt behavioral measure of escape (e.g., Ellsworth et
 25 al., 1972; Wisman & Koole, 2003). Specifically, we assessed whether putrescine would cause
 26 participants (who were under the impression the study was finished) to walk away more quickly
 27 over a predetermined distance of 80 meters.

28 29 **4.1. Method**

30 31 **4.1.1. Participants and Procedure**

32
 33 Forty-five participants (21 females and 24 males; $M_{age} = 27.51, SD = 9.72$) completed the study
 34 on campus. We filled three empty felt-tip pens, each with one of the three compounds
 35 (putrescine, ammonia, or water). To fill each pen, 10ml of liquid odor was injected onto the
 36 pen's fiber rod inside the pen. The pens were then re-assembled and left to stand upside down for
 37 24 hours in order to allow the liquid to soak into the fiber rod. Just before the start of the
 38 experiment, scent blotters were marked with the scent marker pens and stored in separate
 39 sealable containers.

40
 41 Participants were approached on a fixed spot on the campus and asked if they had time to
 42 participate in a brief scent test of approximately ten minutes. Participants were tested
 43 individually and randomly assigned to one of three conditions (putrescine, ammonia, or water).
 44 The experimenter, blind to the conditions, presented one of the three containers to the
 45 participant, who rated the scent on intensity ("This scent is strong"; 1 = strongly disagree and 5
 46 = strongly agree), repugnance ("This scent is repugnant"; 1 = strongly disagree and 5 = strongly

1 agree), and familiarity (“This scent is familiar”; 1 = strongly disagree and 5 = strongly agree).
2 After finishing and being thanked for their participation, a second experimenter blind to the
3 condition and hypotheses of the experiment and out of sight of the participants— used a standard
4 stopwatch to time how many seconds it took participants to walk away over a distance of 80
5 meters (pre-measured before the experiment began). The recorded time constituted our
6 dependent variable. After they reached this distance, participants were re-approached, fully
7 debriefed and thanked again.

8 9 **4.2. Results and Discussion**

10 11 **4.2.1. Hedonic Evaluations**

12
13 Consistent with Experiment 1, separate one-way between-subjects ANOVAs revealed that there
14 was no significant difference between ammonia and putrescine on repugnance, $F(1, 28) = 2.30$, p
15 $= .14$, $\eta^2 = .07$, and familiarity, $F(1, 28) = 0.04$, $p = .75$, $\eta^2 = .01$. However, ammonia was rated
16 as relatively more intense ($M = 4.73$; $SD = 0.46$) compared to putrescine ($M = 4.27$; $SD = 0.70$; p
17 $= .04$; see Table 3). Once again, the results reported below were not altered when we entered the
18 intensity (nor the other hedonic) ratings into the analyses as covariates. We also note that the
19 results are similar whether participants rate how “intense” or “strong” the scent smells (see
20 Experiment 3 below).

21 22 **4.2.2. Escape Behavior**

23
24 To test our hypothesis that putrescine elicited an escape motivation, we compared our scent
25 conditions in a one-way ANOVA, using gender as a covariate². The results yielded a significant
26 effect of the scent prime on the time it took to walk 80 meters, $F(2, 41) = 19.03$, $p < .001$, $\eta^2 =$
27 $.48$. The only significant differences occurred between putrescine ($M = 56.40$ seconds; $SD =$
28 4.19) and ammonia ($M = 59.93$, $SD = 5.04$), and between putrescine and the neutral scent prime
29 ($M = 60.00$, $SD = 4.42$; both $ps < .005$; see Figure 1). Thus, putrescine caused participants to
30 walk away more quickly, supporting our assumption that putrescine evoked a stronger
31 motivation to escape. Experiment 3 was conducted to replicate this finding, and furthermore to
32 test whether putrescine elicited implicit cognitions related to escape and threat.

33 34 **5. Experiment 3: The effect of putrescine on escape behavior and thoughts**

35
36 The procedure for Experiment 3 was similar to Experiment 2’s. First, we asked participants to
37 evaluate the scents on the different dimensions (repugnance, familiarity, intensity). In addition,
38 we gauged participants’ implicit threat-related associations using a word stem-completion task.
39 Specifically, this task measured the implicit accessibility of thoughts related to “escape” and
40 “threat.” We predicted that only putrescine would increase the accessibility of these cognitions.
41 Finally, we assessed whether putrescine would cause participants to walk away more quickly
42 over a predetermined distance of 60 meters.

² Because previous research has shown that men and women tend to walk at different speeds (Chumanov, Wall-Scheffler, & Heiderscheit, 2008), the results of Experiments 2 and 3 included gender as a covariate. In addition, we analyzed the results of Experiments 2 and 3 with gender as a separate factor and this did not alter the significance of the results.

5.1. Method

5.1.1. Participants and Procedure

Sixty participants (32 females and 28 males, $M_{\text{age}} = 21.57$, $SD = 1.12$) completed the study on campus. Individuals were approached just outside campus on a path sloping downhill and asked if they had time to participate in a brief scent test for about 15 minutes.

Participants were randomly assigned to one of the three scent conditions, then they rated the scent on intensity, repugnance, and familiarity (“This scent is intense”; 1 = strongly disagree and 9 = strongly agree), repugnance (“This scent is repugnant”; 1 = strongly disagree and 9 = strongly agree), and familiarity (“This scent is familiar”; 1 = strongly disagree and 9 = strongly agree). Then, to assess cognitions relevant to the concepts of “escape” and “threat,” participants completed the word-stem completion task, a widely used and well-established measurement that gauged the thought accessibility of these two concepts (Arndt et al., 1997; Greenberg et al., 1994; Lozito & Mulligan, 2010; Migo et al., 2010). Participants were asked to complete 30 word fragments, 20 of which were neutral (e.g., B_ NK could be BANK or BUNK) in terms of any particular theme, five of which could be words related to “escape” (e.g., the fragment RU_ could be completed as RUN or RUB, the latter a neutral word), and another five could be completed with a word related to “threat” (e.g., _ _ RROR could be TERROR or MIRROR). We summed the number of escape- ($M = 2.73$, $SD = 1.07$) and threat-related words ($M = 1.90$, $SD = .66$) that participants completed to assess the thought accessibility of these concepts. Finally, participants were again timed by a second experimenter, who was blind to the conditions and the hypotheses, for how long it took them to walk away over a distance of 60 meters (Due to natural constraints a slightly shorter distance than in Experiment 2).

5.2. Results and Discussion

5.2.1. Hedonic Evaluations

Separate one-way between-subjects ANOVAs revealed no difference between the chemosensory primes on repugnance, $F(1, 38) = .35$, $p = .56$, $\eta^2 = .01$, familiarity, $F(1, 38) = .04$, $p = .85$, $\eta^2 = .001$, and intensity, $F(1, 38) = 0.29$, $p = .59$, $\eta^2 = .008$ (see Table 4). Thus, participants rated ammonia and putrescine similarly to one another on intensity, repugnance, and familiarity. Again, the results reported below were did not differ when we entered the hedonic evaluations into the analyses as covariates.

5.2.2. Escape- and Threat-Related Cognitions

To test our hypothesis that putrescine elicited implicit cognitions related to escape and threat, we analyzed the escape and threat word-completion results separately. The results revealed a significant effect of scent prime on escape thought accessibility, $F(2, 57) = 10.90$, $p < .001$, $\eta^2 = .28$ (see Table 5). Putrescine caused participants to complete word stems more frequently with escape related words ($M = 3.45$, $SD = .69$) than both the ammonia ($M = 2.45$, $SD = 1.05$) and the neutral scent ($M = 2.15$, $SD = .99$) primes (both $ps < .005$). Similarly, the scent primes affected

1 the accessibility of threat-related thoughts, $F(2, 57) = 8.39, p < .001, \eta^2 = .23$. Putrescine led to
 2 more threat word-stem completions ($M = 2.55, SD = .94$) than ammonia ($M = 1.73, SD = .64$)
 3 and the neutral scent ($M = 1.68, SD = .65$; all $ps < .005$).

5 5.2.3. Escape Behavior

6
 7 Like Experiment 2, the analyses showed a significant effect of chemosensory primes on walking
 8 speed, $F(2, 56) = 9.11, p < .001, \eta^2 = .24$ (see Figure 2). The pattern of results again showed that
 9 putrescine ($M = 33.38, SD = 2.99$) caused people to walk more quickly than ammonia ($M =$
 10 $35.92, SD = 3.38$) and the neutral scent prime ($M = 37.67, SD = 3.13; p < .05$). Again, no
 11 difference was found between the ammonia and the neutral scent condition ($p = .87$).

12
 13 Experiment 3 revealed that putrescine elicited implicit cognitions of escape and threat. In
 14 addition, Experiment 3 replicated the finding that putrescine increased walking speed. Thus,
 15 taken together, the results of Experiments 2 and 3 indicated that putrescine motivated (automatic)
 16 escape behavior. An important feature of the settings in Experiments 2 and 3 was that
 17 participants were outdoors and in a context that facilitated the possibility that they could distance
 18 themselves from the scent.

20 6. Experiment 4: The effects of putrescine on defensive responses toward an out-group

21
 22 Experiment 4 sought to extend our understanding of the effects of putrescine in two important
 23 respects. First, we tested the hypothesis that non-conscious (unobtrusive) exposure to putrescine
 24 could elicit threat management responses. As we highlighted in the Introduction, this possibility
 25 is consistent with evidence that scent primes, even when presented at sub-threshold levels, can
 26 influence brain activation (Sobel et al., 1999), learning (Koster et al., 2002), and physiological
 27 state (Stern & McClintock, 1998). This applies similarly to aversive scent primes, which for
 28 example, have the ability to alter skin conductance (Jacquot et al., 2004), social preferences (Li
 29 et al., 2007), and cognitive performance (Epple & Herz, 1999) in ways that correspond to
 30 supraliminal exposure to aversive stimuli (Sela & Sobel, 2010). Thus, we predicted that
 31 subliminal presentation of putrescine would be capable of activating threat responses.

32
 33 Second, Experiment 4 focused on the fight rather than the flight component of alarm
 34 responses. Consistent with previous research showing that implicit threat cues increase
 35 intolerance toward out-group members (Holbrook et al., 2011) and defensive responses
 36 (Blanchard et al., 2001; Wheatley & Haidt, 2005), we hypothesized that putrescine would
 37 increase defensiveness toward an out-group member, in a situation where there was no
 38 immediate opportunity to escape (Cesario et al., 2010). Like Experiment 1, we conducted this
 39 experiment in a laboratory setting. After priming the participants with one of the scents, they
 40 filled out a standard positive and negative affect scale that gauged their mood. Although, our
 41 pilot study (see Table 1) and some research (e.g., Knasko, 1993) revealed that aversive scent
 42 primes do not alter mood on a conscious level, we intended to rule out the possibility that the
 43 subliminal primes influenced participants' feelings at a conscious level. After that, they read
 44 about an out-group member—a foreign student who criticized the participants' value system—
 45 and were asked to evaluate the target. This evaluation was designed to assess how much hostility
 46 participants felt toward the target.

6.1. Method

6.1.1. Participants and Procedure

Sixty-nine participants (39 females and 30 males, $M_{\text{age}} = 24.00$, $SD = 8.38$) were run in our lab individually, in different cubicles (randomized) to avoid carryover effects of scents. Furthermore, participants were booked at least 30 minutes apart in order to ventilate the rooms between sessions. Upon arrival, participants were given the first of two questionnaire packets to complete. This first questionnaire consisted of demographic questions and a number of filler items. We then randomly assigned participants to their condition by marking one of the three liquid scents (putrescine, ammonia, water) to the top of each page (0.5 ml) of the second questionnaire participants were given. In the putrescine and ammonia conditions, this amounted to a very subtle scent prime that was not meant to be detected. At the conclusion of the experiment, we funnel debriefed participants to determine whether they noticed or smelled anything unusual during the study. None of them reported being aware of the scents.

The second questionnaire assessed participants' mood, and our dependent variables. First, to rule out the possibility that our results could be explained by generalized affect, participants began the second part of the questionnaire by completing the 20-item Positive and Negative Affect Scale (PANAS; Tellegen et al., 1988). This scale measured the extent to which each of 10 positive affect descriptors ($\alpha = .86$) and 10 negative affect descriptors ($\alpha = .85$) reflected how they felt at that moment (1 = very slightly or not at all, 5 = extremely). We computed the average positive affect ($M = 3.31$, $SD = .68$) and negative affect ($M = 1.61$, $SD = .59$) scores for everybody.

This was followed by the description and evaluation of the out-group member (Greenberg et al., 2001; Navarrete et al., 2004; Norenzayan et al., 2007). Specifically, participants read an essay supposedly written by a college student from the Middle East who was visiting the United Kingdom to study English. In this essay, the student went on to criticize Western values, predicting its eventual decline (see Norenzayan et al., 2007). Participants were then asked to evaluate the author and his message by responding to four questions on a 9-point Likert scale ("To what extent do you like the author"; "To what extent would you like to be friends with the author"; "How much would you oppose the author teaching your (future) children"; and "How much do you want the ideas of the author to be publicized"; 1 = very much, 9 = not at all). We derived an overall out-group hostility index ($M = 5.82$, $SD = 1.63$) by averaging all items together ($\alpha = .77$), such that larger values indicated greater hostility. Finally, we measured motivation to escape the situation by timing how long it took participants to complete the second (scented) questionnaire followed by a standard demographic questionnaire (91% of the participants were native to England, 3% Greece, 4% Ireland, and 1% to the United States).

6.2. Results and Discussion

6.2.1. Ancillary Analyses

1 A one-way ANOVA tested whether the chemosensory primes elicited different levels of self-
 2 reported affect across the three conditions. However, the primes had no impact on positive affect
 3 $F(2, 66) = 1.87, p > .16$, nor negative affect, $F(2, 66) = .36, p > .70$. Moreover, the analyses
 4 below were no different when we used these affect measures as covariates, showing that any
 5 effect of our primes on out-group defense was not mediated by mood.

6 7 **6.2.2. Out-group Defense**

8
 9 As predicted, we found a significant effect of scent prime on defensiveness toward the author of
 10 the essay, $F(2, 66) = 11.83, p < .001, \eta^2 = .26$ (see Figure 3). Post-hoc analyses found that
 11 putrescine led to greater hostility ($M = 6.98, SD = 1.42$) compared to ammonia ($M = 5.05, SD =$
 12 1.54) and the neutral conditions ($M = 5.43, SD = 1.30$; both $ps < .005$). There was no significant
 13 difference between the ammonia and control conditions, $p > .6$.

14
 15 Experiment 4 supported the hypothesis that non-conscious exposure to putrescine evoked
 16 defensive responses toward an out-group member, and this effect was not due to conscious
 17 awareness of the scents, mood, or to the motivation to escape the aversive scent primes³.
 18 Although these results suggest that the scent primes elicited an odor percept (non-consciously),
 19 future studies may wish to control the precise intensities of the stimulus odors that are presented
 20 (e.g., using an olfactometer).

21 22 **7. General Discussion**

23
 24 This research was designed to test the hypothesis that putrescine could serve as a warning signal
 25 that mobilizes protective responses to deal with threats. In four experiments, we found support
 26 for this idea: conscious and non-conscious exposure to putrescine elicited distancing and
 27 defensive reactions (e.g., fight and flight responses). Putrescine increased vigilance (Experiment
 28 1), heightened the accessibility of escape- and threat-relevant cognitions (Experiment 3), and
 29 increased the speed participants walked away from the location of the scent (Experiments 2 and
 30 3). Experiment 4 created a situation where immediate escape was not likely and gave participants
 31 the opportunity to evaluate an out-group member. Subtle exposure to putrescine produced greater
 32 defensiveness toward the out-group member, suggesting an aggressive readiness in participants
 33 (Cesario et al., 2010). As a whole, the findings indicate that even brief exposure to putrescine
 34 mobilizes threat management responses designed to cope with environmental threats.

35
 36 These are the first results to show that a specific chemical compound (putrescine) can be
 37 processed as a threat signal. Thus far, nearly all the evidence for threat chemosignals has come
 38 from those that are transmitted by body sweat (de Groot et al., 2012; Pause et al., 2012).
 39 Moreover, these are among the first studies that show that a specific chemical compound can
 40 cause overt behavior in humans (Wysocki & Preti, 2004). Furthermore, an advantage of isolating
 41 putrescine in threat management processes is that it may help in determining which sensory and
 42 brain pathways are involved in chemosensory threat detection and processing. For instance,
 43 research suggests that the central nucleus of the amygdala projects to the midbrain

³ When the amount of time participants took to complete the questionnaire was used as a covariate, the results remained significant, $F(2, 65) = 13.13, p < .001, \eta^2 = .29$.

1 periaqueductal gray, the hypothalamus and the brainstem, which together coordinate to prepare
2 fight-or-flight responses to threatening stimuli (Misslin, 2003). We speculate that putrescine
3 activates a similar neurological pathway. Future research could include physiological
4 measurements (e.g., systolic blood pressure, heart rate) to test the thesis that the observed effects
5 of putrescine are modulated by processes originating in the sympathetic nervous system.
6

7 An important direction for future research will be to understand the precise nature of the
8 threat produced by putrescine (e.g., microbial, predatory). Our view is that putrescine is relevant
9 to both of these domains, though the immediate context should determine which type of threat is
10 more primary. Recent work on trace amine receptors (TAARs) has the potential to shed light on
11 some of these mechanisms, as the activation of different receptors may function to detect specific
12 threats, such as predators and pathogens (Li & Liberles, 2015; Pérez-Gómez et al., 2015). In
13 addition, this research suggests that cadaverine (a compound with a similar chemical structure as
14 putrescine; both are diamines) activates a similar pathway and produces similar escape and
15 avoidance responses (Hussain et al., 2013; Oliveira et al., 2014) in animals. Thus, we render it
16 likely that cadaverine evokes a similar threat response as putrescine (see Li & Liberles, 2015).
17

18 It would also be interesting to examine how putrescine detection affects sensitivity to
19 particular types of threat and whether it produces elevated responses to certain stimuli more than
20 others (e.g., fear- vs. disgust-based sensitivities). For instance, further research could elucidate
21 how putrescine activates sensory acquisition (typically associated with fear experiences) and
22 sensory rejection (associated with disgust) processes (Susskind et al., 2008), and whether
23 exposure to putrescine augments physiological responses (e.g., heart rate, pupil dilation) that
24 typically co-occur with adaptive responses to threats. This type of research would benefit from
25 including individual differences in both disgust and fear sensitivity (Garfinkel et al., 2014; Haidt
26 et al., 1994). By the same token, future work could clarify whether putrescine elicits discrete
27 emotions (e.g., fear versus disgust) or less specific affective states associated with negative
28 valence and high arousal (see also Li & Liberles, 2015; Smeets & Dijksterhuis, 2014). Our
29 findings, which showed that responses to putrescine were automatic, occurred after various
30 lengths of delay (Experiments 1-3), and when presented at sub-threshold levels (Experiment 4),
31 suggested that conscious evaluations are not at the heart of the observed responses to putrescine.
32 This is consistent with our theorizing and ample work showing that chemosensory cues influence
33 psychological and physiological operations outside conscious awareness (for extended reviews,
34 see Sela & Sobel, 2010; Smeets & Dijksterhuis, 2014). However, we hasten to add that more
35 research is needed to specify the exact nature of the effects produced by the sub-threshold
36 priming of putrescine, for instance, by varying the exposure times to putrescine, the delay after
37 the primes, and the intensity of the putrescine stimulus.
38

39 Another important question is how specific threat management responses develop.
40 Within non-olfactory sensory channels, for example, there may be an innate bias for humans to
41 detect certain biologically-relevant stimuli as threatening, such as the sight of snakes and spiders
42 (Ohman & Mineka, 2001). Although controversial in human research, some work suggests that
43 responses to chemosensory stimuli are innate (Hussain et al., 2013; Misslin, 2003; Dielenberg et
44 al., 2001). For instance, Soussignan et al. (1997) showed that soon after birth, butyric acid (a
45 malodorous scent) evoked disgust reactions in neonates, a finding they claim is consistent with
46 an innate predisposition toward ecologically-relevant scents. To test for possibility of innate

1 biases toward threatening chemosensory cues, it would be interesting to examine whether
2 putrescine triggers facial expressions associated with fear in infants. In fact, research indicates
3 that adults do not habituate so readily to the scent of putrescine emitted from rotting flesh
4 (Roberson et al., 2008), suggesting that there might be a bias to respond warily to it.
5

6 Although the innateness of responses to chemosignals is still controversial, humans'
7 ability to incorporate learned information into cultural practices is beyond question (Boyd &
8 Richerson, 2005). Consequently, the magnitude of specific chemosensory threat responses could
9 be different in cultures where people are exposed to putrescine more frequently. Likewise,
10 reactions to putrescine may differ between cultures with different burial practices (e.g.,
11 embalming practices, the duration before burial). These factors should remind us that the context
12 is critical to how people react to putrescine. How olfactory information modulates other sensory
13 inputs (Zhou et al., 2012) is no doubt central to whether it will be interpreted as threatening.
14

15 One alternative theoretical perspective of our findings on the effects of putrescine is
16 Terror Management Theory (TMT; Greenberg et al., 1994). According to this theory, reminders
17 of death are regulated by a "cultural anxiety buffer" that consists of beliefs and values that imbue
18 life with meaning and the promise of immortality. Interestingly, TMT argues that a great deal of
19 the darker side of human behavior (e.g., aggression, out-group prejudice, religious intolerance)
20 stems from the need to maintain and defend the integrity of this cultural anxiety buffer, due to its
21 vital role in managing existential angst. In this view, putrescine could function as a reminder of
22 mortality, and subsequently elicits similar defensive processes, as activated by reminders of
23 death. We do not rule out this possibility, but render it unlikely that chemosensory threats trigger
24 the same type of processes as those that originate from the unique human ability to reflect on the
25 conundrum of life and death (Landau et al., 2007). Nevertheless, examining whether putrescine
26 can be used as a subtle reminder of death, and whether it influences cultural beliefs, values, and
27 practices, would open up fascinating directions of research.
28

29 Most research has shown that humans process threats either visually or audibly, while
30 other animals inhabit the inaccessible world of scents. At the same time, we know that humans
31 are guided by many of the same olfactory processes, especially when they involve fitness-
32 relevant information. We believe that by identifying putrescine as one of these signals, a further
33 understanding of its mechanisms can shed light on more general processes that modulate
34 chemosensory signaling and threat management responses.
35
36

1 **8. Conflict of Interest Statement**

2

3 The authors declare that the research was conducted in the absence of any commercial or
4 financial relationships that could be construed as a potential conflict of interest.

5

6

9. References

- Ackerl, K., Atzmueller, M., & Grammer, K. (2002). The scent of fear. *Neuroendocrinology Letters*, 23, 79-84.
- Arakawa, H., Arakawa, K., & Deak, T. (2010). Sickness-related odor communication signals as determinants of social behavior in rat: a role for inflammatory processes. *Hormones and Behavior*, 57, 330-341.
- Arndt, J., Greenberg, J., Pyszczynski, T., & Solomon, S. (1997). Subliminal exposure to death-related stimuli increases defense of the cultural worldview. *Psychological Science*, 8, 379-385.
- Blanchard, R. J., Flannelly, K. J., & Blanchard, D. C. (1986). Defensive behaviors of laboratory and wild *Rattus norvegicus*. *Journal of Comparative Psychology*, 100, 101-107.
- Blanchard, C. D., Hynd, A. L., Minke, K. A., Minemoto, T., & Blanchard, R. J. (2001). Human defensive behaviors to threat scenarios show parallels to fear-and anxiety-related defense patterns of non-human mammals. *Neuroscience & Biobehavioral Reviews*, 25, 761-770.
- Boissy, A., Terlouw, C., & Le Neindre, P. (1998). Presence of cues from stressed conspecifics increases reactivity to aversive events in cattle: evidence for the existence of alarm substances in urine. *Physiology and Behavior*, 63, 489-495.
- Boyd, R., & Richerson, P. J. (2005). *The origin and evolution of cultures*. Oxford University Press, USA.
- Bradley, M. M., Codispoti, M., Cuthbert, B. N., & Lang, P. J. (2001). Emotion and motivation I: defensive and appetitive reactions in picture processing. *Emotion*, 1, 276-298.
- Cannon, W. B. (1927). The James-Lange theory of emotions: A critical examination and an alternative theory. *The American Journal of Psychology*, 39, 106-124.
- Cesario, J., Plaks, J. E., Hagiwara, N., Navarrete, C. D., & Higgins, E. T. (2010). The Ecology of Automaticity How Situational Contingencies Shape Action Semantics and Social Behavior. *Psychological Science*, 21, 1311-1317.
- Chen, D., & Haviland-Jones, J. (2000). Human olfactory communication of emotion. *Perceptual and Motor Skills*, 91, 771-781.
- Chumanov, E. S., Wall-Scheffler, C., & Heiderscheit, B. C. (2008). Gender differences in walking and running on level and inclined surfaces. *Clinical Biomechanics*, 23, 1260-1268.
- Degel, J., Piper, D., & Köster, E. P. (2001). Implicit learning and implicit memory for odors: the influence of odor identification and retention time. *Chemical Senses*, 26, 267-280.
- de Groot, J. H., Smeets, M. A., Kaldewaij, A., Duijndam, M. J., & Semin, G. R. (2012). Chemosignals communicate human emotions. *Psychological Science*, 23, 1417-1424.
- Dielenberg, R. A., Hunt, G. E., & McGregor, I. S. (2001). 'When a rat smells a cat': The distribution of Fos immunoreactivity in rat brain following exposure to a predatory odor. *Neuroscience*, 104, 1085-1097.
- Dinges, D. F., Orne, M. T., Whitehouse, W. G., & Orne, E. C. (1987). Temporal placement of a nap for alertness: contributions of circadian phase and prior wakefulness. *Sleep*, 10(4), 313-329.
- Dinges, D. F., & Powell, J. W. (1985). Microcomputer analyses of performance on a portable,

- 1 simple visual RT task during sustained operations. *Behavior Research Methods,*
2 *Instruments, & Computers*, 17(6), 652-655.
- 3 Dorrian, J., Sweeney, M., & Dawson, D. (2004). A prior sleep/wake model of fatigue-related
4 accident risk in truck drivers. *Journal of Sleep Research*, 13, A194.
- 5 Ellsworth, P. C., Carlsmith, J. M., & Henson, A. (1972). The stare as a stimulus to flight in
6 human subjects: a series of field experiments. *Journal of Personality and Social*
7 *Psychology*, 21(3), 302.
- 8 Epple, G., & Herz, R. S. (1999). Ambient odors associated to failure influence cognitive
9 performance in children. *Developmental Psychobiology*, 35(2), 103-107.
- 10 Garfinkel, S. N., Minati, L., Gray, M. A., Seth, A. K., Dolan, R. J., & Critchley, H. D. (2014).
11 Fear from the heart: sensitivity to fear stimuli depends on individual heartbeats. *The*
12 *Journal of Neuroscience*, 34(19), 6573-6582.
- 13 Gilbert, P., & Gilbert, J. (2003). Entrapment and arrested fight and flight in depression: An
14 exploration using focus groups. *Psychology and Psychotherapy: Theory, Research and*
15 *Practice*, 76(2), 173-188.
- 16 Gray, J. A., & McNaughton, N. (2003). *The neuropsychology of anxiety: An enquiry into the*
17 *function of the septo-hippocampal system* (No. 33). Oxford University Press.
- 18 Greenberg, J., Arndt, J., Schimel, J., Pyszczynski, T., & Solomon, S. (2001). Clarifying the
19 function of mortality salience-induced worldview defense: Renewed suppression or
20 reduced accessibility of death-related thoughts?. *Journal of Experimental Social*
21 *Psychology*, 37(1), 70-76.
- 22 Greenberg, J., Pyszczynski, T., Solomon, S., Simon, L., & Breus, M. (1994). Role of
23 consciousness and accessibility of death-related thoughts in mortality salience
24 effects. *Journal of Personality and Social Psychology*, 67(4), 627.
- 25 Haidt, J., McCauley, C., & Rozin, P. (1994). Individual differences in sensitivity to disgust: A
26 scale sampling seven domains of disgust elicitors. *Personality and Individual*
27 *Differences*, 16(5), 701-713.
- 28 Holbrook, C., Sousa, P., & Hahn-Holbrook, J. (2011). Unconscious vigilance: Worldview
29 defense without adaptations for terror, coalition, or uncertainty management. *Journal of*
30 *Personality and Social Psychology*, 101(3), 451.
- 31 Horowitz, L. F., Saraiva, L. R., Kuang, D., Yoon, K. H., & Buck, L. B. (2014). Olfactory
32 receptor patterning in a higher primate. *The Journal of Neuroscience*, 34, 12241-12252.
- 33 Hummel, T., & Kobal, G (1992). Differences in human evoked potentials related to
34 olfactory or trigeminal chemosensory activation. *Electroencephalogr. Clin.*
35 *Neurophysiol.*, 84, 84-89.
- 36 Hussain A., Saraiva, L. R., Ferrero, D. M., Ahuja, G., Krishna, V. S., Liberles, S. D., &
37 Korsching, S. I. (2013). High-affinity olfactory receptor for the death associated odor
38 cadaverine. *Proceedings of the National Academy of Sciences*, 48, 19579-19584.
- 39 Jacquot, L., Monnin, J., & Brand, G. (2004). Unconscious odor detection could not be due to
40 odor itself. *Brain Research*, 1002(1), 51-54.
- 41 Knasko, S. C. (1993). Performance, mood, and health during exposure to intermittent
42 odors. *Archives of Environmental Health: An International Journal*, 48(5), 305-308.
- 43 Koster, E. P., Degel, J., & Piper, D. (2002). Proactive and retroactive interference in implicit
44 odor memory. *Chemical Senses*, 27, 191-207.
- 45 Krusemark, E. A., & Li, W. (2012). Enhanced olfactory sensory perception of threat in anxiety:
46 an event-related fMRI study. *Chemosensory Perception*, 5(1), 37-45.

- 1 Landau, M. J., Solomon, S., Pyszczynski, T., & Greenberg, J. (2007). On the compatibility of
2 terror management theory and perspectives on human evolution. *Evolutionary*
3 *Psychology*, 5, 476-519.
- 4 Lee, S. W., & Schwarz, N. (2012). Bidirectionality, mediation, and moderation of metaphorical
5 effects: The embodiment of social suspicion and fishy smells. *Journal of Personality and*
6 *Social Psychology*, 103(5), 737-749.
- 7 Li, Q., & Liberles, S. D. (2015). Aversion and attraction through olfaction. *Current Biology*, 25,
8 R120-R129.
- 9 Li, W., Moallem, I., Paller, K. A., & Gottfried, J. A. (2007). Subliminal smells can guide social
10 preferences. *Psychological Science*, 18(12), 1044-1049.
- 11 Loh, S., Lamond, N., Dorrian, J., Roach, G., & Dawson, D. (2004). The validity of psychomotor
12 vigilance tasks of less than 10-minute duration. *Behavior Research Methods*, 36(2), 339-
13 346.
- 14 Lozito, J. P., & Mulligan, N. W. (2010). Exploring the role of attention during implicit memory
15 retrieval. *Journal of Memory and Language*, 63, 387-399.
- 16 Migo, E. M., Roper, A., Montaldi, D., & Mayes, A. R. (2010). British English norms for the
17 spontaneous completion of three-letter word stems. *Behavior Research Methods*, 42, 470-
18 473.
- 19 Mobbs, D., Marchant, J. L., Hassabis, D., Seymour, B., Tan, G., Gray, M., et al. (2009). From
20 threat to fear: the neural organization of defensive fear systems in humans. *The Journal*
21 *of Neuroscience*, 29, 12236-12243.
- 22 Miller, S. L., & Maner, J. K. (2010). Scent of a woman: men's testosterone responses to
23 olfactory ovulation cues. *Psychological Science*, 21, 276-283.
- 24 Millot, J. L., Brand, G., & Morand, N. (2002). Effects of ambient odors on reaction time in
25 humans. *Neuroscience Letters*, 322(2), 79-82.
- 26 Misslin, R. (2003). The defense system of fear: Behavior and neurocircuitry. *Clinical*
27 *Neurophysiology*, 33, 55-66.
- 28 Mujica-Parodi, L. R., Strey, H. H., Frederick, B., Savoy, R., Cox, D., Botanov, Y.,
29 Tolkunov, D., Rubin, D., & Weber, J. (2009). Chemosensory cues to conspecific
30 emotional stress activate amygdala in humans. *PLoS One*, 4, e6415.
- 31 Navarrete, C. D., Kurzban, R., Fessler, D. M., & Kirkpatrick, L. A. (2004). Anxiety and
32 intergroup bias: Terror management or coalitional psychology?. *Group Processes &*
33 *Intergroup Relations*, 7(4), 370-397.
- 34 Neuberg, S. L., Kenrick, D. T., & Schaller, M. (2011). Human threat management systems: Self-
35 protection and disease avoidance. *Neuroscience & Biobehavioral Reviews*, 35, 1042-
36 1051.
- 37 Norenzayan, A., Dar-Nimrod, I., Hansen, I. G., & Proulx, T. (2007). Mortality salience
38 and religion: Divergent effects on the defense of cultural worldviews for the religious and
39 the non-religious. *European Journal of Social Psychology*, 39, 101-113.
- 40 Ohman, A., & Mineka, S. (2001). Fears, phobias, and preparedness: Toward an evolved module
41 of fear and fear learning. *Psychological Review*, 108, 483-522.
- 42 Olsson, M. J., Lundström, J. N., Kimball, B. A., Gordon, A. R., Karshikoff, B., Hosseini, N., et
43 al. (2014). The scent of disease: Human body odor contains an early chemosensory cue of
44 sickness. *Psychological Science*, 25, 817-823.
- 45 Pause, B. M. (2012). Processing of body odor signals by the human brain. *Chemosensory*
46 *Perception*, 5, 55-63.

- 1 Pause, B. M., Adolph, D., Prehn-Kristensen, A., & Ferstl, R. (2009). Startle response
2 potentiation to chemosensory anxiety signals in socially anxious individuals.
3 *International Journal of Psychophysiology*, 74, 88-92.
- 4 Pérez-Gómez, A., Blyemehl, K., Stein, B., Pyrski, M., Birnbaumer, L., Munger, S. D., et al.
5 (2015). Innate predator odor aversion driven by parallel olfactory subsystems that
6 converge in the ventromedial hypothalamus. *Current Biology*, 25, 1-7.
- 7 Prehn, A., Ohrt, A., Sojka, B., Ferstl, R., & Pause, B. M. (2006). Chemosensory anxiety signals
8 augment the startle reflex in humans. *Neuroscience Letters*, 394, 127-130.
- 9 Prounis, G. S., & Shields, W. M. (2013). Necrophobic behavior in small mammals. *Behavioural*
10 *Processes*, 94, 41-44.
- 11 Rieser, J., Yonas, A., & Wikner, K. (1976). Radial localization of odors by human newborns.
12 *Child Development*, 47, 856-859.
- 13 Roberson, D. W., Neil, J. A., & Bryant, E. T. (2008). Improving wound care simulation with the
14 addition of odor: A descriptive, quasi-experimental study. *OstomyWound Management*,
15 54, 36-43.
- 16 Sekizawa, S. I., & Tsubone, H. (1994). Nasal receptors responding to noxious chemical irritants.
17 *Respiration Physiology*, 96(1), 37-48.
- 18 Sela, L., & Sobel, N. (2010). Human olfaction: A constant state of change-blindness.
19 *Experimental Brain Research*, 205(1), 13-29.
- 20 Smeets, M. A. M., & Dijksterhuis, G. B. (2014). Smelly primes—when olfactory primes do or do
21 not work. *Frontiers in Psychology*, 5, 96.
- 22 Sobel, N., Prabhakaran, V., Hartley, C. A., Desmond, J. E., Glover, G. H., Sullivan, E. V., &
23 Gabrieli, J. D. (1999). Blind smell: brain activation induced by an undetected air-borne
24 chemical. *Brain*, 122(2), 209-217.
- 25 Soussignan, R., Schaal, B., Marlier, L., & Jiang, T. (1997). Facial and autonomic responses to
26 biological and artificial olfactory stimuli in human neonates: Re-examining early hedonic
27 discrimination of odors. *Physiology & Behavior*, 62, 745-758.
- 28 Stern, K., & McClintock, M. K. (1998). Regulation of ovulation by human pheromones. *Nature*,
29 392, 177-179.
- 30 Stevenson, R. J. (2010). An initial evaluation of the functions of human olfaction. *Chemical*
31 *Senses*, 35, 3-20.
- 32 Stevenson, R. J., Boakes, R. A., & Prescott, J. (1998). Changes in odor sweetness
33 resulting from implicit learning of a simultaneous odor-sweetness association: An
34 example of learned synesthesia. *Learning and Motivation*, 29, 113-132.
- 35 Susskind, J. M., Lee, D. H., Cusi, A., Feiman, R., Grabski, W., & Anderson, A. K. (2008).
36 Expressing fear enhances sensory acquisition. *Nature Neuroscience*, 11(7), 843-850.
- 37 Tellegen, A., Watson, D., & Clark, L. A. (1988). Development and validation of brief measures
38 of positive and negative affect: The PANAS scales. *Journal of Personality and Social*
39 *Psychology*, 54, 1063-1070.
- 40 Tybur, J. M., Bryan, A. D., Magnan, R. E., & Hooper, A. E. C. (2011). Smells like safe sex:
41 Olfactory pathogen primes increase intentions to use condoms. *Psychological*
42 *Science*, 22, 478-480.
- 43 Wise, P. M., Canty, T. M., & Wysocki, C. J. (2005). Temporal integration of nasal irritation from
44 ammonia at threshold and supra-threshold levels. *Toxicological Sciences*, 87(1), 223-231.
- 45 Wisman, A., & Koole, S. L. (2003). Hiding in the crowd: can mortality salience promote

- 1 affiliation with others who oppose one's worldviews?. *Journal of Personality and Social*
2 *Psychology*, 84(3), 511-526.
- 3 Wheatley, T., & Haidt, J. (2005). Hypnotic disgust makes moral judgments more severe.
4 *Psychological Science*, 16, 780-784.
- 5 Woody, E. Z., & Szechtman, H. (2011). Adaptation to potential threat: the evolution,
6 neurobiology, and psychopathology of the security motivation system. *Neuroscience &*
7 *Biobehavioral Reviews*, 35(4), 1019-1033.
- 8 Wyatt, T. D. (2009). Fifty years of pheromones. *Nature*, 457, 262-263.
- 9 Wysocki, C. J., & Preti, G. (2004). Facts, fallacies, fears, and frustrations with human
10 pheromones. *The Anatomical Record Part A: Discoveries in Molecular, Cellular, and*
11 *Evolutionary Biology*, 281, 1201-1211.
- 12 Yao, M., Rosenfeld, J., Attridge, S., Sidhu, S., Aksenov, V., & Rollo, C. D. (2009). The ancient
13 chemistry of avoiding risks of predation and disease. *Evolutionary Biology*, 36, 267-281.
- 14 Yeoman, C. J., Thomas, S. M., Miller, M. E., Ulanov, A. V., Torralba, M., Lucas, S., et al.
15 (2013). A multi-omic systems-based approach reveals metabolic markers of bacterial
16 vaginosis and insight into the disease. *PloS One*, 8(2), e56111.
- 17 Yoon, K. H., Ragozczy, T., Lu, Z., Kondoh, K., Kuang, D., Groudine, M., & Buck, L. B. (2015).
18 Olfactory receptor genes expressed in distinct lineages are sequestered in different
19 nuclear compartments. *Proceedings of the National Academy of Sciences*, 112, E2403-
20 E2409.
- 21 Zhou, W., & Chen, D. (2009). Fear-related chemosignals modulate recognition of fear in
22 ambiguous facial expressions. *Psychological Science*, 20, 177-183.
- 23 Zhou, W., Zhang, X., Chen, J., Wang, L., & Chen, D. (2012). Nostril-specific olfactory
24 modulation of visual perception in binocular rivalry. *The Journal of Neuroscience*, 32,
25 17225-17229.
26
27

1 Table 1

2 Hedonic evaluations of putrescine, ammonia, indole, "fart spray," and skatole¹ (Pilot study).

3

4 Scent primes	Putrescine	Ammonia	Indole	Skatole	Fart spray
5					
6 Intensity ²					
7 M	5.98 _b	6.60 _b	5.25 _a	7.23 _c	5.52 _b
8 SD	2.50	2.46	2.15	2.08	2.07
9 Familiarity					
10 M	4.98 _a	5.10 _a	6.88 _b	5.21 _a	4.90 _a
11 SD	2.71	2.95	2.46	2.56	2.69
12 Repugnance					
13 M	5.94 _b	5.94 _b	3.65 _a	6.54 _b	5.31 _b
14 SD	2.65	2.55	1.78	2.94	2.63
15 Positivity					
16 M	2.63 _b	2.69 _b	3.81 _a	2.50 _b	2.67 _b
17 SD	1.55	1.78	2.05	1.87	1.77
18 N	48	48	48	48	48

19
20
21 ¹ "How intense is this scent?", 1 = Not at all and 10 = Very much; "How familiar is this
22 scent?", 1 = Not at all and 10 = Very much; "How repugnant is this scent?", 1 = Not at all and 10
23 = Very much; "How positive does this scent make you feel?", 1 = Not at all and 10 = Very much.

24 ² Different subscripts on a hedonic dimension (within a row) indicate a significant
25 difference of $p < .05$.

26

27

1 Table 2

2 Scent ratings for the chemosensory primes (Experiment 1)

3

4 Chemosensory primes Neutral Ammonia Putrescine

5

6 Intensity

7 M 3.30 4.73 4.27

8 SD 1.81 1.45 1.92

9

10 Familiarity

11 M 6.00 5.10 4.40

12 SD .86 2.25 1.60

13

14 Repugnance

15 M 2.35 5.90 5.65

16 SD 1.46 1.34 1.23

17 N 20 20 20

18

1 Table 3

2 Scent ratings for the chemosensory primes (Experiment 2)

3

4 Chemosensory primes Neutral Ammonia Putrescine

5

6 Intensity

7 M 1.53 4.73 4.27

8 SD .64 .46 .70

9

10 Familiarity

11 M 4.75 1.60 1.67

12 SD .46 .51 .62

13

14 Repugnance

15 M 1.73 4.47 4.80

16 SD .70 .74 .41

17 N 15 15 15

18

1 Table 4

2 Scent ratings for the chemosensory primes (Experiment 3)

3

4 Chemosensory primes Neutral Ammonia Putrescine

5

6 Intensity

7 M 1.85 3.20 3.40

8 SD .99 1.32 .99

9

10 Familiarity

11 M 2.95 2.20 2.15

12 SD .83 .89 .75

13

14 Repugnance

15 M 2.60 3.70 3.50

16 SD .60 .98 1.15

17 N 20 20 20

18

19

1 Table 5

2 The ratings of escape-related and threat-related cognitions for the chemosensory primes

3 (Experiment 3).

4

5 Chemosensory primes	Neutral	Ammonia	Putrescine
-----------------------	---------	---------	------------

6

7 Escape cognitions

8 M 2.15 2.45 3.45

9 SD 0.99 1.05 0.69

10

11 Threat cognitions

12 M 1.68 1.73 2.55


13 SD 0.65 0.64 0.94

14 N 20 20 20

15

1

2



3

4 Figure 1. The number of seconds it took participants to walk 80 meters after exposure to the
5 scent prime (Experiment 2). Asterisks denote that two groups differ at $**p < .005$.

6

7

8





Figure 2. The number of seconds it took participants to walk 60 meters after exposure to the scent prime (Experiment 3). Asterisks denote that two groups differ at $*p < .05$.

1

2



3

4 Figure 3. Mean scores on the worldview defense scale for all three conditions (Experiment 4).
5 Higher scores reflect greater hostility toward the target. Asterisks denote two groups differ at $**p$
6 $< .005$.

7

8

9

10

11