



# University of HUDDERSFIELD

## University of Huddersfield Repository

Charles, C.J., Rout, S.P., Garratt, E.J., Patel, K., Laws, A.P. and Humphreys, Paul

The enrichment of an alkaliphilic biofilm consortia capable of the anaerobic degradation of isosaccharinic acid from cellulosic materials incubated within an anthropogenic, hyperalkaline environment.

### Original Citation

Charles, C.J., Rout, S.P., Garratt, E.J., Patel, K., Laws, A.P. and Humphreys, Paul (2015) The enrichment of an alkaliphilic biofilm consortia capable of the anaerobic degradation of isosaccharinic acid from cellulosic materials incubated within an anthropogenic, hyperalkaline environment. *FEMS Microbiology Ecology*. ISSN 1574-6941 (In Press)

This version is available at <http://eprints.hud.ac.uk/25235/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: [E.mailbox@hud.ac.uk](mailto:E.mailbox@hud.ac.uk).

<http://eprints.hud.ac.uk/>

1 **The enrichment of an alkaliphilic biofilm consortia capable of the anaerobic**  
2 **degradation of isosaccharinic acid from cellulosic materials incubated within an**  
3 **anthropogenic, hyperalkaline environment.**

4 C.J. Charles<sup>a</sup>, S.P. Rout<sup>a</sup>, E.J. Garratt<sup>a</sup>, K. Patel<sup>a</sup>, A.P. Laws<sup>b</sup>, P.N. Humphreys<sup>a\*</sup>.

5 <sup>a</sup> Department of Biological Sciences, School of Applied Sciences, University of  
6 Huddersfield, Queensgate, Huddersfield HD1 3DH, United Kingdom.

7 <sup>b</sup> Department of Chemical Sciences, School of Applied Sciences, University of Huddersfield,  
8 Queensgate, Huddersfield HD1 3DH, United Kingdom.

9 \* Corresponding author. Tel: +441484472771.

10 E-mail address: p.n.humphreys@hud.ac.uk (P.N. Humphreys).

11 Running title = Isolation of an alkaliphilic biofilm consortia.

12 **Abstract**

13 Anthropogenic hyper-alkaline sites provide an environment that is analogous to  
14 proposed cementitious geological disposal facilities (GDF) for radioactive waste. Under  
15 anoxic, alkaline conditions cellulosic wastes will hydrolyse to a range of cellulose  
16 degradation products (CDP) dominated by isosaccharinic acids (ISA). In order to investigate  
17 the potential for microbial activity in a cementitious GDF, cellulose samples were incubated  
18 in the alkaline (~pH 12), anaerobic zone of a lime kiln waste site. Following retrieval, these  
19 samples had undergone partial alkaline hydrolysis and were colonised by a Clostridia  
20 dominated biofilm community, where hydrogenotrophic, alkaliphilic methanogens were also  
21 present. When these samples were used to establish an alkaline CDP fed microcosm, the  
22 community shifted away from Clostridia, methanogens became undetectable and a flocculate

23 community dominated by *Alishewanella sp.* established. These flocs were composed of  
24 bacteria embedded in polysaccharides and protein stabilised by extracellular DNA. This  
25 community was able to degrade all forms of ISA with >60% of the carbon flow being  
26 channelled into extracellular polymeric substance (EPS) production. This study demonstrated  
27 that alkaliphilic microbial communities can degrade the CDP associated with some  
28 radioactive waste disposal concepts at pH 11. These communities divert significant amounts  
29 of degradable carbon to EPS formation, suggesting that EPS has a central role in the  
30 protection of these communities from hyper-alkaline conditions.

31

31

32 **Introduction**

33           The UK's national nuclear waste legacy contains approximately 290 000 m<sup>3</sup> (N.D.A.,  
34 2013) of intermediate level radioactive wastes (ILW) which includes an estimated (~2000  
35 tonnes) (N.D.A., 2010a) of cellulosic materials (wood, paper and cloth) (Humphreys et al.,  
36 2010a). One of the proposed strategies for the disposal of this ILW is a deep geological  
37 disposal facility (GDF) (N.D.A., 2010a) employing a multi-barrier system which is likely to  
38 include a cement based backfill (Chapman and Hooper, 2012). Upon the closure of such a  
39 facility, groundwater ingress combined with corrosion processes will result in the  
40 development of a chemically reducing high pH (pH 12.5) environment (Libert et al., 2011,  
41 N.D.A., 2010b). Under these conditions the cellulose portion of ILW is expected to undergo  
42 chemical, alkaline hydrolysis to form a variety of cellulose degradation products (CDP)  
43 (Knill and Kennedy, 2003, Humphreys et al., 2010a).

44           CDP are comprised of the alpha and beta diastereomers of isosaccharinic acid (ISA),  
45 alongside other small chain organic compounds including acetic acid (Van Loon and Glaus,  
46 1997, Motellier et al., 1998, Knill and Kennedy, 2003). The diastereomers of ISA are of  
47 significance when considering the performance of a GDF as they possess the ability to  
48 enhance the mobility of a range of radionuclides, including nickel, thorium, plutonium and  
49 uranium through complexation (Greenfield et al., 1991, Allard and Ekberg, 2006, Warwick et  
50 al., 2003). In addition, the hemicellulose fraction of cellulosic waste components will also  
51 undergo anoxic, alkaline hydrolysis to form an additional 5-carbon form of ISA, known as  
52 xyloisosaccharinic acid (X-ISA) (Almond et al., 2012). Recent work by Randall et al (2013)  
53 suggests that X-ISA does not have the same complexation properties as the alpha and beta

54 forms of ISA but could, however, represent a source of organic carbon available for microbial  
55 metabolism.

56           Although the harsh geochemical conditions of an ILW-GDF place limitations upon  
57 microbial life it may not prevent microbes from colonising a facility. An investigation of an  
58 anthropogenic analogue of an ILW-GDF at a hyperalkaline contaminated site in Buxton, UK  
59 where ISA is generated in-situ (Rout et al., 2015) has revealed a microbially active site  
60 despite porewaters of up to pH 13 (Burke et al., 2012). The range of microbes present within  
61 the background sediments is diverse, with organisms within the Phyla Bacteroidetes,  
62 Proteobacteria and Firmicutes consistently making up large proportions of the sediment  
63 taxonomic profiles (Burke et al., 2012, Williamson et al., 2013, Bassil et al., 2014). The  
64 subsequent culturing of these sediments has shown that these communities are able to utilise  
65 the alpha form of ISA as a substrate under aerobic, nitrate, iron reducing (Bassil et al., 2014)  
66 and methanogenic conditions (Rout et al., 2015). Sulphate reduction appears to be inhibited at  
67 pH >10 (Bassil et al., 2014), however, the utilisation of ISA under sulphate reducing  
68 conditions has been observed at neutral pH indicating that this limitation is thermodynamic  
69 (Rizoulis et al., 2012, Rout et al., 2014). The heterogeneity of ILW and its compaction in  
70 grout may limit the availability of higher energy terminal electron acceptors such as nitrate  
71 and ferric iron, with the inundating ground water also depleted in these electron acceptors due  
72 to its passage through the microbial thermodynamic ladder (Bethke et al., 2011).  
73 Fermentation processes and subsequent methanogenesis therefore represent the most likely  
74 conditions to dominate an ILW-GDF

75           Microbes in nature can be found in biofilms of mixed syntrophic communities, with  
76 microbial biofilms found in a diverse range of environments (Summons et al., 2015, Urbieta  
77 et al., 2015). The secretion of extracellular polymeric substance (EPS) such as  
78 polysaccharides, proteins, lipids and nucleic acids during biofilm formation assist in bacterial

79 survival and propagation (Flemming and Wingender, 2010) and confer an increased  
80 resistance to environmental stresses such as pH and temperature fluctuations, desiccation and  
81 UV radiation (Ordoñez et al., 2009, Gorlenko et al., 2004, Rodrigues et al., 2006, Jones et al.,  
82 1994, Conrad et al., 2014). When considering the colonisation of an ILW-GDF, the ability of  
83 microbes to migrate and adhere to niche areas such as ungrouted surfaces may allow for both  
84 microbial survival and growth under extreme alkaline conditions (Humphreys et al., 2010b).  
85 The aim of this work was to culture, *in situ*, a biofilm forming consortium capable of  
86 colonising cellulosic materials under anoxic, hyper-alkaline conditions and to determine its  
87 ability to degrade CDP, which represent the primary organic carbon source within an ILW-  
88 GDF.

## 89 **Methods**

### 90 Cellulose cotton preparation

91 In order to prepare the cellulose cotton for incubation, raw woven cotton fabric  
92 (Greige) was treated with NaOH to saponify the natural waxes along with an alkali stable  
93 phosphate ester detergent to emulsify the suspended impurities. Further treatment with NaOH  
94 and phosphonate stabilised H<sub>2</sub>O<sub>2</sub> was carried out to bleach the fabric. The cotton was then  
95 rinsed, neutralised under acetic acid before finally being rinsed, dried and autoclaved at  
96 121°C prior to use.

### 97 Analogue site investigation

98 During May 2014 a 2.2cm Ø borehole was hand drilled to an approximate depth of  
99 0.5m into an area inundated with alkaline leachate at Brook Bottom, Harpur Hill, Buxton, UK  
100 (Figure 1). An inert plastic liner with a perforated lower section was placed into the borehole.  
101 Approximately 5g of sterile treated cellulose cotton was loaded into a nylon mesh bag and

102 placed at the bottom of the borehole. After a period of 3 months the cotton was recovered  
103 along with sediment and porewater samples from the immediate vicinity of the sample. *In*  
104 *situ* pH and Eh values were determined prior to sample recovery using a handheld portable  
105 pH meter with calibrated electrodes and an InLab Redox Micro probe (Mettler Toledo, UK)  
106 tested in accordance with BS ISO 11271:2002 (B.S.I, 2002). All recovered materials were  
107 sealed in airtight containers along with anaerobic gas packs (Anaerogen, Oxoid, UK) for  
108 transport. Sediment and porewater samples were stored at -20°C until analysis and cotton not  
109 used for immediate studies was stored at -20°C in a solution of 140mL of ultrapure water,  
110 10mL of 1M TRIS-HCl (pH7.5) and 250ml of 96% ethanol after an overnight fixation step in  
111 4% paraformaldehyde in phosphate buffered saline.

112 Porewater, cotton and sediment ISA content was determined as previously described  
113 by Rout et al (2015, 2014) against ISA standards in the alpha, beta and xylo conformations  
114 (Almond et al., 2012, Shaw et al., 2012). C1-8 volatile fatty acid (VFA) content of both the  
115 sediment and cotton was determined using a standard extraction method outlined in Eaton et  
116 al (2005) and analysed via GC-FID as described by Rout et al (2014).

### 117 Microscopy

118 Scanning electron microscopy was undertaken using a JEOL JSM-6060LV  
119 microscope (JEOL, USA). Samples were dehydrated using a serial ethanol dilution of 25 %,  
120 50 %, 75% and 100 % for 2 minutes per step then sputter coated via a gold palladium plasma  
121 (CA7625 Polaron, Quorum Technologies Ltd, UK). Fluorescence microscopy was carried out  
122 using an Olympus BX41 laboratory microscope (Olympus, USA). Live dead staining was  
123 carried out using the BAC light Live/dead kit (Life technologies, UK), fluorescein  
124 isothiocyanate (FITC) (Sigma-Aldrich, UK) staining was used for protein and visualisation of  
125 individual bacteria cells and the polysaccharide components was achieved using ethidium

126 bromide and Calcofluor White (Sigma-Aldrich, UK) staining, respectively. For DNase  
127 digestion, microcosm fluid (1mL) was centrifuged at 10,000 xg for 1 minute and re-  
128 suspended in ultrapure water (1mL). A 10 fold dilution of this was then subjected to digestion  
129 by DNase using a DNase 1 kit (Sigma-Aldrich, UK).


### 130 Microcosm

131 In order to investigate ISA degradation, approximately 1g of colonised cotton was  
132 washed with 10mL N<sub>2</sub> purged sterile PBS under an inert environment to remove any transient  
133 microorganisms. The washed cotton was then added to a continuously stirred microcosm  
134 containing 175mL of pre-reduced 10% CDP and 90% mineral media (B.S.I, 2005) at pH 11  
135 and 20°C that had been purged with nitrogen and maintained with a nitrogen headspace to  
136 ensure anoxic conditions. CDP was produced as previously described by Rout et al (2014).  
137 The microcosm was brought up to a final volume of 250mL by feeding 25mL of CDP every 2  
138 weeks with the pH adjusted using 4M NaOH every 7 days. After this period the cotton was  
139 removed and the microcosm was switched to a 10% waste/feed cycle with CDP every 2  
140 weeks. The microcosm was maintained with a nitrogen atmosphere and all reagents were  
141 reduced prior to use with disodium sulfide nonahydrate (Sigma-Aldrich, UK) and sodium  
142 dithionite (Fisher, UK) as per BS ISO 14853:2005 (B.S.I, 2005) and stored under nitrogen.  
143 Resazurin redox indicator (Fisher, UK) present within the mineral media provided an  
144 indication of anaerobic conditions within the microcosm and all manipulations of the  
145 microcosm were carried out under a stream of nitrogen to maintain anoxic conditions.  
146 Sufficient time (50 weeks) was allowed for the microcosm chemistry to stabilise and also to  
147 allow for the washout of any transient microorganisms. The microcosm was sampled every 2  
148 days over 2 feed/waste cycles to determine the ISA and VFA content. For each sample period  
149 microcosm fluid (1mL) was taken, centrifuged at 10,000 xg for 1 minute and the supernatant  
150 filter sterilised using a 0.45µm syringe filter (Sartorius, UK) and stored at -20°C prior to



151 analysis. The gas headspace (75ml starting volume) was sampled every 2 days with the  
152 composition determined via gas chromatography using Agilent 6850 gas chromatograph  
153 (Hewlett Packard, UK) fitted with a HP-Plot/Q+ PT column and thermal conductivity  
154 detection (TCD). Headspace gas (100µL) was removed using a lockable gas syringe from the  
155 microcosm and passed through the column under the following conditions: initial temperature  
156 of 60°C for 2 minutes, followed by an increase to 120°C at a ramp rate of 30°C min<sup>-1</sup> with a  
157 detector temperature of 250°C. Gas headspace pressure was measured using a digital  
158 manometer (TPI, UK) before gas sample periods.

159 Microcosm fluid (1mL) containing the suspended flocs was taken on days 0, 7 and 14  
160 and spun at 10,000 xg for 1 minute for ATP/biomass detection using a 3M™ Clean-Trace™  
161 Biomass Detection Kit and Luminometer employing a modified method (3M, UK). The pellet  
162 was washed once with pH 4 PBS and reconstituted in pH 7 PBS to remove interference from  
163 excess alkalinity and salts. Following analysis, CFU mL<sup>-1</sup> and dry weight biomass (DW)  
164 were calculated against a standard curve of *E.coli* K12 concentrations. In addition, a set of  
165 control microcosms amended with 50µg mL<sup>-1</sup> chloramphenicol were prepared and were  
166 sampled as per the test microcosms. The controls served as an abiotic comparison for the  
167 elimination of sorption and precipitation events. All data was processed in Microsoft Excel  
168 with calculated means and associated standard error shown in all relevant results. Carbon  
169 flow calculations were undertaken using balanced equations 1 and 2 for the fermentation of  
170 ISA to acetate and hydrogen.



173 Preparation of 16S rDNA clone libraries

174 Total genomic DNA was extracted from the cotton and microcosm using a Powersoil  
175 DNA extraction kit (Mo-BIO, Carlsbad, US) with the following modifications. For the cotton  
176 approximately 0.25g was washed with pH 7 PBS and loaded into a glass bead tube with  
177 100 $\mu$ L  $\beta$ - mercaptoethanol and the bead beating step extended to 1 hour in order to overcome  
178 dampening effects introduced by the material. For genomic DNA extraction from the  
179 microcosm, 25mL of fluid was centrifuged at 5000 xg for 15min and the pellet re-suspended  
180 in 25mL pH 4 PBS. The sample was then centrifuged again at 5000 xg for 15 minutes and re-  
181 suspended in 2mL of pH 7 PBS. 1mL of the concentrated sample was transferred to a 1.5mL  
182 tube and centrifuged again at 10,000 xg for 1 minute, after which the supernatant was  
183 removed and the cell pellet re-suspended in the reaction fluid provided in the glass bead tubes  
184 of the Powersoil kit. The resulting mixture was then transferred back to a glass bead tube and  
185 bead beaten with 100 $\mu$ L  $\beta$ -mercaptoethanol for an increased time of 20 minutes to overcome  
186 clogging due to the EPS and then run as per the supplier's instruction. These modifications  
187 were found to increase the yield and purity of DNA obtained from both samples by removing  
188 excess salts, inhibiting nucleases and neutralising the samples.

189 Purified genomic DNA was quantified and quality checked by spectroscopic methods  
190 and used as a template to amplify the 16s rRNA gene. A ~1500bp fragment of the Eubacterial  
191 16S rRNA gene was amplified using broad specificity primers pA and pH (Edwards et al.,  
192 1989) and a ~750bp fragment of the archaeal 16S rRNA gene was amplified using primers Ar  
193 and Af (Gantner et al., 2011). PCR reactions were carried out using BIOMIX red master mix  
194 (BIOLINE, UK) with PCR fragments purified via a Qiaquick PCR purification kit (Qiagen,  
195 UK) and visualised using a 1.0% agarose TAE gel with SYBR® Safe staining (Life  
196 technologies, UK). PCR products were ligated into the standard cloning vector PGEM-T easy  
197 (Promega, US) and transformed into *E.coli* JM109 competent cells (Promega, US).  
198 Transformed cells were grown on Luria Bertani (LB) agar containing 100 $\mu$ g mL<sup>-1</sup> ampicillin

199 overlaid with 40 $\mu$ L of 100mM IPTG and 40 $\mu$ L of 40mg mL<sup>-1</sup> X-GAL (5-bromo-4-chloro-3-  
200 indolyl- $\beta$ -D-galactopyranoside) in N’N dimethylformamide for blue-white colour screening.  
201 Insert containing colonies were transferred to 96 well plates containing LB agar with 150mg  
202 mL<sup>-1</sup> ampicillin and sequenced using Sanger sequencing technology (GATC Biotech,  
203 Germany). Inserts were amplified using a T7 forward primer and the resulting 16S rRNA  
204 gene sequences aligned using the multiple sequence alignment package MUSCLE  
205 (www.ebi.ac.uk/Tools/services/web/toolform.ebi?tool=muscle) and chimera checked using  
206 the UCHIME component of the Mothur suite, where chimeric sequences were omitted from  
207 the analysis (Schloss et al., 2009). Sequences were analysed against the NCBI database using  
208 Basic Local Alignment Search Tool (MegaBLAST) utilising the 16S ribosomal RNA  
209 sequences for Bacteria and Archaea (Altschul et al., 1997). Phylogenetic families were then  
210 determined at a 95% confidence level by comparison with the Ribosomal Database Project  
211 (Cole et al., 2009).

#### 212 Nucleotide accession numbers

213 The 16S rRNA sequence data from the colonised cotton has been submitted to  
214 GenBank under accession numbers KP263977 - KP264111 and the microcosm sequences  
215 under the numbers KP728118 - KP728176.

## 216 **Results**

### 217 Chemical and physiological cotton analysis

218 The pH in the vicinity of the cotton samples was between pH 11.5 and 12 and redox  
219 measurements were found to be negative in both the associated sediment (-77mV) and  
220 porewaters (-66mV). Both the alpha and beta forms of ISA were extracted from the cotton  
221 (>0.5mg (g dry wt)<sup>-1</sup>), the sediment (>0.5mg (g dry wt)<sup>-1</sup>)) and porewater (7.64mg L<sup>-1</sup> alpha,

222 6.82mg L<sup>-1</sup> beta) (Table 1) indicating in-situ alkaline cellulose hydrolysis (Knill and  
223 Kennedy, 2003).

224 The surfaces of the colonised cotton showed areas of EPS indicative of biofilm  
225 formation and surface associated mineral precipitates (Figure 2B and Figure S1) with  
226 individual viable bacterial cells being visible on some fibres (Figure 2C and D).

#### 227 16S rDNA profile of colonised cotton

228 The cotton's Eubacterial clone library (Table S1) was dominated by the Order  
229 Clostridiales which represented 58% of the clones obtained (n=67, Figure 3A). Of these  
230 Clostridia, 33 sequences most closely matched organisms from the family Clostridiaceae 2,  
231 where 13 sequences most closely matched *Clostridium formicaceticum* strain DSM 92 (95%  
232 sequence similarity) and a further 10 to *Anaerovirgula multivorans* strain SCA (97%  
233 sequence similarity). The remaining 9 clones most closely matched sequences belonging to  
234 the genus *Alkaliphilus*, of which 8 were closely related to *Alkaliphilus oremlandii* strain  
235 OhILAs (91-93% sequence similarity) and 1 related to *Alkaliphilus transvaalensis* strain  
236 SAGM1 (98% sequence similarity). The remaining clones of the Clostridia were represented  
237 by sequences most closely related to organisms from the families *Clostridium insertae sedis*  
238 *XI* (3 sequences) and *Clostridium insertae sedis XIV* (2 sequences).

239 The remainder of the clone library was made up of a diverse range of taxonomic  
240 families (Table S1) including: representatives from Brucellaceae, primarily related to  
241 *Ochrobactrum anthropi* strain ATCC 49188 (96-99% sequence similarity);  
242 Corynebacteriaceae, dominated by *Corynebacterium marinum* strain D7015 (98-99%  
243 sequence match similarity); and the Bacillaceae 1, dominated by *Bacillus pseudofirmus* strain  
244 OF4 (89-99% sequence similarity).

245 The Archaeal clone library (Table S2) was dominated (93%) by sequences most  
246 closely matching *Methanobacterium alcaliphilum* strain NBRC 105226 (99% sequence  
247 similarity) (n=68, Figure 3B). The remaining sequences were most closely related to  
248 *Methanomassiliicoccus luminyensis* strain B10 (4 sequences 89% sequence match) and  
249 *Methanosarcina mazei* Go1 (99% sequence similarity).

#### 250 CDP driven microcosms

251 The microcosm demonstrated significant degradation of ISA at pH 11.0 over 2  
252 waste/feed cycles (Figure 4) with first order rate constants of  $3.33 \times 10^{-2} \text{ day}^{-1}$  ( $\text{SE} \pm 2.0 \times 10^{-2}$ )  
253  $^2$ ) for alpha,  $9.36 \times 10^{-2} \text{ day}^{-1}$  ( $\text{SD} \pm 2.2 \times 10^{-2}$ ) for beta and  $6.78 \times 10^{-2} \text{ day}^{-1}$  ( $\text{SE} \pm 2.85 \times 10^{-2}$ )  
254 for X-ISA. Acetate was the only VFA detected and gradually accumulated in the system  
255 reaching a peak of 2.06 mmoles ( $\text{SE} \pm 0.2$ ), similarly hydrogen gas accumulated in the  
256 headspace over the course of the feed cycle reaching 1.00 mmoles ( $\text{SE} \pm 0.04$ ). Neither  
257 carbon dioxide nor methane was detected in the headspace of the microcosm, however,  
258 soluble inorganic carbon increased within the system (data not shown) with the pH after each  
259 cycle having an average pH of 10.80 ( $\text{SE} \pm 0.4$ ). The CDP fed microcosm inoculated with the  
260 colonised cotton was dominated by polymicrobial flocs with fluorescence microscopy  
261 showing microbial cells embedded in an EPS composed of protein, polysaccharide and  
262 extracellular DNA (eDNA) (Figure 5).

263 Measurement of the ATP concentration of the microcosm showed that cell density  
264 increased over the feed/waste cycles (Table S4) indicating that a portion of the organic  
265 carbon was used for the generation of both cell biomass and EPS. Carbon flow calculations  
266 (Rittmann and Mccarty, 2001) based on the degradation of ISA showed 23.7% of the carbon  
267 was converted to acetate and 12.1% converted to carbonate from energy generating  
268 processes, 0.5% was converted to cell biomass and a further 63.7% was theorised to be

269 involved in processes relating to EPS production. The yield of dry cell biomass was 0.012 mg  
270 (mg ISA)<sup>-1</sup> degraded, the system could not be stoichiometrically balanced due to the  
271 unknown composition of the flocculate EPS material. Comparison of the samples amended  
272 with chloramphenicol showed no ISA degradation and the production of acetate and  
273 hydrogen was not detected (Figure S2) indicating that ISA degradation was via microbial  
274 activity rather than chemical processes or sorption.

#### 275 Microcosm clone library

276 The microcosm microbial populations demonstrated a significant shift away from that  
277 associated with the emplaced cotton samples, with Archaeal taxa no longer being detectable  
278 and the Eubacterial population no longer dominated by the Clostridiales. The environmental  
279 and physiological constraints imposed within the microcosm resulted in a population  
280 dominated by clones of *Alishewanella jeotgali* strain MS1 (99% sequence similarity) from  
281 the family Alteromonadaceae (Table S3, Figure 6). The remaining clones included  
282 representatives of the family Bacillaceae, most closely matching *Bacillus pseudofirmus* strain  
283 OF4 (98% sequence similarity) and *Alkaliphilus crotonatoxidans* strain B11-2 (98% sequence  
284 similarity) of the family Clostridiaceae 2.

#### 285 **Discussion**

286 Previous authors noted the presence of an organic electron donor within the soils at  
287 Harpur Hill that allowed for electron flow into nitrate and iron reducing processes at depth  
288 (Burke et al., 2012). The generation of CDPs from the site's soil organic matter has been  
289 demonstrated (Rout et al., 2015) and in this study the addition of cotton cellulose resulted in  
290 its partial alkaline hydrolysis to CDPs with the concentration of alpha and beta ISA in the  
291 porewater and sediments being higher than those measured by Rout et al (2015). This  
292 supports the concept that the hyper-alkaline conditions created at this site are capable of

293 generating CDP. The presence of acetate, a common end product of ISA fermentation (Rout  
294 et al., 2015, Bassil et al., 2014, Rout et al., 2014), in the porewater, sediment and cotton  
295 indicated an active anaerobic microbial community in the immediate proximity of the cotton  
296 even though the ambient pH was between pH 11.5 and 12.

297 Cotton fibres were covered with large areas of EPS indicative of biofilm formation  
298 (Figure 2A and B) with individual cells being only rarely visible (Figure 2C). This is a  
299 marked contrast to the colonisation of cotton incubated in a landfill site under neutral  
300 anaerobic conditions reported by McDonald et al (2012) where fibres were heavily colonised  
301 with cells and exhibited the characteristic pits and grooves associated with microbial  
302 cellulose hydrolysis. The reduced colonisation of the cotton under the hyperalkaline  
303 conditions present at the site are further illustrated by the live/dead staining of the cotton  
304 (Figure 2D) which revealed a low density of live cells on the individual cotton fibres and  
305 within the surrounding biofilm material. Previous work by Grant et al (2002) demonstrated  
306 the ability of alkaliphilic microorganisms to form a biofilm upon the surface of the  
307 cementitious materials presumably to provide a degree of protection from the alkaline  
308 stresses imposed by the local environment. This formation of EPS as a response to  
309 hyperalkaline conditions is replicated in these microcosm studies where a polymicrobial,  
310 eDNA stabilised floc based population developed (Figure 5A). The importance of EPS  
311 generation in this system is illustrated by the fact that >60% of the available carbon is  
312 diverted to EPS formation, a finding similar to the carbon distribution in biofilm systems  
313 reported by Jahn and Nielsen (1998).

314 The microbial flocs were composed of an EPS containing protein, polysaccharides  
315 and eDNA. Polysaccharides are a common component of EPS and moderate a range of  
316 bacterial biofilm properties including adhesion, cell aggregation, cohesive nature, protection  
317 as well as the sorption of organic compounds and inorganic ions (Flemming and Wingender,

318 2010). Imaging of the polysaccharide component revealed its distribution throughout the  
319 flocculate with large globular like structures (Figure 5A). DNase treatment caused the loss of  
320 these structures resulting a less compact structure of cells associated with polysaccharide,  
321 indicating a relationship between the eDNA and the distribution of the polysaccharide  
322 components (Figure 5B). The role of eDNA within biofilms appears to serve a number of  
323 functions (Dominiak et al., 2011), in this case it is likely to aid the structure and function of  
324 the flocculate community (Gloag et al., 2013). The presence of eDNA within the flocculate  
325 structure is also likely to act as a phosphate store for the constituent microbial consortia  
326 (Dell'anno and Danovaro, 2005). Calcium ions are abundant at the site and as such the  
327 interaction between eDNA and these ions is likely to promote cell aggregation and biofilm  
328 formation within these alkaliphilic cultures (Das et al., 2014). This is illustrated by the fact  
329 that treatment of the flocs with DNase resulted in the loss of flocculate stability (Figure 5A  
330 and B). Imaging of the protein component of the flocs showed large concentrated areas of  
331 protein within the flocculate (Figure 5C). Protein serves a wide range of functions within  
332 biofilm including the permitting of redox activity, protection from environmental conditions,  
333 enzymatic reactions and sorption of organic compounds sorption and inorganic ions  
334 (Flemming and Wingender, 2010).

335         The presence of the cotton cellulose within the sediments selected for organisms of  
336 the Order Clostridia which contrasts with previous investigations of the background  
337 sediments where a larger degree of taxonomic diversity was observed (Bassil et al., 2014,  
338 Williamson et al., 2013), presumably due to greater diversity of energy sources and  
339 colonisation from surrounding pasture land. Of the Clostridiaceae 2 species identified,  
340 *Clostridium formicaceticum* has broad spectrum carbohydrate fermentation capabilities  
341 (Andreesen et al., 1970), but was not previously associated with alkaline conditions. This  
342 contrasts with species from the genera *Anaerovirgula* and *Alkaliphilus* which have all been



343 previously associated with alkaline sites (Fisher et al., 2008, Takai et al., 2001, Pikuta et al.,  
344 2006).

345 The Archaeal population associated with the cotton was dominated by  
346 hydrogenotrophic, alkaliphilic *Methanobacterium* sp. showing sequence similarity to  
347 *Methanobacterium alcaliphilum* (Worakit et al., 1986). These findings are in agreement with  
348 clone libraries generated from microcosms previously developed from sediment samples  
349 from the same site (Rout et al., 2015). Although these organisms are able to utilise acetate as  
350 a growth factor (Kotelnikova et al., 1998, Wu et al., 1992), they are incapable of acetoclastic  
351 methanogenesis which accounts for the accumulation of acetic acid in extracts from the  
352 cotton and surrounding sediment and porewaters. In addition a small number of sequences  
353 showing similarity to *Methanomassiliicoccus luminyensis* (Dridi et al., 2012) and  
354 *Methanosarcina* sp. (Maestrojuan et al., 1992) were also detected.

355 The microbial population established in the microcosm was much less diverse than  
356 that present on the cotton samples with the almost complete removal of Clostridia and the  
357 total loss of methanogens from the system. This resulting fermentative system was dominated  
358 (95% of clones) by organisms most closely related to *Alishewanella* sp., which was a minor  
359 component (3% of clones) of the population present on the colonised cotton. This facultative  
360 anaerobic genus is most commonly associated with fermented seafood, but has also been  
361 isolated from landfill soils (Jung et al., 2012, Kim et al., 2009, Kim et al., 2010, Kolekar et  
362 al., 2013). Its ability to grow in alkaline conditions up to pH 12 has also been reported (Kim  
363 et al., 2009, Tarhriz et al., 2012, Kim et al., 2010), and its ability to degrade a range of  
364 substrates appears to have enhanced its ability to thrive within the CDP driven microcosm.  
365 The ability to form biofilms and pellicles has been reported in *Alishewanella jeotgali* which  
366 may indicate a pivotal role for the *Alishewanella* sp. in the formation and maintenance of the  
367 bacterial aggregates within the microcosm (Jung et al., 2012) (Figure 5A).

368           A range of degradation rate constants for the various forms of ISA (alpha, beta and  
369 xylo) were observed in the derived microcosms. The rate constant of beta ISA degradation  
370 was similar to that reported by Rout et al (2015) at pH 11, whilst the rate constant of alpha  
371 ISA degradation was greatly reduced, potentially due to the reduced role of key genera such  
372 as *Alkaliphilus* (Rout et al., 2015). This is the first time that a microbial degradation rate  
373 constant for xylo ISA has been published.

374           The loss of methanogens from the microcosm cannot be entirely attributed to  
375 the pH, since a pH\_11.0 methanogenic microcosm has been successfully established using  
376 sediments from the Buxton site (Rout et al., 2015). In that case a similar range of  
377 methanogens were observed to that identified here associated with the cotton but with a  
378 Eubacterial population dominated by *Alkaliphilus*. The lack of Clostridia species specifically  
379 *Alkaliphilus* sp. within the microcosm formed from the colonised cotton appears to have  
380 retarded the ability of the associated methanogenic population to become established. Tight  
381 adherence to the cotton fibres and a possible differences in redox potential between the  
382 internal biofilm environment and the enrichment media may have also contributed to the poor  
383 transition of the methanogens and Clostridia species leading to an *Alishewanella* dominated  
384 system (Sridhar and Eiteman, 1999, Stuart et al., 1999).

385           The presence of cotton fibres with the hyper alkaline analogue site at Harpur Hill  
386 provided both a source of CDP to drive anoxic metabolism and a surface for microbial  
387 colonisation. Subsequent sub culturing indicated that the cotton provided a surface for the  
388 adherence of a narrow range of Clostridiaceae 2 species and promoted the development of a  
389 floc based alkaliphilic population dominated by *Alishewanella* sp. able to degrade CDP up to  
390 a pH of 11.0. Although methanogenic populations were detected on the cotton fibres, they  
391 were unable to make the transition to floc based suspended growth.

392 **Funding**

393 The work here was partially supported by both Radioactive Waste Management Ltd  
394 (RWM Ltd) and the C14-BIG project (EPSRC, EP/I036354/1).

395 **Acknowledgements**

396 The authors would like to thank Peter Finan for his assistance with preparing cotton samples  
397 for emplacement.

398 **References**

- 399 Allard, S. & Ekberg, C. Complexing properties of  $\alpha$ -Isosaccharinate: stability constants,  
400 enthalpies and entropies of Th-complexation with uncertainty analysis. *Journal of*  
401 *Solution Chemistry* 2006; **35**:1173-1186.
- 402 Almond, M., Shaw, P. B., Humphreys, P. N., Chadha, M. J., Niemelä, K. & Laws, A. P.  
403 Behaviour of xyloisosaccharinic acid and xyloisosaccharino-1, 4-lactone in aqueous  
404 solutions at varying pHs. *Carbohydrate Research* 2012; **363**:51-57.
- 405 Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman,  
406 D. J. Gapped BLAST and PSI-BLAST: a new generation of protein database search  
407 programs. *Nucleic Acids Research* 1997; **25**:3389-3402.
- 408 Andreesen, J. R., Gottschalk, G. & Schlegel, H. G. *Clostridium formicoaceticum* nov. spec.  
409 isolation, description and distinction from *C. aceticum* and *C. thermoaceticum*.  
410 *Archives of microbiology* 1970; **72**:154-174.
- 411 B.S.I. BS ISO 11271:2002 Soil quality — Determination of redox potential — Field method.  
412 British Standards Institute, London, UK. 2002. BS ISO 11271:2002.
- 413 B.S.I. BS ISO 14853:2005 Plastics-Determination of the ultimate anaerobic biodegradation  
414 of plastic materials in an aqueous system- Method by measurement of biogas  
415 production. British Standards Institute, London, UK. 2005. BS ISO 14853:2005.
- 416 Bassil, N. M., Bryan, N. & Lloyd, J. R. Microbial degradation of isosaccharinic acid at high  
417 pH. *The ISME Journal* 2014; **9**:310-320.
- 418 Bethke, C. M., Sanford, R. A., Kirk, M. F., Jin, Q. & Flynn, T. M. The thermodynamic ladder  
419 in geomicrobiology. *American Journal of Science* 2011; **311**:183-210.
- 420 Burke, I. T., Mortimer, R. J., Palaniyandi, S., Whittleston, R. A., Lockwood, C. L., Ashley,  
421 D. J. & Stewart, D. I. Biogeochemical reduction processes in a hyper-alkaline  
422 leachate affected soil profile. *Geomicrobiology Journal* 2012; **29**:769-779.
- 423 Chapman, N. & Hooper, A. The disposal of radioactive wastes underground. *Proceedings of*  
424 *the Geologists' Association*. 2012; **123**:46-63.
- 425 Cole, J. R., Wang, Q., Cardenas, E., Fish, J., Chai, B., Farris, R. J., Kulam-Syed-Mohideen,  
426 A. S., Mcgarrell, D. M., Marsh, T., Garrity, G. M. & Tiedje, J. M. The Ribosomal  
427 Database Project: improved alignments and new tools for rRNA analysis. *Nucleic*  
428 *Acids Research* 2009; **37**:D141-D145.

- 429 Conrad, R., Ji, Y., Noll, M., Klose, M., Claus, P. & Enrich-Prast, A. Response of the  
430 methanogenic microbial communities in Amazonian oxbow lake sediments to  
431 desiccation stress. *Environmental Microbiology* 2014; **16**:1682-1694.
- 432 Das, T., Sehar, S., Koop, L., Wong, Y. K., Ahmed, S., Siddiqui, K. S. & Manefield, M.  
433 Influence of calcium in extracellular DNA mediated bacterial aggregation and biofilm  
434 formation. *PloS one* 2014; **9**:e91935.
- 435 Dell'anno, A. & Danovaro, R. Extracellular DNA plays a key role in deep-sea ecosystem  
436 functioning. *Science* 2005; **309**:2179.
- 437 Dominiak, D. M., Nielsen, J. L. & Nielsen, P. H. Extracellular DNA is abundant and  
438 important for microcolony strength in mixed microbial biofilms. *Environmental*  
439 *Microbiology* 2011; **13**:710-721.
- 440 Dridi, B., Fardeau, M.-L., Ollivier, B., Raoult, D. & Drancourt, M. *Methanomassiliicoccus*  
441 *luminyensis* gen. nov., sp. nov., a methanogenic archaeon isolated from human faeces.  
442 *International Journal of Systematic and Evolutionary Microbiology* 2012; **62**:1902-  
443 1907.
- 444 Eaton, A. D., Clesceri, L. S., Rice, E. W. & Greenberg, A. E. (eds.) 2005. *Standard Methods*  
445 *for the Examination of Water and Wastewater 21st Edition*, Washington, US:  
446 APHA/AWWA/WEF.
- 447 Edwards, U., Rogall, T., Blöcker, H., Emde, M. & Böttger, E. C. Isolation and direct  
448 complete nucleotide determination of entire genes. Characterization of a gene coding  
449 for 16S ribosomal RNA. *Nucleic Acids Research* 1989; **17**:7843-7853.
- 450 Fisher, E., Dawson, A. M., Polshyna, G., Lisak, J., Crable, B., Perera, E., Ranganathan, M.,  
451 Thangavelu, M., Basu, P. & Stolz, J. F. Transformation of inorganic and organic  
452 arsenic by *Alkaliphilus oremlandii* sp. nov. strain OhILAs. *Annals of the New York*  
453 *Academy of Sciences* 2008; **1125**:230-241.
- 454 Flemming, H. C. & Wingender, J. The biofilm matrix. *Nature Reviews Microbiology* 2010;  
455 **8**:623-33.
- 456 Gantner, S., Andersson, A. F., Alonso-Sáez, L. & Bertilsson, S. Novel primers for 16S  
457 rRNA-based archaeal community analyses in environmental samples. *Journal of*  
458 *Microbiological Methods* 2011; **84**:12-18.
- 459 Gloag, E. S., Turnbull, L., Huang, A., Vallotton, P., Wang, H., Nolan, L. M., Mililli, L.,  
460 Hunt, C., Lu, J. & Osvath, S. R. Self-organization of bacterial biofilms is facilitated  
461 by extracellular DNA. *Proceedings of the National Academy of Sciences* 2013;  
462 **110**:11541-11546.
- 463 Gorlenko, V., Tsapin, A., Namsaraev, Z., Teal, T., Tourova, T., Engler, D., Mielke, R. &  
464 Nealsen, K. *Anaerobranca californiensis* sp. nov., an anaerobic, alkalithermophilic,  
465 fermentative bacterium isolated from a hot spring on Mono Lake. *International*  
466 *Journal of Systematic and Evolutionary Microbiology* 2004; **54**:739-743.
- 467 Grant, W. D., Holtom, G. J., O'Kelly, N., Malpass, J., Rosevear, A., Watkiss, P. &  
468 Widdowson, D. 2002. Microbial degradation of cellulose-derived complexants under  
469 repository conditions. AEA Technology and University of Leicester for UK Nirex  
470 Ltd. AEAT/ERRA-0301.
- 471 Greenfield, B. F., Moreton, A. D., Spindler, M. W., Williams, S. J. & Woodwark, D. R. The  
472 effects of the degradation of organic materials in the near field of a radioactive waste  
473 repository. *MRS Online Proceedings Library* 1991; **257**.

474 Humphreys, P., Laws, A. & Dawson, J. A review of cellulose degradation and the fate of  
475 degradation products under repository conditions. SERCO / TAS / 002274 / 001. Serco  
476 contractors report for the Nuclear Decommissioning Authority, UK 2010a.

477 Humphreys, P. N., West, J. M. & Metcalfe, R. 2010b. Microbial effects on repository  
478 performance. Quintessa contractors report for the Nuclear Decommissioning  
479 Authority, UK. QRS-1378Q-1.

480 Jahn, A. & Nielsen, P. H. Cell biomass and exopolymer composition in sewer biofilms.  
481 *Water Science and Technology* 1998; **37**:17-24.

482 Jones, B. E., Grant, W. D., Collins, N. C. & Mwatha, W. E. 1994. Alkaliphiles: diversity and  
483 identification. *Bacterial Diversity and Systematics*. Springer.

484 Jung, J., Chun, J. & Park, W. Genome sequence of extracellular-protease-producing  
485 *Alishewanella jeotgali* isolated from traditional Korean fermented seafood. *Journal of*  
486 *Bacteriology* 2012; **194**:2097.

487 Kim, M.-S., Jo, S. K., Roh, S. W. & Bae, J.-W. *Alishewanella agri* sp. nov., isolated from  
488 landfill soil. *International Journal of Systematic and Evolutionary Microbiology*  
489 2010; **60**:2199-2203.

490 Kim, M. S., Roh, S. W., Nam, Y. D., Chang, H. W., Kim, K. H., Jung, M. J., Choi, J. H.,  
491 Park, E. J. & Bae, J. W. *Alishewanella jeotgali* sp. nov., isolated from traditional  
492 fermented food, and emended description of the genus *Alishewanella*. *International*  
493 *Journal of Systematic and Evolutionary Microbiology* 2009; **59**:2313-6.

494 Knill, C. J. & Kennedy, J. F. Degradation of cellulose under alkaline conditions.  
495 *Carbohydrate Polymers* 2003; **51**:281-300.

496 Kolekar, Y. M., Konde, P. D., Markad, V. L., Kulkarni, S. V., Chaudhari, A. U. & Kodam, K.  
497 M. Effective bioremoval and detoxification of textile dye mixture by *Alishewanella*  
498 sp. KMK6. *Applied Microbiology and Biotechnology* 2013; **97**:881-889.

499 Kotelnikova, S., Macario, A. J. & Pedersen, K. *Methanobacterium subterraneum* sp. nov., a  
500 new alkaliphilic, eurythermic and halotolerant methanogen isolated from deep granitic  
501 groundwater. *International Journal of Systematic Bacteriology* 1998; **48**:357-367.

502 Libert, M., Bildstein, O., Esnault, L., Jullien, M. & Sellier, R. Molecular hydrogen: An  
503 abundant energy source for bacterial activity in nuclear waste repositories. *Physics*  
504 *and Chemistry of the Earth, Parts A/B/C* 2011; **36**:1616-1623.

505 Maestrojuan, G. M., Boone, J. E., Mah, R. A., Menaia, J. A., Sachs, M. S. & Boone, D. R.  
506 Taxonomy and halotolerance of mesophilic *Methanosarcina* strains, assignment of  
507 strains to species, and synonymy of *Methanosarcina mazei* and *Methanosarcina*  
508 *frisla*. *International Journal of Systematic Bacteriology* 1992; **42**:561-567.

509 McDonald, J. E., Houghton, J. N., Rooks, D. J., Allison, H. E. & McCarthy, A. J. The  
510 microbial ecology of anaerobic cellulose degradation in municipal waste landfill sites:  
511 evidence of a role for fibrobacters. *Environmental Microbiology* 2012; **14**:1077-1087.

512 Motellier, S., Richet, C. & Merel, P. Analysis of cellulose degradation products by capillary  
513 electrophoresis. *Journal of Chromatography A* 1998; **804**:363-370.

514 N.D.A. Geological Disposal: An introduction to the generic disposal system safety case.  
515 Nuclear Decommissioning Authority. Harwell, Didcot, Oxfordshire, UK. 2010a.

516 N.D.A. Near-field Evolution Status Report Nuclear Decommissioning Authority. Harwell,  
517 Didcot, Oxfordshire, UK 2010b. NDA/RWMD/033.

518 N.D.A. Radioactive Wastes in the UK: A Summary of the 2013 Inventory. Nuclear  
519 Decommissioning Authority. Harwell, Didcot, Oxfordshire, UK. 2013.

520 Ordoñez, O. F., Flores, M. R., Dib, J. R., Paz, A. & Farías, M. E. Extremophile culture  
521 collection from Andean lakes: extreme pristine environments that host a wide  
522 diversity of microorganisms with tolerance to UV radiation. *Microbial Ecology* 2009;  
523 **58**:461-473.

- 524 Pikuta, E. V., Itoh, T., Krader, P., Tang, J., Whitman, W. B. & Hoover, R. B. *Anaerovirgula*  
525 *multivorans* gen. nov., sp. nov., a novel spore-forming, alkaliphilic anaerobe isolated  
526 from Owens Lake, California, USA. *International Journal of Systematic and*  
527 *Evolutionary Microbiology* 2006; **56**:2623-9.
- 528 Randall, M., Rigby, B., Thomson, O. & Trivedi, D. Assessment of the effects of cellulose  
529 degradation products on the behaviour of europium and thorium. Report prepared on  
530 behalf of NDA by National Nuclear Laboratory. 2013. NNL (12) 12239 Part A, Issue  
531 4.
- 532 Rittmann, B. E. & Mccarty, P. L. *Environmental biotechnology*, McGraw Hill New  
533 York.2001
- 534 Rizoulis, A., Steele, H., Morris, K. & Lloyd, J. The potential impact of anaerobic microbial  
535 metabolism during the geological disposal of intermediate-level waste. *Mineralogical*  
536 *Magazine* 2012; **76**:3261-3270.
- 537 Rodrigues, D. F., Goris, J., Vishnivetskaya, T., Gilichinsky, D., Thomashow, M. F. & Tiedje,  
538 J. M. Characterization of *Exiguobacterium* isolates from the Siberian permafrost.  
539 Description of *Exiguobacterium sibiricum* sp. nov. *Extremophiles* 2006; **10**:285-294.
- 540 Rout, S. P., Charles, C. J., Garratt, E. J., Laws, A. P., Gunn, J. & Humphreys, P. N. Evidence  
541 of the generation of isosaccharinic acids and their subsequent degradation by local  
542 microbial consortia within hyper-alkaline contaminated soils, with relevance to  
543 intermediate level radioactive waste disposal. *PloS one* 2015; **10**:e0119164.
- 544 Rout, S. P., Radford, J., Laws, A. P., Sweeney, F., Elmekawy, A., Gillie, L. J. & Humphreys,  
545 P. N. Biodegradation of the alkaline cellulose degradation products generated during  
546 radioactive waste disposal. *PloS one* 2014; **9**:e107433.
- 547 Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B.,  
548 Lesniewski, R. A., Oakley, B. B., Parks, D. H. & Robinson, C. J. Introducing mothur:  
549 open-source, platform-independent, community-supported software for describing and  
550 comparing microbial communities. *Applied and Environmental Microbiology* 2009;  
551 **75**:7537-7541.
- 552 Shaw, P. B., Robinson, G. F., Rice, C. R., Humphreys, P. N. & Laws, A. P. A robust method  
553 for the synthesis and isolation of  $\beta$ -gluco-isosaccharinic acid ((2R,4S)-2,4,5-  
554 trihydroxy-2-(hydroxymethyl)pentanoic acid) from cellulose and measurement of its  
555 aqueous pKa. *Carbohydrate Research* 2012; **349**:6-11.
- 556 Sridhar, J. & Eiteman, M. A. Influence of redox potential on product distribution in  
557 *Clostridium thermosuccinogenes*. *Applied Biochemistry and Biotechnology* 1999;  
558 **82**:91-101.
- 559 Stuart, S. L., Woods, S. L., Lemmon, T. L. & Ingle, J. D. The effect of redox potential  
560 changes on reductive dechlorination of pentachlorophenol and the degradation of  
561 acetate by a mixed, methanogenic culture. *Biotechnology and Bioengineering* 1999;  
562 **63**:69-78.
- 563 Summons, R. E., Schubotz, F., Hays, L. E., Meyer-Dombard, D., Gillespie, A. & Shock, E.  
564 Stable isotope labeling confirms mixotrophic nature of streamer biofilm communities  
565 at alkaline hot springs. *Name: Frontiers in Microbiology* 2015; **6**:42.
- 566 Takai, K., Moser, D. P., Onstott, T. C., Spoelstra, N., Pfiffner, S. M., Dohnalkova, A. &  
567 Fredrickson, J. K. *Alkaliphilus transvaalensis* gen. nov., sp. nov., an extremely  
568 alkaliphilic bacterium isolated from a deep South African gold mine. *International*  
569 *Journal of Systematic and Evolutionary Microbiology* 2001; **51**:1245-56.
- 570 Tarhriz, V., Nematzadeh, G., Vahed, S. Z., Hejazi, M. A. & Hejazi, M. S. *Alishewanella*  
571 *tabrizica* sp. nov., isolated from Qurugöl Lake. *International Journal of Systematic*  
572 *and Evolutionary Microbiology* 2012; **62**:1986-1991.

- 573 Urbietta, M. S., González-Toril, E., Bazán, Á. A., Giaveno, M. A. & Donati, E. Comparison  
574 of the microbial communities of hot springs waters and the microbial biofilms in the  
575 acidic geothermal area of Copahue (Neuquén, Argentina). *Extremophiles* 2015:1-14.
- 576 Van Loon, L. R. & Glaus, M. A. Review of the kinetics of alkaline degradation of cellulose  
577 in view of its relevance for safety assessment of radioactive waste repositories.  
578 *Journal of Environmental Polymer Degradation* 1997; **5**:97-109.
- 579 Warwick, P., Evans, N., Hall, T. & Vines, S. 2003. Complexation of Ni(II) by  $\alpha$ -  
580 isosaccharinic acid and gluconic acid from pH 7 to pH 13. *Radiochimica*  
581 *Acta/International journal for chemical aspects of nuclear science and technology*.
- 582 Williamson, A. J., Morris, K., Shaw, S., Byrne, J. M., Boothman, C. & Lloyd, J. R. Microbial  
583 reduction of Fe(III) under alkaline conditions relevant to geological disposal. *Applied*  
584 *Environmental Microbiology* 2013; **79**:3320-3326.
- 585 Worakit, S., Boone, D. R., Mah, R. A., Abdel-Samie, M.-E. & El-Halwagi, M.  
586 *Methanobacterium alcaliphilum* sp. nov., an H<sub>2</sub>-utilizing methanogen that grows at  
587 high pH values. *International Journal of Systematic Bacteriology* 1986; **36**:380-382.
- 588 Wu, W.-M., Jain, M. K., De Macario, E. C., Thiele, J. H. & Zeikus, J. G. Microbial  
589 composition and characterization of prevalent methanogens and acetogens isolated  
590 from syntrophic methanogenic granules. *Applied Microbiology and Biotechnology*  
591 1992; **38**:282-290.
- 592

592

593 Table 1: Analysis of porewater, sediment and cotton retrieved from sample borehole.

594

Source	pH	eH	Acetate	$\alpha$ -ISA	$\beta$ -ISA
Porewater (mg L <sup>-1</sup> )	11.92	-66.00	208.90	7.64	6.82
Sediment (mg (g dry wt) <sup>-1</sup> )					
1)	11.50	-77.00	127.24	1.01	0.54
Cotton (mg (g dry wt) <sup>-1</sup> )	N/S	N/S	141.16	2.34	0.85


595 \*N/S-Not sampled

596



596

597




598

599 Figure 1: Overview of hyperalkaline contaminated site and position of emplaced

600 cotton within bore hole 1 (ECBH-1)

601


601



602

603 Figure 2: Microscopy investigation of the cellulose cotton. [A] Sterile cotton. [B]  
604 Cellulose cotton from the borehole showing biofilm formation. [C] Close up of  
605 individual fibre showing individual cells, EPS aggregates and mineral precipitate. [D]  
606 Live/dead image of individual cotton fibre.

607




608

609

610 Figure 3: 16S rRNA gene clone libraries of the colonised cotton. [A] Eubacterial  
 611 (n=67). [B] Archaeal (n=68). Phylogenetic families were assigned to clones through a  
 612 MegaBLAST database search.

613


613



614

615 Figure 4: Chemistry of the CDP driven pH 11 microcosm over two waste/feed cycles  
616 using colonised cellulose cotton as an inoculation source. [A] Alpha, beta and xylo  
617 isosaccharinic acid degradation profile. [B] Hydrogen and acetate production profile.

618



619


620 Figure 5: Microscopy investigation into the morphology of the pH 11 microcosm. [A]  
 621 Ethidium bromide and Calcofluor white stain of bacterial flocculate showing individual  
 622 cells and eDNA (red) and extracellular polysaccharides (blue). [B] DNase digest of  
 623 flocculates stained with ethidium bromide and Calcofluor white. [C] FITC stain of

624 bacterial flocculate showing areas containing protein (green). [D] Live/dead image of

625 bacterial flocculate.

626

626



627

628 Figure 6: Eubacterial (n=59) 16s rRNA gene clone library of pH 11 CDP driven  
629 microcosm.

630

631