



University of HUDDERSFIELD

University of Huddersfield Repository

Edward, Karen-Leigh and Ousey, Karen

Exploring resilience for people with type 2 diabetes who have a wound

Original Citation

Edward, Karen-Leigh and Ousey, Karen (2015) Exploring resilience for people with type 2 diabetes who have a wound. *Wounds UK*, 11 (2). pp. 28-31. ISSN 1746-6814

This version is available at <http://eprints.hud.ac.uk/25513/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>

Exploring resilience for people with type 2 diabetes who have a wound

KEY WORDS

- ▶ Diabetes
- ▶ Type 2
- ▶ Wounds
- ▶ Wound healing
- ▶ Resilience
- ▶ Self-management

Comorbidities related to diabetes such as chronic wounds can increase the need for surgical procedures, with at least 10% of all patients undergoing surgery having diabetes. Being diagnosed with diabetes can be seen as a risk factor for developing wound healing problems. **Discussion:** This short report highlights the potential positive influences gained from providing resilience education and self-management education to people with type 2 diabetes, potentially enhancing self-managing abilities and reducing poor wound healing. **Summary:** Modern wound care practice is centred on symptom reduction and working with pathology; however, working with people to enhance their personal resilience and promoting positive psychological adaptation can impact positively on their mental health.

As a chronic condition diabetes is endemic, where the prevalence of diabetes in England in 2013 was 3.2 million people (with an estimated 630,000 undiagnosed cases) (Holman et al, 2014) and in Australia affects approximately one million people (Australian Bureau of Statistics, 2014). Concomitant factors such as obesity, comorbidity and poor homeostatic control place a significant number of people with diabetes at risk of developing both micro and macro complications (Huxley et al, 2006), resulting in accelerated mortality and economic burden (Lee et al, 2013). The complications arising from having a diabetes diagnosis are most evident in hospital admissions where, regardless of the cause for admission, people with diabetes experience a prolonged length of stay. For those who have a wound, the complications of diabetes escalate. For example, neuropathy and ischaemia can lead to foot ulceration, unstable blood sugars, and increased risk of infection, potentially resulting in surgery for lower-limb amputations. Unsurprisingly, at least 10% of all patients undergoing surgery have diabetes. In consideration of the World Health Organisation (WHO) position on attending to the mental health and physical needs of mental health consumers (WHO, 2013), care for people living with serious mental illness still needs to improve in this area. When chronic conditions are comorbid (such as mental

illness, diabetes and diabetes-related wounds), quality of life can be reduced. Quality of life is associated with the individual's perception of their own life and in this context the importance of working with people in building their personal self-righting capabilities — i.e. resilience — is heightened (Edward et al, 2010). The purpose of this short communication is to elucidate the potential positive influences from enhancing resilience and self-management for people with type 2 diabetes who have a diabetes-related or influenced wound towards positive psychological adaptation. Since this article was a review of the literature and no primary research was undertaken ethical approval was not required.

UNDERSTANDING THE COST OF DIABETES

There are approximately 347 million people living with diabetes worldwide, with a predicted increase to almost 552 million people by 2030 (WHO, 2014). The WHO also predicts that diabetes will be the 7th leading cause of death by 2030. Type 2 diabetes accounts for between 85 and 95 per cent of cases in high-income regions, but is estimated to be much higher in regions of lower income. In the UK, a point prevalence estimate of 3.9% people with diabetes has been suggested, with this rising to above 15% in certain ethnic minority origin communities.

KAREN-LEIGH EDWARD
Associate Professor of Nursing Research, Nursing Research Unit Faculty of Health Sciences, Australian Catholic University; Director/Chair, St Vincent's Private Hospital Melbourne, Australia

KAREN OUSEY
Reader Advancing Clinical Practice, Ramsden Building, School of Human and Health Sciences, University of Huddersfield, Huddersfield

Similarly, recent information about the incidence of diabetes in Australia according to the Australian Institute of Health and Welfare (AIHW, 2014) revealed over 4% of the population had diabetes, an increase from 1.5% prevalence in 1989. In 2012, 8.2 million pharmaceutical scripts were claimed for diabetes, with diabetes healthcare expenditure accounting for 2.3% of all health expenditure in Australia. The Australian hospital statistics 2012–13 (AIHW, 2014) revealed that there were 2.5 million admissions that involved surgery; of these 2 million were elective and 67% of these were undertaken in a private hospital. Potentially preventable hospitalisations were identified as relating to conditions that theoretically would not result in hospitalisation if adequate and timely non-hospital care was received. These included such conditions as appendicitis, dehydration, asthma and diabetes. Diabetes complications resulting in hospitalisations during the 2012–13 period in Australia accounted for over 200,000 hospital separations and is also listed in the top 20 common principal diagnoses for emergency admissions involving surgery in both public and private hospitals during the 2012–13 period. Alarming, reported type 2 diabetes in all diagnoses (either as principal diagnosis or a comorbidity) has jumped from approximately 500,000 diagnoses in the period 2008–9 to over 1 million in 2012–13 (AIHW, 2014). Diabetes is associated with adverse events following surgery, which include: increased length of stay, morbidity rates and inpatient costs. They are well recognised, such that the excess bed days were recently estimated to be 45% greater for surgical patients with diabetes, and of significance, the surgical mortality rate is up to 50% higher than that of the non-diabetic population (Frisch et al, 2010).

STRESS AND WOUND MANAGEMENT

The immunosuppressive effects of diabetes and its impact on healthy wound healing (Shupp and Moffatt, 2014) are of consideration for people with diabetes. Often when a person with diabetes is inflicted with an acute wound it can develop into a chronic or infected wound due to poor healing as part of their condition. The burden of chronic wounds is well known, including the impact on both physical and psychological wellbeing due to the multiple effects of pain, reduced mobility, poor circulation and stigma. The pathophysiology of healing in both acute and chronic wounds is still an area of scientific examination, as little is known

about the interplay of mental state on wound healing. The available evidence suggests that a person's perception of their wounds contributes to healing, and wound healing can be negatively impacted by stress (Gouin and Kiecolt-Glaser, 2011). A recent systematic review with meta-analysis (Walburn et al, 2009) reported that stress was significantly associated with impaired healing ($r=0.42$; $P<0.01$). When people have an active involvement in the care of their wounds, however, outcomes are often improved (Broadbent and Koschwanez, 2012). Active participation of consumers in healthcare services, healthcare developments and individual care can lead to improved healthcare outcomes. Use of health-related education to build knowledge and mastery can offer a vehicle for patients to become more engaged in their care, improve clinical outcomes and enhance personal resilience.

THE IMPORTANCE OF BUILDING RESILIENCE

Modern health practice is centred on symptom reduction and working with pathology; however, working with people's resilience has not been a consideration. In light of the noticeable absence of a focus on psychological resilience as a method of intervention, and a scarcity of research on resilience in relation to resistance to the stress of chronic conditions, the findings of a study undertaken by Edward et al (2009) has the potential to make a substantive contribution to existing knowledge on the topic. The results of Edward et al's study (2009) identified how being resilient was experienced by people through sharing their experience with others and realising they were not alone, and an acceptance of self, others and the situation were integral to going beyond the negative impact of being in poor health. Edward (2013) later offers a model for conceptualising and building resilience in people with chronic conditions (such as mental illness, cardiac conditions and diabetes). In practice this is underpinned by the premise that when individuals believe life events and outcomes are manageable, learned helplessness is avoided and active attempts are made by the individual to overcome aversive situations, opening the possibility of moving forward and achieving resilience (Edward, 2013; 2014). By being resilient, individuals have the power to adjust, resist stress and potentially thrive in the face of adversity (Edward, 2005b). For healthcare professionals, exploration of resilience included in

“The complications arising from having a diabetes diagnosis is most evident in hospital admissions where regardless of the cause for admission, people with diabetes experience a prolonged length of stay.”

REFERENCES

- Australian Bureau of Statistics (2014) *Australian Health Survey: First Results, 2011–12* Available at: <http://www.abs.gov.au/ausstats/abs@.nsf/Lookup/4364.0.55.001main+features12011-12> (accessed 4.06.2015)
- Australian Institute of Health and Welfare (2014) *Australian Hospital Statistics 2012–13*. Available at: <http://www.aihw.gov.au/publication-detail/?id=60129546922> (accessed 4.06.2015)
- Broadbent E, Koschwanez HE (2012) The psychology of wound healing. *Curr Opin Psychiatry* 25(2):135–40
- Cole-King A, Harding KG (2001) Psychological factors and delayed healing in chronic wounds. *Psychosom Med* 63(2):216–20
- Edward KL (2005a) The phenomenon of resilience in crisis care mental health clinicians. *Int J Ment Health Nurs* 14(2):142–48
- Edward KL (2005b) Resilience: A Protector From Depression. *Journal of the American Psychiatric Nurses Association* 11(4):241–43
- Edward KL (2013) Chronic illness and wellbeing: using nursing practice to foster resilience as resistance. *Br J Nurs* 22(13):741–2,744,746
- Edward KL (2014) Empowering cardiac patients using resilience as a practice strategy. *British Journal of Cardiac Nursing* 9(3):110
- Edward KL, Rasmussen B, Munro I (2010) Nursing care of clients treated with atypical antipsychotics who have a risk of developing metabolic instability and/or type 2 diabetes. *Arch Psychiatr Nurs* 24(1):46–53
- Edward KL, Welch A, Chater K (2009) The phenomenon of resilience as described by adults who have experienced mental illness. *J Adv Nurs* 65(3):587–95
- Finestone HM, Alfeeli A, Fisher WA (2008) Stress-induced physiologic changes as a basis for the biopsychosocial model of chronic musculoskeletal pain: a new theory? *Clin J Pain* 24(9):767–75
- Frisch A, Chandra P, Smiley D et al (2010) Prevalence and clinical outcome of hyperglycemia in the perioperative period in noncardiac surgery. *Diabetes Care* 33(8):1783–8
- Gouin JP, Kiecolt-Glaser JK (2011) The impact of psychological stress on wound healing: methods and mechanisms. *Immunol Allergy Clin North Am* 31(1):81–93
- the assessment phase of the health care continuum could offer a basis for clinical interventions that can assist the person to move beyond the difficulties of their condition. The International Diabetes Federation (IDF, 2012) recognised that psychological wellbeing and working with people and people's strength were important goals of medical care, and psychosocial factors are relevant to a wide range of considerations in diabetes management.

SELF-MANAGEMENT TO REDUCE WOUND COMPLICATIONS

Chronic conditions such as diabetes require a considerable amount of patient involvement for effective self-management, improving knowledge and for making healthy lifestyle choices. For people living with type 2 diabetes, diabetes education has been standard care. However, it is clear that the provision of knowledge alone does not change behaviour and care goals need to be achievable (Snow et al, 2014). Evidence related to effective education programmes focusing on reducing wound complications for people with diabetes emphasises the importance for control of blood glucose levels, regular screening, immediate self-reporting of any wound changes to health professionals and the use of a simple daily self-care routine. Also, education that entails one-on-one support from a healthcare professional trained in wound care practices and principles, in conjunction with an individualised program reflecting the recipient's specific and unique experiences is recommended as best practice (Rankin et al, 2012).

The available research regarding the superiority of which education approach to adopt, however, appears undecided and this may be potentially due to the heterogeneity and complexity of this patient population (Rygg et al, 2012). Encouragingly, recent evidence supports a significant positive impact in the ability to self-care through education, with improvements in outcomes compared to baseline post education as high as 84% (Kafaie et al, 2012). This finding reinforces the assertion made by some researchers for the need to focus efforts on enhancing self-management of wound care and potentially enhancing resilience, and adaptation rather than conveying information that may not be specific for self-management (Rosenthal, 2013; Haas et al, 2013). The IDF (L'Heveder and Nolan, 2013) recognised that diabetes self-management is critical in enabling individuals to learn the skills to be effective self-managers; however, they suggest that

these programmes should consider putting a greater emphasis on the promotion of positive behaviour change. This has an important focus, as knowledge about the intrinsic and extrinsic factors that increase self-management capabilities has the potential to guide appropriate self-management choices with regard to wound care with the back-drop of having the chronic condition of type 2 diabetes. As such, practitioners who care for people with type 2 diabetes who have a wound should ensure that self-management programmes include information and related activities for both diabetes and wound management and should be seen as one programme rather than two separate conditions.

PSYCHOLOGICAL STRESS AND IMPAIRED HEALING

A range of clinical studies (Loo et al, 2007; Li et al, 2007; Schrementi et al, 2008) have identified that psychological stress is associated with impaired healing or dysregulation of a biomarker associated with wound healing across different clinical and experimental wounds in both cutaneous and mucosal tissue types. Intervention studies have provided evidence of an association between stress and healing (Holden-Lund, 1988), with stress being associated with slower or delayed wound healing in stressed older adults, adults with leg wounds, people with diabetes and surgical patients (Finestone et al, 2008). Cole-King and Harding (2001) examined the relationship between the healing of chronic wounds and anxiety and depression using the Hospital Anxiety and Depression Scale (HADS). Psychological and clinical wound assessments were conducted, with the investigators and participants blinded to the results of the other assessments. The relationship between the healing of chronic wounds and anxiety and depression was statistically significant: delayed healing was associated with a higher mean Hospital Anxiety and Depression Scale score ($P=0.03$), demonstrating that symptoms of both depression and anxiety were associated with chronic wound healing. Walburn et al (2009) in their systematic review examining the effect of stress on wound healing concluded that the size of the relationship between stress and wound healing was estimated to be $r=-0.42$, classified as a medium effect size, suggesting that it may be of significance clinically as well as statistically. Resilience as resistance to the stress associated with both wound care and diabetes management

can offer a new way of working with service users/consumers/patients. That is, by being resilient, individuals have the power to adjust, resist stress and potentially thrive in the face of adversity (Edward 2005a; 2005b; 2013, 2014).

Wounds are common in people with diabetes due to impaired healing as part of the diabetic condition. With the ever-increasing priority of healthcare providers internationally to reduce length of hospitalisations, it can be assumed there will be an increase in the amount of people with diabetes and a wound who are cared for in the community. A large number of these people may present with a further complication of mental illness such as depression or anxiety. This therefore highlights the importance of effective support, development of resilient strategies, which enables the individual to manage their diabetes while allowing them to manage stress or anxiety that may be a secondary outcome of having a wound. It is recognised that psychosocial factors, such as anxiety and depression, are associated with delayed healing of wounds, and that when patients are actively involved in their care, outcomes are improved. Psychologically, stress may increase the likelihood of patients making cognitive errors or negative appraisals. A prolonged fight- or flight-response may result from these negative feelings that affect biological and behavioural responses,

which may have a negative effect on HbA1C (glycated haemoglobin). The notion of developing resilience is essential if patients with mental health problems and their families/carers are able to transcend the potential negative impacts of living with comorbid chronic conditions (mental illness, diabetes and a wound).

SUMMARY

When people with wounds are able to accept the wound may impinge on normal day-to-day activities and develop strategies that manage this temporary change in lifestyle, their stress levels may be kept to a minimum. Complications of diabetes can impact on the normal trajectory of wound healing, for example, neuropathy and ischemia, which may lead to foot ulceration; unstable blood sugars; and an increased risk of infection that can lead to lower-limb amputation. The ability to develop resilient behaviours has the potential to facilitate a positive outcome for quality of life and subjective wellbeing. **WUK**

ACKNOWLEDGMENTS

The authors wish to acknowledge the continued support of each author affiliation through a memorandum of understanding between organisations. No funding was received in the generation of this review.

- Haas L, Maryniuk M, Beck J et al (2012) Standards Revision Task Force. National standards for diabetes self-management education and support. *Diabetes Care* 36(Suppl 1):S100–8
- Holden-Lund C (1988) Effects of relaxation with guided imagery on surgical stress and wound healing. *Res Nurs Health* 11(4):235–44
- Holman N, Young B, Gadsby R (2014) What is the current prevalence of diagnosed and yet to be diagnosed diabetes in the UK? *Diabet Med* 31(5):510–1
- Huxley R, Barzi F, Woodward M (2006) Excess risk of fatal coronary heart disease associated with diabetes in men and women: meta-analysis of 37 prospective cohort studies. *BMJ* 332(7533):73–8
- International Diabetes Federation (2012) *Global Guideline for Type 2 Diabetes*. Available at: <http://www.idf.org/sites/default/files/IDF-Guideline-for-Type-2-Diabetes.pdf> (accessed 4.06.2015)
- Kafaie P, Noorbala MT, Soheilikhah S, Rashidi M (2012) Evaluation of Patients' Education on Foot Self-Care Status in Diabetic Patients. *Iran Red Crescent Med J* 14(12):829–32
- L'Heveder R, Nolan T (2013) *International Diabetes Federation*. 101(3): 349–51
- Li J, Chen J, Kirsner R (2007) Pathophysiology of acute wound healing. *Clin Dermatol* 25:9–18
- Lee CM, Colagiuri R, Magliano DJ et al (2013) The cost of diabetes in adults in Australia. *Diabetes Res Clin Pract* 99(3):385–90
- Loo WT, Sasano H, Chow LW (2007) Pro-inflammatory cytokine, matrix metalloproteinases and TIMP-1 are involved in wound healing after mastectomy in invasive breast cancer patients. *Biomed Pharmacother* 61(9):548–52
- Rankin D, Cooke DD, Elliott J et al (2012) Supporting self-management after attending a structured education programme: a qualitative longitudinal investigation of type 1 diabetes patients' experiences and views. *BMC Public Health* 12:652
- Rosenthal S (2013) A workshop on prevention and management of diabetic foot complications. *Wound Care Canada* 11(2):26–31
- Rygg LØ, Rise MB, Grønning K, Steinsbekk A (2012) Efficacy of ongoing group based diabetes self-management education for patients with type 2 diabetes mellitus. A randomised controlled trial. *Patient Educ Couns* 86(1):98–105
- Schrementi ME, Ferreira AM, Zender C, DiPietro LA (2008) Site-specific production of TGF-beta in oral mucosal and cutaneous wounds. *Wound Repair Regen* 16(1):80–6
- Shupp J, Moffatt, L (2014) From mental health to the host microbiome: Looking at the forest not just the trees in wound healing *The FASEB Journal*, 28(1): Suppl. 337.1.
- Snow R, Sandall J, Humphrey C (2014) Use of clinical targets in diabetes patient education: qualitative analysis of the expectations and impact of a structured self-management programme in Type 1 diabetes. *Diabet Med* 31(6):733–8
- Walburn J, Vedhara K, Hankins M, Rixon L, Weinman J (2009) Psychological stress and wound healing in humans: a systematic review and meta-analysis. *J Psychosom Res* 67(3):253–71
- World Health Organization (2013) *Noncommunicable Diseases and Mental Health. Global Action Plan for the Prevention and Control of NCDs 2013–2020*. Available at: http://www.who.int/nmh/events/ncd_action_plan/en/ (accessed 10.06.2015)
- World Health Organization (2014) *Diabetes Programme*. Available at: <http://www.who.int/diabetes/en/> (accessed 10.06.2015)