



University of HUDDERSFIELD

University of Huddersfield Repository

Kipling, Kevin, McCluskey, Serena, Boduszek, Daniel, Kirshbaum, Marilyn and Garbutt, Gerard

Supervised exercise for older women treated for breast cancer: results from a pilot randomised controlled trial

Original Citation

Kipling, Kevin, McCluskey, Serena, Boduszek, Daniel, Kirshbaum, Marilyn and Garbutt, Gerard (2015) Supervised exercise for older women treated for breast cancer: results from a pilot randomised controlled trial. *Journal of Sports Sciences*, 33 (1). s85-s93. ISSN 0264-0414

This version is available at <http://eprints.hud.ac.uk/26451/>

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

<http://eprints.hud.ac.uk/>


Supervised exercise for older women treated for breast cancer.

Results from a pilot randomised controlled trial.

Kipling, K.N.^{1*}, McCluskey, S.¹, Boduszek, D.¹ Kirshbaum, M.N.² and Garbutt, G.^{1,3} (2015).

¹University of Huddersfield; ²Charles Darwin University, Australia; ³Pennine Acute Hospitals Trust.

*Corresponding author: kevin.kipling@hud.ac.uk



Introduction

Breast cancer is now the most common cancer in the UK. Almost two-thirds of women diagnosed earlier this decade with breast cancer are now likely to survive their disease for at least twenty years.

Exercise and breast cancer research

There is compelling evidence of the benefits of exercise in younger cancer survivors and older populations, however, evidence from older breast cancer survivors, is limited, despite the higher incidence of diagnosis and lower survival rates in this population.

Aims

To investigate whether a 12-week supervised exercise intervention with older women (>60 years) during adjuvant therapy for breast cancer, improved function (12-min walk), body composition (air displacement plesythmography), quality of life (European Organisation for the Research and Treatment of Cancer) and physical activity (PA) levels (Scottish PA Questionnaire) and if these could be sustained over a 12-month period.



Methods

A pilot randomised controlled trial

35 female breast cancer patients were recruited (mean = 67 years; SD ± 5.02) Supervised exercise intervention (n=16) or a control group (n=19). The exercise programme consisted of both aerobic and resistance exercises at RPE 3-4. Outcomes measures were assessed at baseline, 3, 6 and 12 months. Statistical analyses were conducted using descriptive statistics, mixed between-within ANOVA and repeated measures ANOVA.




Results


Attrition rates to the study were good (12.5%-intervention, 26%-control, 20%-overall) with no adverse events reported. Adherence to the supervised exercise sessions was high (>85%). Although no statistically significant interaction terms were detected between groups for any outcome measures at all four time points, both intervention and control groups significantly increased walking distance (P < 0.01, ES = .78) and physical activity levels (P < 0.05, ES = .30) over 12 months. Positive trends in favour of the intervention group was observed for body composition during the intervention period.




Comparison of body fat between groups over 12 months




Distance walked in 12-mins between groups over 12-months



Self reported PA levels between groups over 12-months



Comparison of fat free mass between groups over 12 months



Summary and Conclusions

Recruitment onto a supervised exercise intervention with older breast cancer survivors (BCS) was feasible with high adherence rates without adverse events. Future studies should incorporate larger sample sizes and consider longer interventions to evaluate sustained positive health benefits and behavioural change. This is important with BCS now living much longer to further examine the effects of physical activity in this under researched population.