



TICK POPULATION IN GOATS AND SHEEP IN ŠABAC

Ivan Pavlović¹, Snežana Ivanović¹, Aleksandar Dimitrić², Mensur Vegara³,
Ana Vasić⁴, Slavica Živković⁵, Bojana Mijatović⁵

¹Scientific Veterinary Institute of Serbia, V. Toze 14, 11000 Belgrade, Serbia

²Agrimatco DOO, Narodnog Fronta 73/I, 21000 Novi Sad, Serbia

³Department of International Environment and Development Studies Norwegian University of Life Science (NMBU) Campus As, Postboks 5003, No-1432 Aas, Norway

⁴Faculty of Veterinary Medicine, Belgrade University, Bul. Oslobođenja 18, 11000 Belgrade, Serbia

⁵Agricultural School PKB, Pančevački Put 39, 11000 Krnjača-Belgrade, Serbia

Received 14 June 2015; Received in revised form 4 January 2016; Accepted 25 January 2016

ABSTRACT

During our examination performed in the period from 2010 to 2012, we collected ticks from 52 flocks of sheep and 38 goat flocks. Ticks infestation occurred in 15.97% (214/1340) of sheep and 16.93% (107/632) of goats. The result showed the presence of *Ixodes ricinus*, *Rhipicephalus sanguineus*, *R. bursa*, *Dermacentor marginatus*, *D. pictus*, *Haemaphysalis punctata* and *Ha. inermis*. Additional to determination of tick species during the research, the sex ratio and the monthly influence of microclimate conditions (temperature, relative humidity and precipitation quantity) on the dynamics of populations of ticks were followed. Obtained results indicate the importance of the impact of climatic factors on the population dynamics of some species of ticks as well as the dynamics and abundance of different sexes within established species of ticks.

Key words: ecological factors, goat, sheep, ticks

INTRODUCTION

Ticks are obligate haematophagous ectoparasites which have multiple adverse effects on the host organism. A particular problem is that they spread diseases to humans, domestic and wild animals, which can be reservoirs, vectors and/or transient hosts for the tick-borne pathogens (5, 12, 14, 20, 23, 35, 36, 37, 41).

Most of the tick species of the family *Ixodidae* are egsophilic and during the search for the host they inhabit open habitats. During this development, ticks go through four life stages. These stages are egg, larvae (or seed tick), nymph, and adult (1, 5). *Ixodidae* ticks mainly belong to three host life cycle

group of ticks, while during each developmental stage it feeds once on a separate host and will remain attached to the host for several days (8, 43).

An important feature of the life cycle of ticks is the diapause (10, 15). Throughout diapause, a tick goes into hibernation, during which the metabolism slows (49, 50). Diapause allows ticks to survive adverse conditions, such as extreme temperatures, drought and lack of food (19).

We have taken into account the epidemiological significance of ticks, because they carry many disease pathogens that belong to the group of natural focal infections, characterized by endemic and seasonal occurrence (27, 41). That was the starting point of our investigation of domestic animals tick fauna in certain areas of Serbia. In our paper we present the tick fauna of sheep and goats from the area of Šabac (Serbia) in the 2010-2012 period.

MATERIAL AND METHODS

Sheep and goat breeds play important role in the Šabac area. Usually, they are kept in small herds of 20 to 25 animals. From early spring to late autumn,

Corresponding author: Dr. Ivan Pavlović, PhD

E-mail address: dripavlovic58@gmail.com

Present address: Scientific Veterinary Institute of Serbia, V. Toze 14, 11000 Belgrade, Serbia

Copyright: © 2016 Pavlović I. This is an open-access article published under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Competing Interests: The authors have declared that no competing interests exist.

Available Online First: 5 February 2016

Published on: 15 March 2016

<http://dx.doi.org/10.1515/macvetrev-2016-0077>

they are kept on pastures and graze on any land that is not being cultivated. It is common that flocks of goats and sheep are found on the same pastures. During the winter, goats and sheep are kept in stables. During the course of our study, we collected ticks from 52 flocks of sheep and 38 goat flocks. Ticks were collected monthly during the period from the exit to the pasture to the retreat to the barn (March to October). All specimens were placed into glass specimen bottles which had a piece of hard paper inserted bearing the name of locality, name of host and date and hour of collection. The tick species and sex/gender were identified by morphometric characteristics (24, 25, 47).

The data obtained were analyzed using Chi-square test (χ^2) to determine if the tick species and the prevalence of infection with ticks depended on the host animal species. For all analyses, the confidence level was kept at 95%.

Our studies were carried out in the period from March 2010 to November 2012 in the municipality of Šabac, which is located at 44° 46' northern latitude and 19° 41' east longitude and is in the western part of Serbia. Morphologically, the municipality of Šabac has three natural zones. North of the town there is a vast plain area known as Mačva where lowland humus is the dominant soil type. The second morphological unit constitutes the western part of the territory of the municipality of Šabac, which is characterized by a hilly relief - the Pocerina area, where the plain area Mačva gradually turns into a hilly area down to the Cer Mountain, where the relief and forest caused degradation and evolution of lowland soils into brown forest soil. The third morphological unit covers the southeastern part and

pressure occurs in the coldest month of January, while the lowest in April. In this area winds blow from all quadrants. On windless periods as waste 1/3 the frequency of occurrence of winds. The mean air temperature in Šabac is around 11,7°C. Humidity is on average 78-92% in winter and 51-63% in summer. The annual precipitation is about 435 l/m². Insolation increases from January to July and then decreases until December (official data by the Hydrometeorological Service of Serbia).

The climatic vegetation zones in this area correspond to the vegetation that is usually found on the territory of Serbia, such as forests of Hungarian oak (*Quercus confestim*), oak oak (*Quercus*) and beech (*Fagetum montanuum*), while in the underwater terrain we commonly encounter poplar (*Populus alba*) and white willow (*Salicium alba*). With regards to bushes and low trees the most common are: mandrel (*Prunus spinosa*), hawthorn (*Scataegus monogyna*), wild service tree (*Sorbus aria*) and maple (*Acer compestere*). From the terrestrial plants, most common are: meadow fescue (*Festuca pratensis*), meadow grass (*Poa pratensis*), couch grass (*Agropyrum repens*), couch grass without awns (*Bromus inermis*), sheep fescue (*Festuca ovina*), French ryegrass (*Arrhenatherum elatius*), red clover (*Trifolium pratense*), common vetch (*Vicia sativa*) and meadow pea (*Lathyrus sativa*) (21, 22).

RESULTS

During our examination we examined 1.340 sheep and 632 goats (Table 1).

Table 1. Number of examined sheep and goats in the period 2010-2012

host	Examined animals / years			Total
	2010.	2011.	2012.	
Sheep	502	486	352	1.340
Goats	215	246	171	632

it represents Posavina - the river valley along the right bank of the Sava River water intersected by former backwaters. There the most common type of soil is sandy black soil, which appears as a young Pedogenic creation and is most frequently found in the bottomlands of major rivers (2).

The municipality of Šabac has a continental climate. On the north side, Šabac is widely open to the Srem and therefore in terms of climate is strongly influenced by the Panonian continental climate. This climate is characterized by cold winters and warm, dry summers. The highest air

Tick infestation has occurred in 15.97% (214/1340) of sheep and 16.93% (107/632) of goats. The results showed presence of *Ixodes ricinus*, *Rhipicephalus sanguineus*, *R. bursa*, *Dermacentor marginatus*, *D. pictus*, *Haemaphysalis punctata* and *Ha. inermis*. Their prevalence is presented in Table 2.

Relative abundance analysis revealed that the *I. ricinus* was absolutely dominant species in sheep, followed by *Rhipicephalus bursa*, *R. sanguineus*, *Haemaphysalis punctata*, *Dermacentor marginatus* and *D. pictus* and *Ha. inermis*. Similar fauna of ticks

Table 2. Percent of sheep and goats infected with ticks in the period 2010-2012

Tick species	Examined animals					
	sheep's			goats		
	total	infected	%	total	infected	%
<i>Ixodes ricinus</i>	1340	214	15.97	632	107	16.93
<i>Rhipicephalus sanguineus</i>	1340	112	8.35	632	34	0.53
<i>Rhipicephalus bursa</i>	1340	209	15.59	632	101	15.98
<i>Dermacentor marginatus</i>	1340	98	7.31	632	19	3.00
<i>Dermacentor pictus</i>	1340	87	6.49	632	7	1.10
<i>Haemaphysalis punctata</i>	1340	101	7.53	632	21	3.32
<i>Haemaphysalis inermis</i>	1340	4	0,29	632	2	0.31

was found in goats, since the herds of goats graze at the same pasture areas with herds of sheep.

The chi-square test showed there was a difference present between sheep and goats with regards to the prevalence and intensity of infection with *R. sanguineus*, *Ha. punctata*, *D. marginatus* and *D. pictus* (Fig. 1).

Ixodes ricinus reached maximum abundance in May, in which we also found the maximum occurrence of the species *Dermacentor pictus*. In June, the population peak is observed for the species *Rhipicephalus sanguineus* and *Rhipicephalus bursa*, which are the most common types both in July and August. In September, we saw an increase in the




Figure 1. Percent of sheep and goats infected with ticks in the period 2010-2012

Population dynamics of the established species of ticks can be observed every year from March to October. No sampling took place from November to February, therefore their presence in nature cannot be confirmed. During the three years of examination, in the month of March the following types of ticks were found: *Ixodes ricinus*, *Rhipicephalus sanguineus*, *Dermacentor marginatus* and *Haemaphysalis punctata*. In April, we observed the occurrence of the following species: *Dermacentor pictus*, *Rhipicephalus bursa* and *Haemaphysalis inermis*. In April, species that reached maximum numbers were *Dermacentor marginatus*, *Haemaphysalis punctata* and *Haemaphysalis inermis*. Species

population of two species of ticks: *Ixodes ricinus* and *Dermacentor marginatus*, while in October we observed the emergence of the following species: *Haemaphysalis punctata* and *Ha. Inermis*, and *Rhipicephalus sanguineus* and *R. bursa*.

According to the Chi-square test for averages of the three year sampling periods, there is no significant difference between population dynamics of tick species infection with sheep and goats.

Overall, the female-male ratio during the course of our study was 61.02% : 38.98% in favor of females, with a higher percentage of females established in all three years of research. The ratio of males and females of the same species is




Figure 2. Population dynamics of the established species of ticks by months in the period 2010-2012

also interesting. Of the seven species established, a greater number of males than females (65.08% : 34.92%) occurred only in the species *Rhipicephalus bursa*, while for the other six species we established a larger number of females. For the two most commonly found species *Rhipicephalus sanguineus* and *Ixodes ricinus*, this ratio was 69.50%: 30.50% and 63.42%: 36.58%, respectively in favor of females.

DISCUSSION

Comparison of the obtained results with findings in other regions of Serbia indicated that there is a great similarity in the established tick species. Examination performed in the Belgrade area established *I. ricinus*, *R. sanguineus*, *D. pictus* and *D. marginatus* as the most abundant species (12, 31, 32, 40, 44).

Results obtained during the examination of the tick fauna of sheep and goats in the northeastern, eastern and western part of Serbia showed that *Ixodes ricinus* and *Dermacentor marginatus* are the dominant tick species in those areas (31, 34, 35, 44, 46). Examination performed in sheep in the Prizren district (Kosovo) during 1991 (39) pointed to the presence of the same tick species, including *Hy. savignyi*, *Ha. inermis*, *Boophilus calcaratus* and *Ornithonius lachorenis*. The latest examination conducted during 2013 in the Kumanovo area

(Macedonia) established that most abundant in sheep were *Ixodes ricinus*, followed by *Dermacentor marginatus*, *Rhipicephalus sanguineus*, *R. bursa*, *Haemaphysalis punctata* and *D. pictus* (45). During all the examinations, *Ixodes ricinus* was the most abundant tick species in sheep and goats. The found species of ticks are most common in sheep and goats in the regions of the Western Balkans, Mediterranean and Central Europe (14, 16, 25, 27, 28, 37, 38, 48).

Climate conditions have a great influence on the population dynamics of ticks. Population dynamics of ticks is related to the impact of climate factors like air temperature, relative humidity and rainfall (32, 49).

Our results confirmed the results of the studies carried out in northeast, eastern and south-eastern Serbia (29, 33, 34) which established that *I. ricinus* was by far the most abundant species with the largest number of specimens collected in the spring at a temperature of about 15°C, relative humidity of 76.00% and rainfall of 81.11 l/m². March marks the start of, the increase in the number of *Haemaphysalis (punctata and inermis)* species, while similar results were obtained during examination in France (26), Romania (28) and in Macedonia (45). April is the month of greatest abundance of *D. marginatus*, *Haemaphysalis punctata* and *Ha. inermis* which reached their peak at a temperature of 9.01°C, relative humidity of 75.66% and rainfall of 35.80 and 36.06 mm/m².

This data is in correlation with the results of other examinations of the seasonal dynamics of ticks in Europe (3, 6, 9, 11, 17, 29, 37). Similar values were obtained during research in Serbia (12, 41), France (26), Italy (27, 38) and in a Berlin forest (11). The authors point out that the low temperature, high humidity and rainfall significantly affect the life cycle of ticks, particularly the *I. ricinus* species (20, 28, 46). Same results were obtained worldwide e.g. in coastal areas of New York (7) and in South Africa (18, 20). Both studied the temperature below which the activity of ticks is completely stopped and the temperature at which it expressed its full activity and found a significant relationship of tick activities and the degree of reduction of temperature for adult forms. This agrees with the data published about ticks in various part of Russia, Central Europe and the West Balkans (4, 12, 13, 14, 20, 21, 25, 26, 29, 31).

The female abundance of established tick species has been in correlation with previously established population dynamics. The females of *Ixodes ricinus* species were present from March to October, with a peak population in May and June. Females of two species of the genus *Rhipicephalus* (*sanguineus* and *bursa*) have been found most often in the summer months - June and July. Findings of the females of species *Dermacentor marginatus* and *Haemaphysalis punctata* were most common in April and May, while sporadic finding of females of *Dermacentor pictus* and *Haemaphysalis inermis* species was attached to the spring months. This population dynamics of female ticks is characteristic for this microclimate (3, 8, 9, 15).

Males of the species *Ixodes ricinus* were found from March to October, with the spring peak population in May and autumnal in September which corresponds to the values obtained in our earlier research (12, 17). Males of the species *Rhipicephalus sanguineus* were established from March to October, a species *Rhipicephalus bursa* from April to September with a population peak of both species in June, which also corresponds to values for this geographical area (28, 30, 32, 38). Males of the two species of the genus *Dermacentor* (*marginatus* and *pictus*) were usually found from April to June, while the males of *Haemaphysalis punctata* species were established from April to June. A small number of males of *Haemaphysalis inermis* was found only in April which corresponds to values of research in this area and in Central Europe and the Mediterranean basin (16, 20, 35, 42, 47).

CONCLUSION

Being the vectors and reservoirs for many endemic tickborne pathogens, the tick transmit diseases that cause health disturbance in domestic animals and humans in affected areas.

The occurrence of tick born diseases in this species is usual during the exposure time period. The population dynamics and the climatic factors that influence the tick population need to be studied in order to predict the critical points and implement adequate protection measures in animals with the final goal of disease prevention and control.

ACKNOWLEDGEMENT

Paper is accomplished as a part of scientific and technological project BT 31053 of the Ministry of Education, Science and Technical Development of Republic of Serbia

REFERENCES

- Anderson, J.F., Magnarelli, L.A. (2008). Biology of ticks. *Infect Dis Clin North Am.* 22 (2): 195-215. <http://dx.doi.org/10.1016/j.idc.2007.12.006> PMID:18452797
- Anonimus (2015). Geografski Atlas. Republički geodetski zavod. Beograd.
- Babenko, L.V. (1974). Sutočnie kolebanija aktivnosti golodnih nimf *Ixodes ricinus* L. i *Ixodes persulcatus* P. SCH (Parazitiformes: Ixodidae). *Med. Parazitol. Parazit. Bolezn.* 42, 520-527.
- Babenko, L.V., Arumova, E.A., Bush, M.A., Skadinsh, E.A. (1977). O sootonošenii polov v prirodnihih populaciah imago *Ixodes ricinus* L. i *Ixodes perculacatus* P. SCH (Ixodidea, Ixodidae). *Med. Parazitol. Parazit. Bolezn.* 46, 294-301.
- Belozerow, V.N. (1982). Diapausa and biological rhythms in ticks. In: F.D.Obenchain and R.Galun (Ed.), *Physiology of ticks*, 1st edition (pp.469-500). Oxford: Pergamon Press. <http://dx.doi.org/10.1016/B978-0-08-024937-7.50018-4>
- Carrol, J.F., Kramer, M. (2003). Winter activity of *Ixodes scapularis* (Acari: Ixodidae) and the operation of deer-targeted tick control devices in Mryland. *J.Med.Entomol.* 40, 238-244. <http://dx.doi.org/10.1603/0022-2585-40.2.238>
- Clark, D.D. (1995). Lower temperature limits for activity of several Ixodid ticks (Acari: Ixodidae): effects of body size and rate of temperature change. *J.Med.Entomol.* 32, 449 - 452. <http://dx.doi.org/10.1093/jmedent/32.4.449> PMID:7650705

8. Černy, V., Daniel, M., Rosicky, B. (1974). Some features of developmental cycle of the tick *Ixodes ricinus* L. (Acarina, Ixodidae). *Folia Parasitol.* 21, 85-87.
9. Černy, V., Szymanski, S., Dusbabek, F., Daniel, M., Hozakova, E. (1982). Survival of unfed *Dermacentor ricitulatus* adults under natural conditions. *Wiadom. Parasitol.* 28, 27-31.
10. Daniel, M. (1978). Microclimate as a determining element in the distribution of ticks and their developmental cycles. *Folia Parasitol.* 25, 91-94.
11. Dautel, H., Dippel, C., Kammer, D., Werkhausen, A., Kahl, O. (2008). Winter activity of *Ixodes ricinus* in a Berlin forest. *Int. Med. Microbiol.* 298, 50-54. <http://dx.doi.org/10.1016/j.ijmm.2008.01.010>
12. Dimitrić, A. (1999). Fauna i ekologija krpelja (Acari:Ixodidae) kao prenosioca metazoonoza. MSc thesis, Faculty of Veterinary Medicine University in Belgrade, Serbia
13. Duffy, D.C., Campbell, S.R. (1994). Ambient air temperature as a predictor of activity of adult *Ixodes scapularis* (Acari: Ixodidae). *J. Med. Entomol.* 31, 178-180. <http://dx.doi.org/10.1093/jmedent/31.1.178> PMID:8158624
14. Dumitrache, M.O., Gherman, C.M., Cozma, V., Mircean, V., Györke, A., Sándor, A.D., Mihalca, A.D. (2012). Hard ticks (Ixodidae) in Romania: surveillance, host associations, and possible risks for tick-borne diseases. *Parasitol.Res.* 110, 2067-2070. <http://dx.doi.org/10.1007/s00436-011-2703-y> PMID:22033737
15. Dyk, V., Boučkova, L. (1968). Die Temperature-Feuchte-Relation in der Aktivitat des gemeinen Holzbocks. *Angew Parasitol.* 9, 83-87.
16. Estrada-Pe-a, A., Martinez Avilez, M., Mu-oz Reoyo, M.J. (2011). A population model to describe the distribution and sesonal dynamics of the ticks *Hyalomma marginatus* in the Mediterranean basin. *Transbound. Emerg. Dis.* 58, 213-223. <http://dx.doi.org/10.1111/j.1865-1682.2010.01198.x> PMID:21223534
17. Fourie, L.J., Horak, I.G., Mrais, L. (1988). The seasonal abundance of adult Ixodid ticks on Merino sheep in the South-Western Orange Free State. *J.S.A.V.A.* 59,191-194.
18. Fourie, L.J., Horak, I.G. (1991). The seasonal activity of adult Ixodid ticks on Angora goats in the South-Western Orange Free State. *J.S.A.V.A.* 62, 104-106.
19. Harlan, H.J., Foster, W.A. (1990). Micrometeorologic factors affecting field host-seeking activity of adult *Dermacentor variabilis* (Acari : Ixodidae). *J. Med. Entomol.* 27, 471-479. <http://dx.doi.org/10.1093/jmedent/27.4.471> PMID:2388223
20. Hornok, S. (2009). Allochronic seasonal peak activites of *Dermacentor* ond *Haemaphysalis* spp. Under continental climate in Hungary. *Vet. Parasitol.* 163, 366-369. <http://dx.doi.org/10.1016/j.vetpar.2009.03.048> PMID:19410373
21. Janković, M. (1973). Biljni svet prirodnih ekosistema SR Srbije. SANU, referati naučnog skupa »Čovek i životna sredina«, Beograd. PMID:PMC1423154
22. Janković, M., Pantić, N., Mišić, V., Diklić, N., Gajić, M. (1984). Vegetacija SR Srbije. SANU, Beograd. PMID:PMC1009372
23. Jongejan, F., Uilenberg, G. (2004). The global importance of ticks. *Parasitol.* 129, Suppl. S3-14.
24. Kapustin, F.U. (1955). Atlas parazitov krovi životnih i klešćei iksodid. Gasudarstvenoe izdetejlstvo seljskohazajstvenoi literaturi. Moskva PMID:13313366
25. Kolonin, G.V. (2009). Fauna of ixodid ticks of the world (Acari:Ixodidae), Moscow. <http://www.kolonin.org/3.html>.
26. L'Hostis, M., Dumon, H., Dorchies, B., Biosdron, F., Gorenflot, A. (1995). Seasonal incidence and ecology of the ticks *Ixodes ricinus* (Acari: Ixodidae) on grazing pastures in Western France. *Exp. Appl. Acarol.* 19, 211-220. <http://dx.doi.org/10.1007/BF00130824> PMID:7641568
27. Maroli, M., Khouzy, C., Frusteri, L., Manila, G. (1996). Distribution of dogs ticks (*Rhipicephalus sanguineus* Latreille, 1806) in Italy: a public health problem. *Ann. Inst. Super. Sanita.* 23, 387-397.
28. Mihalca, A.D., Dumitrache, M.O., Magdaş, C., Gherman, C.M., Domşa, C., Mircean, V., Ghira I. V., Pocora, V., Ionescu, D.T., Sikó Barabási, S., Cozma, V., Sándor, A.D. (2012). Synopsis of the hard ticks (Acari: Ixodidae) of Romania with update on host associations and geographical distribution. *Exp. Appl. Acarol.* 58, 183-206. <http://dx.doi.org/10.1007/s10493-012-9566-5> PMID:22544174
29. Milutinović, M. (1992). Ekološka istraživanja krpelja (Acarina, Ixodidae, Ixodia) Srbije, PhD disertation, Biology faculty University in Belgrade, Serbia
30. Milutinović, M., Petrović, Z., Bobić, B., Pavlović, I. (1996). Ecological notes on ticks collected in West Serbia Yugoslavia. *Parasitol.Hung* 29/30: 67-74.
31. Milutinović, M., Aleksić-Bakrač, N., Pavlović, I. (1997). Ticks (Acari: Ixodidae, Argasidae) in the Belgrade area. *Acta. Entom. Serb.* 2, 77-85.
32. Milutinović, M., Aleksić-Bakrač, N., Pavlović, I. (1998). Faunistic and ecological notes on ticks (Acari: Ixodidae, Argasidae) in the extended area of Belgrade. *Mag. Allator. Lapja* 120, 434-436.

33. Milutinović, M., Aleksić-Bakrač, N., Pavlović, I. (1998). Reserch of tick population (Acari: Ixodidae) in Eastern part of Serbia. *Ars Vet.* 14, 227-234.
34. Milutinović, M., Petrović, Z., Mišćević, Z. (1987). Fauna i ekologija krpelja (Acarina, Ixodoidea, Ixodidae) severoistočnog dela SR Srbije. *Zbornik V jugoslovenski kongres infektologov*, (pp.140-145), Ljubljana, Yugoslavia.
35. Mišćević, Z., Milutinović, M., Biševac, Lj. (1990). Tick fauna (Acarina, Ixodoidea, Ixodidae) of northeast Serbia with special emphasis on the species *Ixodes ricinus*. *Acta Vet.* 40, 143-150.
36. Nieder, M., Bojkovski, J., Pavlović, I., Savić, B., Elezović, M., Silaghi, C. (2013). Studies on the occurrence of granulocytic anaplasmosis in cattle and on biodiversity of vectors (ixodid ticks) in Serbia. *Zbornik kratkih sadržaja 18.godišnjeg savetovanja doktora veterinarske medicine Republike Srpske sa međunarodnim učešćem*, pp. 25, Teslić, Republika Srpska, BiH
37. Papazahariadou, M.G., Saridomichelakis, M.N., Koutinas, A.F., Papadopoulos, E.G., Leontides, L. (2003). Tick infestation of dogs in Thessaloniki, northern Greece. *Med. Vet. Entomol.* 17, 110-113. <http://dx.doi.org/10.1046/j.1365-2915.2003.00404.x> PMID:12680933
38. Pavlović, I., Kulišić, Z., Nešić, D., Romanić, S. (1995). Ectoparasites of sheep and goats in Prizren district. *Proceeding of 3rd Internat Conference Sheep and Goat Production*, (pp.101-105), Ohrid, Macedonia
39. Pavlović, I., Milutinović, M., Kulišić, Z., Dimitrić, A., Pavlović, V. (1999). Prisustvo artropoda od biomedicinskog značaja na zelenim površinama grada Beograda. *Zbornik radova II Gradske konferencije o suzbijanju štetnih artropoda i glodara sa međunarodnim značajem*, (pp. 81-87), Belgrade, SRYugoslavia
40. Pavlović, I., Milutinović, M., Petković, D., Terzin, V., Terzin, D. (2002). Epizootiological research of canine babesiosis in the Belgrade district. *J. Protozool. Res.* 12 (1-2): 10-15.
41. Pavlović, I., Savić, B., Ivetić, V., Radanović, O., Žutić, M., Jakić-Dimić, D., Bojkovski, J. (2009). The effect of parasitic infections to production results of sheep. *IV Balkan Conf Animal Sci BALNIMALCON 2009, Proceeding of Challenges of the Balkan Animal industry and the Role of science and Cooperation*, (pp. 389-391), Stara Zagora, Bulgaria
42. Pavlović, I., Petković, D., Kukovska, V., Stamenković, V., Jovčevski, S., Pavlović, M., Jovčevski, St., Elezović, M. (2012). Most important food-borne disease of dogs caused by ticks and its control. *Proceeding of 3rd International Scientific Meeting Days of Veterinary Medicine*, (pp. 34-37), Ohrid, Macedonia
43. Pavlovic, I., Ivanovic, S., Zujovic, M. (2013). Tick fauna of goat and sheep in Belgrade area. *Sci Works Series C Vet Med LIX* (1), 51-53.
44. Pavlović, I., Jovčevski, S., Jovčevski, St., Kukovska, V., Dimitrić, A. (2014). Tick fauna of sheep and cattle at Kumanovo arae (Macedonia). *Lucr.Ştii. Med.Vet. XLVII* (3), 88-95.
45. Petrović, Z., Milutinović, M., Pavlović, I. (1996). Istraživanja krpelja (Acari: Ixodidae, Argasidae) u Jugoslaviji. In: Z.Petrović (Ed.) *Akademik Čedomir P. Simić, Naučni skup posvećen 100-godišnjici rođenja* (pp. 96-101). Beograd: SANU Odeljenje mecidinskih nauka i Jugoslovensko društvo parazitologa
46. Pomerancev, B.L. (1950). Fauna SSSR. Paukoobraznie. Iksodidovie kleščei (Ixodidae). *Akadema Nauk SSSR, Moskva-Leningrad*
47. Rivosecchi, L., Khoury, C., Lezzerini, C., Dell Uomo, G. (1980). Osservazioni su *Rhipicephalus sanguineus* (Ixodidae) nella periferia di Roma. *Riv. Parasitol.* 41, 273-276.
48. Tovornik, D. (1976). Seasonal and diurnal periodicity of the tick *Ixodes ricinus* L. in the Pannonian tick-borne encephalitis focus (Stara Ves). *JAZU, Razred za medicinske znanosti* 99-103.
49. Zahler, M., Gothe, R. (1995). Effect of temperature and humidity on egg hatch, moulting and longevity of larvae and nymphs of *Dermacentor reticulatus* (Ixodidae). *Appl. Parasitol.* 36, 53 -65. PMID:7780450