

# A THEORY OF LENGTH FOR NOETHERIAN MODULES

Tor H. Gulliksen

## Introduction.

In this paper we shall introduce a theory of length for Noetherian modules over an arbitrary ring (with identity), assigning to each Noetherian module  $M$  an ordinal number  $l(M)$  which will briefly be called the length of  $M$ , see § 2 for definition.  $l(M)$  is finite if and only if  $M$  has a finite composition-series, in which case  $l(M)$  equals the length of the composition-series. Thus we are working with a generalization of the classical theory of length.

$l(M)$  carries important information about  $M$ . Being an ordinal,  $l(M)$  can be expressed as a polynomial in  $\omega$  with integral coefficients and ordinal exponents,  $\omega$  denoting the first non-finite ordinal. This polynomial - the Cantor normal form of  $l(M)$  - has properties similar to the properties of the Hilbert-Samuel polynomials in local algebra. First of all, its degree coincides with the Krull dimension of  $M$  (2.3), the Krull dimension being interpreted as an ordinal as in Krause [5]. Moreover, if  $\alpha$  is an ordinal, then the coefficient of the term of degree  $\alpha$  is additive on the category of Noetherian modules of Krull dimension not greater than  $\alpha$  (2.7).

In § 1 we fix the notation concerning ordinal numbers and the Krull ordinal of a partially ordered Noetherian set.

§ 2 contains general results concerning the length function  $M \mapsto l(M)$ . Although  $l$  is not additive in general, 2.1 gives the following satisfactory substitute for additivity: if

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

is an exact sequence of Noetherian modules, then we have

$$l(M'') + l(M') \leq l(M) \leq l(M') \oplus l(M'')$$

Moreover we have (2.11):

$$l(M' \oplus M'') = l(M') \oplus l(M'').$$

Here  $\oplus$  is used ambiguously to denote the Hessenberg natural sum of ordinals, cf. § 1, and the direct sum of modules.

In general there does not exist a good notion of composition series in terms of which  $l(M)$  can be defined. However, we show in 2.12 that if  $M$  has countable Krull dimension, then there exists a chain of non-zero submodules of  $M$  which is of ordinal type  $l(M)$ .

Unlike the case with factor modules of  $M$  (2.3), not every ordinal less than  $l(M)$  is the length of a submodule of  $M$ . In fact if  $N$  is a submodule of  $M$  then each of the coefficients in the polynomial  $l(N)$  is less than or equal to the corresponding coefficient in the polynomial  $l(M)$ . In particular,  $l(N)$  can only take a finite number of values (2.9).

In § 3 we obtain more precise results by assuming that all modules be finitely generated over a commutative Noetherian ring. In this case we can give an interpretation of the set of exponents in the polynomial  $l(M)$ , in terms of  $\text{Ass } M$  (3.2). We also give a complete description of the possible lengths of the submodules of  $M$ .

In Bass [1]  $o(M)$  denotes the supremum of the ordinal types of descending chains of non-zero submodules of  $M$ . In 3.4 we show that also  $o(M)$  can be expressed in terms of  $l(M)$ . We have the relation

$$o(M) = \min(\omega_1, l(M))$$

$\omega_1$  being the first non-countable ordinal.

§ 1 Notation and basic definitions.

If  $W$  is a set of ordinal numbers, we let  $\sup W$  denote the least ordinal which is greater than or equal to every element in  $W$ . In particular we put  $\sup \emptyset = 0$ . If  $\beta_1, \dots, \beta_k$  are ordinals, we let  $\sum_{i=1}^k \beta_i$  denote their sum in the following order

$$\beta_1 + \dots + \beta_k$$

Letting  $\omega$  denote the ordinal type of the natural numbers, any ordinal  $\alpha$  can be written

$$* \quad \alpha = \sum_{i=1}^k \omega^{\alpha_i} n_i$$

where  $n_1, \dots, n_k$  are non-negative integers and the exponents form a decreasing sequence of ordinals, i.e.

$$j < i \Rightarrow \alpha_i < \alpha_j \quad \text{for all } i, j$$

The representation (\*) will be called the Cantor normal form of  $\alpha$ . If  $n_1 \neq 0$  the corresponding exponent  $\alpha_1$  will be called the degree of  $\alpha$  and will be denoted by  $\deg \alpha$ . It is convenient to define  $\deg 0 = -1$ . The Cantor normal form is unique in the following sense: Let  $\alpha$  and  $\beta$  be ordinals with Cantor normal forms  $\sum_{i=1}^k \omega^{\alpha_i} n_i$  and  $\sum_{i=1}^k \omega^{\alpha_i} m_i$  respectively. Then we have  $\alpha = \beta$  if and only if  $n_i = m_i$  for all  $i$ . If  $n_i \leq m_i$  for all  $i$ , then this fact will be expressed by writing  $\alpha \ll \beta$ . Finally we define the direct sum (Hessenberg natural sum) of  $\alpha$  and  $\beta$  as follows

$$\alpha \oplus \beta := \sum_{i=1}^k \omega^{\alpha_i} (n_i + m_i).$$

A justification for this notation is contained in 2.11.

Let  $S$  be a non-empty partially ordered set which is Noetherian, i.e. every non-empty subset has a maximal element. Let Ord denote the class of ordinal numbers. By the ordinal map on  $S$  we mean the map

$$\lambda: S \rightarrow \text{Ord}$$

defined by

$$\lambda(x) = \sup\{\lambda(y) + 1 : x < y\}$$

The Krull ordinal of  $S$  will be denoted  $\kappa(S)$  as in [1].  $\kappa(S)$  can be expressed in terms of the ordinal map as follows

$$\kappa(S) = \sup\{\lambda(x) : x \in S\}.$$

## § 2 The length of Noetherian modules.

Let  $M$  be a Noetherian (left)module over a ring (with identity) and let  $S(M)$  be the set of all submodules of  $M$  ordered by inclusion. The Krull ordinal of  $S(M)$  will be called the length of  $M$  and will be denoted by  $l(M)$ . The degree of the ordinal  $l(M)$ , cf. § 1, will be called the dimension of  $M$  and will be denoted  $d(M)$ . By the Krull dimension of  $M$  we will mean the ordinal  $KdimM$  as defined in Krause [5] and equivalently in [2]. We shall see in theorem 2.3 below that  $d(M) = KdimM$ .

2.1 Theorem. Let  $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$  be an exact sequence of Noetherian modules. Then we have

$$l(M'') + l(M') \leq l(M) \leq l(M') \oplus l(M'')$$

In particular we have

$$d(M) = \max(d(M'), d(M'')).$$

Proof: The last equality clearly follows from the two inequalities. We will start by proving the first inequality. Let  $P$  be the partially ordered set obtained from  $S(M')$  and  $S(M'')$  by identifying the unique maximal element in  $S(M')$  with the unique minimal element in  $S(M'')$ . Let  $\lambda^+$  and  $\lambda'$  be the ordinal maps on  $P$  and  $S(M')$  respectively. It is easily shown by induction that

$$\lambda^+(N) = \kappa(S(M'')) + \lambda'(N) \quad \text{for all } N \in S(M')$$

Hence

$$\kappa(P) = \kappa(S(M'')) + \kappa(S(M')) = l(M'') + l(M').$$

Since we have an order preserving injection  $P \rightarrow S(M)$ , it is easily shown that  $\kappa(P) \leq \kappa(S(M))$ . Hence

$$l(M'') + l(M') \leq l(M).$$

We shall now prove the second inequality in 2.1. Let  $\lambda''$ ,  $\lambda$  and  $\lambda'$  denote the ordinal maps on  $S(M'')$ ,  $S(M)$  and  $S(M')$  respectively. We will define a map

$$\lambda^* : S(M) \rightarrow \underline{\text{Ord}}$$

as follows. Let  $N \in S(M)$ . Put

$$\lambda^*(N) = \lambda'(N \cap M') \oplus \lambda''(N + M'/M')$$

I claim that  $\lambda^*$  is strictly order reversing. Indeed, let  $N_1 \subseteq N_2$  be submodules of  $M$ . Clearly we have

$$\lambda^*(N_1) \geq \lambda^*(N_2)$$

Assume that we have equality. We are going to show that  $N_1 = N_2$ .

For  $i = 1, 2$  put

$$\alpha_1 = \lambda'(N_1 \cap M') \quad \text{and} \quad \beta_1 = \lambda''(N_1 + M'/M')$$

We have

$$\alpha_1 \geq \alpha_2, \quad \beta_1 \geq \beta_2 \quad \text{and} \quad \alpha_1 \oplus \beta_1 = \alpha_2 \oplus \beta_2$$

These three relations are easily seen to imply

$$\alpha_1 = \alpha_2 \quad \text{and} \quad \beta_1 = \beta_2.$$

Since  $\lambda'$  and  $\lambda''$  are strictly order reversing we have

$$N_1 \cap M' = N_2 \cap M' \quad \text{and} \quad N_1 + M'/M' = N_2 + M'/M'.$$

It follows that  $N_1 = N_2$ . Since  $\lambda^*$  is strictly orderreversing, it is easily shown by induction that

$$\lambda(N) \leq \lambda^*(N) \quad \text{for all } N \in S(M).$$

Hence

$$l(M) = \kappa(S(M)) = \lambda((o)) \leq \lambda^*((o)) = l(M') \oplus l(M'')$$

**2.2 Remark.** It is possible to generalize the notion of length to non-Noetherian modules  $M$ , by letting  $l(M)$  be the supremum of all ordinals  $\kappa(S)$  where  $S$  runs through the set of all Noetherian subsets of  $S(M)$ . With this generalized notion the previous theorem would still be valid, except for the first of the two inequalities which has to be replaced by the following weaker inequality

$$\max(l(M''), l(M')) \leq l(M).$$

**2.3 Theorem.** Let  $M$  be a non-zero Noetherian module. Then we have

- (i) Every ordinal less than  $l(M)$  is the length of a proper factor module of  $M$ . Conversely, if  $N$  is a non-zero submodule of  $M$  then  $l(M/N) < l(M)$ .
- (ii)  $d(M) = K \dim M$ .

Proof: (i) Let  $\beta$  be an ordinal less than  $l(M)$ , and let  $\lambda$  be the ordinal map on  $S(M)$ . Letting  $0_M$  denote the zero-submodule in  $M$  we have  $\lambda(0_M) = l(M) > \beta$ . Hence we can find a submodule  $N \subseteq M$  such that  $\lambda(N) = \beta$ , so  $l(M/N) = \beta$ . Conversely, if  $N$  is a non-zero submodule of  $M$ , then by 2.1  $l(M/N) < l(M)$ .

(ii) We will first show that  $Kdim M \leq d(M)$  using induction on  $d(M)$ . If  $l(M) \leq 0$  then  $M$  has finite length, so clearly  $Kdim M = d(M)$ . Let  $\alpha$  be a non-zero ordinal and assume that the inequality is valid whenever  $d(M) < \alpha$ . Now assume that  $d(M) = \alpha$ . Assume that  $Kdim M > \alpha$ . Then there exists a descending chain

$$M = M_0 \supset M_1 \supset \dots$$

such that  $Kdim M M_i/M_{i+1} \geq \alpha$  for  $i \geq 0$ . By the induction hypothesis we have  $d(M_i/M_{i+1}) \geq \alpha$ . Hence  $l(M_i/M_{i+1}) \geq \omega^\alpha$ . By 2.1 we have  $l(M) \geq \omega^\alpha \omega = \omega^{\alpha+1}$ . So  $d(M) \geq \alpha+1$  which is a contradiction. We conclude that  $Kdim M \leq \alpha$ .

We will now show that  $d(M) \leq Kdim M$  using induction on  $Kdim M$ . If  $Kdim M \leq 0$  then  $M$  has finite length, hence  $d(M) = Kdim M$ . Put  $Kdim M = \alpha > 0$ . Assume that  $d(M) \geq \alpha+1$ . Then  $l(M) \geq \omega^{\alpha+1}$ .

By (i) we can find a submodule  $M_1 \subset M$  such that  $l(M/M_1) = \omega^\alpha$ . By 2.1 it follows that

$$\omega^{\alpha+1} \leq l(M) \leq l(M_1) \oplus \omega^\alpha$$

Hence  $l(M_1) \geq \omega^{\alpha+1}$ . Now we can find a submodule  $M_2 \subset M_1$  such that  $l(M_1/M_2) = \omega^\alpha$ . Repeating the argument we can find a descending sequence  $M = M_0 \supset M_1 \supset M_2 \supset \dots$  such that  $d(M_i/M_{i+1}) = \omega^\alpha$  for  $i \geq 0$ . Hence  $d(M_i/M_{i+1}) = \alpha$ . We may assume by induction that

$\text{Kdim}(M_i/M_{i+1}) \geq \alpha$  . Hence  $\text{Kdim} M \geq \alpha+1$  , which is a contradiction. We conclude that  $d(M) \leq \alpha$  . ■

2.4 Corollary. To each ordinal  $\alpha$  there exists a Noetherian, commutative ring  $R$  such that  $l(R) = \alpha$  .

Proof There exists a commutative, Noetherian ring  $R_\alpha$  such that  $\text{Kdim} R_\alpha \geq \alpha$  , cf. [2] or [3]. Hence  $l(R_\alpha) \geq \omega^\alpha \geq \alpha$  . By 2.3(i) there exists an ideal  $\mathcal{O}$  in  $R_\alpha$  such that  $l(R_\alpha/\mathcal{O}) = \alpha$  . ■

In [2] a module  $M$  is called  $\alpha$ -critical if  $M$  has Krull-dimension equal to  $\alpha$  and every proper factor-module has Krull-dimension less than  $\alpha$  . The following corollary is an immediate consequence of 2.3:

2.5 Corollary. Let  $M$  be a Noetherian module. Then the following statements are equivalent:

- (i)  $M$  is  $\alpha$ -critical.
- (ii)  $l(M) = \omega^\alpha$  .

2.6 Definition Let  $M$  be a Noetherian module and let  $\alpha$  be any ordinal. The coefficient of the term of degree  $\alpha$  in the Cantor normal form of  $l(M)$  is a non-negative integer which will be denoted by  $\mu_\alpha(M)$  .

2.7 Lemma Let  $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$  be an exact sequence of Noetherian modules. Put  $\alpha = \text{Kdim} M$  . Then we have

$$\mu_\alpha(M) = \mu_\alpha(M') + \mu_\alpha(M'')$$

Proof By 2.3  $\alpha$  equals the degree of  $l(M)$  , hence the equality follows from 2.1. ■

2.8 Lemma and definition. Let  $M$  be a Noetherian module of dimension  $\alpha \neq 0$ . Then there exists a unique maximal submodule of  $M$  of dimension less than  $\alpha$ , which will be denoted by  $M_*$ . Put

$$l(M) = \omega^{\alpha}n + \beta$$

where  $n \neq 0$  and  $\beta < \omega^{\alpha}$  then we have  $l(M_*) = \beta$  and  $l(M/M_*) = \omega^{\alpha}n$ .

Proof Since  $M$  is Noetherian, the existence of  $M_*$  is clear in view of 2.1. By 2.3 we can choose a submodule  $N$  in  $M$  such that  $l(M/N) = \omega^{\alpha}n$ . Using 2.6 we obtain  $\mu_{\alpha}(N) = 0$ , hence  $\text{Kdim } N < \alpha$ , so  $N \subseteq M_*$ . Moreover it follows from 2.1 that

$$\omega^{\alpha}n + l(N) \leq l(M) \leq \omega^{\alpha}n \oplus l(N) = \omega^{\alpha}n + l(N)$$

Hence

$$\omega^{\alpha}n + l(N) = l(M) = \omega^{\alpha}n + \beta$$

so  $l(N) = \beta$ . It suffices to show that  $N = M_*$ . Since

$$\mu_{\alpha}(M/M_*) = \mu_{\alpha}(M)$$

we have

$$l(M/M_*) = \omega^{\alpha}n + \gamma$$

for some  $\gamma$ . Using 2.1 on the exact sequence

$$0 \rightarrow M_*/N \rightarrow M/N \rightarrow M/M_* \rightarrow 0$$

we obtain

$$\omega^{\alpha}n + \gamma + l(M_*/N) \leq l(M/N) = \omega^{\alpha}n.$$

Hence we have  $l(M_*/N) = 0$  so  $M_* = N$ . ■

2.9 Theorem. Let  $M$  be a Noetherian module and consider the following sets of ordinals:

$$A(M) := \{\beta : l(M) = \gamma + \beta \text{ for some ordinal } \gamma\}$$

$$lS(M) := \{l(N) : N \subseteq M\}$$

$$C(M) := \{\beta : \beta \ll l(M)\}$$

Then we have

$$A(M) \subseteq lS(M) \subseteq C(M) .$$

Proof We will first prove that  $A(M) \subseteq lS(M)$  . Let  $\gamma$  and  $\beta$  be ordinals such that  $l(M) = \gamma + \beta$  . We are going to show the existence of a submodule  $N \subseteq M$  such that  $l(N) = \beta$  .

Let  $\alpha$  be the degree of  $\beta$  . We may write

$$l(M) = \gamma' + \omega^{\alpha}m + \beta'$$

where  $\deg \beta' < \alpha$  and where each term in the Cantor normal form of  $\gamma'$  has degree greater than  $\alpha$  . Clearly there exists an integer  $n \leq m$  such that

$$\beta = \omega^{\alpha}n + \beta'$$

By repeated application of the operation  $*$  in 2.8 we obtain a submodule  $N_1 \subseteq M$  such that

$$l(N_1) = \omega^{\alpha}m + \beta'$$

By 2.3 we can find a submodule  $N \subseteq N_1$  such that

$$l(N_1/N) = \omega^{\alpha}(m-n)$$

Using 2.1 on the exact sequence

$$0 \rightarrow N \rightarrow N_1 \rightarrow N_1/N \rightarrow 0$$

we obtain

$$\omega^{\alpha}(m-n) + l(N) \leq l(N_1) \leq \omega^{\alpha}(m-n) \oplus l(N) = \omega^{\alpha}(m-n) + l(N)$$

Hence

$$\omega^{\alpha(m-n)} + l(N) = l(N_1) = \omega^{\alpha m + \beta'}$$

So

$$l(N) = \omega^{\alpha n + \beta'} = \beta .$$

To prove the relation  $lS(M) \subseteq C(M)$ , let  $N$  be any submodule of  $M$ . We are going to show that  $l(N) \ll l(M)$ , i.e.  $\mu_{\alpha}(N) \leq \mu_{\alpha}(M)$  for all  $\alpha$ . We will use induction on the dimension of  $M$ . If  $Kdim M = 0$ , then  $N$  and  $M$  have finite length, and the inequalities are satisfied in this case.

We will now assume that  $Kdim M > 0$ . By the obvious induction hypothesis it follows that

$$(1) \quad \mu_{\alpha}(N \cap M_*) \leq \mu_{\alpha}(M_*) \quad \text{for all } \alpha .$$

Moreover, it follows from 2.7 that

$$(2) \quad \mu_{\alpha}(M_*) = \mu_{\alpha}(M) \quad \text{for all } \alpha \neq Kdim M$$

$$(3) \quad \mu_{\alpha}(M_*) = 0 \quad \text{for } \alpha = Kdim M .$$

There are two cases:

(i)  $Kdim N < Kdim M$ . In this case we have  $N = N \cap M_*$ .

Hence by (1), (2) and (3) we have

$$\mu_{\alpha}(N) \leq \mu_{\alpha}(M) \quad \text{for all } \alpha .$$

(ii)  $Kdim N = Kdim M$ . In this case we have  $N_* = N \cap M_*$ .

For  $\alpha \neq Kdim M$  we have

$$\mu_{\alpha}(N) = \mu_{\alpha}(N_*) \leq \mu_{\alpha}(M_*) = \mu_{\alpha}(M)$$

For  $\alpha = Kdim M$  it follows from 2.7 that

$$\mu_{\alpha}(N) = \mu_{\alpha}(M) - \mu_{\alpha}(M/N) \leq \mu_{\alpha}(M)$$

2.10 Remark Jategaonkar shows in [4] that, given any ordinal  $\alpha$ , there is a principal right ideal domain  $R$  whose proper right ideals are linearly ordered of order type  $\omega^\alpha$ . Considering  $R$  as a right module it is easily seen that we have  $A(R) = lS(R)$ . In 3.2 below we shall see that if  $M$  is a Noetherian module over a commutative ring, then we have  $lS(M) = C(M)$ . This, combined with 2.4, shows that  $A(M)$  is not equal to  $lS(M)$  in general.

The inclusion  $lS(M) \subseteq C(M)$  expresses that if  $N$  is a submodule of  $M$ , then each of the coefficients in the Cantor normal form of  $l(N)$  is less than or equal to the corresponding coefficient in the Cantor normal form of  $l(M)$ . This will be referred to as the principle of coefficientwise comparison.

2.11 Proposition Let  $M$  be a Noetherian module, and let  $M_1$  and  $M_2$  be submodules such that  $M = M_1 + M_2$ . Then the sum is direct if and only if

$$l(M) = l(M_1) \oplus l(M_2)$$

Proof We will first show that

$$l(M') \oplus l(M'') = l(M' \oplus M'')$$

The inequality  $\geq$  follows immediately from 2.1. We are going to show the opposite inequality by induction on  $l(M'')$ . For  $l(M'') = 0$  there is nothing to prove. Now let  $l(M'') > 0$ , and let  $l(M')$ ,  $l(M'')$  and  $l(M' \oplus M'')$  be denoted by  $\alpha'$ ,  $\alpha''$  and  $\alpha$  respectively. Letting  $\beta$  be a variable running over the ordinals less than  $\alpha''$  we have  $\alpha'' = \sup(\beta+1)$ . For each value of  $\beta$  we can find (2.3) a proper factor module  $\bar{M}''$  of  $M''$  such that  $l(\bar{M}'') = \beta$ . Since  $M' \oplus \bar{M}''$  is a proper factor-module of  $M' \oplus M''$ , it follows

by the obvious induction hypothesis that

$$l(M') \oplus l(\bar{M}'') \leq l(M' \oplus \bar{M}'') < l(M' \oplus M'') = \alpha$$

Hence

$$(\alpha' \oplus \beta) + 1 \leq \alpha$$

This gives

$$\alpha' \oplus \alpha'' = \alpha' \oplus (\sup(\beta+1)) = \sup((\alpha' \oplus \beta)+1) \leq \alpha,$$

which was to be shown.

Let us now assume that

$$l(M) = l(M_1) \oplus l(M_2)$$

It remains to show that  $M_1 \cap M_2 = 0$ . We have an exact sequence

$$0 \rightarrow M' \cap M'' \rightarrow M' \oplus M'' \rightarrow M \rightarrow 0$$

Using 2.1 we obtain

$$l(M) + l(M' \cap M'') \leq l(M' \oplus M'') = l(M') \oplus l(M'') = l(M)$$

Hence  $l(M' \cap M'') = 0$  so  $M' \cap M'' = 0$ . ■

**2.12 Proposition** Let  $M$  be a Noetherian module. Assume that  $Kdim M$  is countable. Then there exists a well ordered chain of non-zero submodules of  $M$  of ordinal type equal to  $l(M)$ .

Proof We will use induction on  $l(M)$ . Put  $\alpha := Kdim M$ . If  $l(M)$  is finite, then the proposition is obvious. Hence we may assume that  $\alpha \geq 1$ . We will first treat the case where  $l(M) = \omega^\alpha$ . Since  $\alpha$  is countable, we can find a non-decreasing sequence of ordinal numbers less than  $\alpha$

$$\beta_1 \leq \beta_2 \leq \dots \leq \beta_n \leq \dots$$

such that

$$\omega^\alpha = \omega^{\beta_1} + \omega^{\beta_2} + \dots + \omega^{\beta_n} + \dots$$

We are going to construct a filtration of non-zero submodules

$$M = M_0 \supset M_1 \supset M_2 \supset \dots$$

such that

$$l(M_{i-1}/M_i) = \omega^{\beta_i} \quad \text{for } i \geq 1.$$

We put  $M_0 := M$ . Now let  $i \geq 1$  and assume that  $M_0, \dots, M_{i-1}$  has been constructed. By the principle of coefficientwise comparison (2.10) any non-zero submodule of  $M$  has length equal to  $\omega^\alpha$ , hence

$$l(M_{i-1}) > \omega^{\beta_i}$$

Thus by 2.3 we can find a non-zero submodule  $M_i \subset M_{i-1}$  such that

$$l(M_{i-1}/M_i) = \omega^{\beta_i},$$

and the construction is complete.

By the induction hypothesis  $M_{i-1}/M_i$  contains a chain consisting of non-zero submodules and having ordinal type equal to  $\omega^{\beta_i}$ . Clearly these chains induce a chain in  $M$  of ordinal type  $\omega^\alpha$ .

In the general case we can write

$$l(M) = \omega^\alpha n + \beta$$

where  $n \neq 0$  and  $\beta < \omega^\alpha$ . By the first part of the proof we may assume that  $l(M) > \omega^\alpha$ . By 2.3 we can find a non-zero submodule  $N \subset M$  such that  $l(M/N) = \omega^\alpha$ . Using 2.1 on the exact sequence

$$0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0$$

we obtain

$$\omega^\alpha + l(N) \leq l(M) \leq \omega^\alpha \oplus l(N) = \omega^\alpha + l(N)$$

Hence

$$l(M) = \omega^\alpha + l(N).$$

By the induction hypothesis,  $M/N$  and  $N$  contain chains of ordinal type  $\omega^\alpha$  and  $l(N)$  respectively. Two such chains clearly induce a chain in  $M$  of ordinal type  $l(M)$ .  $\blacksquare$

§3 Noetherian modules over commutative rings.

In this section all modules are assumed to be finitely generated over a commutative Noetherian ring  $R$ . The results depend heavily on the assumption that  $R$  be commutative.

3.1 Lemma Let  $M$  be a module with length

$$l(M) = \omega^{\alpha} n + \gamma$$

where  $n \neq 0$  is a natural number and  $\gamma < \omega^{\alpha}$ . Let  $k$  be an integer such that  $0 \leq k \leq n$ . Then  $M$  contains a submodule  $N$  such that  $l(N) = \omega^{\alpha} k$ .

Proof By ascending induction on  $k$  we are going to construct submodules

$$0 = N_0 \subset \dots \subset N_k \subset \dots \subset N_n$$

such that  $l(N_k) = \omega^{\alpha} k$ . Assume that  $1 \leq k \leq n$  and that  $N_0, \dots, N_{k-1}$  has been constructed. By 2.7 we have

$$u_{\alpha}(M/N_{k-1}) = (n-k+1) \neq 0$$

Hence  $\text{Kdim } M/N_{k-1} = \alpha$ , so there exists a prime ideal  $\mathfrak{p}$  in  $\text{Ass}(M/N_{k-1})$  such that  $\text{Kdim } R/\mathfrak{p} = \alpha$ . In view of 2.5 we have  $l(R/\mathfrak{p}) = \omega^{\alpha}$ . There exists an injection of  $R/\mathfrak{p}$  into  $M/N_{k-1}$ . The image of  $R/\mathfrak{p}$  in  $M/N_{k-1}$  pulls back to a submodule in  $M$  which we will denote by  $N_k$ . Thus we have an exact sequence

$$0 \rightarrow N_{k-1} \rightarrow N_k \rightarrow R/\mathfrak{p} \rightarrow 0$$

By (2.1) we obtain  $l(N_k) = l(N_{k-1}) + l(R/\mathfrak{p}) = \omega^{\alpha} k$ . ■

3.2 Theorem Let  $M$  be a Noetherian module over a commutative ring  $R$ , and let the length  $l(M)$  have Cantor normal form

$$l(M) = \omega^{\alpha_k n_k} + \dots + \omega^{\alpha_1 n_1}$$

where  $n_1 \dots n_k \neq 0$ . Then

(i)  $M$  is an essential extension of a direct sum of submodules  $N_i$  such that  $l(N_i) = \omega^{\alpha_i n_i}$  ( $1 \leq i \leq k$ ).

(ii)  $\{l(N) : N \subseteq M\} = \{\beta \ll l(M)\}$

(iii)  $\{\alpha_1, \dots, \alpha_k\} = \{Kdim R/\mathfrak{p} : \mathfrak{p} \in Ass M\}$

Proof (i). Using 2.8 and the previous lemma we see that  $M$  contains submodules  $N_i$  such that  $l(N_i) = \omega^{\alpha_i n_i}$  for  $1 \leq i \leq k$ . Put  $N := \sum_{i=1}^k N_i$ . Using 2.11 in combination with the principle of coefficientwise comparison (2.10) one easily shows that this sum is direct and that  $l(N) = l(M)$ . The last relation shows that  $M$  is an essential extension of  $N$ .

(ii). With the notation introduced in 2.9 we are going to show  $lS(M) = C(M)$ . Since the inclusion  $\subseteq$  was established in 2.9 we need only take care of the opposite inclusion. Let  $\beta$  be an arbitrary ordinal such that  $\beta \ll l(M)$ . We can write

$$\beta = \omega^{\alpha_k b_k} + \dots + \omega^{\alpha_1 b_1}$$

where  $b_i \leq n_i$  for  $1 \leq i \leq k$ . By 3.1 we can find submodules  $L_i \subseteq N_i$  such that  $l(L_i) = \omega^{\alpha_k b_k}$ . Put  $L := \sum_{i=1}^k L_i$ . Clearly this sum is direct, so by 2.11 we obtain  $l(L) = \beta$ .

(iii). We shall first prove the inclusion  $\subseteq$ . Let  $\alpha$  be one of the members in the set  $\{\alpha_1, \dots, \alpha_k\}$ . By (possibly repeated) application of the  $*$ -operation in 2.8 to  $M$ , we obtain a submodule  $N \subseteq M$  with Krull dimension  $\alpha$ . Hence there is a prime ideal  $\mathfrak{p} \in Ass N \subseteq Ass M$  such that  $Kdim R/\mathfrak{p} = \alpha$ . Conversely,

let  $\mathfrak{p}$  be a prime ideal in  $\text{Ass } M$  such that  $\text{Kdim } R/\mathfrak{p} = \alpha$ . Then  $M$  contains an isomorphic copy of  $R/\mathfrak{p}$  having length equal to  $\omega^\alpha$ . By the principle of coefficientwise comparison (2.10),  $\alpha$  is one of the exponents  $\alpha_1, \dots, \alpha_k$  in the Cantor normal form of  $l(M)$ . ■

**3.3 Definition** As in [1] we let  $o(M)$  denote the supremum of the ordinal types of descending chains of non-zero submodules of  $M$ .

We close this section by expressing  $o(M)$  in terms of  $l(M)$ .

**3.4 Theorem** Let  $M$  be Noetherian module over a commutative ring. Then we have

$$o(M) = \min(\omega_1, l(M)) ,$$

where  $\omega_1$  denotes the first non-countable ordinal.

Proof Let us first treat the case where  $l(M) < \omega_1$ . In this case  $\text{Kdim } M$  is countable. It follows from 2.12 that  $o(M) \geq l(M)$ . On the other hand it is easily seen that we (in general) have  $o(M) \leq l(M)$ . Hence

$$o(M) = l(M)$$

which proves the theorem in this case.

Let us now treat the case where  $l(M) \geq \omega_1$ . Let  $\beta$  be an arbitrary ordinal less than  $\omega_1$ . By 2.3 there exists a submodule  $N_\beta \subset M$  such that  $l(M/N_\beta) = \beta$ . By 2.12  $M/N_\beta$  has a descending chain of non-zero modules of ordinal type  $\beta$ , hence such a chain also exists in  $M$ . This gives  $o(M) \geq \beta$ , so  $o(M) \geq \omega_1$ . On the other hand, by 1.1 in [1] every chain in  $M$  is countable, so

$o(M) \leq \omega_1$  . This gives

$$o(M) = \omega_1$$

and the proof is now complete. ■

#### References.

- [1] H. Bass, Descending chains and the Krull ordinal of commutative Noetherian rings. J. Pure and Appl. Algebra 1 (1971) 347-360.
- [2] R. Gordon, J.C. Robson, Krull dimension, critical modules and monoform modules. To appear.
- [3] T.H. Gulliksen, The Krull ordinal, coprof and Noetherian localizations of large polynomial rings. To appear in Amer.J.Math.
- [4] A.V. Jategaonkar, A counter-example in ring theory and homological algebra. J.Algebra 12 (1969) 418-440.
- [5] G. Krause, On the Krull-dimension of left Noetherian left Matlis-rings, Math.Z. 118(1970) 207-214.