

PRIMARY RESEARCH**Open Access**

TWIST1, A novel androgen-regulated gene, is a target for NKX3-1 in prostate cancer cells

Turid Eide^{1*}, Håkon Ramberg¹, Carlotta Glackin³, Donald Tindall⁴ and Kristin Austlid Taskén^{1,2}

Abstract

Background: TWIST1 plays a key role in EMT-mediated tumor invasion and metastasis. Since bone metastasis is a hallmark of advanced prostate cancer and is detected in at least 85% of patients who die of this disease, it is of great importance to understand the regulation of the cellular signaling pathways involved in the metastatic process.

Methods: Prostatic cell lines were analyzed using real time RT-PCR, chromatin immunoprecipitations (ChIP) and transfection of siRNA's and reporter constructs.

Results: We report in this paper that TWIST1 is an androgen-regulated gene under tight regulation of NKX3-1. Androgens repress the expression of TWIST1 via NKX3-1, which is a prostate-specific tumor suppressor that is down-regulated in the majority of metastatic prostate tumors. We show that NKX3-1 binds to the TWIST1 promoter and that NKX3-1 over-expression reduces the activity of a TWIST1 promoter reporter construct, whereas NKX3-1 siRNA up-regulates endogenous TWIST1 mRNA in prostate cancer cells.

Conclusion: Our finding that NKX3-1 represses TWIST1 expression emphasizes the functional importance of NKX3-1 in regulating TWIST1 expression during prostate cancer progression to metastatic disease.

Keywords: Prostate cancer, LNCaP cells, TWIST1, NKX3-1

Background

Metastasis is the leading cause of morbidity and mortality among men with prostate cancer (PCa). Metastasis is a complex multistep process controlled by distinct genes and signaling pathways in each step. Epithelial-mesenchymal transition (EMT), a critical event for morphogenetic movements during formation of parietal endoderm during gastrulation, may represent the initial phase of metastasis. Furthermore, EMT seems to induce stem-like properties of epithelial cells [1].

A pivotal step of EMT is the loss of E-cadherin [2]. TWIST1, a master regulator of mesodermal development and a key mediator in the metastatic process, represses the expression of E-cadherin [3]. Furthermore, TWIST1 depletion reduces the expression of N-cadherin in PC3 cells, a metastatic prostate cancer cell line, suggesting that TWIST1 supports EMT in prostate cancer [4]. In support of this, a positive correlation between the level of TWIST1 and prostate cancer metastasis has been reported [5].

NKX3-1 is a critical gene associated with early stage of prostate tumorigenesis as down-regulation of NKX3-1 is observed in both prostatic intraepithelial neoplasia (PIN) and adenocarcinomas [6]. Whereas high levels of TWIST1 is expressed in prostate cancer metastasis, low levels of NKX3-1 expression is observed in most prostate cancer metastasis examined [7]. NKX3-1 encodes a homeobox gene that is switched on during embryonic development of prostate tissue and is one of the earliest markers of luminal prostate epithelium [8]. NKX3-1 expression is strictly regulated by androgens and also appears to mark a sub-population of prostate stem cells [9,10].

We report in this paper that TWIST1 is a novel androgen-regulated gene whose expression is tightly controlled by NKX3-1.

Results**TWIST1 is regulated by androgens**

The expression of TWIST1 mRNA increased 4-fold in LNCaP cells incubated with 0.5 nM R1881 for 72 hours relative to untreated cells (ctr.) and reached more than a 10-fold induction at 10 nM R1881 (Figure 1A).

* Correspondence: turid.eide@ous-hf.no

¹Department of Tumor Biology, Institute for Cancer Research, Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway
Full list of author information is available at the end of the article




Figure 1 Androgen up-regulates TWIST1 expression. **A.** LNCaP cells were stimulated with increasing doses of the synthetic androgen R1881, and the relative level of TWIST1 compared to LNCaP cells left untreated in charcoal stripped fetal calf serum and phenol red free medium (CSS) was quantified using sqRT-PCR. Data are presented as mean \pm SD (n=3) relative to unstimulated cells (control). A paired t-test was performed and a two-tailed p-value <0.05 is indicated with a *. **B.** LNCaP cells were stimulated with increasing doses of the synthetic androgen R1881 or left untreated in CSS for 3 days before total protein was extracted. A representative Western blot probed with anti human TWIST1 antibody and anti PRKAR1A antibody as loading control is shown.

Corresponding regulation of TWIST1 mRNA was also demonstrated in RWPE-1, a benign prostate cell line (data not shown). Up-regulation of TWIST1 expression was also observed at the protein level in LNCaP cells stimulated with increasing concentrations of R1881 (Figure 1B).

To determine if AR mediates the effect of androgens on TWIST1 expression, LNCaP cells were transfected with a siRNA targeting AR (AR) or a non-silencing siRNA as control (ctr.). Down-regulation of AR expression (Figure 2, lower panel) reduced the mRNA level of TWIST1 by 80% relative to control (Figure 2A, upper panel) supporting the concept that TWIST1 is an AR target gene in LNCaP cells. Down-regulation of AR also reduced the level of NKX3-1 (Figure 2, lower panel).


TWIST1 is an NKX3-1 target gene

In an effort to find binding sites for AR (ARE) or androgen-regulated transcription factors, we examined the human TWIST1 genomic sequence using MatInspector [11]. Three putative binding sites for NKX3-1 designated BS1, BS2 and BS3 (Figure 3A) were identified. In addition we observed several potential AREs. Chromatin immunoprecipitation analyses (ChIP) using chromatin isolated

from LNCaP cells stimulate with 10 nM R1881 (+R1881) for 24 hours or unstimulated cells as control (-R1881) resulted in amplification of BS3. We were not able to detect binding to either BS1 or BS2 (data not shown, n=3). A 5-fold increase in NKX3-1 binding activity to the TWIST1 promoter upon R1881 stimulation was observed (Figure 3B). Additionally, PCR amplification of non-targeted DNA (genomic contig of chromosome 12) was used as negative control (data not shown).

Next, a reporter construct containing 5kb of the mouse TWIST1 promoter (pGL3-TWIST-Luc) was co-transfected with an NKX3-1 expression vector. Over-expression of NKX3-1 reduced the promoter activity of TWIST1 by 50%, indicating that NKX3-1 represses the TWIST1 promoter activity (Figure 3C). No effect of NKX3-1 over-expression was observed on cells transfected with pGL3 lacking the TWIST1 promoter (pGL3).

Finally, the effect of NKX3-1 on endogenous TWIST1 mRNA level was studied by transfecting LNCaP cells with siRNA targeting NKX3-1. A 5-fold increase in TWIST1 mRNA level was observed when NKX3-1 expression was down-regulated (Figure 3D). The impact of R1881 stimulation, NKX3-1 overexpression and siRNA on NKX3-1



expression in LNCaP cells are summarized in Figure 3E. Both R1881 stimulation as well as overexpression of NKX3-1 led to a significant increase in NKX3-1 expression, while siRNA against NKX3-1 reduced the expression markedly.

Discussion

In this study we show both that TWIST1 mRNA is up-regulated by androgen via AR and that NKX3-1, a well-known androgen-regulated gene, binds the upstream

regulatory region of the TWIST1 gene and represses the expression of TWIST1.

In a recent study, Takayama et al. [12] identified putative direct target genes of AR using ChIP-on ChIP and Cap Analysis Gene Expression (CAGE). One of the genes identified by *in silico* analysis was TWIST1. Our results support this report as increased levels of TWIST1 mRNA was observed in R1881 stimulated LNCaP cells, while reduced expression of TWIST1 was observed in cells transfected with siRNA targeting AR.

Furthermore, our data shows binding of NKX3-1 to the promoter region of TWIST1. We therefore suggest that NKX3-1 mediates an indirect and late effect of androgen stimulation on TWIST1 expression. Interestingly, we show that siRNA targeting AR reduces the level of TWIST1, whereas siRNA targeting NKX3-1 increases TWIST1 expression suggesting that TWIST1 expression is tightly controlled by androgen. NKX3-1 has been shown to function both as an activator and a repressor of transcription, but few target genes have been identified [13-15].

The physical binding of NKX3-1 to the TWIST1 promoter might block the mesenchymal drive of TWIST1, until NKX3-1 expression is down-regulated or lost in PIN or adenocarcinoma lesions. Loss of NKX3-1 expression has been observed in ~20% of PIN lesions, ~40% of advanced prostate tumors and up to 80% of metastatic prostate cancer [16]. Androgen deprivation therapy as the most widely used treatment for advanced prostate cancer is likely to abolish androgen-stimulation of NKX3-1, leading eventually to down-regulation of repressor protein and de-repression of TWIST1's metastatic potential.

In an attempt to identify genes whose regulation are altered by NKX3-1, Song et al. [17] performed gene expression profiling analyses on micro dissected glands from NKX3-1-deficient prostate tissues during prostate cancer progression. They observed similarities between the expression profile of the micro dissected glands and constitutive activated AKT-transgenic mice as well as PTEN-deficient mice, suggesting that the PTEN-AKT-NKX3-1 axis serve as a major molecular path of prostate tumorigenesis. Li and Zhou [18] showed that activation of the AKT pathway by TWIST1 is critical for the sustention of cancer stem cell-like traits generated by EMT, again suggesting a link between loss of NKX3-1 expression, relive of TWIST1 expression and eventually activation of AKT pathway.

Conclusions

We report in this paper that TWIST1 is an androgen-regulated gene, tightly regulated by NKX3-1. We show that NKX3-1 binds to the TWIST1 promoter and that NKX3-1 over-expression reduces the activity of a TWIST1 promoter reporter construct, whereas NKX3-1 siRNA up-regulated endogenous TWIST1 mRNA in prostate cancer




Figure 3 (See legend on next page.)

(See figure on previous page.)

Figure 3 NKX3-1 mediates androgen-regulation of TWIST1. **A.** Three putative NKX3-1-binding sites (BS1, BS2 and BS3) are shown as boxes in the TWIST1 promoter (from NCBI MapViewer). **B.** Chromatin immunoprecipitation was performed using chromatin isolated from LNCaP cells stimulate with 10 nM R1881 (+R1881) for 24 hours or left untreated (-R1881) and using a monoclonal antibody against NKX3-1 (NKX3-1). sqPCR results using primers targeting BS3 are shown in B. The results of the BS3 primer set are shown adjusted to the levels of chromatin immunoprecipitated with IgG as a negative control. Data are presented as mean \pm SD (n=3). **C.** LNCaP cells were transfected with pGL3-TWIST-Luc (pGL3-TWIST) and a human NKX3-1 expression plasmid or an empty vector (pCR3) as well as pCMV β -gal. As control, LNCaP cells were transfected with a reporter plasmid lacking the TWIST promoter region (pGL3) and the human NKX3-1 expression vector or an empty expression plasmid (pCR3). The luciferase activities were normalized against the corresponding β -galactosidase activities and the results are shown as mean \pm SD (n=3). A paired t-test was performed and a two-tailed p value <0,05 is indicated with a *. **D.** LNCaP cells were transfected with 100 nM of siRNA targeting NKX3-1 or a non-silencing siRNA (ctr.) and total RNA was harvested 72 hours after transfection. TWIST1 mRNA expression was examined using sqRT-PCR. The data are shown as mean \pm SD (n=3) relative to the levels of LNCaP cells transfected with a non-silencing control siRNA. **E.** Representative Western blots showing expression levels of NKX3-1 in (from top) LNCaP cells stimulated with 10 nM of R1881, transfected with NKX3-1 overexpression vector or targeted by siRNA against NKX3-1. α -tubulin is shown as loading control.

cells. Our finding that NKX3-1 represses TWIST1 expression emphasizes the functional importance of NKX3-1 in regulating TWIST1 expression during prostate cancer progression to metastatic disease.

Methods

Cell culture

LNCaP and RWPE-1 cells were purchased from ATCC (Rockville, MD) and cultured in RPMI 1640 medium containing 10% fetal calf serum (FCS) or Keratinocyte-SFM medium from Invitrogen (Carlsbad, CA, USA) supplemented with 2.5 μ g Epidermal Growth Factor (EGF) and 25 mg Bovine Pituitary Extract (BPE), respectively, and stimulation with synthetic androgen R1881 was performed as previously described [19].

Semi-quantitative real time RT-PCR (sqRT-PCR)

Total RNA was isolated using Trizol™ from Invitrogen (Carlsbad, CA), and 100 ng of total RNA was used in a one-step RT-PCR reaction (QIAGEN Quantitect SYBR Green RT-PCR kit) that was performed using an MJ Research DNA Engine Opticon Continuous Fluorescence Detection System (MJ Research Inc., Waltham, MA). RT-PCR cycles were performed as previously described by Ramberg et al. [20]. G6PD was used for normalization. The $\Delta\Delta$ Ct formula was used as described in the protocol from Applied Biosystems (Foster City, CA). All the PCR-products were verified by sequencing.

Primer sets used in sqRT-PCR

G6PD (NM_000402); Left: tgcatagagccagatagc and right: acagggaggagatgtggttg, NKX3-1 (NM_006167); Left: gagacgtggcagagacc and right: ttctgctgctgcttaggg, AR (NM_000044); Left: gcgatcttcaccaatgta and right: cattcggacactggctgt, TWIST1 (NM_000474); Left: ctctcgtctgaggatgg and right: ctctctctggaacaatgaca.

Immunoblotting

Protein extraction followed by Western blot analysis was performed as previously described by Kvissel et al. [19]. Primary antibodies used were the following: anti-TWIST1 (H-81, sc-15393), anti-AR (N-20, sc-816), both from Santa Cruz Biotechnology (Santa Cruz, CA). The anti-NKX3-1 antibody was kindly provided by Professor Fahri Saatcioglu at Department of Molecular Biosciences, University of Oslo, Norway. For loading control we used anti-PRKARIA (610609, BD Transduction Laboratories) or anti- α -tubulin antibody from Sigma (St. Louis, MO).

Transfection and luciferase assay

LNCaP cells were cultured at 300.000 cells per 6-well dish and transfected with a luciferase reporter plasmid including 5 kb of the mouse TWIST1 promoter- pGL3-TWIST (kindly provided by Steven Kendall and Carlotta Glackin, Beckman Research Institute, City of Hope, USA) or pGL3 as negative control using Lipofectamine 2000 (INVT11668019, Invitrogen) according to the manufacturer's protocol. For overexpression of NKX3-1, a commercial transfection-ready TRUE clone (sc116287, ORIGENE, Rockville, MD, USA) was purchased. Cells were co-transfected with pCMV β -gal (Clontech) to monitor the transfection efficiency. After 72 hours, luciferase and β -galactosidase activity were measured as previously described [20].

Preparation and transfection of synthetic small interfering RNA (siRNA)

siRNA targeting the following sequence within the androgen receptor mRNA: 5'-AAAAGCCCATCGTAGAG GCCCA-3' was purchased from Dharmacon (Dharmacon Inc. (Lafayette, CO). A non-silencing siRNA (cat.no D-001810-10-20, Dharmacon) was included in all siRNA experiments. siRNA against NKX3-1 was purchased from QIAGEN (2-for-silencing) and included the following sequences: NKX3-1-1: 5'-CAGGCTATCATATATACTGTA-3', NKX3-1-2: 5'-ACGCTATAAGACTAAGCGAAA-3'.

All siRNAs were transfected into cells using Dharma-*TECT*[™] 3 transfection reagent according to the manufacturer's protocol (cat.no T-2003, Dharmacon).

Chromatin immunoprecipitation

ChIP assays were carried out according to the QuickChIP protocol (Imgenex, San Diego, CA) with a crosslinking time of 10 minutes using 1% formaldehyde for the LNCaP cells and using an anti human NKX3-1 monoclonal antibody (cat.no 35–9700, Zymed Laboratories Inc., San Francisco, CA). Following DNA purification (QuickChIP DNA Purification kit), sqPCR was performed using QIAGEN Quantitect SYBR Green RT-PCR kit with the following primers:

NKX3-1 BS3; Left: cccagttacacttgatgcagta and right: tccctggtgagatcatacatc. Negative control primers from human chromosome 12 genomic contig; Left: atggtgctactgggatct and right: tgccaaagcctagggaaga. Chromatin immunoprecipitated with IgG was used as a negative control. The $\Delta\Delta C_t$ formula was used as described in the protocol from Applied Biosystems (Foster City, CA).

Abbreviations

PCa: Prostate Cancer; EMT: Epithelial-Mesenchymal Transition; PIN: Prostatic Intraepithelial Neoplasia; AR: Androgen Receptor; ChIP: Chromatin Immunoprecipitation Analyses.

Competing interests

The authors declare that they have no competing interest.

Authors' contributions

TE took part in designing the study and conducted the experiments, collected and analyzed the data, and wrote the manuscript. HR conducted the Western blot analysis of androgen stimulation of TWIST1 and took part in reviewing of the manuscript. CG contributed with the mouse TWIST1-promoter plasmid and took part in reviewing of the manuscript. DJT and KAT both conceived and coordinated the study, and participated in the discussion of all analyses and critical revision of the manuscript. All authors have read and approved the final draft of the manuscript.

Acknowledgements

We thank the Department of Urology at the Mayo Clinic and Foundation in Rochester, Minnesota, USA and Department of Urology at the Oslo University Hospital, Aker. Furthermore, we thank Lucy Smith (Mayo Clinic, Rochester, USA) for skilled technical assistance and Fahri Saatcioglu (University of Oslo, Norway) for providing us with anti human NKX3-1 antibody. This work was supported by The Norwegian Research Council, Health Region South East, The Norwegian Cancer Society, University of Oslo and Oslo University Hospital, Aker. DJT was supported by NIH grants CA121277, CA91956, CA125747 and the T.J. Martell Foundation.

Author details

¹Department of Tumor Biology, Institute for Cancer Research, Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway. ²Faculty of Medicine, University of Oslo, Oslo, Norway. ³Division of Molecular Medicine, Beckman Research Institute of the City of Hope, Duarte 91010-3011CA, USA.

⁴Department of Urology, Mayo Clinic and Foundation, Rochester 55905MN, USA.

References

1. Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, et al: The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* 2008, **133**:704–715.
2. Birchmeier W, Behrens J: Cadherin expression in carcinomas: role in the formation of cell junctions and the prevention of invasiveness. *Biochim Biophys Acta* 1994, **1198**:11–26.
3. Yang J, Mani SA, Donaher JL, Ramaswamy S, Itzykson RA, Come C, et al: Twist, a master regulator of morphogenesis, plays an essential role in tumor metastasis. *Cell* 2004, **117**:927–939.
4. Alexander NR, Tran NL, Rekapally H, Summers CE, Glackin C, Heimark RL: N-cadherin gene expression in prostate carcinoma is modulated by integrin-dependent nuclear translocation of Twist1. *Cancer Res* 2006, **66**:3365–3369.
5. Kwok WK, Ling MT, Lee TW, Lau TC, Zhou C, Zhang X, et al: Up-regulation of TWIST in prostate cancer and its implication as a therapeutic target. *Cancer Res* 2005, **65**:5153–5162.
6. Bethel CR, Faith D, Li X, Guan B, Hicks JL, Lan F, et al: Decreased NKX3.1 Protein expression in focal prostatic atrophy, prostatic intraepithelial neoplasia, and adenocarcinoma: association with gleason score and chromosome 8p deletion. *Cancer Res* 2006, **66**:10683–10690.
7. Gelmann EP, Bowen C, Bubendorf L: Expression of NKX3.1 In normal and malignant tissues. *Prostate* 2003, **55**:111–117.
8. Bhatia-Gaur R, Donjacour AA, Sciavolino PJ, Kim M, Desai N, Young P, et al: Roles for Nkx3.1 In prostate development and cancer. *Genes Dev* 1999, **13**:966–977.
9. Prescott JL, Blok L, Tindall DJ: Isolation and androgen regulation of the human homeobox cDNA, NKX3.1. *Prostate* 1998, **35**:71–80.
10. Shen MM, Wang X, Economides KD, Walker D, Abate-Shen C: Progenitor cells for the prostate epithelium: roles in development, regeneration, and cancer. *Cold Spring Harb Symp Quant Biol* 2008, **73**:529–538.
11. Quandt K, Frech K, Karas H, Wingender E, Werner T: MatInd and MatInspector: new fast and versatile tools for detection of consensus matches in nucleotide sequence data. *Nucleic Acids Res* 1995, **23**:4878–4884.
12. Takayama K, Tsutsumi S, Katayama S, Okayama T, Horie-Inoue K, Ikeda K, et al: Integration of cap analysis of gene expression and chromatin immunoprecipitation analysis on array reveals genome-wide androgen receptor signaling in prostate cancer cells. *Oncogene* 2011, **30**:619–630.
13. Carson JA, Fillmore RA, Schwartz RJ, Zimmer WE: The smooth muscle gamma-actin gene promoter is a molecular target for the mouse bagpipe homologue, mNkx3-1, and serum response factor. *J Biol Chem* 2000, **275**:39061–39072.
14. Simmons SO, Horowitz JM: Nkx3.1 Binds and negatively regulates the transcriptional activity of Sp-family members in prostate-derived cells. *Biochem J* 2006, **393**:397–409.
15. Abate-Shen C, Shen MM, Gelmann E: Integrating differentiation and cancer: the Nkx3.1 Homeobox gene in prostate organogenesis and carcinogenesis. *Differentiation* 2008, **76**:717–727.
16. Bowen C, Bubendorf L, Voeller HJ, Slack R, Willi N, Sauter G, et al: Loss of NKX3.1 Expression in human prostate cancers correlates with tumor progression. *Cancer Res* 2000, **60**:6111–6115.
17. Song H, Zhang B, Watson MA, Humphrey PA, Lim H, Milbrandt J: Loss of Nkx3.1 Leads to the activation of discrete downstream target genes during prostate tumorigenesis. *Oncogene* 2009, **28**:3307–3319.
18. Li J, Zhou BP: Activation of beta-catenin and Akt pathways by twist are critical for the maintenance of EMT associated cancer stem cell-like characters. *BMC Cancer* 2011, **11**:49.
19. Kvissel AK, Ramberg H, Eide T, Svindland A, Skalhegg BS, Tasken KA: Androgen dependent regulation of protein kinase a subunits in prostate cancer cells. *Cell Signal* 2007, **19**:401–409.
20. Ramberg H, Eide T, Krobort KA, Levy FO, Dizzey N, Bjartell AS, et al: Hormonal regulation of beta2-adrenergic receptor level in prostate cancer. *Prostate* 2008, **68**:1133–1142.

doi:10.1186/1475-2867-13-4

Cite this article as: Eide et al.: TWIST1, A novel androgen-regulated gene, is a target for NKX3-1 in prostate cancer cells. *Cancer Cell International* 2013 **13**:4.