

Geophysical Research Abstracts, Vol. 8, 03063, 2006
SRef-ID: 1607-7962/gra/EGU06-A-03063
© European Geosciences Union 2006



An integrated study of interior layered deposits in Hebes Chasma, Valles Marineris, Mars

E. Hauber (1), K. Gwinner (1), A. Gendrin (2,6), F. Fueten (3), R. Stesky (4), S. Pelkey (2), H. Wulf (1), D. Reiss (1), T. Zegers (5), P. MacKinnon (3), and the HRSC Co-Investigator Team

(1) Institute of Planetary Research, German Aerospace Center (DLR), Berlin, (2) Brown University, Providence, (3) Brock University, St. Catharines, (4) Pangaea Scientific, Brockville, (5) ESTEC, ESA, Noordwijk (Ernst.Hauber@dlr.de/+49.30.67055.402)

Despite more than three decades of analysis, the origin of the Interior Layered Deposits (ILD) in the Valles Marineris (VM) trough system is still unknown. The advance of new remote sensing data obtained by the recent planetary missions Mars Global Surveyor (MGS), Mars Odyssey (MO), and Mars Express (MEX) allow investigation of the morphology and composition in unprecedented detail. This study focusses on Hebes Chasma (HC) in the central VM, which is unique because it contains a huge mesa of ILD in a completely closed depression. We used topographic data from the HRSC camera onboard MEX to analyze the geometry of layering in HM with the Orion structural analysis software. Strike and dip were measured in 50m/px gridded Digital Elevation Models and corresponding orthoimages. These data have a higher spatial resolution than those used in an earlier study. We find that the layers dip gently with up to 12° on the northern ILD wall (wall slope 20°) and with steeper slopes up to 22° on the southern ILD wall (wall slope >30°). The layers always dip in the down-slope direction. These results are in agreement with our earlier results in HC and with our similar ILD studies in western Candor and Ophir Chasmata, also based on HRSC topography. A mineralogic map indicating concentrations of polyhydrated sulfates, kieserite, and oxides was produced from OMEGA spectral data. It shows that these alteration minerals are only observed in low-lying areas, which are not covered by landslides. We consider a lacustrine origin of the ILD in HC as unlikely. The down-slope dipping of ILD layers is in agreement with a draping process, e.g., pyroclastic fall deposits from an W-E trending volcanic vent. The occurrence of alteration minerals

only in very deep portions of HC also argues against a deposition in a deep body of standing water. Groundwater, not meteoric water, might have played a major role in rock alteration, as elsewhere on Mars. Alternatively, it can not be excluded that sulfates and oxides belong to ancient altered deposits, which were exhumed during HC formation.