



## Geophysical Research Letters

### RESEARCH LETTER

10.1002/2013GL059159

#### Key Points:

- Airborne in situ measurements in the UTLS were performed with HALO
- Stratospheric tracer correlations are used to quantify mixing in the UTLS
- The stratospheric content relative to the tropopause is investigated

#### Correspondence to:

T. Jurkat,  
tina.jurkat@dlr.de

#### Citation:

Jurkat, T., et al. (2014), A quantitative analysis of stratospheric HCl, HNO<sub>3</sub>, and O<sub>3</sub> in the tropopause region near the subtropical jet, *Geophys. Res. Lett.*, 41, 3315–3321, doi:10.1002/2013GL059159.

Received 30 DEC 2013

Accepted 3 MAR 2014

Accepted article online 12 MAR 2014

Published online 14 MAY 2014

## A quantitative analysis of stratospheric HCl, HNO<sub>3</sub>, and O<sub>3</sub> in the tropopause region near the subtropical jet

T. Jurkat<sup>1</sup>, C. Voigt<sup>1,2</sup>, S. Kaufmann<sup>1,2</sup>, A. Zahn<sup>3</sup>, M. Sprenger<sup>4</sup>, P. Hoor<sup>2</sup>, H. Bozem<sup>2</sup>, S. Müller<sup>2</sup>, A. Dörnbrack<sup>1</sup>, H. Schlager<sup>1</sup>, H. Bönisch<sup>5</sup>, and A. Engel<sup>5</sup>

<sup>1</sup>Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany, <sup>2</sup>Institut für Physik der Atmosphäre, Johannes Gutenberg-Universität Mainz, Mainz, Germany, <sup>3</sup>Institute for Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany, <sup>4</sup>Institute for Atmospheric and Climate Science, Swiss Federal Institute of Technology Zurich, Zurich, Switzerland, <sup>5</sup>Institute for Atmospheric and Environmental Sciences, Goethe Universität Frankfurt, Frankfurt, Germany

**Abstract** The effects of chemical two-way mixing on the Extratropical Transition Layer (ExTL) near the subtropical jet are investigated by stratospheric tracer-tracer correlations. To this end, in situ measurements were performed west of Africa (25–32°N) during the Transport and Composition of the Upper Troposphere Lower Stratosphere (UTLS)/Earth System Model Validation (TACTS/ESMVal) mission in August/September 2012. The Atmospheric chemical Ionization Mass Spectrometer sampling HCl and HNO<sub>3</sub> was for the first time deployed on the new German High Altitude and Long range research aircraft (HALO). Measurements of O<sub>3</sub>, CO, European Centre for Medium-Range Weather Forecasts (ECMWF) analysis, and the tight correlation of the unambiguous tracer HCl to O<sub>3</sub> and HNO<sub>3</sub> in the lower stratosphere were used to quantify the stratospheric content of these species in the ExTL. With increasing distance from the tropopause, the stratospheric content increased from 10% to 100% with differing profiles for HNO<sub>3</sub> and O<sub>3</sub>. Tropospheric fractions of 20% HNO<sub>3</sub> and 40% O<sub>3</sub> were detected up to a distance of 30 K above the tropopause.

### 1. Introduction

Dynamical and chemical processes modify the ozone (O<sub>3</sub>) budget of the upper troposphere/lowermost stratosphere (UT/LMS) and lead to locally variable O<sub>3</sub> trends. In this region, O<sub>3</sub> has steep gradients and acts as a strong greenhouse gas with a positive radiative forcing (RF) depending on its location with respect to the tropopause. Only few models explicitly discuss the uncertainty of mixing parametrization and its effect on the RF. They emphasize that small changes in O<sub>3</sub> result in large changes in RF at the tropopause [Riese et al., 2012]. However, understanding of UT/LMS O<sub>3</sub> trends still remains incomplete [World Meteorological Organization (WMO), 2010].

Similar to O<sub>3</sub>, nitric acid (HNO<sub>3</sub>) shows steep gradients at the tropopause with low concentrations in the upper troposphere and a steady increase in the lower stratosphere. After downward transport into the troposphere, the predominant removal process is particle interaction with subsequent wash out or dry deposition [Murphy et al., 1993]. In addition to the stratospheric contribution, upper tropospheric HNO<sub>3</sub> can be formed in situ from biomass burning plumes, lightning, or aviation. The relative contribution of each source is globally highly variable and therefore difficult to quantify. Next to photochemical production and dry and wet deposition, mixing is an important part of irreversible distribution of tracers, driven by horizontal strain and vertical shear.

The Extratropical Transition Layer is a region of air with mixed tropospheric and stratospheric characteristics centered around the thermal tropopause and has been described extensively using tracer-tracer correlations of carbon monoxide (CO), O<sub>3</sub>, and water vapor (H<sub>2</sub>O) [Fischer et al., 2000; Hoor et al., 2004; Hegglin et al., 2009]. Tropospheric intrusions transporting low ozone from lower latitudes are known to ventilate the lower stratosphere above the subtropical jet [Pan et al., 2009]. During summer, the weaker isentropic potential vorticity (PV) gradient in the region of the subtropical jet (STJ) allows for a stronger mixing [Haynes and Shuckburgh, 2000].

Due to similar source regions at altitudes above 20 km and long chemical lifetimes in the lower stratosphere, in the order of months, it has been suggested that the correlation of the stratospheric tracer hydrogen chloride (HCl) with O<sub>3</sub> can be used to quantify stratospheric O<sub>3</sub> in the upper troposphere [Marcy et al.,

2004]. While it is the main chlorine reservoir gas in the stratosphere, HCl has a low tropospheric background, thus being an ideal tracer to study quantitative mixing of trace gases with both tropospheric and stratospheric sources.

Due to the steep gradients at the tropopause, highly resolved measurements of  $\text{HNO}_3$ ,  $\text{O}_3$ , and HCl are needed. To this end, we performed in situ measurements of HCl and  $\text{HNO}_3$  with a newly developed Atmospheric chemical Ionization Mass Spectrometer (AIMS) during the first atmospheric science HALO mission TACTS/ESMVal in August/September 2012. High spatial and temporal resolution of the trace gas measurements allow a good characterization of mixing near the tropopause.


This paper discusses the benefits and limitations of using the relation between stratospheric tracers to study mixing. By comparison of HCl/ $\text{HNO}_3$  and HCl/ $\text{O}_3$  in the stratosphere with an air mass probed in the vicinity of the subtropical jet, we describe the state of mixing and discuss the location of a mixing layer [Fischer *et al.*, 2000] relative to the dynamical and the thermal tropopause. While Marcy *et al.* [2004] discuss transport of stratospheric  $\text{O}_3$  into the upper troposphere, we extend the method to altitudes above the tropopause to determine the tropospheric content in the stratosphere. Additionally, we include a second mainly stratospheric tracer  $\text{HNO}_3$  in our analysis and compare the different mixing states in a region of enhanced isentropic transport. A quantitative composition analysis in the Extratropical Transition Layer (ExTL) is performed.

## 2. The Atmospheric Chemical Ionization Mass Spectrometer

The Atmospheric chemical Ionization Mass Spectrometer (AIMS) is a newly designed CIMS instrument, operated with an electrical discharge source and in-flight calibration. In contrast to earlier measurements using radioactive sources for ionization [Marcy *et al.*, 2005; Jurkat *et al.*, 2010],  $\text{SF}_5^-$  reagent ions are produced by an electrical discharge between a gold needle tip and a circular pin hole at the upstream part of the flow reactor. The reagent ions are instantaneously mixed in the flow tube with the sample flow generating stable product ions by fluoride transfer. Product and reagent ions are detected at a rate of 1 Hz by a linear quadrupole and a channeltron housed inside a differentially pumped vacuum chamber. Sensitivity of the instrument and possible absorption effects on the walls are monitored using an in-flight online calibration with isotopically labeled  $\text{SO}_2$ , which was also measured but will not be shown here. The relative sensitivity and stability for HCl and  $\text{HNO}_3$  are likewise monitored and are stable within named measurement errors over the whole flight on 4 September 2012. For an absolute calibration, two pressure- and temperature-controlled permeation ovens are used to generate standards for HCl and  $\text{HNO}_3$  calibration. The backward facing trace gas inlet mounted on the central fuselage of the aircraft is equipped with a 1/2" heated perfluoroalkoxy (PFA) tube ( $T = 40^\circ\text{C}$ ). A high bypass flow of 4.7 slm maintained at ambient pressure is achieved by an additional scroll pump. A smaller flow of 1.1 slm enters the mass spectrometer through a PFA pressure control valve. Background measurements are hourly performed by flushing the inlet with dry synthetic air. The  $1\sigma$  detection limit of the instrument for both HCl and  $\text{HNO}_3$  was between 10 and 15 parts per trillion by volume (pptv) during the campaign with a running mean over 20 data points and for atmospheric  $\text{H}_2\text{O} < 100$  ppmv. A systematical error is estimated to be between 18 and 25%. The reagent ions additionally allow detection of nitrous acid (HONO) [Voigt *et al.*, 2010; Jurkat *et al.*, 2011] and chlorine nitrate ( $\text{ClONO}_2$ ) [Marcy *et al.*, 2005], which are not considered here. In addition, the instrument can be used to measure high concentrations of tropospheric HCl,  $\text{HNO}_3$ , and  $\text{SO}_2$  in volcanic plumes [Voigt *et al.*, 2014].  $\text{O}_3$  mixing ratios were measured at a rate of 10 Hz with high precision and reproducibility of 0.5% and 1.5% using a combination of a chemiluminescence sensor and an UV photometer [Zahn *et al.*, 2012]. The instrument measuring CO is based on the three-channel tunable diode laser instrument for atmospheric research used during SPURenstofftransport in der Tropopausenregion (SPURT) [Hoor *et al.*, 2004]. With a new quantum cascade laser setup, we achieved a short-term precision of 0.9 ppbv at 1.5 s integration time and an upper limit for the total uncertainty of 2.3 ppbv without any corrections applied.


## 3. In Situ Measurements of HCl and $\text{HNO}_3$ in the Tropopause Region

The TACTS/ESMVal mission comprised 13 flights from latitudes of  $68^\circ\text{N}$  to  $65^\circ\text{S}$  and  $25^\circ\text{E}$  to  $74^\circ\text{W}$  with more than 120 flight hours. The aim of the campaign was to study the transport and composition of the UT/LMS region and provide observations for global climate model validation.



**Figure 1.** (a) Flight track of HALO on 4 September 2012 and 10-day backward trajectories ending on the flight track, color coded with the distance to the thermal tropopause (DTH). In the vicinity of the subtropical jet at 26°N (grey-shaded area), the backward trajectories show a change in air mass origin from stratospheric (DTH > 0) to tropospheric values when the aircraft flew from high to low latitudes. The black contour lines show mean zonal wind velocities of 20 and 25 m/s during the 10 day period. (b) Trace gas mixing ratios of HCl (red), HNO<sub>3</sub> (green), O<sub>3</sub> (blue), and CO (black) measured on 4 September 2012 together with the HALO cruise altitude. The grey-shaded area corresponding to Figure 1a marks the mixed air mass. (c) Potential temperature ( $\theta$ , black) derived from onboard HALO measurements together with calculated measured concentration in potential vorticity (PV, yellow) from operational ECMWF data, the distance to the thermal (DTH\_WMO, red), and dynamical tropopause (DTH\_2PV, blue), respectively. (d) The color code shows horizontal wind velocities (white  $\leq 2.5$  m/s to light green 32 m/s, in steps of 2.5 m/s) as a curtain along the flight track derived from ECMWF analysis. Grey contours are potential temperature isolines, in black the flight altitude and in light grey the dynamical tropopause (2 PVU) are shown. At approximately 12:00 UT, the aircraft entered a region with enhanced horizontal wind velocities.

In this paper we concentrate on one flight performed on 4 September 2012 going from Oberpfaffenhofen, Germany, to Sal, Cape Verde. In Figure 1a, the flight track and 10-day backward trajectories run with the Lagrangian Analysis Tool LAGRANTO [Wernli and Davies, 1997] based on operational ECMWF data are shown. Using these trajectory calculations, we derive the most recent contact to the stratosphere for air masses sampled below the tropopause. Figures 1b and 1c show the time series of HCl, HNO<sub>3</sub>, O<sub>3</sub> and CO




**Figure 2.** Scatterplots of 10 s averaged mole fractions of CO vs O<sub>3</sub>, HCl vs O<sub>3</sub>, and HCl vs HNO<sub>3</sub>. A tight correlation ( $R^2 = 0.8$  and  $0.84$ ) is observed for all stratospheric tracers in the unperturbed lower stratosphere (red, O<sub>3</sub> > 280 ppbv). The stratospheric fit curves with slopes  $m_1$  and  $m_2$  are colored in blue. Data marked in green correspond to the grey-shaded area in Figure 1 and were collected in the vicinity of the subtropical jet. Black squares represent the median of 30 ppbv O<sub>3</sub> and 0.1 ppbv HNO<sub>3</sub> bins for O<sub>3</sub> < 280 ppbv.

tropopause in the vicinity of the STJ reveals the largest potential temperature difference up to 57 K, with the dynamical tropopause 38 K below the aircraft and the thermal tropopause at a distance of 19 K above. To study the effects of mixing of tropospheric and stratospheric air, we use the tropopause definitions and tracer-tracer correlations of HCl to O<sub>3</sub> and HCl to HNO<sub>3</sub>.

#### 4. Stratospheric Tracer-Tracer Coordinates

Tracer-tracer coordinates are commonly used to study the chemical transition from troposphere to stratosphere. While some studies apply CO as a tropospheric marker and O<sub>3</sub> as a stratospheric marker [Zahn et al., 2000; Hoor et al., 2004], we focus on the relationship of two stratospheric tracers [Marcy et al., 2004]. To support our analysis we use the tropospheric tracer CO. Figure 2 shows scatterplots of CO to O<sub>3</sub>, HCl to O<sub>3</sub> and HCl to HNO<sub>3</sub> for the flight on 4 September 2012. While the branching of the CO-O<sub>3</sub> correlation and the large CO variability between 100 and 280 ppb O<sub>3</sub> indicate mixing between tropospheric and stratospheric air, the scatterplots of the stratospheric tracers show a compact linear relationship. To separate the characteristic, unperturbed stratospheric background from the mixing layer, slopes for different O<sub>3</sub> ranges have been derived. By computing the slopes for lower O<sub>3</sub> cutoff values between 100 and 350 ppbv in steps of 50 ppbv, a change of the slope of 15% for the fit of all data points as compared to the fit for O<sub>3</sub> > 280 ppbv was observed. The change in the slope will be used here as an independent criterion to derive the upper

together with flight altitude, potential temperature ( $\theta$ ), vertical distance to the thermal (DTH\_WMO) and the dynamical tropopause (DTH\_2PV), measured in potential temperature and the potential vorticity along the flight path. The dynamical tropopause is represented by the 2 PVU-isoline in the extratropics and the 380 K-isentrope in the tropics. The thermal tropopause is calculated according to the WMO definition based on the standard lapse rate. Within the 8.5 h flight both tropospheric and stratospheric air was encountered, from 20 K below the tropopause to 57 K above the tropopause, and potential temperatures between 300 and 382 K. Tropospheric background mixing ratios of HCl encountered at 7:50 UT and after 13:20 UT are below the detection limit of 15 pptv. This lack of HCl has been observed during most flights and confirms the exclusive stratospheric origin of upper tropospheric HCl in the absence of recent volcanic eruptions. In the following discussion we focus on the trace gas relation in the grey-shaded area. Here PV is dropping from stratospheric (6 PVU) to tropospheric values (1 PVU). Figure 1d shows the horizontal wind velocities (color coded) in a vertical curtain plot along the flight route. The STJ is represented by high horizontal wind velocities with maximum values of 32 m/s in the center of the jet. The high horizontal wind speeds are extending deep into the stratosphere where HALO sampled at the upper edge of these extensions. Isentropes are crossing the dynamical tropopause indicating a potential pathway of cross-tropopause trace gas transport. Comparison of the thermal and the dynamical



**Figure 3.** The relative stratospheric contributions  $O_{3, \text{strat}}^*$  and  $HNO_{3, \text{strat}}^*$  and the 5 K percentiles (median (black squares), 10th and 90th (dark blue lines)) plotted against the distance to the (a, b) dynamical and (c, d) thermal tropopause are shown. Colored in light grey are data of the entire flight. The dark grey-colored data represent the mixed air mass above the tropopause. Data below the thermal tropopause have been color coded with the trajectory based analysis of the most recent contact to the stratosphere (in hours). Light green colored is air with the most recent contact to the stratosphere  $\geq 240$  h.

boundary of the mixing layer equivalent to the lower boundary of the unperturbed lower stratosphere. The boundary is consistent with a broadening of the stratospheric branch in the CO-O<sub>3</sub> tracer-tracer space below 280 ppbv O<sub>3</sub>. The linear fits for the unperturbed stratospheric branch (data with O<sub>3</sub> > 280 ppbv) yield high correlation coefficients of  $R^2 = 0.8$  and  $0.84$ , with a slope of  $m_1 = 8.1 \times 10^{-4}$  for the HCl and O<sub>3</sub> correlation and a slope of  $m_2 = 0.239$  for the fit of HCl and HNO<sub>3</sub>.

The data within the grey-shaded area in Figure 1 are green colored in Figure 2. They show a systematic lower ratio of HCl and O<sub>3</sub> and HCl and HNO<sub>3</sub> than the corresponding ratio on the extrapolated stratospheric fit curve (blue fit line). Additionally, the fits for the correlations of tracers sampled in the mixing layer have significantly higher slopes of  $1 \times 10^{-3}$  for HCl/O<sub>3</sub> and  $0.324$  for HCl/HNO<sub>3</sub>.

In the following discussion, X represents the mole fraction of either O<sub>3</sub>, HNO<sub>3</sub>, or HCl in the mixed (index M) air mass, and the stratospheric (index S) and tropospheric (index T) end-members, respectively. The index *strat* is used for the stratospheric content  $X_{\text{strat}}$  of the observed mole fraction  $X_M$  in the mixed air mass, thus  $X_M = X_{\text{strat}} + X_{\text{trop}}$ . We assume linear mixing of air along a mixing line between a stratospheric end-member ratio  $HCl_S/X_S$  located on the stratospheric fit curve and an upper tropospheric end-member ratio  $HCl_T/X_T$  with  $HCl_T = 0$ . Thus, we derive the stratospheric content  $X_{\text{strat}}$  based on *Marcy et al.* [2004]:

$$X_{\text{strat}} = \frac{HCl_M X_S}{HCl_S} \quad (1)$$

and  $X_{\text{strat}}^* = X_{\text{strat}}/X_M$  is the relative stratospheric content. To calculate  $X_{\text{strat}}^*$ , we use the stratospheric end-member at the lower end of the stratospheric fit curve. The choice of the end-member implies that the stratospheric part in the mixed air mass originated from the lower boundary of the unperturbed stratosphere. For our calculation we use the stratospheric end-member at  $O_{3,S} = 280$  ppbv and  $HNO_{3,S} = 0.63$  ppbv, closest to the ratios in the mixed air mass in the tracer-tracer space, with  $HCl_S/O_{3,S} = 7.9 \times 10^{-4}$  and  $HCl_S/HNO_{3,S} = 0.318$ .

### 5. Two-Way Transport Across the Tropopause

The vertical profiles of  $O_{3, \text{strat}}^*$  and  $HNO_{3, \text{strat}}^*$  with respect to the dynamical (Figures 3a and 3b) and the thermal tropopause (Figures 3c and 3d) are shown. For a clearer picture of the profiles and to minimize the influence of small-scale variability, the median, the 10th, and 90th percentiles are calculated for 5 K bins in the vertical.

The mixed air mass marks a distinct transition layer between pure tropospheric (0%  $X_{\text{strat}}^*$ ) and pure stratospheric air (100%  $X_{\text{strat}}^*$ ). The stratospheric content ranges between 20% and 100% for  $O_{3, \text{strat}}^*$  and 30% and

100% for  $\text{HNO}_3^*_{3,\text{strat}}$ . It is decreasing with decreasing distance from the tropopause. The air mass was sampled mostly above the dynamical but centered around the thermal tropopause at potential temperatures between 320 K and 370 K. Thus, mixed air with significant tropospheric content was detected above the thermal and dynamical tropopause and mixed air with stratospheric content below the thermal tropopause. For  $\text{O}_3^*_{3,\text{strat}}$  and  $\text{HNO}_3^*_{3,\text{strat}}$  below the thermal tropopause, different transport time scales, derived from trajectory analysis, can be distinguished: air with recent contact to the stratosphere corresponds to a higher stratospheric content of 60%–80% while air with the last contact to the stratosphere at more than 240 h before the measurement contains only 20%–30% of stratospheric  $\text{O}_3$  and  $\text{HNO}_3$ . By using the HCl-tagged air, we can unambiguously distinguish the stratospheric contribution to the upper tropospheric  $\text{HNO}_3$  and  $\text{O}_3$  budget with additional information of the time of tropopause crossing from backward trajectory calculations. Thus, we see mixing in the vicinity of the STJ affecting the ExTL up to 30–40 K above the low extratropical thermal tropopause. The difference in the mixing state between the two profiles arises from the different sources and background concentrations of upper tropospheric  $\text{O}_3$  as compared to upper tropospheric  $\text{HNO}_3$  and the difference in gradients across the tropopause. Tropospheric air with low  $\text{O}_3$  is mixed into the stratosphere with generally high  $\text{O}_3$  mole fractions. After mixing the stratospheric air exhibits a reduced  $\text{O}_3$  concentration and is therefore diluted with respect to  $\text{O}_3$ . Due to different gradients of  $\text{O}_3$  and  $\text{HNO}_3$ , the stratospheric air mass experiences a stronger dilution of the  $\text{O}_3$  concentration than dilution of  $\text{HNO}_3$  concentration.

## 6. Discussion

Using tracer-tracer correlations to study mixing demands an analysis of the stability and lifetime of the trace gases compared to the mixing time scales. The following discussion focuses on the instrumental and atmospheric parameters affecting the analysis. Generally, the deviation of  $\text{HCl}/\text{O}_3$  from the stratospheric ratio may have three independent reasons which may all influence the ratio: (1) mixing of tropospheric with stratospheric air, (2) solution of HCl and  $\text{HNO}_3$  in liquid water droplets, and (3) heterogeneous reactions on cirrus cloud ice particles, e.g., trapping of  $\text{HNO}_3$ . HCl activation on cirrus clouds may cause a reduction in the concentration of up to 50% but only for high particulate surface areas [Thornton *et al.*, 2007]; these are rarely observed in the stratosphere. Trapping and burial of  $\text{HNO}_3$  in tropical cirrus cloud particles, potentially leading to a small depletion in the gas phase with subsequent vertical redistribution, have been observed in the tropical tropopause. A climatology and overview are given in Voigt *et al.* [2006]. In the present case, the slope of the two trace gas correlations is shifted and tilted when compared to the stratospheric slope but it is still a relative compact mixing line with correlation coefficients of  $R^2 = 0.89$  and  $0.9$ . In case of chemical or microphysical removal of either component, the compactness would have suffered. Therefore, we argue that mixing is mainly responsible for the observed shift in the ratios of the trace gases.

The criteria for the choice of the stratospheric end-members leave some uncertainty to the method. To compare the range of potential end-members, we investigated the variability in the stratosphere and in the troposphere; here  $\text{O}_3$  is used as an example. The slope of the stratospheric fit curve represents an upper limit of potential stratospheric end-members of  $\text{HCl}_5/\text{O}_{3,5} = 8.1 \times 10^{-4}$ . The lower limit is given for  $\text{O}_3$  at 280 ppbv with  $\text{HCl}_5/\text{O}_{3,5} = 7.9 \times 10^{-4}$ . The choice of this end-member is also supported by a broadening of the stratospheric branch in the  $\text{CO}/\text{O}_3$  correlation. The respective tropospheric end-members  $\text{O}_{3,T}$  are on average  $104 \pm 26$  ppbv, thus within the expected range for typical upper tropospheric concentrations [Murphy *et al.*, 1993]. We examine the spread of potential tropospheric end-members by deriving the frequency of occurrence of  $\text{O}_{3,T}$  for  $\text{HCl}_T < \text{detection limit}$  and altitudes  $> 6$  km for midlatitude measurements. The distribution was fitted by a gaussian fit with a mean of  $74 \pm 24$  ppbv. Due to a smaller spread, the uncertainty for the calculation of  $X_{\text{strat}}$  is minimized by using a stratospheric end-member on the stratospheric fit curve.

A lower limit of stratospheric  $\text{O}^*_{3,\text{strat}}$  and  $\text{HNO}^*_{3,\text{strat}}$  that can be derived from this method is given by the detection limit of the HCl measurements. A detection limit of 15 pptv gives an uncertainty of 16 ppbv  $\text{O}_{3,\text{strat}}$  and 0.05 ppbv  $\text{HNO}_{3,\text{strat}}$ . The relative uncertainty is largest at the lowest mixing ratios.

The slope  $m_1$  of the stratospheric fit is higher than former measurements [Marcy *et al.*, 2004; Jurkat *et al.*, 2010], but the slopes agree within the named uncertainties of both measurements. The deviation may also partly evolve from a latitude dependence of the ratio considering that Marcy *et al.* [2004] sampled at subtropical latitudes where  $\text{O}_3$  is produced at a higher rate. The slopes  $m_1$  and  $m_2$  and therefore  $\text{HCl}_5/\text{O}_{3,5}$

and  $\text{HCl}_5/\text{HNO}_{3,5}$  for tropical regions may be smaller which would result in an underestimation of the stratospheric content  $X_{\text{strat}}$ .

## 7. Summary and Outlook

In this paper we present a new method to quantify the stratospheric content of  $\text{HNO}_3$  and  $\text{O}_3$  in a mixing layer above and below the tropopause with stratospheric tracer-tracer correlation analysis. High-resolution airborne measurements of  $\text{HCl}$  and  $\text{HNO}_3$  in the UT/LMS have been conducted with the AIMS instrument onboard the new research aircraft HALO during the TACTS/ESMVal Mission. The mixed layer was observed at low concentrations of  $\text{HCl}$ ,  $\text{HNO}_3$ , and  $\text{O}_3$  up to 30 K above the thermal and dynamical tropopause in the ExTL. The profiles of  $\text{O}_3$  and  $\text{HNO}_3$  show an increasing tropospheric content with decreasing distance from the tropopause. We propose to use the ratios  $\text{HCl}/\text{O}_3$  and  $\text{HCl}/\text{HNO}_3$  as a good proxy to study quantitatively the effects of mixing in the ExTL. Additionally, upper tropospheric  $\text{O}_3$  and  $\text{HNO}_3$  from local or remote sources can be differentiated from downward transported stratospheric  $\text{O}_3$  and  $\text{HNO}_3$ . Future analysis will focus on regional and seasonal variability of mixing and transport of these and other stratospheric tracers in the ExTL. Our measurements extend data from remote sensing instruments on satellites in the lower stratosphere to lower altitudes and better resolutions.

### Acknowledgments

TACTS/ESMVal was funded by the DFG SPP-HALO 1294 and the DLR-Project ESMVal. Part of the work was funded by HGF contract VH-NG-309. Further, we thank the DLR flight department for their excellent support during the campaign.

The Editor thanks two anonymous reviewers for their assistance in evaluating this paper.

### References

- Fischer, H., F. G. Wienhold, P. Hoor, O. Bujok, C. Schiller, P. Siegmund, M. Ambaum, H. A. Scheeren, and J. Lelieveld (2000), Tracer correlations in the northern high latitude lowermost stratosphere: Influence of cross-tropopause mass exchange, *Geophys. Res. Lett.*, *27*(1), 97–100, doi:10.1029/1999GL010879.
- Haynes, P., and E. Shuckburgh (2000), Effective diffusivity as a diagnostic of atmospheric transport: 1. Stratosphere, *J. Geophys. Res.*, *105*(D18), 22,777–22,794, doi:10.1029/2000JD900093.
- Hegglin, M. I., C. D. Boone, G. L. Manney, and K. A. Walker (2009), A global view of the extratropical tropopause transition layer from atmospheric chemistry experiment fourier transform spectrometer  $\text{O}_3$ ,  $\text{H}_2\text{O}$ , and  $\text{CO}$ , *J. Geophys. Res.*, *114*, D00811, doi:10.1029/2008JD009984.
- Hoor, P., C. Gurk, D. Brunner, M. I. Hegglin, H. Wernli, and H. Fischer (2004), Seasonality and extent of extratropical TST derived from in-situ  $\text{CO}$  measurements during SPURT, *Atmos. Chem. Phys.*, *4*, 1427–1442, doi:10.5194/acp-4-1427-2004.
- Jurkat, T., C. Voigt, F. Arnold, H. Schlager, H. Aufmhoff, J. Schmale, J. Schneider, M. Lichtenstern, and A. Dörnbrack (2010), Airborne stratospheric ITCIMS measurements of  $\text{SO}_2$ ,  $\text{HCl}$ , and  $\text{HNO}_3$  in the aged plume of volcano Kasatochi, *J. Geophys. Res.*, *115*, D00L17, doi:10.1029/2010JD013890.
- Jurkat, T., C. Voigt, F. Arnold, H. Schlager, J. Kleffmann, H. Aufmhoff, D. Schäuble, M. Schaefer, and U. Schumann (2011), Measurements of  $\text{HONO}$ ,  $\text{NO}$ ,  $\text{NO}_y$  and  $\text{SO}_2$  in aircraft exhaust plumes at cruise, *Geophys. Res. Lett.*, *38*, L10807, doi:10.1029/2011GL046884.
- Marcy, T., R. Gao, M. Northway, P. Popp, H. Stark, and D. Fahey (2005), Using chemical ionization mass spectrometry for detection of  $\text{HNO}_3$ ,  $\text{HCl}$ , and  $\text{ClONO}_2$  in the atmosphere, *Int. J. Mass Spectrom.*, *243*(1), 63–70, doi:10.1016/j.ijms.2004.11.012.
- Marcy, T. P., et al. (2004), Quantifying stratospheric ozone in the upper troposphere with in situ measurements of  $\text{HCl}$ , *Science*, *304*(5668), 261–265, doi:10.1126/science.1093418.
- Murphy, D. M., D. W. Fahey, M. H. Proffitt, S. C. Liu, K. R. Chan, C. S. Eubank, S. R. Kawa, and K. K. Kelly (1993), Reactive nitrogen and its correlation with ozone in the lower stratosphere and upper troposphere, *J. Geophys. Res.*, *98*(D5), 8751–8773, doi:10.1029/92JD00681.
- Pan, L. L., W. J. Randel, J. C. Gille, W. D. Hall, B. Nardi, S. Massie, V. Yudin, R. Khosravi, P. Konopka, and D. Tarasick (2009), Tropospheric intrusions associated with the secondary tropopause, *J. Geophys. Res.*, *114*, D10302, doi:10.1029/2008JD011374.
- Riese, M., F. Ploeger, A. Rap, B. Vogel, P. Konopka, M. Dameris, and P. Forster (2012), Impact of uncertainties in atmospheric mixing on simulated UTLS composition and related radiative effects, *J. Geophys. Res.*, *117*, D16305, doi:10.1029/2012JD017751.
- Thornton, B. F., et al. (2007), Chlorine activation near the midlatitude tropopause, *J. Geophys. Res.*, *112*, D18306, doi:10.1029/2006JD007640.
- Voigt, C., H. Schlager, H. Ziereis, B. Kärcher, B. P. Luo, C. Schiller, M. Krämer, P. J. Popp, H. Irie, and Y. Kondo (2006), Nitric acid in cirrus clouds, *Geophys. Res. Lett.*, *33*, L05803, doi:10.1029/2005GL025159.
- Voigt, C., et al. (2010), In-situ observations of young contrails: Overview and selected results from the CONCERT campaign, *Atmos. Chem. Phys.*, *10*(18), 9039–9056.
- Voigt, C., P. Jessberger, T. Jurkat, S. Kaufmann, R. Baumann, H. Schlager, N. Bobrowski, G. Guffrda, and G. Salerno (2014), Evolution of  $\text{SO}_2$ ,  $\text{HCl}$ ,  $\text{HNO}_3$  and  $\text{CO}_2$  in the volcanic plume from Etna, *Geophys. Res. Lett.*, doi:10.1002/2013GL058974.
- Wernli, H., and H. C. Davies (1997), A lagrangian-based analysis of extratropical cyclones. I: The method and some applications, *Q. J. R. Meteorol. Soc.*, *123*(538), 467–489, doi:10.1002/qj.49712353811.
- WMO, W. M. O. (2010), Scientific assessment of ozone depletion: 2010, *WMO Rep. 52*, World Meteorol. Org., Geneva, Switzerland.
- Zahn, A., et al. (2000), Identification of extratropical two-way troposphere-stratosphere mixing based on CARIBIC measurements of  $\text{O}_3$ ,  $\text{CO}$ , and ultrafine particles, *J. Geophys. Res.*, *105*(D1), 1527–1535, doi:10.1029/1999JD900759.
- Zahn, A., J. Weppner, H. Widmann, K. Schlote-Holubek, B. Burger, T. Kühner, and H. Franke (2012), A fast and precise chemiluminescence ozone detector for eddy flux and airborne application, *Atmos. Meas. Tech.*, *5*(2), 363–375, doi:10.5194/amt-5-363-2012.