



Pyridine-promoted dediazonation of aryl diazonium tetrafluoroborates: Application to the synthesis of SF₅-substituted phenylboronic esters and iodobenzenes

George Iakobson¹, Junyi Du², Alexandra M. Z. Slawin² and Petr Beier^{*1}

Full Research Paper

Open Access

Address:

¹Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Flemingovo nám. 2, 166 10 Prague 6, Czech Republic and ²EaStCHEM School of Chemistry, University of St Andrews, St Andrews, KY16 9ST, United Kingdom

Email:

Petr Beier^{*} - beier@uochb.cas.cz

* Corresponding author

Keywords:

borylation; diazonium salts; iodination; pyridine; sulfur pentafluorides

Beilstein J. Org. Chem. **2015**, *11*, 1494–1502.

doi:10.3762/bjoc.11.162

Received: 25 March 2015

Accepted: 05 August 2015

Published: 26 August 2015

Associate Editor: V. Gouverneur

© 2015 Iakobson et al; licensee Beilstein-Institut.

License and terms: see end of document.

Abstract

Pyridine promotes dediazonation of aryl diazonium tetrafluoroborates. The formed aryl radicals were trapped with B₂pin₂, iodine, or tetrahydrofuran to afford boronic esters, iodobenzenes and benzenes, respectively. The application to the synthesis of (pentafluorosulfanyl)phenylboronic esters, iodo(pentafluorosulfanyl)benzenes and (pentafluorosulfanyl)benzene is shown.

Introduction

Pentafluorosulfanyl-containing compounds have been known for more than half a century [1–4]; however, for a long time they remained a relatively underdeveloped class of compounds [5,6]. The main reason for the slow development of the chemistry of SF₅-containing compounds was the lack of availability of key building blocks. However, in recent years, the scientific community has been witnessing a renewed interest in this functional group. Synthetic methods towards aliphatic SF₅-containing compounds are based on free radical addition of SF₅Cl or SF₅Br to unsaturated compounds [7–9], whereas aromatic derivatives are available either by the Umemoto's two-step synthesis from diaryl disulfides or benzenethiols [10–12], or by the reaction of nitrophenyl disulfides with elemental fluorine [13–16]. Aromatic and heteroaromatic SF₅ compounds are

mostly prepared by the derivatization of commercial nitro-(pentafluorosulfanyl)benzenes [14,17–27] and approaches from SF₅-aliphatics have also been studied [28–30]. The unique combination of properties the SF₅ group imparts includes high chemical, thermal, and metabolic stability, strong electron-acceptor property, and high lipophilicity. Furthermore, applications of SF₅ compounds in catalysis [31,32], life-science [6,18,33–38], and material sciences [5,19,38,39] are emerging.

Arylboronic acids and arylboronates represent versatile building blocks in organic synthesis [40]. They have found wide applications in transition metal-catalyzed cross-coupling reactions [41,42]. These boron compounds are accessed mainly by the reactions of arylmagnesium or aryllithium species with trialkyl-

boronates [43,44], Pd- or Cu-catalyzed borylations of aryl halides using $B_2\text{pin}_2$, H-Bpin [45–50] or $R_2\text{N}\text{-BH}_2$ [51], direct borylations via aromatic C–H bond activations [52–58], Lewis acid catalyzed electrophilic borylations of electron-rich arenes [59–62], and Sandmeyer-type borylation of arylamines or diazonium salts with $B_2\text{pin}_2$ [63–67], $\text{B}_2(\text{OH})_4$ [68] or $R_2\text{N}\text{-BH}_2$ [69]. Several attempts were made to synthesize the SF_5 -phenylboronates. Patent literature describes the synthesis of 3- or 4-(pentafluorosulfanyl)phenylboronates or boronic acids from SF_5 -bromobenzenes via lithiation or magnesiation. These approaches suffer from low yields and other drawbacks [70,71]. For lithiation of the aryl bromide, $t\text{-BuLi}$ had to be used and the formation of Grignard reagents is inefficient. On the other hand, Shibata and co-workers have recently reported the synthesis of 3,5-bis(pentafluorosulfanyl)phenylboronic acid from the corresponding aryl bromide, trimethyl borate and $i\text{PrMgBr}$ [32]. Finally, Joliton and Carreira have recently shown efficient Ir-catalyzed C–H borylation of several 1-substituted-3-(pentafluorosulfanyl)benzenes and applied the products of borylation to the Pd-catalyzed Suzuki–Miyaura reaction with aryl bromides or iodides. However, the reaction is limited to borylations in position five of 1-substituted-3-(pentafluorosulfanyl)benzenes [72].

Straightforward access to SF_5 -phenylboronic acids or boronates would be highly desirable since it would allow easy installation of the SF_5 -phenyl group by the subsequent Suzuki–Miyaura reaction. Nitro-(pentafluorosulfanyl)benzenes are the primary industrial SF_5 -aromatics, therefore the easiest access to SF_5 -phenylboronates appears to be starting from readily available SF_5 -substituted anilines or diazonium salts rather than SF_5 -containing halobenzenes. Herein, we report a new protocol for efficient borylation, iodination and hydrodediazonation of SF_5 -phenyldiazonium tetrafluoroborates in the presence of pyridine. The generality of the borylation and iodination reactions was demonstrated on several examples.

Results and Discussion

At the onset of our investigation, Sandmeyer-type borylation of 3- and 4-(pentafluorosulfanyl)anilines (**1a** and **1b**, respectively) to pinacolboronates **2a** and **2b** according to Wang and co-workers was studied [64,65] (Table 1). The borylation of **1a** took place in a reasonable yield in the presence of catalytic amounts of benzoyl peroxide (BPO, Table 1, entry 1), while for **1b**, heating without any additives was preferable; however, the yield of **2b** was only moderate (Table 1, entry 3).

For a detailed investigation of the borylation reaction, the diazonium tetrafluoroborates **3a** and **3b** were prepared and isolated according to Doyle conditions [73]. High yields (above 90%) of both isomers of the diazonium tetrafluoroborates **3a** and **3b**


Table 1: Synthesis of boronates **2** from aniline derivatives **1**^a.

Entry	1	Additive	Temp. (°C)	2 , Yield (%) ^b
1	1a , 3-SF ₅	BPO	rt	2a , 70
2	1b , 4-SF ₅	BPO	rt	2b , traces
3	1b , 4-SF ₅	—	80	2b , 55 ^c

^aReaction conditions: **1** (1 mmol), *t*-BuONO (1.5 mmol), $B_2\text{pin}_2$ (1.1 mmol), additive (2 mol %), MeCN (4 mL). ^bIsolated yield. ^c82% purity.

were obtained on 0.3–6 g scale (Table 2). Crystal structures of **3a** (CCDC 1009848) and **3b** (CCDC 1009849) were determined confirming the nature of the products. During the course of our studies, Okazaki and co-workers reported the synthesis of **3b** in 84% yield under similar conditions and have shown its reactivity in various cross-coupling reactions with varied degree of success. The most efficient cross-coupling reactions were the Heck reactions with alkenes, a biaryl homocoupling reaction, an azo coupling to electron-rich arenes, and a dediazonation with TMSN_3 in an ionic liquid medium [74,75].

An efficient borylation of aryldiazonium tetrafluoroborates with NHC-Pd catalysts was reported recently [63]. When applied to **3a** and **3b** using $\text{Pd}(\text{OAc})_2$ and NHC ligand precursors **L**, the borylated products **2a** and **2b** were isolated in only moderate yields (Table 2, entries 1 and 2). However, it was found that the Pd catalyst was not required for an efficient reaction. Alkali metal acetates are known to facilitate decomposition of aryldiazonium salts by the formation of diazoacetates and diazo anhydrides, which decompose to aryl radicals [76]. These additives were used in Meerwein arylation of isopropenyl acetate [77]. In our case, two-fold excess of sodium acetate in acetonitrile afforded **2a** and **2b** in good and moderate yields, respectively (Table 2, entries 3 and 4). The conditions reported by Yu and co-workers [66] (Table 2, entry 5) provided a mixture with starting **3b** as the major component. While the diazonium salts **3a** and **3b** were found to be stable in acetonitrile, we observed slow decomposition in methanol and in the presence of $B_2\text{pin}_2$ under strictly Pd-free conditions (new Teflon stirring bar and glassware), the borylation took place with low conversion (Table 2, entry 6) [78]. In pyridine, however, the decomposition of **3b** was very fast and a vigorous evolution of nitrogen was observed affording the borylated product in a moderate yield together with a mixture of SF_5 -pyridines in ca. 10% GC–MS yield (Table 2, entry 7). Finally, the use of a

Table 2: Synthesis of diazonium tetrafluoroborates **3^a** and boronates **2^b**.

Entry	3 (mmol)	Reagents (equiv), solvent	Temp. (°C)	Time (h)	2 , Yield (%) ^c
1	3a (1)	Pd(OAc) ₂ (0.01), L (0.02), THF	rt	18	2a , 43
2	3b (1)	Pd(OAc) ₂ (0.01), L (0.02), THF	rt	18	2b , 57
3	3a (1)	NaOAc (2), MeCN	-50 to rt	3	2a , 73
4	3b (1)	NaOAc (2), MeCN	-50 to rt	3	2b , 49
5	3b (1)	CuBr (0.05), MeCN/H ₂ O (3:1)	rt	330	2b , 23
6	3a (1)	—, MeOH	rt	24	2a , 32
7	3b (1)	—, pyridine ^d	-30 to rt	2	2b , 55
8	3b (1)	pyridine (4), MeCN ^d	-30 to rt	2	2b , 77
9	3b (3)	pyridine (4), MeCN ^d	-30 to rt	2	2b , 77
10	3a (2)	pyridine (4), MeCN ^d	-30 to rt	2	2a , 80

^aReaction conditions: **1** (1–28 mmol), BF₃·OEt₂ (2.1 equiv), t-BuONO (1 equiv), CH₂Cl₂ or Et₂O (3 mL/1 mmol of **1**), 30 min. ^bReaction conditions: **3** (1–3 mmol), B₂pin₂ (1 equiv), reagents, solvent (2 mL/1 mmol of **3**) under N₂. ^cIsolated yield. ^dThe reaction was conducted under air.


4-fold excess of pyridine in acetonitrile was found to be optimal. Conducting the reaction on a gram scale proceeded without a notable loss of efficiency and the reaction can be performed in air (Table 2, entries 8–10).

Pyridine is known to induce decomposition of aryl diazonium salts. Zollinger and Abramovitch studied the interaction of pyridine with aryl diazonium tetrafluoroborates and suggested the formation of diazopyridinium salts which homolytically decompose to aryl radicals, nitrogen and a pyridinium tetrafluoroborate radical [79,80]. Tanaka and co-workers have used the combination of PhN₂⁺BF₄⁻ and pyridine for arylations of silylenol ethers [81]. They have observed the formation of large amounts of phenylpyridines. In our case, we detected SF₅-phenylpyridines in trace amounts only during borylation using pyridine as a solvent. The major products in the borylation reaction apart from **2a** and **2b** were found to be F-Bpin and pyridine-BF₃ complexes. Both compounds are easily hydrolyzable but they were observed by NMR of the crude reaction mixture and compared to the synthesized authentic samples. On the other hand, we were not able to observe a pyridine-B₂pin₂ complex by ¹¹B NMR in CD₃CN or CDCl₃. The borylation was extended to several other aryl diazonium tetrafluoroborates showing that both electron-donor and electron-acceptor substituted phenyldiazonium tetrafluoroborates undergo efficient


borylation with an equimolar amount of B₂pin₂ (Scheme 1); however, *ortho*-substituted phenyldiazonium salts were found to be either not efficient substrates (**3f**) or completely unreactive (**3g**), presumably due to a large steric demand of B₂pin₂.

The mechanism of this borylation reaction remains to be elucidated. Based on experimental results and literature precedent, we propose the following free-radical mechanism (Scheme 2). Aryldiazonium salt **3** reacts with pyridine to form aryl diazopyridinium **4** which decomposes to an aryl radical, a pyridinium tetrafluoroborate radical and nitrogen. The aryl radical reacts with B₂pin₂ to form the borylated product **2** and the Bpin radical (likely to be stabilized by pyridine) [82]. The byproducts pyridine-BF₃ and F-Bpin are formed by the reaction of the pyridinium tetrafluoroborate radical and the Bpin radical.

Subjecting the diazonium salt **3i** to the borylation conditions gave further indirect evidence for the formation of aryl radicals (Scheme 3). Full conversion of **3i** was observed affording a mixture of products **2i**, **2i'**, **5i** (two isomers of unknown configuration) and **6i** in 16:31:28:25 GC–MS ratio. The presence of these products can be explained only by the formation of the substituted SF₅-phenyl radical which undergoes borylation to **2i**, hydrogen atom transfer followed by borylation to **2i'**, intramolecular cyclization to **5i** or hydrogen abstraction to **6i**.



Scheme 1: Borylation of arydiazonium tetrafluoroborates **3**. Reaction conditions: **3** (1 mmol), B_2pin_2 (1 mmol), pyridine (4 mmol), MeCN (2 mL), 2 h.




Scheme 2: Proposed reaction mechanism.

Starting from aniline derivative **1b**, a one pot diazotization–borylation sequence using different acids afforded the corresponding borylated product **2b** in good yields (Table 3).

Suzuki–Miyaura cross-coupling reactions of boronates **2a** and **2b** with aryl iodides using a simple system without any optimization proceeded in satisfactory yields considering the electron-deficient character of the boronates and consequently less efficient transmetalation step (Scheme 4).

Table 3: One pot diazotization-borylation of **1b**^a.




Entry	Acid (equiv)	Time (h)	2b , Yield (%) ^b
1 ^c	<i>p</i> -TsOH·H ₂ O (1)	15	51
2	aq HBF ₄ (1.7)	1	81
3	aq HCl (1.7)	1	78


^aReaction conditions: **1b** (1 mmol), acid, *t*-BuONO (1 mmol), MeCN (3 mL), B₂pin₂ (1.0–1.1 mmol), pyridine (4 mmol). ^bIsolated yield.

^cReaction temperature for steps 1 and 2 was rt.


Transformation to SF₅-phenylboronic acid **8b** and potassium trifluoroboroborates **9** was straightforward under standard conditions (Scheme 5). Similar potassium SF₅-phenyltrifluoroboroborates were found to be highly reactive with a variety of aryl bromides and iodides in the presence of catalytic amounts of



Scheme 3: Reaction of diazonium salt **3i** under borylation conditions.




Scheme 4: Suzuki–Miyaura reaction of boronates **2a** and **2b** with aryl iodides. Reaction conditions: **2** (1 mmol), ArI (1.1 mmol), Pd(PPh₃)₄ (5 mol %), LiOH·H₂O (4 mmol), 1,4-dioxane (2 mL), water (1 mL), 3.5 h.



Scheme 5: Synthesis of boronic acid **8b** and trifluoroborates **9**. Reaction conditions for the synthesis of **8b**: **2** (2 mmol), NaIO₄ (8 mmol), THF (8 mL), H₂O (2 mL), rt, 1 h. Reaction conditions for the synthesis of **9**: **2** (2 mmol), KHF₂ (10 mmol), MeOH (8 mL), H₂O (3.6 mL), rt, 20 min.

PdCl₂(dppf)·CH₂Cl₂ or Pd(OAc)₂ [72]. The recently published synthesis of arylboronic acids from anilines or aryl diazonium tetrafluoroborates using B₂(OH)₄ [68] applied to **3b** provided **8b** in only 25% ¹⁹F NMR yield.

To extend the synthetic utility of the pyridine-mediated derivatization of aryl diazonium tetrafluoroborates we investigated the reaction with iodobenzene and iodine as efficient scavengers of aryl radicals [83,84]. A competitive experiment starting from **3a** and equimolar amounts of B₂pin₂ and iodobenzene in the presence of pyridine (4 equiv) in MeCN afforded a mixture of **2a** (72% yield) and 1-iodo-3-(pentafluorosulfanyl)benzene (**10a**, 23% yield). Additionally, a reaction of **3b** with PhI (4 equiv) in the absence of B₂pin₂ gave 1-iodo-4-(pentafluorosulfanyl)benzene (**10b**) in 35% yield. Both experiments point to the formation of aryl radicals during the reaction and suggest a possibility to conduct practical aromatic iodination. Indeed, the iodination reaction with I₂ proved to be more efficient than with PhI (Scheme 6). In contrast to borylation, iodination with I₂ shows a higher sensitivity to electronic properties of substituents on the aromatic ring. Electron-acceptor substituted aryl diazonium compounds are excellent substrates while those with electron-



Scheme 6: Iodination of aryl diazonium tetrafluoroborates **3**. Reaction conditions: **3** (1 mmol), I₂ (1.1 mmol), pyridine (4 mmol), MeCN (3 mL), 2 h.
^aYields in the presence of collidine instead of pyridine.

donor groups react much less efficiently. The substitution of pyridine with collidine (*2,4,6*-trimethylpyridine) gave similar yields. Unlike borylations, the iodination reactions were not sensitive to *ortho* substitutions. In the case of **3i**, compound **10i** was the sole product; no product of hydrogen atom transfer or cyclization was observed, demonstrating that the reaction with I₂ is much faster than with B₂pin₂. Importantly, the yields of SF₅-phenyl iodides **10a** and **10b** using our two-step diazotization–iodination method significantly exceed those obtained by classical Sandmeyer reaction (ca. 80% yield over two steps compared to 63% for **10a** and 50% for **10b** by one-pot Sandmeyer procedure requiring 10 fold excess of KI) [14]. The side-product in the iodination of **3** was bis(pyridine)iodonium tetrafluoroborate, which can be easily isolated from the reaction mixture by precipitation upon addition of diethyl ether. This iodonium salt was first synthetized by Barluenga [85] and later used for mild iodination of alkenes, alkynes and aromatics [85–87]. Its formation can be explained by the reaction of the pyridinium tetrafluoroborate radical (Scheme 2), pyridine and I₂ or iodine radical. Bromination of **3b** with Br₂ was attempted under conditions similar to iodination but the reaction was slow and inefficient resulting in a mixture of products with expected 1-bromo-4-(pentafluorosulfanyl)benzene as a minor product. With equimolar NBS instead of bromine, the reaction is much cleaner but slow; after overnight at ambient temperature the bromo product was isolated in 35% yield.

Finally, hydrodediazonation using tributyltin hydride or THF was tested and THF proved to be a more efficient hydrogen atom donor. The addition of excess pyridine to MeCN/THF solution of diazonium tetrafluoroborates **3a** or **3b** led to an efficient hydrodediazonation and the formation of (pentafluorosulfanyl)benzene (**6**). Deuteriation experiments established that the hydrogen atom in the product comes exclusively from THF and not from pyridine or MeCN (Table 4). The observed deuterium enrichment using THF-*d*₈ was around 80%. Thermal decomposition of aryl diazonium salts prepared from immobilized triazene precursors and the formation of deuterated aromatics using THF-*d*₈ was reported [88]. We explain the reduced yield of **6-D** (48% yield) and the formation of tar products by hydrogen atom abstraction from **3** or **6** and subsequent polymerization. No significant amounts of double deuterated products were detected. The kinetic isotope effect was determined from intermolecular competition experiment using **3b** and a 1:1 mixture of THF and THF-*d*₈ giving KIE = 5.5 (**6**:**6-D** ratio determined by GC–MS) and combined yield of 62%. This means that the hydrogen abstraction is much faster than the deuterium abstraction and suggests the C-H(D) bond formation as the rate-limiting step. For unambiguous identification of the rate-limiting step, individual rate constants *k*_H and *k*_D in two parallel reactions would have to be determined [89]. Dihydro-

Table 4: Hydrodediazonation of **3a** and **3b** with THF^a.

3a (3-SF ₅)		6 (6-D)		
3b (4-SF ₅)		11		
Entry	3 (mmol)	D source	D enrichment ^b	6 , Yield (%) ^c
1	3a (0.5)	—	n/a	6 , 75
2	3b (0.5)	—	n/a	6 , 70
3	3b (0.5)	THF- <i>d</i> ₈ (0.5 mL)	77–82	6-D , 48
4	3b (0.25)	C ₅ D ₅ N (4 mmol)	0	6 , 70
5	3b (0.25)	CD ₃ CN (0.75 mL)	0	6 , 67

^aReaction conditions: **3** (0.25–0.5 mmol), pyridine (4 equiv), THF (1 mL/1 mmol of **3**), MeCN (3–4 mL/1 mmol of **3**), 2 h. ^bBased on GC–MS. ^cBased on ¹⁹F NMR using 1-nitro-4-(pentafluorosulfanyl)-benzene as an internal standard.

furan (**11**) and pyridinium tetrafluoroborate were identified as byproducts of the dediazonation reactions. Similarly to the previous processes, we presume the formation of aryl diazonium **4** and its decomposition to an aryl radical and the pyridinium tetrafluoroborate radical. The aryl radical abstracts a hydrogen atom from THF forming **6** and a THF radical. The THF radical then transfers the hydrogen atom to the pyridinium tetrafluoroborate radical giving dihydrofuran (**11**) and pyridinium salt.

Conclusion

In conclusion, a novel dediazonation–borylation methodology was developed based on the reaction of aryl diazonium tetrafluoroborates with pyridine and B₂pin₂ to give aryl pinacolborates. Particular emphasis was on the synthesis of SF₅-phenyl boronates where our methodology represents a considerable improvement in reaction efficiency compared to previously published syntheses. Furthermore, no transition metals are needed and mild reaction conditions are used. The borylation is applicable to a variety of aryl diazonium tetrafluoroborates with electron-donor or acceptor groups while *ortho*-substituted substrates are less reactive. A mechanism involving aryl radicals is suggested. The Suzuki–Miyaura reaction of SF₅-phenyl boronates with aryl iodides provided the cross-coupling biaryl products. In analogy to the borylation reaction, iodination of aryl diazonium tetrafluoroborates with pyridine and iodine resulted in aryl iodides. An efficient reaction was observed with electron-acceptor substituted aromatic compounds even with *ortho*-substituted derivatives. In the case of SF₅-substituted

iodobenzenes, the method is much more efficient than the classical Sandmeyer reaction starting from SF₅-containing aniline derivatives. Finally, hydrodediazonation of SF₅-phenyldiazonium tetrafluoroborates by hydrogen atom abstraction from THF in the presence of pyridine provided (pentafluorosulfanyl)benzene.

Supporting Information

Synthesis and characterization of all products, copies of ¹H, ¹³C, and ¹⁹F NMR spectra of newly synthesized products, and X-ray crystallographic files of the compounds **3a** and **3b**.

Supporting Information File 1

Experimental part.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-11-162-S1.pdf>]

Supporting Information File 2

Crystal structure of compound **3a**.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-11-162-S2.cif>]

Supporting Information File 3

Crystal structure of compound **3b**.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-11-162-S3.cif>]

Acknowledgement

Financial support from the Academy of Sciences of the Czech Republic (Research Plan RVO: 61388963) and the Grant Agency of the Czech Republic (P207/12/0072) is gratefully acknowledged.

References

- Clifford, A. F.; El-Shamy, H. K.; Emeleus, H. J.; Haszeldine, R. N. *J. Chem. Soc.* **1953**, 2372–2375. doi:10.1039/jr9530002372
- Hoffmann, F. W.; Simmons, T. C.; Beck, R. B.; Holler, H. V.; Katz, T.; Koshar, R. J.; Larsen, E. R.; Mulvaney, J. E.; Rogers, F. E.; Singleton, B.; Sparks, R. S. *J. Am. Chem. Soc.* **1957**, 79, 3424–3429. doi:10.1021/ja01570a029
- Dresdner, R. D.; Mao, T. J.; Young, J. A. *J. Am. Chem. Soc.* **1959**, 81, 574–577. doi:10.1021/ja01545a027
- Sheppard, W. A. *J. Am. Chem. Soc.* **1960**, 82, 4751–4752. doi:10.1021/ja01502a083
- Savoie, P. R.; Welch, J. T. *Chem. Rev.* **2015**, 115, 1130–1190. doi:10.1021/cr500336u
- Altomonte, S.; Zanda, M. *J. Fluorine Chem.* **2012**, 143, 57–93. doi:10.1016/j.jfluchem.2012.06.030
- Aït-Mohand, S.; Dolbier, W. R., Jr. *Org. Lett.* **2002**, 4, 3013–3015. doi:10.1021/o1026483o
- Dolbier, W. R., Jr.; Aït-Mohand, S.; Schertz, T. D.; Sergeeva, T. A.; Cradlebaugh, J. A.; Mitani, A.; Gard, G. L.; Winter, R. W.; Thrasher, J. S. *J. Fluorine Chem.* **2006**, 127, 1302–1310. doi:10.1016/j.jfluchem.2006.05.003
- Vida, N.; Pastýřková, T.; Klepetářová, B.; Beier, P. *J. Org. Chem.* **2014**, 79, 8906–8911. doi:10.1021/jo501562z
- Umemoto, T.; Garrick, L. M.; Saito, N. *Beilstein J. Org. Chem.* **2012**, 8, 461–471. doi:10.3762/bjoc.8.53
- Lummer, K.; Ponomarenko, M. V.; Röschenthaler, G.-V.; Bremer, M.; Beier, P. *J. Fluorine Chem.* **2014**, 157, 79–83. doi:10.1016/j.jfluchem.2013.10.009
- Kanishchev, O. S.; Dolbier, W. R., Jr. *Angew. Chem., Int. Ed.* **2015**, 127, 282–286. doi:10.1002/ange.201409990
- Bowden, R. D.; Greenhall, M. P.; Moillet, J. S.; Thomson, J. *Chem. Abstr.* **1997**, 126, 199340. The preparation of fluorinated organic compounds, WO 9705106, 13 Feb 1997.
- Bowden, R. D.; Comina, P. J.; Greenhall, M. P.; Kariuki, B. M.; Loveday, A.; Philp, D. *Tetrahedron* **2000**, 56, 3399–3408. doi:10.1016/s0040-4020(00)00184-8
- Chambers, R. D.; Spink, R. C. H. *Chem. Commun.* **1999**, 883–884. doi:10.1039/A901473J
- Sipyagin, A. M.; Bateman, C. P.; Matsev, A. V.; Waterfeld, A.; Jilek, R. E.; Key, C. D.; Szulczeński, G. J.; Thrasher, J. S. *J. Fluorine Chem.* **2014**, 167, 203–210. doi:10.1016/j.jfluchem.2014.07.031
- Sheppard, W. A. *J. Am. Chem. Soc.* **1962**, 84, 3064–3072. doi:10.1021/ja00875a006
- Crowley, P. J.; Mitchell, G.; Salmon, R.; Worthington, P. A. *Chimia* **2004**, 58, 138–142.
- Kirsch, P.; Bremer, M.; Heckmeier, M.; Tarumi, K. *Angew. Chem., Int. Ed.* **1999**, 38, 1989–1992. doi:10.1002/(SICI)1521-3773(19990712)38:13/14<1989::AID-ANIE198>3.0.CO;2-K
- Frischmuth, A.; Unsinn, A.; Groll, K.; Stadtmüller, H.; Knochel, P. *Chem. – Eur. J.* **2012**, 18, 10234–10238. doi:10.1002/chem.201201485
- Beier, P.; Pastýřková, T.; Vida, N.; Iakobson, G. *Org. Lett.* **2011**, 13, 1466–1469. doi:10.1021/o12001478
- Beier, P.; Pastýřková, T.; Iakobson, G. *J. Org. Chem.* **2011**, 76, 4781–4786. doi:10.1021/jo200618p
- Beier, P.; Pastýřková, T. *Tetrahedron Lett.* **2011**, 52, 4392–4394. doi:10.1016/j.tetlet.2011.06.011
- Pastýřková, T.; Iakobson, G.; Vida, N.; Pohl, R.; Beier, P. *Eur. J. Org. Chem.* **2012**, 2123–2126. doi:10.1002/ejoc.201200127
- Vida, N.; Beier, P. *J. Fluorine Chem.* **2012**, 143, 130–134. doi:10.1016/j.jfluchem.2012.04.001
- Iakobson, G.; Pošta, M.; Beier, P. *Synlett* **2013**, 24, 855–859. doi:10.1055/s-0032-1318452
- Wang, C.; Yu, Y.-B.; Fan, S.; Zhang, X. *Org. Lett.* **2013**, 15, 5004–5007. doi:10.1021/o14023326
- Sergeeva, T. A.; Dolbier, W. R., Jr. *Org. Lett.* **2004**, 6, 2417–2419. doi:10.1021/o10491991
- Dolbier, W. R., Jr.; Zheng, Z. *J. Fluorine Chem.* **2011**, 132, 389–393. doi:10.1016/j.jfluchem.2011.03.017
- Dolbier, W. R., Jr.; Zheng, Z. *J. Org. Chem.* **2009**, 74, 5626–5628. doi:10.1021/jo9007699
- Lee, J.-W.; List, B. *J. Am. Chem. Soc.* **2012**, 134, 18245–18248. doi:10.1021/ja3096202
- Yang, Y.-D.; Lu, X.; Tokunaga, E.; Shibata, N. *J. Fluorine Chem.* **2012**, 143, 204–209. doi:10.1016/j.jfluchem.2012.06.007

33. Chia, P. W.; Brennan, S. C.; Slawin, A. M. Z.; Riccardi, D.; O'Hagan, D. *Org. Biomol. Chem.* **2012**, *10*, 7922–7927. doi:10.1039/c2ob26402a
34. Sun, L.; Li, J.; Bera, H.; Dolzhenko, A. V.; Chiu, G. N. C.; Chui, W. K. *Eur. J. Med. Chem.* **2013**, *70*, 400–410. doi:10.1016/j.ejmech.2013.10.022
35. Stump, B.; Eberle, C.; Schweizer, W. B.; Kaiser, M.; Brun, R.; Krauth-Siegel, R. L.; Lentz, D.; Diederich, F. *ChemBioChem* **2009**, *10*, 79–83. doi:10.1002/cbic.200800565
36. Altomonte, S.; Baillie, G. L.; Ross, R. A.; Riley, J.; Zanda, M. *RSC Adv.* **2014**, *4*, 20164–20176. doi:10.1039/c4ra01212g
37. Yang, Y.-D.; Tokunaga, E.; Akiyama, H.; Saito, N.; Shibata, N. *ChemMedChem* **2014**, *9*, 913–917. doi:10.1002/cmcd.201400059
38. Nakayama, H.; Nishida, J.-i.; Takada, N.; Sato, H.; Yamashita, Y. *Chem. Mater.* **2012**, *24*, 671–676. doi:10.1021/cm202650u
39. Winter, R.; Nixon, P. G.; Gard, G. L.; Graham, D. J.; Castner, D. G.; Holcomb, N. R.; Grainger, D. W. *Langmuir* **2004**, *20*, 5776–5781. doi:10.1021/la040011w
40. Hall, D. G., Ed. *Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine*; Wiley-VCH: Weinheim, Germany, 2005.
41. Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483. doi:10.1021/cr00039a007
42. Miyaura, N. *Top. Curr. Chem.* **2002**, *219*, 11–59. doi:10.1007/3-540-45313-X
43. Brown, H. C.; Cole, T. E. *Organometallics* **1983**, *2*, 1316–1319. doi:10.1021/om50004a009
44. Brown, H. C.; Srebnik, M.; Cole, T. E. *Organometallics* **1986**, *5*, 2300–2303. doi:10.1021/om00142a020
45. Ishiyama, T.; Murata, M.; Miyaura, N. *J. Org. Chem.* **1995**, *60*, 7508–7510. doi:10.1021/jo00128a024
46. Murata, M.; Watanabe, S.; Masuda, Y. *J. Org. Chem.* **1997**, *62*, 6458–6459. doi:10.1021/jo970963p
47. Murata, M.; Oyama, T.; Watanabe, S.; Masuda, Y. *J. Org. Chem.* **2000**, *65*, 164–168. doi:10.1021/jo991337q
48. Fürstner, A.; Seidel, G. *Org. Lett.* **2002**, *4*, 541–543. doi:10.1021/o10171463
49. Zhu, W.; Ma, D. *Org. Lett.* **2006**, *8*, 261–263. doi:10.1021/o1052633u
50. Kleberg, C.; Dang, L.; Lin, Z.; Marder, T. B. *Angew. Chem., Int. Ed.* **2009**, *48*, 5350–5354. doi:10.1002/anie.200901879
51. Euzenat, L.; Horlant, D.; Ribourdouille, Y.; Duriez, C.; Alcaraz, G.; Vaultier, M. *Chem. Commun.* **2003**, 2280–2281. doi:10.1039/B306874A
52. Chen, H.; Schlecht, S.; Semple, T. C.; Hartwig, J. F. *Science* **2000**, *287*, 1995–1997. doi:10.1126/science.287.5460.1995
53. Ishiyama, T.; Takagi, J.; Ishida, K.; Miyaura, N.; Anastasi, N. R.; Hartwig, J. F. *J. Am. Chem. Soc.* **2002**, *124*, 390–391. doi:10.1021/ja0173019
54. Ishiyama, T.; Takagi, J.; Hartwig, J. F.; Miyaura, N. *Angew. Chem., Int. Ed.* **2002**, *41*, 3056–3058. doi:10.1002/1521-3773(20020816)41:16<3056::AID-ANIE3056>3.0.CO;2-#
55. Preshlock, S. M.; Plattner, D. L.; Maligres, P. E.; Krska, S. W.; Maleczka, R. E., Jr.; Smith, M. R., III. *Angew. Chem., Int. Ed.* **2013**, *52*, 12915–12919. doi:10.1002/anie.201306511
56. Dai, H.-X.; Yu, J.-Q. *J. Am. Chem. Soc.* **2012**, *134*, 134–137. doi:10.1021/ja2097095
57. Kawamorita, S.; Miyazaki, T.; Ohmiya, H.; Iwai, T.; Sawamura, M. *J. Am. Chem. Soc.* **2011**, *133*, 19310–19313. doi:10.1021/ja208364a
58. Ros, A.; Estepa, B.; López-Rodríguez, R.; Álvarez, E.; Fernández, R.; Lassaletta, J. M. *Angew. Chem., Int. Ed.* **2011**, *50*, 11724–11728. doi:10.1002/anie.201104544
59. Ingleson, M. J. *Synlett* **2012**, *23*, 1411–1415. doi:10.1055/s-0031-1291147
60. De Vries, T. S.; Prokofjevs, A.; Vedejs, E. *Chem. Rev.* **2012**, *112*, 4246–4282. doi:10.1021/cr200133c
61. Del Grosso, A.; Singleton, P. J.; Muryn, C. A.; Ingleson, M. J. *Angew. Chem., Int. Ed.* **2011**, *50*, 2102–2106. doi:10.1002/anie.201006196
62. Del Grosso, A.; Helm, M. D.; Solomon, S. A.; Caras-Quintero, D.; Ingleson, M. J. *Chem. Commun.* **2011**, *47*, 12459–12461. doi:10.1039/c1cc14226g
63. Ma, Y.; Song, C.; Jiang, W.; Xue, G.; Cannon, J. F.; Wang, X.; Andrus, M. B. *Org. Lett.* **2003**, *5*, 4635–4638. doi:10.1021/o1035857q
64. Mo, F.; Jiang, Y.; Qiu, D.; Zhang, Y.; Wang, J. *Angew. Chem., Int. Ed.* **2010**, *49*, 1846–1849. doi:10.1002/anie.200905824
65. Qiu, D.; Jin, L.; Zheng, Z.; Meng, H.; Mo, F.; Wang, X.; Zhang, Y.; Wang, J. *J. Org. Chem.* **2013**, *78*, 1923–1933. doi:10.1021/jo3018878
66. Yu, H.; Zhang, J.; Wang, X.; Ye, J. *Synlett* **2012**, *23*, 1394–1396. doi:10.1055/s-0031-1290960
67. Yu, J.; Zhang, L.; Yan, G. *Adv. Synth. Catal.* **2012**, *354*, 2625–2628. doi:10.1002/adsc.201200416
68. Erb, W.; Hellal, A.; Albini, M.; Rouden, J.; Blanchet, J. *Chem. – Eur. J.* **2014**, *20*, 6608–6612. doi:10.1002/chem.201402487
69. Marciasini, L. D.; Richy, N.; Vaultier, M.; Pucheault, M. *Adv. Synth. Catal.* **2013**, *355*, 1083–1088. doi:10.1002/adsc.201200942
70. Sherrington, J. Organoboron compound comprising sulphur pentafluoride. WO 123749 A1, Dec 29, 2005.
71. Stamford, A. W.; Cumming, J. N. Pentafluorosulfur imino heterocyclic compounds as BACE-1 inhibitors, composition, and their use. WO 044184 A1, April 14, 2011.
72. Joliton, A.; Carreira, E. M. *Org. Lett.* **2013**, *15*, 5147–5149. doi:10.1021/o14025666
73. Doyle, M. P.; Bryker, W. J. *J. Org. Chem.* **1979**, *44*, 1572–1574. doi:10.1021/jo01323a048
74. Okazaki, T.; Laali, K. K.; Bunge, S. D.; Adas, S. K. *Eur. J. Org. Chem.* **2014**, 1630–1644. doi:10.1002/ejoc.201301538
75. Okazaki, T.; Laali, K. K.; Reddy, A. S. *J. Fluorine Chem.* **2014**, *165*, 91–95. doi:10.1016/j.fluchem.2014.06.021
76. Galli, C. *Chem. Rev.* **1988**, *88*, 765–792. doi:10.1021/cr00087a004
77. Molinaro, C.; Mowat, J.; Gosselin, F.; O'Shea, P. D.; Marcoux, J.-F.; Angelaud, R.; Davies, I. W. *J. Org. Chem.* **2007**, *72*, 1856–1858. doi:10.1021/jo062483g
78. Zhao, C.-J.; Xue, D.; Jia, Z.-H.; Wang, C.; Xiao, J. *Synlett* **2014**, *25*, 1577–1584. doi:10.1055/s-0033-1339118
79. Abramovitch, R. A.; Saha, J. G. *Tetrahedron* **1965**, *21*, 3297–3303. doi:10.1016/S0040-4020(01)96951-0
80. Loewenschuss, H.; Wahl, G. H., Jr.; Zollinger, H. *Helv. Chim. Acta* **1976**, *59*, 1438–1448. doi:10.1002/hclca.19760590505
81. Sakakura, T.; Hara, M.; Tanaka, M. *J. Chem. Soc., Perkin Trans. 1* **1994**, 283–288. doi:10.1039/P19940000283
82. Köster, R.; Bellut, H.; Ziegler, E. *Angew. Chem., Int. Ed.* **1967**, *6*, 255. doi:10.1002/anie.196702551
83. Friedman, L.; Chlebowski, J. *J. Org. Chem.* **1968**, *33*, 1636–1638. doi:10.1021/jo01268a070
84. Wojnarovits, L.; LaVerne, J. A. *Radiat. Phys. Chem.* **1996**, *47*, 99–101. doi:10.1016/0969-806X(95)00089-G
85. Barluenga, J.; González, J. M.; Campos, P. J.; Asensio, G. *Angew. Chem., Int. Ed.* **1985**, *24*, 319–320. doi:10.1002/anie.198503191

86. Barluenga, J.; Rodriguez, M. A.; Campos, P. J. *J. Org. Chem.* **1990**, *55*, 3104–3106. doi:10.1021/jo00297a027
87. Barluenga, J.; Gonzalez, J. M.; Garcia-Martin, M. A.; Campos, P. J.; Asensio, G. *J. Org. Chem.* **1993**, *58*, 2058–2060. doi:10.1021/jo00060a020
88. Wanderheiden, S.; Bulat, B.; Zevaco, T.; Jung, N.; Bräse, S. *Chem. Commun.* **2011**, *47*, 9063–9065. doi:10.1039/c1cc12950c
89. Simmons, E. M.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2012**, *51*, 3066–3072. doi:10.1002/anie.201107334

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions:
(<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
doi:10.3762/bjoc.11.162