



**Greenwich Academic Literature Archive (GALA)**  
– the University of Greenwich open access repository  
<http://gala.gre.ac.uk>

---

*Citation for published version:*

Stevenson, Philip C., D'Cunha, Reju F. and Grzywacz, David (2010) Inactivation of baculovirus by isoflavonoids on chickpea (*Cicer arietinum*) leaf surfaces reduces the efficacy of nucleopolyhedrovirus against *Helicoverpa armigera*. *Journal of Chemical Ecology*, 36 (2). pp. 227-235. ISSN 0098-0331 (Print), 1573-1561 (Online) (doi:10.1007/s10886-010-9748-8)

*Publisher's version available at:*

<http://dx.doi.org/10.1007/s10886-010-9748-8>

---

**Please note that where the full text version provided on GALA is not the final published version, the version made available will be the most up-to-date full-text (post-print) version as provided by the author(s). Where possible, or if citing, it is recommended that the publisher's (definitive) version be consulted to ensure any subsequent changes to the text are noted.**

*Citation for this version held on GALA:*

Stevenson, Philip C., D'Cunha, Reju F. and Grzywacz, David (2010) Inactivation of baculovirus by isoflavonoids on chickpea (*Cicer arietinum*) leaf surfaces reduces the efficacy of nucleopolyhedrovirus against *Helicoverpa armigera*. London: Greenwich Academic Literature Archive.  
Available at: <http://gala.gre.ac.uk/3454/>

---

Contact: [gala@gre.ac.uk](mailto:gala@gre.ac.uk)

*This is the Author's Accepted Manuscript version of a paper published in JOURNAL OF CHEMICAL ECOLOGY by Springer International Publishing on 10 February 2010. The final publication is available at Springer via <http://dx.doi.org/10.1007/s10886-010-9748-8>*

1     **INACTIVATION OF BACULOVIRUS BY ISOFLAVONOIDS ON CHICKPEA**  
2           **(*Cicer arietinum*) LEAF SURFACES REDUCES THE EFFICACY OF**  
3           **NUCLEOPOLYHEDROVIRUS AGAINST *Helicoverpa armigera*.**

4  
5           PHILIP C. STEVENSON<sup>1&2</sup>, REJU F. D'CUNHA<sup>1</sup>, & DAVID GRZYWACZ<sup>1\*</sup>

6  
7           <sup>1</sup> Natural Resources Institute, University of Greenwich, Chatham, Kent, ME4 4TB, UK.

8           <sup>2</sup>Jodrell Laboratory, Royal Botanic Gardens, Kew, Surrey, TW9 3DS, UK.

9  
10          Chickpea isoflavonoids inhibit *Helicoverpa armigera* NPV

11  
12          Corresponding author; [D.Grzywacz@gre.ac.uk](mailto:D.Grzywacz@gre.ac.uk) tel. +44-1634883360

13  
14          **ABSTRACT**-Biological pesticides based on nucleopolyhedroviruses (NPVs) can provide  
15          an effective and environmentally benign alternative to synthetic chemicals. On some  
16          crops, however, the efficacy and persistence of NPVs is known to be reduced by plant  
17          specific factors. The present study investigated the efficacy of *Helicoverpa armigera*  
18          NPV (HearNPV) for control of *H. armigera* larvae and showed that chickpea reduced the  
19          infectivity of virus occlusion bodies (OBs) exposed to the leaf surface of chickpea for at

20 least one hr. The degree of inactivation was greater on chickpea than on previously  
21 reported on cotton and the mode of action is different to that of cotton. The effect was  
22 observed for larvae that consumed OBs on chickpea leaves but also occurred when OBs  
23 were removed after exposure to plants and inoculated on to artificial diet, indicating that  
24 inhibition was leaf surface related and permanent. Despite their profuse exudation from  
25 trichomes on chickpea leaves and low pH, organic acids – primarily oxalic and malic acid  
26 – caused no inhibition. When HearNPV was incubated with biochanin A and sissotrin,  
27 however, two minor constituents of chickpea leaf extracts, the OB activity was reduced  
28 significantly. These two isoflavonoids increased in concentration by up to 3 times within  
29 one hr of spraying the virus suspension onto the plants and also when spraying only  
30 carrier, indicating induction was in response to spraying and not a specific response to the  
31 HearNPV. Although inactivation by the isoflavonoids did not account completely for the  
32 level of effect recorded on whole plants this work constitutes evidence for a novel  
33 mechanism of NPV inactivation in legumes. Expanding the use of biological pesticides  
34 on legume crops will be dependent upon the development of suitable formulations for  
35 OBs to overcome plant secondary chemical effects.

36

37 **Key Words-** Baculovirus, Biopesticide, Nucleopolyhedrovirus, *Helicoverpa armigera*,  
38 Chickpea, Induced resistance, Plant leaf chemistry, Isoflavonoid.

39

40 INTRODUCTION

41

42 *Helicoverpa armigera* (Hubn.) is a major crop pest in Asia, Africa and Australasia  
43 attacking a wide range of important crops including cotton, maize, tomato, peppers,  
44 chilies, and legumes such as chickpea and pigeonpea (Gowda, 2005; King, 1994). Its  
45 status as arguably the world's most important agricultural pest can be attributed to its  
46 wide geographical and host range coupled with its ability to develop high levels of  
47 resistance to chemical insecticides (Armes et al., 1992b; Kranthi et al., 2002). The  
48 baculovirus biopesticide *Helicoverpa armigera* nucleopolyhedrovirus (HearNPV) is an  
49 ecologically benign alternative to chemical insecticides that is effective and can  
50 overcome problems of chemical insecticide resistance (Moscardi, 1999; Grzywacz et al.,  
51 2005). HearNPV is now commercially produced in Australia, Thailand, India and China  
52 for control of *H. armigera* (Buerger et al., 2007, Sun and Peng, 2007, Singhal, 2004).  
53 However, the utility of baculoviruses for insect pest management is compromised by the  
54 fact that some host plants adversely influence the severity of viral disease in insects and  
55 so reduce pest control efficacy (Felton and Duffey, 1990; Duffey et al., 1995; Hoover et  
56 al., 1998a; Cory and Hoover, 2006). It has for some time been recognized that *Heliothis*  
57 *zea* NPV, a closely related baculovirus, performed poorly on some crops such as cotton  
58 (Young and Yearian, 1974; Forschler et al., 1992), a phenomena linked to the direct  
59 action of glandular secretions in reducing the persistence of occlusion bodies (OBs) the  
60 infective stage of the virus (Young and Yearian, 1977; Ellerman and Entwistle, 1985).  
61 OBs are a protective crystalline protein matrix in which virions are embedded during  
62 transmission and in hostile environments (Hunter-Fujita et al., 1998). The maintenance  
63 of OB integrity is crucial to viral persistence outside the host and for initiating infections  
64 in new host insects. Host plant effects on biological pesticides are not restricted to

65 baculoviruses, as plants such as cotton have been shown to reduce the efficacy of other  
66 biopesticides, especially *Bacillus thuringiensis* (Kushner and Harvey, 1962; Johnson,  
67 1982; Ali et al., 2004). Inhibition of NPV infections on cotton has also been attributed to  
68 high peroxidase activity and subsequent free radical generation which was associated  
69 with an increase in the sloughing off of midgut cells that are the point of entry for the  
70 NPV virions, thereby reducing virus-induced mortality (Hoover et al., 1998a; Hoover et  
71 al., 1998b; Hoover et al., 2000). While the use of HearNPV has been shown to be  
72 effective on chickpea (Jayaraj et al., 1987; Rabindra et al., 1992; Cherry et al., 2000) field  
73 trials have indicated OB persistence and activity to be much lower on chickpea leaf  
74 surfaces than on other crops such as tomato (Rabindra et al., 1994), suggestive of some  
75 degree of adverse interaction on chickpea. Chickpea produces copious glandular  
76 secretions rich in organic acids and the leaf surface can subsequently have a very low pH  
77 (<3) (Rembold and Weigner, 1990; Stevenson and Aslam, 2006). This could make it a  
78 challenging host plant for biopesticide use because earlier work on *Lymantria dispar*  
79 NPV has shown that larvae can be less susceptible to OBs when inoculated on highly  
80 acidic (pH 3.8-4.6) oak foliage rather than other less acidic aspen foliage (Keating and  
81 Yendol, 1987) an effect associated with low pH and high levels of organic acids (Keating  
82 et al., 1989).

83         The present study was undertaken to investigate the efficacy of HearNPV on  
84 chickpea in comparison with tomato, a known favorable host (Forschler et al., 1992;  
85 Farrar et al., 2000), and cotton, a host plant known to impair OB infectivity, to better  
86 understand what plant factors affect virus efficacy with a view to developing better  
87 recommendations for the efficacy of NPV-based insecticides on legume crops and to

88 assist in the development of a suitable formulation for OBs for use on crops such as  
89 chickpea.

90

91

## MATERIALS AND METHODS

92

93       Virus. The virus strain (NRI#0210) was provided by Professor R.J. Rabindra of  
94 Tamil Nadu Agricultural University, India, and stored at -80°C. This strain is typical in  
95 activity of strains of HearNPV used in biopesticides products in India having a mean  
96  $LC_{50}$  of  $2.78 \times 10^3$  OB ml<sup>-1</sup> for neonate larvae similar to that reported by others including  
97 Somasekar et al. (1993) and had been used previously in field trials on chickpea in India  
98 (Cherry et al., 2000). It was multiplied up in third instars of *H. armigera* then harvested  
99 and purified using a standard NPV purification protocol (Hunter-Fujita et al., 1998). The  
100 virus was enumerated using a standard Neubauer haemocytometer and phase contrast  
101 microscope at X400 magnification (Wigley, 1980). The identity of the source and  
102 progeny of the virus was checked using a standard DNA restriction analysis protocol for  
103 NPVs with EcoR1 (Hunter-Fujita et al., 1998).

104       Insects. The insects for the bioassays were derived from a culture of *H. armigera*  
105 provided by the NERC Centre for Ecology and Hydrology at Oxford which had been  
106 maintained there for a number of years. The insects were reared at  $26 \pm 2$  °C with a  
107 relative humidity of  $50 \pm 5\%$  and a 14:10 hr light:dark regime. Larvae were reared in  
108 groups in 250 ml plastic pots on an artificial wheatgerm casein diet until the second instar  
109 and then individually in 30 ml plastic pots on wheatgerm diet using a method previously  
110 described (Armes et al., 1992a).

111 Plants. The plants used in the study were cotton (*Gossypium hirsutum*,) variety  
112 Ankur 651 (Ankur Seeds Ltd. Nagpur, India), chickpea (*Cicer arietinum*) variety ICC  
113 11322 provided by ICRISAT, Hyderabad, India and tomato (*Lycopersicon esculentum*)  
114 ‘Moneymaker’ variety. All were grown in plastic pots on John Innes no. 2 potting  
115 compost at  $28 \pm 2^\circ\text{C}$  in a glasshouse with a 14:10 hr light:dark cycle and a relative  
116 humidity of 60%. Plants were used at 5 weeks old. The surface area of leaves was  
117 measured using a Quantimet 520-image analyser (Leica Microsystems Cambridge Ltd.,  
118 UK). Thus, the concentration of different compounds in a sample could be equated to an  
119 area of leaf surface to ensure that insects were presented with naturally occurring  
120 concentrations during feeding bioassays. These data together with the chemical analysis  
121 were used to calculate chemical concentration of leaf extracts in terms of unit area so that  
122 surface contamination bioassays could be calibrated to match concentrations found on  
123 leaf surfaces.

124 Viral Bioassays. To assess OB activity both leaf dip and surface contamination  
125 neonate larval bioassays were used under standard larval rearing conditions,  $26^\circ\text{C}$  with a  
126 14/10 hour light dark cycle. In the leaf dip assays a standard methodology was used  
127 (Evans and Shapiro, 1997). The HearNPV stock suspensions were prepared as fivefold  
128 dilution series in 50 ml of 0.02 % Triton X-100 immediately prior to use in bioassays.  
129 The leaves were cut from the plant at the stem and dipped in the HearNPV dilutions.  
130 Control leaves were dipped in 0.02 % Triton X-100 only. After dipping, the stem of the  
131 treated leaves was mounted in molten agar in 250 ml round plastic containers, either one  
132 cotton leaf, two tomato leaves and six compound chickpea leaves were used per container  
133 ; fifty neonate larvae less than 18 hours old were used for each treatment with 25 being

134 placed in each container. Larvae were allowed to feed on the leaves for 24 h, after which  
135 they were transferred to 25 ml individual pots and reared individually on clean artificial  
136 diet, the mortality was recorded after 5 and 7 days. To ascertain OB activity separately  
137 from leaf surfaces OB treatments the mass surface contamination bioassay was employed  
138 (McKinley, 1985; Jones, 2000). Again fivefold series dilutions of OBs in distilled water  
139 were prepared and then dispensed as 75  $\mu$ l aliquots onto the surface of artificial diet in  
140 30ml plastic pots, spread evenly by tilting and left to dry. Two larvae were added to each  
141 pot, reared for 7 days under standard conditions and mortality counted on days 5 & 7.  
142 Fifty larvae were used for each treatment replicate. All assays were replicated 5-7 times  
143 with each assay including a control and a stock solution positive control and the results  
144 were subjected to probit analysis (Finney, 1971) in SPSS. Comparisons of LC<sub>50</sub> were  
145 performed on log transformed data, to equalize variances, using ANOVA procedure in  
146 SIGMASTAT software and treatment means were compared using LSD test. In some  
147 bioassays where means differed by several orders of magnitude transforming the data did  
148 not normalize variances so the non-parametric Kruskal-Wallis test with Tukey multiple  
149 comparison procedure was adopted.

150 Effect of exposure of HearNPV to cotton, tomato and chickpea leaf surfaces. To  
151 study plant surface chemistry and its effect on HearNPV, OBs suspended in distilled  
152 water were applied to the leaf surfaces on whole plants at a concentration of  $3 \times 10^7$  OB  
153  $\text{ml}^{-1}$  in 0.02% triton using a hydraulic hand sprayer and applied at a rate sufficient to  
154 evenly wet the leaves. The plants used in experiments were after application of OB  
155 maintained in the laboratory at 26°C under the 14/10 hour light dark cycle and the virus  
156 was then left on the leaves for 1 or 24 hr after which OBs were recovered using a



157 standard washing technique in water containing 0.1% sodium dodecyl sulphate for one  
158 hr (Jones 1988). The samples and the OBs concentrated by centrifugation at 2500g at  
159 5°C for 30 min (Hunter Fujita et al., 1998a). The supernatant was discarded, and the OBs  
160 were re-suspended in distilled water then stored at -20°C prior to counting and bioassay.  
161 This procedure was found to have no significant effect on the LC<sub>50</sub> of virus and recovery  
162 of OBs from leaf surfaces was ascertained to be >95%; similar to that reported by other  
163 workers using this technique (McKinley, 1985; Jones, 1988).

164           Analysis of organic acids in methanol extract of chickpea leaf surface by GC-MS.  
165 The surfaces of 50 leaves were extracted in methanol 300ml and analyzed by GC-MS.  
166 Purification of organic acid fraction was carried out according to Stumpf and Burris  
167 (1979). The residue was resuspended in pyridine (50µl) (Sigma-Aldrich) with a glutaric  
168 acid internal standard (1mg ml<sup>-1</sup>) (Sigma-Aldrich). Ten min before injection 25 µl of N,  
169 O-bis (tri-methylsilyl)-acetamide (Supelco) was added; the vial shaken and left to stand at  
170 room temperature for 5 min before injection. GC-MS was carried out on a Hewlett  
171 Packard HP6890 GC linked to an Ion detector (HP 5973 Mass Selective Detector)  
172 operated in Electron Ionisation (EI) mode. A fused silica capillary column (30 m x 0.25  
173 mm i.d., coating 0.25µm) coated with non-polar HP-5MS (5% Phenyl Methyl Siloxane,  
174 Agilent 1909 IS-433) was used with a split/splitless injector and helium as a carrier gas  
175 (0.5kg cm<sup>-2</sup>). The oven temperature was held at 60°C for 2 min and then raised to 250°C  
176 at 6°C per min. Compounds were identified by comparing EI-MS and GC retention  
177 indices with synthetic standards under the same operating conditions. A set of organic  
178 acid standards as reported to occur on chickpea leaf surfaces (Rembold and Weigner,  
179 1990) was prepared in sterile distilled water, derivatised and analysed as described above.


180 Effect of organic acids present on the chickpea leaf surface on the infectivity of  
181 OBs against *H. armigera* neonates. Organic acids (Sigma Aldrich, USA) were mixed  
182 together, at the concentration present on leaf surface as determined above, in 10 ml of  
183 sterile distilled water. A sample of HearNPV ( $1 \times 10^{10}$ OB) was added to the organic acid  
184 solution and then left in a rotator at 30 rpm for one hr. OBs were then recovered by  
185 centrifuging at 2500g for 30 minutes then re-suspended in 5 ml of distilled water and  
186 counted. Serially diluted suspensions of OBs in distilled water were bioassayed alongside  
187 a control OB suspension not exposed to the organic acids.

188 HPLC analysis of chickpea leaf extracts after spraying with OB suspension. To  
189 determine the effect of HearNPV OBs on the chickpea leaf chemistry, a suspension of  $3$   
190  $\times 10^7$  OB ml<sup>-1</sup> in 0.02% Triton was sprayed onto to the leaf surfaces of whole plants using  
191 a hydraulic hand sprayer sufficient to evenly wet the leaves. Control plants were sprayed  
192 with 0.02% Triton. The leaves were excised within 5 min or after 1, 4 or 24 h after  
193 spraying and surface extracted in methanol for 40 sec, and the extracts filtered (Whatman  
194 No. 1), and evaporated to dryness under reduced pressure. The dried extracts were  
195 redissolved in 1 ml of 100% HPLC grade methanol for analysis. Aliquots (10 ul) were  
196 injected onto a reverse-phase column (Spherisorb 5ODS analytical column, 4.6 mm i.d. x  
197 250 mm) and eluted at 1 ml/min using the gradient 90% A: 10% B at t = 0 min to 50% A:  
198 50% B at t =20 mins to 20% A: 80% B at t = 25 mins to 100% B at t = 30 mins and 90%  
199 A: 10% B at t = 37 mins (A is 2% acetic acid and B is 2% acetic acid in acetonitrile).

200 Isolation of leaf surface compounds and their effect on the activity of HearNPV  
201 OBs against *H. armigera* larvae. Compounds **1** and **2** were isolated by repetitive HPLC as  
202 described above and fractions were collected manually at approximately 22 and 29 min.

203 The combined fractions were evaporated under reduced pressure and weighed. LC-MS  
204 was carried out on a Thermo-Finnigan LC/MS/MS system consisting of a ‘Surveyor’  
205 autosampling LC system interfaced to a LCQ Classic quadrupole ion trap mass  
206 spectrometer. Chromatographic separation was performed on a 150 mm × 4.6 mm i.d. (5  
207 µm particle size) Phenomenex Luna C18 column using a linear mobile phase gradient of  
208 1 ml min<sup>-1</sup> flow rate with water (A): MeOH (B): 5% Acetic Acid in MeOH (C). Initial  
209 conditions were 80% A, 0% B and 20% C changing to 0% A, 80% B and 20% C at t = 20  
210 min and maintained at these conditions to t = 25 min. Injection volume was 10 µl and  
211 data analysis was performed using Xcalibur 1.2 software. The ion trap MS was fitted with  
212 an Atmospheric Pressure Chemical Ionisation (APCI) source operated under standard  
213 conditions; i.e. vaporiser temperature 450 °C, needle current 5 mA, heated capillary  
214 temperature 150 °C, sheath and auxiliary nitrogen gas pressure 80 and 20 psi, and the  
215 source voltages tuned for the optimal transmission of protonated rutin. The ion trap was  
216 set to monitor ions from m/z 125-1200 with collision energy of 45 %. Authentic samples  
217 of genistein, daidzein, pratensein, biochanin A and formononetin (Aldrich-Sigma) were  
218 co-chromatographed with methanol leaf extracts of chickpea leaf surface that had been  
219 sprayed with HearNPV (suspended in 0.02% Triton X-100) and indicated that **2** was  
220 biochanin A. Compound **1** had a similar UV spectrum to **2** but eluted earlier (22 min)  
221 indicating a more polar nature and suggesting a glycoside. An aliquot of **1** that had been  
222 isolated from the leaf extracts as described above was analysed by LC-MS and recorded a  
223 molecular ion signal in positive mode [M + H]<sup>+</sup> at m/e = 447 indicating the molecular  
224 weight of 446 and a molecular formula C<sub>22</sub>H<sub>22</sub>O<sub>10</sub>. Comparison of the mass spectrum  
225 with the library confirmed the structure to be biochanin A 7-O-glucoside (sissotrin) with

226 good match in the lower range ( $m/e = 100-300$ ) of the spectrum. For example, the signal  
227 observed at  $[M + H]^+$   $m/e = 285$  indicated loss of a glucose moiety  $[M - 162 + H]^+$  and  
228 corresponded to biochanin A with a base peak at  $m/e = 270$  correlating to the loss of  
229 glucose and a methyl from the methoxy at C-4' and a further fragment at  $m/e = 253$   
230 correlating to  $[M - 162 - OCH_3]^+$  with the loss of the methoxy group. Subsequent co-  
231 chromatography using an authentic standard of sissotrin from natural products collection  
232 at Royal Botanic Gardens, Kew, confirmed this identification.



233 Compounds **1** and **2** were used subsequently in bioassays to evaluate their effects on  
234 HearNPV.

235 The surface area of the leaves was measured as described above. A 200  $\mu\text{l}$  aliquot  
236 of sissotrin ( $25 \mu\text{g ml}^{-1}$ ) in methanol containing the equivalent sissotrin from 1250  $\text{mm}^2$  of  
237 chickpea leaf surface and equal to the surface area of artificial diet in a 30ml container  
238 was placed onto the diet surface and allowed to evaporate. The control diets were treated  
239 with 200  $\mu\text{l}$  methanol. HearNPV concentrations on a five-fold dilution scale were  
240 prepared in distilled water. A control dose containing only distilled water was also  
241 prepared. An aliquot of each virus concentration was dispensed in a volume of 75  $\mu\text{l}$  onto  
242 the surface of the diet and allowed to dry after which 10 neonate larvae were released into  
243 each of the 5 pots. Larvae were allowed to feed for 24 hr and then were transferred to

244 clean artificial diet pots at a rate of two per pot and reared under standard conditions and  
245 mortality recorded after 7 days. The experiment was replicated three times

246 Effect of biochanin A on the efficacy of HearNPV against *H. armigera* larvae.  
247 Biochanin A (Sigma Aldrich, USA) was diluted to 500, 250, 100 and 10 ppm in distilled  
248 water and was also tested against HearNPV. A 200 µl aliquot of biochanin A at 500, 250,  
249 100 or 10 ppm was spread over the surface of artificial diet. Control pots were treated  
250 with same amount of biochanin A. Bioassays were carried out as described above for  
251 sissotrin with 50 larvae treatment<sup>-1</sup> and the experiment was again replicated three times.

252

## 253 RESULTS

254 Effect of cotton, tomato and chickpea plants on HearNPV against *H. armigera*  
255 larvae using a leaf dip bioassay method. The leaf dip bioassay showed that exposure of  
256 HearNPV on chickpea leaf could impair HearNPV activity. The LC<sub>50</sub> values (Fig 1) for  
257 the different plants were significantly different ( $F = 14.6$ ,  $df = 2,20$ ,  $P = <0.001$ ) and the  
258 LC<sub>50</sub> for HearNPV on chickpea was of  $3.96 \times 10^4$  OB ml<sup>-1</sup> was significantly higher than  
259 that on tomato ( $2.65 \times 10^3$  OB ml<sup>-1</sup>) and cotton ( $9.36 \times 10^3$  OB ml<sup>-1</sup>). The result on  
260 tomato was not different from the mean LC<sub>50</sub> of this virus strain obtained on artificial diet  
261 which was  $2.78 \times 10^3$  OB ml<sup>-1</sup>. The bioassays of HearNPV OBs exposed to tomato,  
262 cotton and chickpea leaf surfaces also showed highly significant differences after 1 hr ( $H$   
263 = 10.851,  $df = 3$ ,  $P = 0.017$ ) and 24 hr ( $H=11.033$ ,  $df = 3$ ,  $P = 0.012$ ) (Fig 2); OBs on  
264 chickpea were markedly less infectious than OBs on tomato or cotton which did not  
265 differ significantly from the LC<sub>50</sub> of unexposed control OBs. Thus, exposure of OBs to  
266 the surface of chickpea for 1 and 24 hr resulted in inactivation even after OBs were

267 removed from the leaf surface. The LC<sub>50</sub> values of HearNPV OBs exposed to chickpea  
268 for 1 and 24 hr did not differ significantly, indicating that the observed inactivation  
269 reaches its maximum effect within one hr and exposure beyond that does not further  
270 affect OB infectivity.

271

272 Analysis of organic acids in methanol extract of chickpea leaf surface by GC-MS.  
273 The leaf surfaces of chickpea extracted with 100% methanol contained oxalic, malonic,  
274 malic, citramalic and citric acid (Fig 3). The compounds with retention times 13.47-13.48  
275 and 16.01 min were silane impurities while those at 24.80-24.81 min were sugars.  
276 Glucose-6-phosphate, oxalacetate, succinic and fumaric acids were not found in any of  
277 the solvent extracts despite having been identified earlier by Rembold et al. (1980).

278 Effect of organic acids present on the chickpea leaf surface on the efficacy of  
279 HearNPV against *H. armigera* neonates. The mean LC<sub>50</sub> values of HearNPV exposed to  
280 organic acids and for untreated HearNPV using a surface contamination bioassay system  
281 to neonates of *H. armigera* were 8.05 x 10<sup>2</sup> OB ml<sup>-1</sup> and 6.16 x 10<sup>2</sup> OB ml<sup>-1</sup> respectively  
282 and were not significantly different ( *t* = 0.484, *P* = 0.762).

283 HPLC analysis of chickpea leaf surfaces after spraying with NPV. Chickpea  
284 plants were sprayed with HearNPV in a 0.02% Triton X-100 suspension (to optimize  
285 spreading) and surface extracted in methanol within 5 min and after 1, 4 and 24 hr. After  
286 1 hr there was a more than four-fold increase in the concentration of **1** to 22 µg cm<sup>-2</sup>  
287 compared with unsprayed leaf surfaces (5 µg cm<sup>-2</sup>) in which the presence of **1** is  
288 constitutive. After 2, 4 and 24 hr the concentration of **1** was similar to pre spray quantities  
289 and remained there up to 24 hr. Analysis of control plants that were sprayed with 0.02%

290 Triton only also showed higher levels of **1** after 1hr indicating that the process of  
291 spraying in the absence of virus was itself sufficient to induce the production of this  
292 compound and was not induced by the presence of the HearNPV.

293 Effect of sissotrin on the efficacy of HearNPV against *H. armigera* larvae. The  
294 mean  $LC_{50}$  after exposure of HearNPV to sissotrin for 1 hr at a concentration equivalent  
295 to that found on the leaf surface after spraying was  $1.23 \times 10^4$  OB  $ml^{-1}$  and was  
296 significantly higher than untreated HearNPV at  $2.30 \times 10^3$  OB  $ml^{-1}$  ( $F = 44.24$ ,  $df = 1,4$ ,  $p$   
297  $= 0.003$ ). However, this increase in  $LC_{50}$  for sissotrin treated HearNPV are small  
298 compared to the  $LC_{50}$  values when HearNPV OBs were exposed to chickpea plant  
299 surface for 1 hr suggesting that sissotrin does reduce the efficacy of HearNPV but does  
300 not account for all the inhibition observed when HearNPV was applied to the leaf.

301 The mean  $LC_{50}$ s of HearNPV after exposure to different concentrations of  
302 biochanin A are shown in Fig.4. There was a significant difference ( $F = 4.16$ ,  $df = 4, 10$ ,  
303  $p = 0.031$ ) between the treatments and it was shown using least significant difference  
304 tests that mean  $LC_{50}$  values for HearNPV exposed to biochanin A were not significantly  
305 different from each other but were significantly greater than the untreated sample,  
306 indicating that biochanin A even at concentrations as low as 10 ppm. As with sissotrin,  
307 however, the effect of biochanin A does not explain fully the 5-fold increase in  $LC_{50}$  seen  
308 in HearNPV after exposure on chickpea plants suggesting that other factors must be  
309 involved.

## 310 DISCUSSION

311 This study showed that the efficacy of HearNPV OBs was inhibited considerably  
312 more on chickpea than on cotton and that the effect was caused, at least in part, by

313 surface isoflavonoids and not by organic acids. This was surprising since chickpea leaf  
314 surfaces have pH of <3 due the presence of organic acids (Rembold and Weigner, 1990),  
315 and there is a well known association between low pH with NPV inactivation (Ignoffo  
316 and Garcia, 1966). This study has also demonstrated that the inactivation of OBs on  
317 leaves is caused by their direct interaction with surface chemicals since OBs that had  
318 been exposed to the leaf surface were still inactive once removed and thus differs from  
319 the mechanism of peroxidase inactivation reported previously for cotton (Hoover et al.,  
320 1998a; 1998b.). The present work does not support an earlier proposition that the  
321 reduced efficacy of HearNPV on chickpea could be related to a slower feeding rate of *H.*  
322 *armigera* on chickpea, thus reducing the rate of OB ingestion (Rabindra et al., 1992).  
323 Sissotrin accumulated on the leaf surface at least for a short period of time after plants  
324 that were sprayed with the OB suspension in 0.02% Triton or even with the 0.02% Triton  
325 control. This indicates that the process of spraying was sufficient to induce the  
326 production of these compounds and was not induced by the presence of the HearNPV.  
327 Thus the induction of these compounds is not a specific response to the application of  
328 HearNPV but a response to either wetting or the presence of surfactant. The increased  
329 secretion of biologically active antimicrobial compounds by chickpea in response to  
330 wetting would be biologically explicable as chickpea is subject to the damaging fungal  
331 diseases such as *Botrytis* grey mould during periods of heavy dew or precipitation (Pande  
332 et al., 2005).

333 Plant chemicals have previously been shown to inhibit OB dissolution by binding  
334 irreversibly to OB structural proteins (Schultz and Keating, 1991), a mechanism that is  
335 enhanced at least for orthodihydroxy moieties in the presence of peroxidases and



336 polyphenoloxidases, particularly in damaged plant tissues (Felton and Duffey, 1990). The  
337 present data do not shed light on the mechanism by which isoflavonoids impair NPV  
338 infectivity. Further work to understand this would be useful since the inactivation  
339 mechanism reported here may impact on other biological pesticides such as Bt or  
340 entomopathogenic fungi, given that chickpea isoflavonoids are toxic to numerous  
341 organisms including viruses, bacteria, fungi and insects (Aslam et al., 2009; Getti et al.,  
342 2007; Ito et al., 2003; Simmonds and Stevenson, 2001; Stevenson and Haware, 1999;  
343 Stevenson et al., 1997). The identification of a new group of compounds affecting OBs,  
344 however, adds to the existing literature on this topic and the importance of the finding is  
345 highlighted by the LC<sub>50</sub>s of OBs exposed on leaf surfaces being 3-5 orders of magnitude  
346 greater than that reported in cotton in both the present and earlier studies (Young and  
347 Yearian, 1974; Forschler et al., 1992). It is not known if this mechanism is present or as  
348 profound in all chickpea varieties. However, selective breeding for disease resistance  
349 (Pande et al., 2005) may have resulted in varieties with more biologically active  
350 compounds and may explain the high OB inactivation reported here.

351 This study showed that HearNPV OBs were inactivated when consumed on  
352 cotton leaf material, but showed no sign of inactivation when bioassayed on diets after  
353 exposure on and then removal from cotton; a result that concurs with those of Hoover et  
354 al. (1998a; 1998b). However, there was no evidence of the OB inactivation by ionic  
355 cotton gland secretions reported previously (Ellerman and Entwistle, 1985) on Ankur  
356 651, the cotton variety tested here. This may again be explained by varietal differences in  
357 the chemistry of Ankur 651 and the Deltapine varieties studied earlier. Some Indian

358 cotton are reportedly more detrimental to OB infectivity than chickpea (Rabindra et al.,  
359 1994).

360 While sissotrin and biochanin A have a significant inactivating action, the  
361 magnitude of inactivation by these compounds did not fully account for the effects  
362 observed on leaf surface assays. Therefore, other chemicals are likely to contribute to this  
363 inactivation and further work will be required to identify these.

364 In considering the results reported here it may be surprising that HearNPV is  
365 effective as a biopesticide on chickpea (Jayaraj et al., 1987; Rabindra et al., 1989; Cherry  
366 et al., 2000; Ahmed and Chandel, 2004). However, on some crops 90% of *H. armigera*  
367 larvae killed by HearNPV sprayed onto plants acquire the infection within one hr of  
368 application (D Murray, pers. comm.). The interaction of HearNPV with chickpea may  
369 also be influenced by the variety of chickpea. Cowgill and Bhagwat (1996) for example  
370 reported a field trial in which HearNPV was more effective at killing *H. armigera* when  
371 applied to the *H. armigera* susceptible genotype (ICCC 37) of chickpea than on a *H.*  
372 *armigera* resistant genotype (ICC 506). This may have been due to differences in their  
373 chemistry since the production of isoflavonoids in chickpeas is known to vary between  
374 cultivars at least in association with resistance to plant pathogens such as *Botrytis* and  
375 *Fusarium* (Stevenson et al., 1997).

376 Additives, including milk powder, casein, molasses and Robin blue dye are  
377 reported to improve HearNPV performance on chickpea (Rabindra et al., 1989) and  
378 although it has been assumed that they improved UV stability (Rabindra and Jayaraj,  
379 1988) given the present findings, it is possible that some additives may also contribute to

380 improving OB efficacy by inhibiting chemical inactivation of OBs or by encouraging  
381 feeding and rapid viral acquisition before the OB inactivation processes have taken effect.

382

383 Acknowledgement - We gratefully acknowledge the Higher Education Funding Council  
384 for England for funding and the generous advice and assistance of Prof. David Hall and  
385 Dudley Farman in the chemical analysis.

386

## 387 REFERENCES

388

389 ALI, M. I., YOUNG, S. Y., and McNEW, R. C. 2004. Host plant Influence on activity of  
390 *Bacillus thuringiensis* Berliner against lepidopterous pests of crops. *J. Entomol.*

391 *Sci.* 39:311-317.

392 AHMED, R., and CHANDEL, S. 2004. Farmers field evaluation of IPM module against  
393 *H. armigera* infesting chickpea. *Arch. Phytopath. Plant Prot.* 37:133-137.

394 ARMES, N. J., BOND, G. S., and COOTER, R. J. 1992a. The laboratory culture and  
395 development of *Helicoverpa armigera*. *Natural Resources Institute Bulletin*, 57,  
396 Natural Resources Institute, Chatham, ME4 4TB, UK. pp 18.

397 ARMES, N. J., JADHAV, D. R., BOND G. S., and KING, A. B. S. 1992b. Insecticide  
398 resistance in *Helicoverpa armigera* in South India. *Pestic. Sci.* 34:355-364.

399 ASLAM, S. N., STEVENSON, P. C., KOKUBUN, T., and HALL, D. R. 2009.

400 Antibacterial and antifungal activity of cicerfuran and related 2-arylbenzofurans  
401 and stilbenes. *Microbiol. Res.* 164: 91-195.

402 BUERGER, P., HAUXWELL, C., and MURRAY, D. 2007. Nucleopolyhedrovirus  
403 introduction in Australia. *Virologica Sinica*. 22:173-179.

404 CHERRY, A.J., RABINDRA, R.J., PARNELL, M.A., GEETHA, N., KENNEDY, J.S.  
405 and GRZYWACZ, D. 2000. Field evaluation of *Helicoverpa armigera*  
406 nucleopolyhedrovirus formulations for control of the chickpea pod-borer, *H.*  
407 *armigera* (Hubn.), on chickpea (*Cicer arietinum* var. Shoba) in southern India.  
408 *Crop Prot.* 19:51-60.

409 CORY, J.S., and HOOVER, K. 2006. Plant-mediated effects in insect-pathogen  
410 interactions. *Trends Ecol. Evol.* 21:278-286.

411 COWGILL, S. E., and BHAGWAT, V. R. 1996. Comparison of efficacy of chemical  
412 control and *Helicoverpa* NPV for the management of *Helicoverpa armigera*  
413 (Hubner) on resistant and susceptible chickpea. *Crop Prot.* 15:241-246.

414 DUFFEY, S. S., HOOVER, K., BONNING, B., and HAMMOCK, B. D. 1995. The  
415 impact of host plant on the efficacy of baculoviruses, in R.M. ROE and R.J.  
416 KUHR (eds.). *Rev. Pest. Toxicol.* 3:137-275.

417 ELLERMAN, C. J., and ENTWISTLE, P. F. 1985. Inactivation of a nuclear  
418 polyhedrosis-virus on cotton by the substances produced by the cotton leaf surface  
419 glands. *Ann. Appl. Biol.* 106:83-92.

420 EVANS, H. F., and SHAPIRO, M. 1997. Viruses. Pp. 17-53 in L.A.Lacey (ed.). *Manual*  
421 *of techniques in Insect Pathology.* Academic Press Ltd, London, UK, .

422 FELTON, G. W., and DUFFEY, S. S. 1990. Inactivation of baculovirus by quinones  
423 formed in insect- damaged plant-tissues: *J. Chem. Ecol.* 16:1221-1236.

424 FARRAR, R. R., and RIDGWAY, R. L. 2000. Host plant effects on the activity of  
425 selected nuclear polyhedrosis viruses against corn earworm and beet armyworm  
426 (Noctuidae: Lepidoptera). Environ. Entomol. 29:108-115.

427 FINNEY, D. J. 1971. Probit Analysis, London, Cambridge University Press.

428 FORSCHLER, B. T., YOUNG, S. Y., and FELTON, G. W. 1992. Diet and the  
429 susceptibility of *Helicoverpa zea* (Noctuidae: Lepidoptera) to a nuclear  
430 polyhedrosis virus. Environ. Entomol. 21:1220-1223

431 GETTI, G. T. M., ASLAM, S. N., HUMBER, D. P., STEVENSON, P. C., and CHEKE,  
432 R. A. 2006. The effect of cicerfuran, an aryl benzofuran from *Cicer bijugum*, and  
433 related benzofurans and stilbenes on *Leishmania aethiopica*, *L. tropica* and *L.*  
434 *major*. Planta Med. 72:907-911.

435 GOWDA, C. L. L. 2005. *Helicoverpa* – The Global Problem. Pp. 1-6 in H.C. Sharma,  
436 (ed.) *Heliothis/Helicoverpa Management – Emerging Trends and Strategies for*  
437 *Future Research*. Oxford & IBH Pub. Co. Pvt. Ltd., New Delhi.

438 GRZYWACZ, D., RICHARDS, A., RABINDRA, R. J., SAXENA, H., and RUPELA, O.  
439 P. 2005. Efficacy of biopesticides and natural plant products for  
440 *Heliothis/Helicoverpa* control. Pp. 371-389 in H.C. Sharma, (ed.).  
441 *Heliothis/Helicoverpa Management – Emerging Trends and Strategies for Future*  
442 *Research*. Oxford & IBH Pub. Co. Pvt. Ltd., New Delhi, India.

443 HOOVER, K., STOUT, M. J., ALANIZ, S. A., HAMMOCK, B. D., and DUFFEY, S. S.  
444 1998a. Influence of induced plant defences in cotton and tomato on the efficacy of  
445 baculoviruses on noctuid larvae. J. Chem. Ecol. 24: 253-271.

446 HOOVER, K., YEE, J. L., SCHULTZ, C. M., ROCKE, D. M., HAMMOCK, B. D., and  
447 DUFFEY, S.S. 1998b. Effects of plant identity and chemical constituents on the  
448 efficacy of a baculovirus against *Heliothis virescens*. *J. Chem. Ecol.* 24:221-252.

449 HOOVER, K., WASHBURN J. O., and VOLKMAN L. E. 2000. Midgut-based resistance  
450 of *Heliothis virescens* to baculovirus infection mediated by phytochemicals in  
451 cotton. *J. Insect Physiol.* 6:999-1007,

452 HUNTER-FUJITA, F. R., ENTWISTLE, P. F., EVANS, H. F., and CROOK, N. E. 1998.  
453 Working with viruses. Pp. 391-454 in HUNTER FUJITA, F.R., ENTWISTLE,  
454 P.F., EVANS, H.F. and CROOK, N.E. (eds). *Insect Viruses and Pest*  
455 *Management*. Wiley, Chichester, UK.

456 IGNOFFO, C. M., and GARCIA, C. 1966. The relation of pH to the activity of inclusion  
457 bodies of a *Heliothis nuclear polyhedrosis*. *J. Invertebr. Pathol.* 8:426-427.

458 ITO, C., ITOIGAWA, M., KANEMATSU, T., RUANGRUNGSI, N., MUKAINAKA,  
459 T., TOKUDA, H., NISHINO, H., and FURUKAWA, H. 2003. Isoflavonoids from  
460 *Dalbergia olivari*. *Phytochem.* 64:1265-1268.

461 JAYARAJ, S., RABINDRA, R. J., and SANTHARAM, G. 1987. Control of *Heliothis*  
462 *armigera* (Hubner) on chickpea and lablab bean by nuclear polyhedrosis virus.  
463 *Indian J. Agric. Sci.* 57:738-741.

464 JOHNSON, D. R, 1982. Suppression of *Heliothis* spp. on cotton by using *Bacillus*  
465 *thuringiensis*, baculovirus *Heliothis* and two feeding adjuvants. *J. Econ. Entomol.*  
466 75:207-210.

467 JONES, K.A. 1988. Studies on the persistence of *Spodoptera littoralis* nuclear  
468 polyhedrosis virus on cotton in Egypt, PhD dissertation, University of Reading,  
469 UK.

470 JONES, K.A. 2000. Bioassays of entomopathogenic viruses, pp 95 -140, in A. Navon,  
471 and K.R.S.Ascher (eds.) Bioassays of Entomopathogenic Microbes and  
472 Nematodes. CAB International, Oxon, UK.

473 KEATING, S. T., and YENDOL W. G. 1987. Influence of selected host plants on gypsy-  
474 moth (*Lepidoptera*, *Lymantriidae*) larval mortality caused by a baculovirus.  
475 *Environ. Entomol.* 16:459-462.

476 KEATING, S.T., McCARTHY, W.J., and YENDOL W.G. 1989. Gypsy Moth  
477 (*Lymantria dispar*) larval susceptibility to a baculovirus by selected nutrients,  
478 hydrogen ions (pH) and plant allelochemicals in artificial diets. *J. Invertebr.*  
479 *Pathol.* 54:165-174.

480 KING, A. B. S. 1994. *Heliothis/Helicoverpa* (*Lepidoptera*: *Noctuidae*). Pp. 39-106  
481 in G.A. MATHEWS and J.P. TUNSTALL (eds.). *Insect Pests of Cotton*. CAB  
482 International, Oxon, UK.

483 KRANTHI, K. R., JADHAV, D. R., KRANTHI, S., WANJARI, R. R., ALI, S. S., and  
484 RUSSELL, D. A. 2002. Insecticide resistance in five major insect pests of cotton  
485 in India. *Crop Prot.* 21:449-460

486 KUSHNER, D. J., and HARVEY, G. T. 1962. Antibacterial substances in leaves: their  
487 possible role in insect disease *J. Invertebr. Pathol.* 4:155-184.

488 MCKINLEY, D. 1985. Nuclear Polyhedrosis virus of *Spodoptera littoralis* Boisd.  
489 (Lepidoptera, Noctuidae) as an infective agent in its host and related insects, PhD  
490 dissertation, University of London, UK.

491 MOSCARDI F. 1999. Assessment of the application of baculoviruses for control of  
492 Lepidoptera. *Ann. Rev. Entomol.* 44:257-289.

493 PANDE, S., STEVENSON, P. C., NARAYANA-RAO, J., NEUPANE, R. K.,  
494 CHAUDHARY, R. N., GRZYWACZ, D., BOURAI, V. A., and KRISHNA-  
495 KISHORE, G. 2005. Rehabilitation of chickpea in Nepal through integrated  
496 management of *Botrytis gray* mold. *Plant Disease*, 89:1252-1262

497 RABINDRA, R. J., MUTHUSWAMI, M., and JAYARAJ, S. 1994. Influence of host-  
498 plant surface environment on the virulence of nuclear polyhedrosis virus against  
499 *Helicoverpa-armigera* (Hbn) (Lep, noctuidae) larvae: *J. Appl. Entomol.* 118:453-  
500 460.

501 RABINDRA, R. J., SATHIAH, N., and JAYARAJ, S. 1992. Efficacy of nuclear  
502 polyhedrosis virus against *Heliothis armigera* (Hubner) on *Helicoverpa* resistant  
503 and susceptible varieties of chickpea. *Crop Prot.* 11:320-322.

504 RABINDRA, R. J., SATHIAH, J., MUTHAIAH, C., and JAYARAJ, S. 1989.  
505 Controlled droplet application of nuclear polyhedrosis virus with adjuvants and  
506 UV protectants for the control of *Heliothis armigera* (Hubner.) on chickpea. *J.*  
507 *Biol. Control* 3:37-39.

508 RABINDRA, R. J., and JAYARAJ, S. 1988. Evaluation of certain adjuvants for nuclear  
509 polyhedrosis virus (NPV) of *Heliothis armigera* (Hbn) on chickpea. *Indian J.*  
510 *Exp. Biol.* 26:60-62.



511 REMBOLD, H., and WEIGNER, C. 1990. Chemical composition of chickpea, *Cicer*  
512 *arietinum*, exudate. *Z. Naturforsch.* 45:922-923.

513 SCHULTZ, J. C., and KEATING S. T. 1991. Host plant-mediated interactions between  
514 the gypsy moth and a baculovirus. Pp. 489-553 in P. BARBOSA. V.A.  
515 KRISCHIK and C.G. JONES (eds). *Microbial Mediation of Plant-Herbivore*  
516 *Interaction.* John Wiley and Sons, New York, USA.

517 SHANOWER, T. G., ROMEIS, J., and MINJA, E. M.,1999. Insects pests of pigeonpea  
518 and their management. *Ann. Rev. Entomol.* 44:77-96.

519 SINGHAL, V. 2004. Biopesticides in India. Pp. 31-39 in N. KAUSHIK (ed).  
520 *Biopesticides for Sustainable Agriculture, Prospects and Constraints.* TERI press,  
521 Delhi, India.

522 SIMMONDS, M. S. J., and STEVENSON, P. C. 2001. Effects of isoflavonoids from  
523 *Cicer* on larvae of *Helicoverpa armigera*. *J. Chem. Ecol.* 27:965-977.

524 SOMASEKAR, S., JAYAPRAGASAM, M., and RABINDRA, R.J. 1993.  
525 Characterisation of five Indian isolates of the nuclear polyhedrosis virus of  
526 *Helicoverpa armigera*. *Phytoparasitica* 21:333-337.

527 STEVENSON, P. C., TURNER, H., and HAWARE, M. P. 1997. Phytoalexin  
528 accumulation in roots of chickpea seedlings (*Cicer arietinum* L.) associated with  
529 resistance to *Fusarium* wilt caused by *Fusarium oxysporum* f.sp. *ciceri*. *Physiol.*  
530 *Mol. Plant. Pathol.* 50:167-178.

531 STEVENSON, P. C., and HAWARE, M. P. 1999. Maackiain in *Cicer bijugum* Rech. f.  
532 associated with resistance to *Botrytis* grey mould *Biochem. Sys. Ecol.* 27:761-767.

533 STEVENSON, P.C. and ASLAM, S.N. 2006. The Chemistry of the Genus *Cicer* L. Pp.  
534 905-956 in ATTA-UR-RAHMAN (ed). Studies in Natural Products Chemistry  
535 Vol. 33. Bioactive Natural Products (Part M). Elsevier, Amsterdam.

536 STUMPF, D. K., and BURRIS, R. H. 1979. A micro-method for the purification and  
537 quantification of organic acids of the tricarboxylic acid cycle in plant tissues.  
538 *Anal. Biochem.* 95:311-315.

539 SUN, X., and PENG, H. 2007. Recent Advances in control of insect pests by using  
540 viruses in China. *Virologica Sinica* 22:158-162.

541 WIGLEY, P. J. 1980. Counting micro-organisms. Pp. 9-35 in J. KALMAKOFF and J.F.  
542 LONGWORTH (eds.). *Microbial Control of Insect Pests: New Zealand*  
543 Department of Science and Industrial Research, Bulletin 228, Wellington, New  
544 Zealand.

545 YOUNG, S.Y., and YEARIAN, W.C. 1974. Persistence of *Heliothis* NPV on foliage of  
546 cotton soybean and tomato. *Environ. Entomol.* 3:253-255.

547 YOUNG S. Y., and YEARIAN, W. C., 1977. Effect of dew from cotton and soybean  
548 foliage on activity of *Heliothis* nuclear polyhedrosis virus. *J. Invertebr. Pathol.*  
549 29:105-111.

550