

International Journal of Mathematics and Statistics Studies

Vol.2, No.1, pp. 1-13, March 2014

Published by European Centre for Research Training and Development UK ([www.ea-journals.org](http://www.ea-journals.org))

## ON A SURVEY OF UNIFORM INTEGRABILITY OF SEQUENCES OF RANDOM VARIABLES

S. A. Adeosun and S. O. Edeki

<sup>1</sup>Department of Mathematical Sciences, Crescent University, Abeokuta, Nigeria.

<sup>2</sup>Department of Industrial Mathematics, Covenant University, Cannanland, Ota, Nigeria.

---

**ABSTRACT:** *This paper presents explicitly a survey of uniformly integrable sequences of random variables. We also study extensively several cases and conditions required for uniform integrability, with the establishment of some new conditions needed for the generalization of the earlier results obtained by many scholars and researchers, noting the links between uniform integrability and pointwise convergence of a class of polynomial functions on conditional based.*

**KEYWORDS:** Uniform Integrability, Sequences, Boundedness, Convergence, Monotonicity.

---

### INTRODUCTION

Uniform integrability is an important concept in functional analysis, real analysis, measure theory, probability theory, and plays a central role in the area of limit theorems in probability theory and martingale theory. Conditions of independence and identical distribution of random variables are basic in historic results due to Bernoulli, Borel and A.N. Kolmogorov[1]. Since then, serious attempts have been made to relax these strong conditions; for example, independence has been relaxed to pairwise independence.

In order to relax the identical distribution condition, several other conditions have been considered, such as stochastic domination by an integrable random variable or uniform integrability in the case of weak law of large number. Landers and Rogge [2] prove that the uniform integrability condition is sufficient for a sequence of pairwise independence random variables in verifying the weak law of large numbers.

Chandra [3] obtains the weak law of large numbers under a new condition which is weaker than uniform integrability: the condition of Cesàro uniform integrability. Cabrera [4], by studying the weak convergence for weighted sums of variables introduces the condition of uniform integrability concerning the weights, which is weaker than uniform integrability, and leads to Cesàro uniform integrability as a particular case. Under this condition, a weak law of large

numbers for weighted sums of pairwise independent random variables is obtained; this condition of pairwise independence can also be dropped at the price of slightly strengthening the conditions of the weights.

Chandra and Goswami [5] introduce the condition of Cesàro  $\alpha$ -integrability ( $\alpha > 0$ ), and show that Cesàro  $\alpha$ -integrability for any  $\alpha > 0$  is weaker than uniform integrability. Under the Cesàro  $\alpha$ -integrability condition for some  $\alpha > 1/2$ , they obtain the weak law of large numbers for sequences of pairwise independence random variables. They also prove that Cesàro  $\alpha$ -integrability for appropriate  $\alpha$  is also sufficient for the weak law of large numbers to hold for certain special dependent sequences of random variables and h-integrability which is weaker than all these was later introduced by Cabrera [4].

As an application, the notion of uniform integrability plays central role in establishing weak law of large number. The new condition; Cesàro uniform integrability, introduced by Chandra [3] can be used in cases to prove  $L^p$ -convergence of sequences of pairwise independent random variables, and in the study of convergence of Martingales. Other areas of application include the approximation of Green's functions of some degenerate elliptic operators as shown by Mohammed [6].

**Definition 1** The random variables  $X_1, X_2, X_3, \dots, X_n$  are independent if:

$$\mathbb{P}\left(\bigcap_{j=1}^n \{X_j \in A_j\}\right) = \prod_{j=1}^n \mathbb{P}(X_j \in A_j)$$

$\forall n$  - tuples of measurable sets  $(A_1, A_2, A_3, \dots, A_n)$ . A family of random variables  $\{X_i\}_{i \in J}$  is

independent if for each finite set  $J \subset I$ , the family  $\{X_i\}_{i \in J}$  is independent.

**Definition 2** A family of random variables  $\{X_i\}_{i \in J}$  is pairwise independent if  $X_i$  and  $X_j$  are independent whenever  $i \neq j$ .

**Definition 3** Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space. A sequence of random variables  $\{X_n\}$  defined on  $(\Omega, \mathcal{F}, \mathbb{P})$  is said to converge in probability to a random variable  $X$  if for every  $\epsilon > 0$ ,  $\mathbb{P}(|X_n - X| > \epsilon) \rightarrow 0$  as  $n \rightarrow \infty$ . This is expressed as :  $X_n \xrightarrow{P} X$

**Definition 4.** A sequence of random variables  $\{X_n\}$  defined on  $(\Omega, \mathcal{F}, \mathbb{P})$  is said to converge in  $L^p$  to a random variable  $X$  if  $X_n \in L^p$  for all  $n$ ,  $X \in L^p$ , and  $\mathbb{E}|X_n - X|^p \rightarrow 0$  as  $n \rightarrow \infty$ . This is expressed as :  $X_n \xrightarrow{L^p} X$

**Definition 5.** Sequence of real valued random variables  $\{X_n : n \in \mathbb{N}\}$  is uniformly integrable if and only if, for any  $\epsilon > 0$ ,  $\exists K > 0$  such that

$$\mathbb{E}[|X_n| I_{(|X_n| > K)}] \leq \epsilon$$

where  $I_{(|X_n| > K)}$  is an indicator function of the event  $\{|X_n| > K\}$  i.e, the function which is equal one for  $\{|X_n| > K\}$  and zero otherwise, and  $\mathbb{E}[\cdot]$  is an expectation operator. Expectation values are given by integrals for continuous random variables.

## NOTION OF USEFUL INEQUALITIES AND RESULTS OF UNIFORM INTEGRABILITY

In this section, we introduce the basic inequalities and results needed for the conditions and applicability of uniformly integrable sequences of random variables.

### Markov Inequality

If  $Y$  is a random variable such that  $\mathbb{E}|Y|^p < \infty$  for some positive real number  $p$  which may or may not be a whole number, then for any  $\epsilon > 0$ ,

$$\mathbb{P}\{|Y| \geq \epsilon\} \leq \epsilon^{-p} \mathbb{E}|Y|^p.$$

### Chebyshev's Inequality

Let  $X$  be a random variable with finite mean  $\mu$  and finite variance  $\sigma^2$ . Then

$$\mathbb{P}\{|X - \mu| \geq \epsilon\} \leq \frac{\mathbb{E}|X - \mu|^2}{\epsilon^2}$$

This follows immediately by putting  $Y = X - \mu$  and  $p = 2$  in the Markov's inequality above.

### Chebyshev's Sum Inequality

If  $a_1 \geq a_2 \geq a_3 \geq a_4 \geq a_5 \geq \dots \geq a_n$  and  $b_1 \geq b_2 \geq b_3 \geq b_4 \geq b_5 \geq \dots \geq b_n$ , then

$$n \sum_{k=1}^n a_k b_k \geq \left( \sum_{k=1}^n a_k \right) \left( \sum_{k=1}^n b_k \right)$$

Proof: Assume  $a_1 \geq a_2 \geq a_3 \geq a_4 \geq \dots \geq a_n$  and  $b_1 \geq b_2 \geq b_3 \geq b_4 \geq \dots \geq b_n$ , then by rearrangement of inequality, we have that:

$$a_1 b_1 + \dots + a_n b_n = a_1 b_1 + a_2 b_2 + \dots + a_n b_n$$

$$a_1 b_1 + \dots + a_n b_n \geq a_1 b_2 + a_2 b_3 + \dots + a_n b_1$$

$$a_1 b_1 + \dots + a_n b_n \geq a_1 b_3 + a_2 b_4 + \dots + a_n b_2$$

$$a_1 b_1 + \dots + a_n b_n \geq a_1 b_n + a_2 b_1 + \dots + a_n b_{n-1}$$

Now adding these  $n$  inequalities gives:

$$n(a_1 b_1 + \dots + a_n b_n) \geq (a_1 + a_2 + \dots + a_n)(b_1 + b_2 + \dots + b_n)$$

Hence,  $n \sum_{k=1}^n a_k b_k \geq \left( \sum_{k=1}^n a_k \right) \left( \sum_{k=1}^n b_k \right)$  □

## RESULTS OF UNIFORM INTEGRABILITY

**Lemma 2.1** Uniform integrability implies  $L^1$ - boundedness

Let  $\{X_n: n \in \mathbb{N}\}$  be uniformly integrable. Then  $\{X_n: n \in \mathbb{N}\}$  is  $L^1$ - bounded.

Proof: Choose  $K$  so large such that  $\sup \mathbb{E}[|X_n|; \{|X_n| > K\}] \leq 1$ .

Then  $\mathbb{E}|X_n| = \mathbb{E}[|X_n|; \{|X_n| \leq K\}] + \mathbb{E}[|X_n|; \{|X_n| > K\}] \leq K + 1 \quad \square$

**Remark:** The converse of Lemma 2.1 is not true, i.e boundedness in  $L^1$  is not enough for uniform integrability. For a counter example, we present the following:

Let  $U$  be uniformly distributed on  $[0, 1]$  such that:

$$X_n(u) = \begin{cases} n, & \text{for } u \leq \frac{1}{n} \\ 0, & \text{otherwise} \end{cases}$$

Then  $\mathbb{E}[X_n] = n \frac{1}{n} = 1$ , so  $\{X_n\}$  is  $L^1$ -bounded. But for  $K > 0$ ,

$\mathbb{E}[|X_n| I_{\{|X_n| > K\}}] = 1, \forall n \geq K$ , so  $(|X_n|)_{n \in \mathbb{N}}$  is not uniformly integrable.

**Theorem 2.2** Let  $X$  be a random variable and  $\{X_n: n \in \mathbb{N}\}$  be a sequence of random variables. Then the following are equivalent:

- i)  $X_n \in L^1 \forall n, X \in L^1$  and  $X_n \rightarrow X$  in  $L^1$
- ii)  $\{X_n: n \in \mathbb{N}\}$  is uniformly integrable and  $X_n \rightarrow X$  in probability.

Proof: Suppose i) holds. By Chebychev's inequality, for  $\epsilon > 0$ ,

$$\mathbb{P}\{|X_n - X| > \epsilon\} \leq \epsilon^{-1} \mathbb{E}(|X_n - X|) \rightarrow 0$$

So  $X_n \rightarrow X$  in probability. Moreover, given  $\epsilon > 0$ , there exists  $N$  such that  $\mathbb{E}(|X_n - X|) < \frac{\epsilon}{2}$

whenever  $n \geq N$ . Then we can find  $\delta > 0$  so that  $\mathbb{P}(A) \leq \delta$  implies  $\mathbb{E}(|X| 1_A) < \frac{\epsilon}{2}$ ,

$\mathbb{E}(|X| 1_A) < \epsilon, n = 1, 2, 3, 4, \dots, N$ . Then for  $n \geq N$  and  $\mathbb{P}(A) \leq \delta$ ,

$$\mathbb{E}(|X|1_A) \leq \mathbb{E}(|X_n - X|) + \mathbb{E}(|X|1_A) \leq \epsilon.$$

Hence,  $\{X_n: n \in \mathbb{N}\}$  is uniformly integrable. We have shown that i) implies ii).

Suppose on the other hand that ii) holds, then there is a subsequence  $(n_k)$  such that  $X_{n_k} \rightarrow X$  almost surely. So, by Fatou's Lemma,

$$\mathbb{E}(|X|) \leq \liminf_k \mathbb{E}(|X_{n_k}|) < \infty.$$

Now, given  $\epsilon > 0$ , there exists  $K < \infty$  such that, for all  $n$ ,

$$\mathbb{E}(|X_n|1_{(|X_n| \geq K)}) < \frac{\epsilon}{3}, \quad \mathbb{E}(|X|1_{(|X| \geq K)}) < \frac{\epsilon}{3}.$$

Consider the uniformly bounded sequence  $X_n^K = (-K) \wedge X_n \vee K$  and set  $X^K = (-K) \wedge X \vee K$ .

Then  $X_n^K \rightarrow X^K$  in probability, so by bounded convergence, there exists  $N$  such that, for all

$$n \geq N, \quad \mathbb{E}(|X_n^K - X^K|) < \frac{\epsilon}{3}. \text{ But then, for } n \geq N,$$

$$\mathbb{E}(|X_n - X|) \leq \mathbb{E}(|X_n|1_{(|X_n| \geq K)}) + \mathbb{E}(|X_n^K - X^K|) + \mathbb{E}(|X|1_{(|X| \geq K)}) < \epsilon.$$

Therefore, ii) implies i) since  $\epsilon > 0$  was arbitrary  $\square$

Other authors like [7], [8], [9] have also shown that uniform integrability of functions were related to the sums of random variables.

## UNIFORM INTEGRABILITY OF A CLASS OF POLYNOMIALS ON A UNIT INTERVAL

Let  $\mathbb{P}_{n,K}(X): [0,1] \rightarrow [0,1]$  denote the probability of exactly  $k$  successes in an  $n$  independent Bernoulli trials with probability  $x$  success by any trial. In other words:

$$\mathbb{P}_{n,K}(X) = \mathbb{P}_r\{b(n, x) = K\} = \binom{n}{k} x^k (1-x)^{n-k}$$

and, for integers  $r, s \geq 1$ , we define the family of functions  $\{S_{n,r,s}\}_{n=1}^{\infty}$  by

$$S_{n,r,s}(x) = \sqrt{n} \sum_{k=0}^n \mathbb{P}_{rn,rk}(x) \mathbb{P}_{sn,sk}(x) \quad (1)$$

The family of polynomials arise in the context of statistical density estimation based on Bernstein polynomials. Specifically, the case  $r = s = 1$  has been considered by many authors [10], [11], & [12] while the case  $r = 1$  and  $s = 2$  was considered by [13]. These same authors have considered issues linked to uniform integrability and pointwise convergence of  $\{S_{n,1,1}\}$  and  $\{S_{n,1,2}\}$ . However, the generalization to any  $r, s \geq 1$  has not been considered. In this section, we will establish the following results.

**Theorem 3.1** Let  $r, s$  be fixed positive integers. Then

- i)  $0 \leq S_{n,r,s}(x) \leq \sqrt{n}$  for  $x \in [0,1]$  and  $S_{n,r,s}(0) = S_{n,r,s}(1) = \sqrt{n}$
- ii)  $\{S_{n,r,s}\}_{n=0}^{\infty}$  is uniformly integrable on  $[0,1]$ .
- iii)  $S_{n,r,s}(x) \rightarrow \frac{\text{gcd}(r,s)}{\sqrt{rs(r+s)2\pi x(1-x)}}$  for  $x \in (0,1)$  as  $n \rightarrow \infty$ .

For the case  $r = s = 1$ , Babu et al [10, Lemma 3.1] contains the proof of iii). Leblanc et al [13, Lemma 3.1] considered when  $r = 1$  and  $s = 2$ . The proof here generalizes (but follows the same line as) these previous results.

In establishing Theorem 3.1, we first show that for all  $0 \leq K \leq n$  and  $x \in [0,1]$ ,

$$\mathbb{P}_{n,r}(x) \geq \mathbb{P}_{2n,2r}(x) \geq \mathbb{P}_{3n,3r}(x) \geq \dots \quad (2)$$

The proof of this inequality is based on a class of completely monotonic functions and hence of general interest [14]. Using completely different methods, Leblanc and Johnson [13] previously showed that  $\{\mathbb{P}_{2^j n, 2^j k}(x)\}_{j=0}^{\infty}$  is decreasing in  $j$  and hence, (2) is a generalization of the earlier result.

**Lemma 3.1** Let  $\{a_k\}_{k=1}^m$  and  $\{b_k\}_{k=1}^m$  be real numbers such that  $a_1 \geq a_2 \geq \dots \geq a_m$  and  $b_1 \geq b_2 \geq \dots \geq b_m$  and let  $\varphi$  denote the digamma function. Define

$$\Phi_{\delta}(x) := \sum_{k=1}^m a_k \varphi(b_k x + \delta), \quad x > 0, \delta \geq 0.$$

If  $\delta \geq \frac{1}{2}$  and  $\sum_{k=1}^m a_k \geq 0$ , then  $\Phi'_{\delta}$  is completely monotonic on  $(0, \infty)$  and hence  $\Phi_{\delta}$  is increasing and concave on  $(0, \infty)$ , see [14] & [15].

**Proof:** Let  $x > 0$ , and  $\delta \geq \frac{1}{2}$ . Then the integral representation of  $\varphi^{(n)}(x)$  is:

$$\varphi^{(n)}(x) = (-1)^{n+1} \int_0^{\infty} \frac{t^n e^{-xt}}{1 - e^{-t}} dt, \quad n = 1, 2, \dots$$

Therefore, for  $n = 1, 2, \dots$ ,

$$\begin{aligned} (-1)^{n+1} \Phi^{(n)}_{\delta}(x) &= (-1)^{n+1} \sum_{k=1}^m a_k b_k^n \varphi^{(n)}(b_k x + \delta) \\ &= \sum_{k=1}^m a_k \int_0^{\infty} \frac{(b_k t)^n e^{-x b_k t}}{e^{-t} (1 - e^{-t})} dt \end{aligned} \quad (3)$$

The assumption  $u = b_k t$  yields

$$(-1)^{n+1} \Phi^{(n)}_{\delta}(x) = \int_0^{\infty} u^{n-1} e^{-ux} \sum_{k=1}^m a_k \eta\left(\frac{u}{b_k}\right) du. \quad (4)$$



where  $\eta(x) = xe^{-\delta x}(1 - e^{-x})^{-1} > 0$ . Calculus shows that, for  $\delta \geq \frac{1}{2}$ ,  $\eta$  is strictly decreasing

on  $(0, \infty)$  and hence, for every  $u > 0$ ,  $\left\{\eta\left(\frac{u}{b_k}\right)\right\}_{k=1}^m$  is decreasing [we note that, if  $b_k = 0$ ,

there is no difficulty in taking  $\eta\left(\frac{u}{b_k}\right) = \eta(\infty) = \lim_{x \rightarrow \infty} \eta(x) = 0$ , since these terms vanish in

(3)]. Since  $\{a_k\}_{k=1}^m$  is also decreasing, Chebyshev's inequality for sums yields:

$$\sum_{k=1}^m a_k \eta\left(\frac{u}{b_k}\right) \geq \frac{1}{m} \left(\sum_{k=1}^m a_k\right) \left(\sum_{k=1}^m \eta\left(\frac{u}{b_k}\right)\right)$$

We see that, if  $\sum_{k=1}^m a_k \geq 0$ , the integrand in (4) is non-negative and hence  $(-1)^{n+1} \phi^{(n)}_{\delta} \geq 0$

on  $(0, \infty)$ . We conclude that  $\phi'_{\delta}$  is completely monotonic on  $(0, \infty)$  and, in particular,  $\phi_{\delta}$  is

increasing and concave on  $(0, \infty)$  whenever  $\delta \geq \frac{1}{2}$  and  $\sum_{k=1}^m a_k \geq 0$ . □

**Lemma 3.2 [14]** Let  $n, k, j$  be integers such that  $0 \leq k \leq n$  and  $j \geq 1$  and define

$$Q_{n,k}(j) = \frac{\binom{(j-1)n}{(j-1)k}}{\binom{jn}{jk}} = \frac{\Gamma((j-1)n+1)\Gamma(j(n-k)+1)}{\Gamma(jn+1)\Gamma((j-1)k+1)\Gamma((j-1)(n-k)+1)} .$$

Then  $Q_{n,k}(j)$  is decreasing in  $j$  and

$$\lim_{j \rightarrow \infty} Q_{n,k}(j) = \left(\frac{k}{n}\right)^k \left(\frac{n-k}{n}\right)^{n-k} .$$

**Proof:** The limit is easily verified using Stirling's formula, thus we need only show that  $Q_{n,k}(j)$  is decreasing in  $j$ . Treating  $Q_{n,k}(j)$  as a continuous function in  $j$  and differentiating we obtain:

$$Q'_{n,k}(j) = Q_{n,k}(j) \{k(q_j(k) - q_j(n)) + (n-k)(q_j(n-k) - q_j(n))\}$$

where  $q_j(x) = \varphi(jx + 1) - \varphi(jx - x + 1)$ . Now, taking  $\delta = 1$ ,  $a_1 = 1$ ,  $a_2 = -1$ , and  $b_1 = j$ , and  $b_2 = j - 1$  in Lemma 3.1, we have that  $q_j(x)$  is increasing on  $(0, \infty)$  and hence  $Q'_{n,k}(j) \leq 0$  for all  $j \geq 1$  since  $Q_{n,k}(j) > 0$  always  $\square$

**Corollary 3.1** Let  $0 \leq k \leq n$ . Then  $\{P_{jn,jk}(x)\}_{j=1}^{\infty}$  is decreasing in  $j$  for every fixed  $x \in [0,1]$ .

**Proof:**  $P_{(j-1)n,(j-1)k}(x) \geq P_{jn,jk}(x)$  if and only if,  $Q_{n,k}(j) \geq x^k(1-x)^{n-k}$  and we have, by Lemma 3.2.

$$Q_{n,k}(j) \geq \left(\frac{k}{n}\right)^k \left(\frac{1-k}{n}\right)^{n-k} = \sup_{x \in [0,1]} x^k(1-x)^{n-k}$$

which completes the proof.

**Proof of Theorem 3.1.** First we note that (i) holds since

$$\begin{aligned} \sum_{k=0}^n P_{rn,rk}(x) P_{sn,sk}(x) &\leq \sum_{k=0}^n P_{rn,rk}(x) \\ &\leq \sum_{k=0}^{rn} P_{rn,k}(x) \\ &= P_{rn,0}(x) + P_{rn,1}(x) + P_{rn,2}(x) + \cdots + P_{rn,rn}(x) \\ &= \binom{rn}{0} x^0 (1-x)^{rn-0} + \cdots + \binom{rn}{rn} x^{rn} (1-x)^0 \\ &= [x + (1-x)]^{rn} = 1 \end{aligned}$$

with equality if and only if  $x = 0,1$ . Similarly, (ii) holds since  $\{S_{n,1,1}\}_{n=1}^{\infty}$  is uniformly integrable on  $[0,1]$  and, by Corollary 3.1, we have  $S_{n,r,s}(x) \leq S_{n,1,1}(x)$  for all  $x \in [0,1]$ .

To prove (iii), let  $U_1, U_2, U_3, \dots, U_n$  and  $V_1, V_2, V_3, \dots, V_n$  be two sequences of independent random variables such that  $U_i$  is Binomial  $(r, x)$  and  $V_i$  is Binomial  $(s, x)$ . Now, define

$W_i = r^{-1}U_i - s^{-1}V_i$  so that  $W_i$  has a lattice distribution with span  $\text{gcd}(r, s)/rs$  [17]. We can write  $S_{n,r,s}(x)$  in terms of the  $W_i$  as:

$$\frac{S_{n,r,s}(x)}{\sqrt{n}} = \sum_{k=0}^n P_{rn,rk}(x)P_{sn,sk}(x) = \mathbb{P}\left(\sum_{i=1}^n \frac{U_i}{r} = \sum_{i=1}^n \frac{V_i}{s}\right) = \mathbb{P}(\sum_{i=1}^n W_i = 0).$$

Now, define the standardized variable  $W^*_i = W_i\sqrt{rs}/\sqrt{(r+s)x(1-x)}$  so that  $\text{Var}(W^*_i) = 1$

and note that these also have a lattice distribution, but with span  $\text{gcd}(r, s)/\sqrt{rs(r+s)x(1-x)}$

. Theorem 3 of Section XV.5 of Feller [16] leads to

$$\lim_{n \rightarrow \infty} \frac{S_{n,r,s}(x)}{\sqrt{n}} = \lim_{n \rightarrow \infty} \mathbb{P}\left(\frac{1}{\sqrt{n}} \sum_{i=1}^n W^*_i = 0\right) = \frac{\text{gcd}(r,s)\phi(0)}{\sqrt{nr s(r+s)x(1-x)}},$$

where  $\phi$  corresponds to the standard normal probability density function. The result now

follows from the fact that  $\phi(0) = \frac{1}{\sqrt{2\pi}}$  □

**Remark 3.1** Since  $Q_{n,k}(j)$  is decreasing, it is obvious that:

$$\binom{(j-1)n}{(j-1)k} \binom{(j+1)n}{(j+1)k} \geq \binom{jn}{jk}^2$$

And since  $Q_{n,k}(j-m+1) \geq Q_{n,k}(j+m)$  for  $m = 1, 2, 3, \dots, j$ . we see that the sequence

$\{A_m\}_{m=1}^j$  define by:

$$A_m = \binom{(j+m)n}{(j+m)k} \binom{(j-m)n}{(j-m)k} \tag{5}$$

is increasing.

Also, Corollary 3.1 trivially leads to a similar family of inequalities for “number of failure”-negative binomial probabilities. As such, let  $H_{n,k}$  be the probability of exactly  $n$  failures ( $n \geq 0$ ) before the  $k$ th success ( $k \geq 1$ ) in a sequence of i.i.d. Bernoulli trials with success probability  $p \in [0,1]$  so that, for  $j = 1,2,3, \dots$ ,

$$H_{jn,jk} = \binom{jn+jk-1}{jk-1} p^{jk} (1-p)^{jn} = \frac{k}{n+k} P_{j(n+k),jk} \quad .$$

Hence, as a direct consequence of Corollary 3.1, we have that  $\{H_{jn,jk}\}_{j=1}^{\infty}$  is also decreasing.

As a consequence of Theorem 3.1, we have, for any function  $f$  bounded on  $[0,1]$ ,

$$\lim_{n \rightarrow \infty} \int_0^1 S_{n,r,s}(x) f(x) dx = \frac{\gcd(r,s)}{\sqrt{rs(r+s)}} \int_0^1 \frac{f(x)}{\sqrt{2\pi(1-x)}} dx \quad (6)$$

In particular, Kakizawa [12], establish (6) for the case  $r = s = 1$

## CONCLUSION

Generally, we conclude by pointing out the usefulness of the results to some other interesting areas such as combinatorial and discrete probability inequalities in terms of monotonicity. In addition, the consequence of Theorem 3.1 is a key tool in assessing the performance of nonparametric density estimators based on Bernstein polynomials.

The work complements previous results in the literature with significance in computational analysis and in applied probability. Also, the result is of special interest in the study of uniform integrability of martingales in terms of pointwise boundedness, and equicontinuity of a certain class of functions.

## REFERENCES

- [1] O.O.Ugbebor, Probability distribution and Elementary limit Theorems, University of Ibadan External studies programme (1991), ISBN 978-2828-82-3
- [2] D. Landers and L. Rogge, “Laws of large numbers for pairwise independent uniformly integrable random variables”, *Math. Nachr.* 130 (1987), 189 - 192.

- [3] T.K. Chandra, “Uniform integrability in the Cesàro sense and the weak law of large numbers”, *Sankhya Ser. A* 51 (3), (1989) 309- 317.
- [4] M. O. Cabrera, “Convergence of weighted sums of random variable and uniform integrability concerning the weights”, *Collect. Math.* 45 (2), (1994), 121 - 132.
- [5] T.K. Chandra and A. Goswami, “Cesàro  $\alpha$ - integrability and laws of large numbers *Int., J. Theoret. Prob.* 16 (3), (2003), 655- 669.
- [6] A. Mohammed, “Uniform Integrability of Approximate Green Functions of some Degenerate Elliptic Operators”, *Pacific Journal of Mathematics*, 199,(2), 2001.
- [7] B.M. Brown, “Moments of a stopping rule related to the central limit theorem”. *Ann. Math. Statist.* 40 (2004), 1236 - 1249.
- [8] R. Yokoyama, “Convergence of moments in the central limit theorem for stationary mixing sequences”, *Pompey.* (1995), 48-55.
- [9] Hiroshi Takahata, “Remark on Uniform Integrability of Functions Related to the Sums of Random Variables”, *Bulletin of Tokyo Gakugei University Ser. IV* 30 ,35-40, 1978.
- [10] G. J. Babu, A. J. Canty and Y. P. Chaubay, “Application of Bernstein polynomials for smooth estimation of a distribution and density function”, *J. Statistical Planning and Inference*, 105 (2002), 377-392.
- [11] Y. Kakizawa, “A bias-corrected approach to density estimation”, *J. Nonparametric Statistics*, 11,(2004), 709-729.
- [12] R. A. Vitale, “A Bernstin polynomial approach to density function estimation”, *Statistical Inference and Related Topics*, 2 (1975), 87-99.
- [13] A. Leblanc and B. C. Johnson, “A family of inequalities related to binomial probabilities”. Department of Statistics, University of Manitoba. Tech. Report, 2006
- [14] A. Leblanc and B. C. Johnson, “On a uniformly integrable family of polynomials defined on the unit interval”, *J. Inequal. Pure and Appl. Math.* 8 (3), (2007).
- [15] M. Abramowitz and I.A. Stegun, “Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables”, *National Bureau of Standards, Applied Mathematics Series* 55, 4th printing, Washington (1965).
- [16] W. Feller, “An Introduction to Probability Theory and its Applications”, Volume II, 2<sup>nd</sup> Ed. John Wiley and Sons, New York (1971).
- [17] H. Alzer and C. Berg, “A class of completely monotonic functions”, II, *Ramanujan J.*, 11 (2), (2006), 225-248.
- [18] S.H. Sung, “Weak law of large numbers for arrays of random variables”, *Statist. Probab. Lett* 42 (3), (1999), 293-298.
- [19] Q.M. Shao, “A comparison theorem on moment inequalities between negatively associated and independent random variables”, *J. Theoret. Probab.* 13 (2), (2000) 343-356.

- [20] V. Kruglov, “Growth of sums of pairwise-independent random variables with finite means”, *Teor. Veroyatn. Primen.* 51 (2), (2006),382-385.
- [21] D. Li, A. Rosalsky, and A. Volodin, “On the strong law of large numbers for sequences of pairwise negative quadrant dependent random variables”, *Bull. Inst. Math. Acad. Sin. (N.S)* 1 (2), (2006), 281-305.
- [22] S.H. Sung, T.C. Hu, and A. Volodin, “On the weak laws for arrays of random variables”, *Statist. Probab. Lett.* 72 (4),(2005),291-298.
- [23] S.H. Sung, S. Lisa wadi, and A. Volodin, “Weak laws of large numbers for arrays under a condition of uniform integrability”, *J. Korean Math. Soc.* 45 (1),(2008), 289-300.
- [24] J. Bae, D. Jun, and S. Levental, “The uniform clt for martingale difference arrays under the uniformly integrable entropy”, *Bull. Korean Math. Soc.* 47(1), (2010),39-51.

#### **ACKNOWLEDGEMENT**

We would like to express our sincere thanks to the anonymous reviewers and referees for their helpful suggestions and valuable comments.