

Volume 8 December 22 2002



TECHNOLOGY AND DEVELOPMENT

(A Journal of Scientific and Technological Research, ISSN. 1115-6775)

Published by :
Federal University of Technology, Yola.

INFORMATION TO CONTRIBUTORS

Articles should contain only original research work of good quality in any one of the above areas which is not published or submitted for publication elsewhere. Special priorities will be given to new scientific discoveries and development of new technologies. High quality review work in above areas may also be considered for publication. A publication charge is required for publication of the paper accepted after proper review. While submitting articles foreign authors must also send a completed agreement in the forms shown in Appendix A. This will waive them the payment of handling charges at the time of submission. While submitting articles authors are encouraged to include also their e-mail address and a diskette containing their article(s). Articles written in English should be submitted in triplicate to the Editor-in Chief at the address given above along with a non-refundable fee of N700. Articles should be typed (preferably with MS Word 2000 and in font 12) on one side of the paper and with double line spacing. All pages of the submitted manuscript must be numbered. The manuscript should be as concise as possible and must not exceed 22 typed pages (except review articles) including tables and figures. Review articles must not exceed 32 typed pages. Each copy of the manuscript should contain the first two pages as follows:

- 1) Page 1: Title (Bold font 14, first letter and only special letters e.g., the first letter of a chemical symbol of an element in capital, the rest in small letters) and authors' names (Bold, font 12) with addresses in italic.
 - 2) Page 2: Title and abstract only. The abstract (font 10) containing only one paragraph and not exceeding 250 words should be typed with double line spacing.
 - 3) Key words of not more than 10 words must be included in page 2.
4. The sections of the rest of the text (in font 12) should be classified as follows:
- i) **I. INTRODUCTION** (the aim and scope of the work should be included at the end of the introduction)
 - ii) **II. EXPERIMENTAL DETAILS**
 - iii) **III. RESULTS**
 - iv) **IV. DISCUSSION**
 - v) **V. CONCLUSION AND RECOMMENDATION**

All tables should be numbered with roman capital letters e.g., **TABLE I**). The title of a table should follow the table number (e.g., **TABLE I**. Experimental and theoretically computed values of phase transition temperatures.....). Tables should be referred in the text as Table I not as **TABLE I**. Materials should follow the corresponding section heading after a one-line-gap. Each section may have subsections. For example: under section '**V. DISCUSSION**' may appear a subsection "**V.1. Technological aspect**".

The material should start in the next line following the subsection (i.e. without any line gap) whereas the text should appear with one line gap after any 'major section'. All new paragraphs should be indented by seven spaces. "**Acknowledgements**" is optional. Only SI units should be used. Journal citations should be according to authors' names with years. References should be listed in alphabetical order of the authors names and in chronological order for the same authors cited several times for different works. For the reference listing follow the style given below. Each table or figure should be accompanied by a caption at the appropriate place. All figures (preferably black and white) should be of high standard. Coloured figures will require additional publication charge. Equations should be written using Equation Editor. All pages including the tables and figures must be numbered starting with the abstract page. It is suggested (but optional) that after the article is accepted by the Editorial Board, the authors should send the final version of the manuscript in a camera ready version or in diskette (in MS WORD 2000 or higher version) to speed up the publication process. A galley proof will be sent to each contributing author, for final check of the manuscript before appearing in the journal. Articles may be delayed significantly if the galley-proof is not received back in time from the author. All correspondences should only be directed to the Editor-in-Chief at the address given above.

References:

Journal:

Blakey, L. J. and Priest, F. G. (1980), 'The occurrence of *Bacillus cereus* in some dried foods including pulse and cereals' *Journal of Applied Bacteriology*, **48**, 297-302.

Books and Manuals:

Hudson, K. B. and Sherwood, L. (1997), *Exploitations in Microbiology*, Prentice Hall, New Jersey, p. 83-90.

Fawole, M. O. and Oso, B. A. (1988), *Laboratory Manual of Microbiology*, Spectrum Books Limited Ibadan, Nigeria, p. 24-56.

Edited Books:

Okaruwa, K. and Dashiell, E. R. (1997), in 'Soybean for Tropics. Research, Production and Utilization', Singh, S. R., Racjie, K. O. and Dashiell, E. R. (Eds.), John Wiley and Sons. New York, p.45.

REPRINTS:

All authors must pay for the publication cost of their article(s) after their papers are accepted. This will enable them two free off-prints. Additional copies (say, N copies) may be ordered at the cost price, $COST = N25 \times N \times p$. Where p = the number of printed pages, after publication. For foreign contributors the cost is, $COST = \$0.35 \times N \times p$ plus shipment cost.

SUBSCRIPTION INFORMATION

The journal is published annually. The annual subscription rates are:

In Nigeria

Individuals: N650.00, Institutions: N1000.00.

Foreign (including air freight):

Institutions in Asia (Excluding Japan): \$40.

Institutions in Japan, Europe and USA: \$65.

Africa (outside Nigeria): Institutions: \$30;

Send orders to:

The Editor-in-Chief, Technology and Development, School of Post Graduate Studies,

Federal University of Technology, Yola, P. M. B. 2076, Adamawa State, Nigeria.

Tel: 234 -75 - 627291. FAX: 234 -75 - e-mail:dkde@futy.edu.ng.

Appendix A: Agreement by foreign contributors

I hereby agree to pay the publication cost (which will not exceed \$70) of our article title “

“ by

for publication in your Journal “Technology and Development” after it is accepted for publication. I understand that this will waive the authors paying the handling fee for review of the article and entitle me to receive five free reprints after it is published.

Signed:

Name:

Position:

Address:

Seal:

Contents:

	Titles & Authors	Page
1.	Treatment of water effluent obtained during crude oil production- Semiu A Kareem	1
2.	Planning successful Fadama development in northern Nigeria- Adamu Idama	6
3.	An innovative technique of rapid analysis of petroleum contamination in kerosene- Dilip Kumar De, M. A. Sheikh and Y. T. Usman	13
4.	Antimicrobial effects of components of the bark extract of neem (<i>Azadirachta indica</i> A. Juss)- Nandita De and E. Ifeoma	23
5.	Antimicrobial spectrum of hexane extract of <i>Commiphora Kerstingii</i> (Burseraceae)- D. Kubmarawa, G. Ajoku and D. A. Okorie	29
6.	Preliminary studies of epicentral map in the western sector of the Gulf of Eden- J.B. Yerima	33
7.	Quality assessment of stored soybean flour (<i>Glycine max.</i> (L). marr)- Nandita De	38
8.	Petrological studies of Cretaceous limestone beds of north east- Dadiya Syncline in the Yola Arm of Upper Benue Trough, Nigeria- S. A. Opeloye, S. S. Dada and G. Obaje	42
9.	The relationship between occupational skills required in manufacturing industries and those acquired in polytechnics- M. Ubale Cyril	49
10.	A Survey of Savanna tree barks for bioactive phytochemicals in Yola metropolis of Adamawa State- J. U. Ijomah, M. Idu, and B. A. O. Okusanya	59
11.	Conductivity and pH values of 0-60 cm soil profiles of two commercial areas in Yola, Adamawa State, Nigeria, using column leaching experiment – E., Adeiza, T.C. Wapanuwari & I.F.H. Andenyang	65

Antimicrobial effects of components of the bark extract of neem (*Azadirachta indica* A. Juss)

Nandita De^{*a} and E. Ifeoma¹

^{*}Department of Microbiology, Federal University of Technology, Yola
Adamawa State, Nigeria.

¹Departemnt of Biological Sciences, Federal University of Technology, Minna
Niger State, Nigeria.

(Originally accepted for publication in Technology and Development, Vol. 7, 2001)

Preliminary phytochemical analysis showed that the leaves and bark of *A. indica* possess saponins, tannins and phenolic compounds. The antimicrobial spectrum of the aqueous, methanol and acetone extracts of leaves and bark of the plant was determined by agar diffusion method. The methanol and acetone extracts of the bark at 10mg/ml concentration showed significant antimicrobial activities against *Staphylococcus aureus*, *Streptococcus pyogenes*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Candida albicans* and *Trichophyton mentagrophytes*. The minimum inhibitory concentration (MIC) values of the methanol extract of the bark (0.3-0.4 mg/ml) were same or lower than those of ampicillin, erythromycin and griseofulvin (0.2-0.5 mg/ml). The minimum bactericidal concentration (MBC) values (0.3-0.9 mg/ml) and the minimum fungicidal concentration values (MFC) (0.3-2.0 mg/ml) of the methanol extract were comparable to those of ampicillin, erythromycin (0.2-0.6 mg/ml) and griseofulvin(0.8-1.0 mg/ml). Isolation of antimicrobial substances from the methanol extract of the bark was studied by thin layer chromatography (TLC) developed by using benzene and methanol mixture (4:6). Two fractions (A, B) were obtained. Fraction A (the upper band in TLC plate) showed significant activity (diameter of zone 4.0-5.0 mm at 2.0 mg/ml concentration) against *C. albicans* and *T. mentagrophytes* when compared to crude extract (diameter of zone 3.0-5.0 mm at 10 mg/ml concentration). So it may be concluded that there is a scientific basis for traditional use of extracts of bark and leaves of *A. indica* in skin infection and the active principles in the extracts may be purified by using TLC method.

Key words: *A. indica*, MIC, MBC, TLC. Plants-microbiology; skin-infections; neem (*Azadirachta A. Juss*) products

I. INTRODUCTION

Antimicrobial compounds are known to be present in the extracts of different tissues of some medicinal plants (Akpatha and Akinrimisi, 1977). Emeruwa (1982) showed that antibacterial substances from *Carica papaya* fruits were bactericidal against *Staphylococcus aureus*, *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Shigella flexnari* and these antibacterial substances were appeared to be protein in nature. Marston *et al.* (1993) isolated xanthenes from the root of *Polygala nyikensis* which possess antifungal activity. Different parts of *Azadirachta indica*, a common tree plant in Nigeria, are used medicinally (Adeserrano, 1982). A study was made here to compare the antimicrobial spectrum of the extracts of *A. indica*

to those of griseofulvin, erythromycin and ampicillin which are conventional antibiotics used against some pathogenic fungi and bacteria.

II. MATERIALS AND METHODS

II.1 Materials

The leaves and bark of *A. indica* were collected at School of Agricultural and Agricultural Technology Farm, Federal University of Technology, Minna. The plant was identified and authenticated by Dr. M.I.S. Ezenwa, School of agriculture and Agricultural Technology, FUT, Minna (Keay and Onochie, 1960 and Ivens *et al.*, 1978). After collection, leaves and bark of *A. indica* were sun dried to a constant weight over seven days and then crushed into powder using a blender (National MX 391M,

Matushita Electric). The powdered samples (50-60 gm) were separately extracted with 200 ml water, acetone and methanol using Soxhlet in vacuo to concentrate the extracts. The concentrated extracts were kept refrigerated pending further use. Each of the dried extracts was separately reconstituted in minimum amount of the extracting solvent and further diluted with glycerol to get the desired concentrations (10 mg/ml for susceptibility test and 0.2–2.0 mg/ml for MIC and MMC tests). The standard antibiotics used for MIC and MMC determination were erythromycin, griseofulvin and ampicillin. The antibiotic solutions were made using sterile distilled water.

II.2 Organisms

The fungi used were *Candida albicans* and *Trichophyton mentagrophytes* while the bacteria used were *Staphylococcus aureus*, *Streptococcus pyogenes*, *Escherichia coli* and *Pseudomonas aeruginosa*. All the cultures were obtained from National Veterinary Research Institute, Vom, Jos, Nigeria. The cultures were maintained on appropriate media- Potato dextrose agar (PDA) for fungi, MacConkey agar (MA) for *E. coli* and Nutrient agar (NA) for other bacterial isolates.

II.3 Preliminary phytochemical analysis

The analysis was done to screen the plant parts for phytochemicals using the method of Sofowora (1982). The phytochemicals tested for include sesquiterpenes, anthraquinones, phenolic compounds, tannins and saponins.

II.4 Determination of antimicrobial activity of the extracts

The fungal isolates were inoculated into PD broth and the cultures obtained after 48 hours were adjusted to a concentration of 10^6 cells per ml by serial dilution and used for the susceptibility test. The bacterial isolates except *E. coli* were inoculated into nutrient broth and the cultures after 24 hours were adjusted to 10^6 cells per ml. In case of *E. coli*, the same procedure was followed except that MacConkey agar surfaces in plates were inoculated with the respective cell suspension and the extracts were tested for their

effect on the organisms by cup plate method of Garrod *et al.* (1981).

II.5 Determination of minimum inhibitory concentration (MIC) and minimum microbicidal concentration (MMC) of the methanol extract of the bark

The methanol extract of the bark of *A. indica* which showed significant antimicrobial activity against *T. mentagrophytes*, *C. albicans*, *S. aureus*, *S. pyogenes* and *P. aeruginosa* was selected for MIC and MMC test.

a) MIC test

0.5 ml of varying concentrations of the reconstituted extract was added to two ml broth (potato dextrose for fungi, nutrient broth for bacteria except in case of *E. coli* where MacConkey broth was used). Then the broth containing extract was inoculated with a loopful of the organism at 10^6 cells/ml concentration. For each of the selected bacterial and fungal species the same procedure was followed using either the extract or standard antibiotic. After incubation (for fungi, 48 hours and for bacteria, 24 hours) the highest dilution of the extract or standard antibiotic that prevented visible growth of the test organism was taken as the MIC (Rotimi and Mosadonmi, 1987).

b) MMC (MFC and MBC) test

For each of the test tubes in the MIC determination which did not show any visible growth, a loopful of broth was aseptically inoculated on sterile agar surface by the streak method. After incubation, the minimum microbicidal concentration (MMC) was determined by the highest dilution at which there was no visible growth on the media.

II.6 Isolation of antimicrobial substances from the methanol extract of the bark

Thin layer chromatography was used for this purpose. Glass TLC plates (20x5 cm), each coated with 25 g of silica gel, were used. Different solvent mixtures were used for the separation of the antimicrobial components in the extract. The solvent mixtures used were (a) chloroform:

Methanol (5:5); (b) chloroform : benzene (5:5); (c) petroleum ether : methanol (5:5) (d) petroleum ether : chloroform (5:5) ; (e) petroleum ether : benzene (5:5); (f) benzene : methanol : water (1:4:5). Using a capillary tube, 10 mg/ml solution of the methanol extract of the bark was spotted on a TLC plate. The plate was then placed in a chromatographic jar containing 30 ml of appropriate solvent mixture. The plate was run for 90 minutes, removed, allowed to dry and put into an iodine jar for 2 minutes to locate the spots. Benzene : methanol solvent mixtures (5:5, 4:6, 3:7) were further used to ensure proper separation. The chromatography procedure was repeated for a broad band (5 ml of 10 mg/ml bark extract) using benzene: methanol (4:6) solvent mixture. The resulting bands (fraction A and fraction (B) and the residue (C) at the base were cut out and each washed with 10 ml of ethanol in a test tube. Using a capillary tube, 0.5 ml of the filtrate was applied on mini TLC plates (microscopic slides coated with 2 g of silica gel) to ensure the purity of these bands. The remaining filtrate obtained from the resultant bands were dried and then reconstituted with 5 ml glycerol. The reconstituted solutions were kept for future use (Vogel, 1980).

11.7 Determination of the antimicrobial activity of the isolated bands from TLC plate

The antimicrobial activity of the reconstituted fractions (A, B and C) at 2.0 mg/ml against some selected microorganisms was then determined by cup plate method and compared with that of the crude extract.

III. RESULTS

The phytochemical analysis showed the presence of phenolic compounds, tannins and saponins in both bark and leaves of *A. indica*. The results are shown in Table 1. Table 2 shows the amount of samples obtained by using 100 g of the different plant parts. The aqueous extracts of all plant parts did not exhibit any antimicrobial activity against the test organisms at the concentrations tested. The zones of inhibition produced by acetone and methanol extracts of the leaves against the organisms were not significant. However, they were quite significant for the bark

(Table 3). The MIC and MMC of the methanol extracts of the bark were comparable to those of some antibiotics generally used for the treatment of infections caused by these organisms. The results are shown in Table 4. The diameter of zones produced by fraction A at 2.0 mg/ml was comparable to the diameter of zones produced by crude extract, at 10.0 mg/ml against *S. aureus*, *S. pyogenes*, *P. aeruginosa*, *C. albicans* & *T. mentagrophytes*. The results are shown in Table 5.

IV. DISCUSSION

The preliminary phytochemical analysis showed that the leaves and bark of *A. indica* contain phenolic compounds and tannins. These phytochemical compounds have been reported by Farnsworth (1990) to inhibit bacterial growth and are capable of protecting certain plants against bacterial infections. The aqueous extracts did not produce any measurable zone of inhibition against the organisms at the tested concentration. When plant materials are ground in water or plant cells are damaged a number of phenolases and hydrolases are released. These enzymes might modulate the activity of active components in the extract. Also it is possible that some phytochemical components are not extracted with water (Akpata and Akinrimisi, 1977). The methanol and acetone extracts of the bark produced significant antimicrobial activity against *S. aureus*, *S. pyogenes*, *P. aeruginosa*, *C. albicans* and *T. mentagrophytes*. The solubility of tannins and phenols in methanol might have been the factor responsible for the significant antimicrobial activity (Wolinsky and Sote, 1982). The extracts of the leaves did not possess any antimicrobial properties at 10 mg/ml against the organisms probably because substances extracted from the leaves had no antimicrobial properties. The separation of the methanol extract of the bark into two distinct components (Fractions A and B) by thin layer chromatography shows that this plant part contains more than one active component. The varying susceptibility of each bacterial and fungal species may be a function of the available binding sites on the bacterial and fungal cells, these are probably surface structures like proteins. Tannins have been shown to form

TABLE 1: Preliminary phytochemical analysis of bark and leaves of *A. indica*

Phytochemical components	Bark	leaves
Sesquiterpenes	-	-
Anthraquinones	-	-
Phenolic compounds (including tannins)	++	++
Saponins	++	++

'++' indicates the presence of component in high amount, '-' indicates the absence of component

TABLE 2: Crude extracts obtained from leaves and bark of *A. indica* using water, methanol and acetone (mean \pm SEM; n=3).

Extracting medium	<i>A. indica</i> (plant part)	extract (g/100g of plant part)
Water	leaf	7.0 \pm 0.21
	bark	8.0 \pm 0.17
Methanol	leaf	6.4 \pm 0.21
	bark	6.0 \pm 0.21
Acetone	leaf	3.2 \pm 0.08
	bark	4.0 \pm 0.20

TABLE 3: Zones of inhibition produced by acetone and methanol extracts of parts of *A. indica*

plant extracts	Zones of inhibition(mm)					
	A	B	C	D	E	F
Methanol extract						
leaf	1.0	1.0	1.0	-	-	-
bark	4.0	5.0	3.0	4.0	4.0	2.0
Acetone extract						
leaf	-	-	-	-	-	-
bark	2.0	2.0	2.0	3.0	2.0	1.0
Aqueous extract						
leaf	-	-	-	-	-	-
bark	-	-	-	-	-	-

Key: A- *S. aureus*, B- *S. pyogenes*, C- *P. aeruginosa*, D- *T. mentagrophytes*, E- *C. albicans*, F- *E. coli*.
 "-" in Table 3 means no activity.

TABLE 4: Determination of MIC and MMC of the methanol extract of the bark of *A. indica* and some standard antibiotics.

Extract & Antibiotics	MIC (mg/ml)					MMC(mg/ml)				
	A	B	C	D	E	A	B	C	D	E
Extract	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	2.0	0.9
Erythromycin	0.3	0.2	0.4	NA	NA	0.6	0.6	0.2	NA	NA
Ampicillin	0.3	0.3	0.2	NA	NA	0.5	0.4	0.3	NA	NA
Griseofulvin	NA	NA	NA	0.3	0.4	NA	NA	NA	1.0	0.7

Key: A- *S.aureus*, B- *S. pyogenes*, C- *P.aeruginosa*, D- *C. albicans*, E- *T.mentagrophytes*, NA- non-applicable.

TABLE 5: Comparison of antimicrobial activity of the separated fractions A, B and the residue C and the crude extract

Fractions & Crude extract	Diameter of zone (mm)				
	A	B	C	D	E
Fraction A (2.0 mg/ml)	4.0	5.0	3.0	4.0	4.0
Fraction B (2.0 mg/ml)	-	4.0	2.0	2.0	3.0
Residue C (2.0 mg/ml)	-	-	-	-	-
Crude extract (10.0 mg/ml)	4.0	5.0	3.0	3.5	4.0

Key: A- *S. aureus*, B- *S. pyogenes*, C- *P.aeruginosa*, D- *C. albicans*, E- *T. mentagrophytes*

irreversible complexes with proline rich proteins. This could interfere with cell function. Table 4 showed that the MIC and MMC of the methanol extract of the bark was comparable to those of erythromycin, griseofulvin and ampicillin. Griseofulvin, an antifungal drug, inhibits cell division and is active against dermatophytes *Trichophyton* and *Candida* sp. Erythromycin and ampicillin are broad spectrum antibiotics.

Ampicillin is known to inhibit sensitive strains of *P. mirabilis* and *E. coli*. Erythromycin act by affecting ribosome function by selective binding to bacterial 50S subunit (Russel and Hugo,1977).The comparable activity of the upper band (A), lower band (B), residue (C) at the base and the crude extract shows that the antimicrobial

activity of the fraction A was quite high (zones of inhibition 4-5 mm at 2.0 mg/ml concentration) compared to the crude one (zones of inhibition 4-5 mm at 10.0 mg/ml concentration). Purification may have eliminated substances that reduce the activity of the active components. Therefore, the information obtained from this study have proved scientifically that the components of the bark of *A. indica* possess significant antimicrobial activity against *S. aureus*, *S. pyogenes*, *P. aeruginosa*, *C. albicans* and *T. mentagrophytes*.

^a To whom all correspondence should be addressed

REFERENCES

- Akpata, E. S. and Akinrimisi, E.O.(1977). Antimicrobial activity of extracts from some African chewing sticks. *Oral Surgery* **44**, 717-722.
- Adeserrano, O.(1982). Growth inhibitory and lymphocytotoxic effect of *A. indica*. *Journal of African Medicinal plants* **5**:6-9.
- Emeruwa, A.C.(1982). Antibacterial substances *Carica papaya* fruit extract. *Journal of Natural Products* **45(2)**: 123-127.
- Farnsworth, N.R. (1982). The role of ethnopharmacology in drug development from plants. John Wiley and Sons, England (Ciba Found. Synp.154) p 2-10
- Garrod, L.P., Lambert, H.P. and O'Crady, I.(1981). Antibiotics and Chemotherapy. Churchill Livingstone, London, p 225.
- Ivens, G.W., Moody, K. and Ejungobi, J.K. (1978). West African Weeds. Oxford University Press, Ibadan, p 225.
- Keay, R.W.J. and Onochie, C.F.A.(1960). Nigerian Trees. Federal Department of Forest Research, Federal Govt. Printer, Lagos, Vol. II. p 428.
- Marston, A., Hamburger, M., Sordat_Diserens, L. and Hostettman, K. (1993). Xanthenes from *Polygala nyikensis*. *Phytochemistry* **33(4)**:809-812.
- Rotimi, V.O. and Mosadonmi, H.A. (1987). The effect of crude extracts of nine african chewing sticks on oral anaerobes. *Journal of Medical Microbiology* **23**: 56-60.
- Russel, A.D. and Hugo, W.B.(1977). Pharmaceutical Microbiology, 5th ed. Blackwell Scientific Publication, Oxford, p 97-119.
- Sofowora, E.A. (1982). Medicinal plants and traditional medicine in Africa. John Wiley and Sons Ltd. New York. p 1-20.
- Vogel, A. (1980). Text book on macro and semimacro qualitative inorganic analysis. 15th ed.(ed. Longman), London.
- Wolinsky, L.E. and Sote, E.O. (1984). The isolation of natural plaque inhibiting substances from Nigerian chewing sticks. *Caries Research* **18**:216-225.