

British Journal of Mathematics & Computer Science 4(24): 3497-3507, 2014 ISSN: 2231-0851



SCIENCEDOMAIN international www.sciencedomain.org

# Using the Average of the Extreme Values of a Triangular Distribution for a Transformation, and Its Approximant via the Continuous Uniform Distribution

H. I. Okagbue<sup>1</sup>, S. O. Edeki<sup>1\*</sup>, A. A. Opanuga<sup>1</sup>, P. E. Oguntunde<sup>1</sup> and M. E. Adeosun<sup>2</sup>

<sup>1</sup>Department of Mathematics, College of Science & Technology, Covenant University, Otta, Nigeria. <sup>2</sup>Department of Mathematics and Statistics, Osun State College of Technology, Esa-Oke, Nigeria.

#### Article Information

DOI: 10.9734/BJMCS/2014/12299 <u>Editor(s):</u> (1) H. M. Srivastava, Department of Mathematics and Statistics University of Victoria, Canada. <u>Reviewers:</u> (1) Anonymous, National Institute of Technology, India. (2) Anonymous, Hangzhou Dianzi University, PR China. (3) Seifedine Kadry, American University of the Middle East, Kuwait. (4) Anonymous, Guru Nanak Dev University, India. Complete Peer review History: <u>http://www.sciencedomain.org/review-history.php?iid=699&id=6&aid=6427</u>

Original Research Article

Received: 26 June 2014 Accepted: 18 August 2014 Published: 09 October 2014

### Abstract

This paper introduces a new probability distribution referred to as a transformed triangular distribution (TTD) by using the average of the extreme values (minimum and maximum) of the triangular distribution. The TTD is being approximated by the continuous uniform distribution. The basic moments of the TTD and those of the continuous uniform distribution are compared respectively, and a relationship established. This can be used in modeling and simulation.

Keywords: Moments, uniform distribution, triangular distribution, transformed distribution, continuous random variable.

Mathematical Subject Classification (2010): 05A10, 81S20, 81S30, 33B20

<sup>\*</sup>Corresponding author: soedeki@yahoo.com;

## **1** Introduction

Triangular distribution is a continuous distribution with a fixed minimum, fixed maximum and most likely value to occur (mode). The most likely value lies between the minimum and maximum values, forming a triangular-shaped distribution which shows that values near the minimum and maximum are less likely to occur [1]. The minimum and the maximum values are called the extreme values. The values can either be symmetrical (the mod e = mean = median) or asymmetrical [2].

The distribution is widely used to approximate Beta distribution [3]. Some of the earliest known written work on the triangular distribution are traceable to the work of [4,5], even though it was Simpson that first mentioned the distribution in his papers [6].

Nevertheless, many authors and researchers have worked on the triangular distribution; some of them are either on the statistical, probabilistic nature of the distribution or in the area of simulation and modeling. Some of the contributions include the characteristics of the distribution [7,8], the product of two identically independent distributed triangular variables [9], on the extension of the triangular distribution and the convolution [10,11], applications in project evaluation and review technique PERT [12-16], non-smooth sailing with respect to asymptotic distribution [23-24], on the negative binomial-triangular distribution [25], advanced simulation and risk modeling [26-28] combination of triangular and exponential distributions [29], the sum of two triangular distributions [30], discrete nature of triangular distribution and non-parametric estimation for probability mass function [31].

The remaining part of the paper is structured as follows: section 2 deals with the basic concepts and properties of the concerned probability distributions, section 3 is on methodology and procedures towards the transformation of the triangular distribution and discussion of results; while in section 4, a concluding remark is made.

## 2 The Basic Concepts of the Probability Distributions

In this section, we present both the properties of the triangular and the continuous uniform distributions to be used in the later part of the work.

#### 2.1 Some Moments and Properties of the Triangular Distribution (TD)

Suppose X is a random variable with parameters a, b and c such that

$$a: a \in (-\infty, \infty)$$
  
$$b: a \le b \le c$$
  
$$c: a < c$$

and a support ,  $a \le x \le b$ 

then, X is said to follow a triangular distribution  $X \sim T(a, b, c)$  with the following properties:

Probability density function (pdf) f(x) of X given as:

$$f(x) = \begin{cases} 0 & , & for \quad x < a \\ \frac{2(x-a)}{(c-a)(b-a)}, & for \quad a \le x \le b \\ \frac{2(c-x)}{(c-a)(c-b)}, & for \quad a \le x \le c \\ 0 & , & for \quad x > c \end{cases}$$
(1)

Cumulative Density Function (CDF) X:

$$F(x) = \begin{cases} 0, & \text{for } x < a \\ \frac{(x-a)^2}{(c-a)(b-a)}, & \text{for } a \le x \le b \\ 1 - \frac{(c-x)^2}{(c-a)(c-b)}, & \text{for } b < x \le c \\ 1, & \text{for } c < x \end{cases}$$
(2)

Mean:

$$E(X) = \frac{a+b+c}{3} \tag{3}$$

Median:

$$X_{Median} = \begin{cases} a + \frac{\sqrt{(c-a)(b-a)}}{\sqrt{2}} & \text{for } b \ge \frac{a+c}{2} \\ c - \frac{\sqrt{(c-a)(c-b)}}{\sqrt{2}} & \text{for } b \le \frac{a+c}{2} \end{cases}$$
(4)

Most likely value

$$X_{Mode} = b \tag{5}$$

Variance:

$$Var(X) = \frac{a^2 + b^2 + c^2 - ac - ab - bc}{18}$$
(6)

Skewness:

$$X_{Skewness} = \frac{\sqrt{2}(a+c-2b)(2a-c-b)(a-2c+b)}{5(a^2+c^2+b^2-ac-ab-bc)^{\frac{3}{2}}}$$
(7)

Kurtosis:

$$X_{Kurtosis} = -3/5 \tag{8}$$

## 2.2 Some Moments and Properties of the Continuous Uniform Distribution

A random variable Y over the interval I = [a,b] is a continuous uniform distribution if it is *equally likely* to assume any value in I. Let f(y) and F(y) be the pdf and CDF of Y respectively, then:

$$f(y) = \begin{cases} \frac{1}{(b-a)} &, a \le y \le b \\ 0 &, \text{ otherwise} \end{cases}$$
(9)

$$F(y) = \begin{cases} 0, & \text{for } y < a \\ \frac{(y-a)}{(b-a)}, & \text{for } a < y \le b \\ 1, & \text{for } c \ge b \end{cases}$$
(10)

Remark 2.1: The following can easily be computed.

Mean:

$$Y_{mean} = \frac{b+a}{2} \tag{11}$$

Mode (can be any value in I) so we choose:

$$Y_{\text{mod}e} = \frac{b+a}{2} \tag{12}$$

Median

$$Y_{median} = \frac{b+a}{2} \tag{13}$$

Variance

$$Var(Y) = \frac{(b-a)^2}{12}$$
 (14)

Coefficient of skewness

$$Y_{c-skewness} = 0 \tag{15}$$

# **3** Methodology and the Modification of the Triangular Distribution

#### 3.1 The Transformation of the Triangular Distribution (TTD)

We replace the most likely value b, with the average of the minimum and maximum of the Triangular distribution in order to modify the Triangular distribution, thus;

$$b = \frac{a+c}{2} \tag{16}$$

The resulting distribution will henceforth be referred to as the Transformed Triangular Distribution (TTD).

#### **3.2 The Resulting Distribution TTD and Its Properties**

Suppose  $X^*$  is the random variable associated with the TTD,  $f^*(x)$  and  $F^*(x)$  as the corresponding pdf and CDF respectively, then by using (9), one can easily obtain the following:

$$f^{*}(x) = \begin{cases} \frac{4(x-a)}{(c-a)^{2}} & , a \le x \le \frac{a+c}{2} \\ \frac{4(c-x)}{(c-a)^{2}} & , \frac{a+c}{2} \le x \le c \\ 0 & , otherwise \end{cases}$$
(17)

and it is easy to show that:

$$F^{*}(x) = \begin{cases} 0, & \text{for } x < a \\ \frac{2(x-a)^{2}}{(c-a)^{2}}, & \text{for } a \le x \le \frac{a+c}{2} \\ 1 - \frac{2(c-x)^{2}}{(c-a)^{2}}, & \text{for } \frac{a+c}{2} < x \le c \\ 1, & \text{for } c \le x \end{cases}$$
(18)

Remark 3.1 validation of the probability density function of  $f^{*}(x)$ 

To validate the pdf of the TTD, we need to show that:

$$\int_{a}^{\frac{a+c}{2}} f^{*}(x)dx = 1$$
(19)

*Proof:* By definition,

$$\int_{a}^{c} f^{*}(x)dx = \int_{a}^{\frac{a+c}{2}} \frac{4(x-a)}{(c-a)^{2}}dx + \int_{\frac{a+c}{2}}^{c} \frac{4(c-x)}{(c-a)^{2}}dx$$
$$= \frac{4}{(c-a)^{2}} \left\{ \int_{a}^{\frac{a+c}{2}} (x-a)dx + \int_{\frac{a+c}{2}}^{c} (c-a)dx \right\}$$
$$= \frac{4}{(c-a)^{2}} \{A+B\}$$
(20)

where

$$A = \left(\frac{x^2}{2} - ax\right)\Big|_{a}^{\frac{a+c}{2}} = \frac{(a-c)^2}{8} \quad \& \quad B = \left(cx - \frac{x^2}{2}\right)\Big|_{\frac{a+c}{2}}^{c} = \frac{(c-a)^2}{8}$$
(21)

Thus,

$$\int_{a}^{c} f^{*}(x)dx = \frac{4}{(c-a)^{2}} \{A+B\}$$
=1
(22)

Since

$$A + B = \frac{(a-c)^2}{4} \& (a-c)^2 \equiv (c-a)^2, \ a \neq c$$
(23)

Showing that  $f^*(x)$  is indeed a valid pdf.

For the rest of the moments and properties we shall often refer to (17), as such (16) in (3) gives: *Mean:* 

$$X_{mean}^{*} = \frac{a + \frac{a + c}{2} + c}{3}$$

$$= \frac{a + c}{2} \ge \frac{b + a}{2}$$
(24)

Thus,

 $X^*_{mean} \ge Y_{mean}$ 

showing that:

$$Y_{mean} \approx X^*_{mean} \tag{25}$$

The Median of TTD using the extreme values:

Substitute equation (16) in (4) gives:

$$X_{median}^{*} = a + \sqrt{\frac{(c-a)(\frac{a+c}{2}-a)}{2}}$$
$$= a + \sqrt{\frac{(c-a)(c-a)}{4}}$$
$$= \frac{a+c}{2} \ge Y_{median}$$
(26)

showing that:

$$Y_{median} \approx X^*_{median}$$
 (27)

Similarly

$$Median = c - \sqrt{\frac{(c-a)(c-(\frac{a+c}{2}))}{2}}$$
$$= c - \sqrt{\frac{(c-a)(c-a)}{4}}$$
$$= \frac{a+c}{2}$$

showing the same result in (27).

The Mode of TTD ( can be any value in I ): In this case of mode, choose b such that:

$$X_{\text{mod}e}^* = b = \frac{a+c}{2}$$

$$\geq Y_{\text{mod}e}$$
(28)

showing that:

$$Y_{mode} \approx X^*_{mode} \tag{29}$$

*The variance of the TTD:* Substitute (16) in (6) gives:

$$Var(X^{*}) = \frac{a^{2} + (\frac{a+c}{2})^{2} + c^{2} - ac - a(\frac{a+c}{2}) - c(\frac{a+c}{2})}{18}$$
$$= \frac{(c-a)^{2}}{24}$$
$$\geq Var(Y)$$
(30)

showing that:

$$Var(Y) \approx Var(X^*)$$
 (31)

*The skewness of the TTD:* From (16)

$$b = \frac{a+c}{2} = 2b = a+c \tag{32}$$

Therefore, substituting (16) and (32) in (7) gives:

$$X_{Skewness}^{*} = \frac{\sqrt{2} \left(2b - 2b\right) \left(2a - c - \left(\frac{a + c}{2}\right)\right) \left(a - 2c + \left(\frac{a + c}{2}\right)\right)}{5 \left(a^{2} + c^{2} + \frac{\left(a + c\right)^{2}}{4} - ac - a\frac{\left(a + c\right)}{2} - c\frac{\left(a + c\right)^{\frac{3}{2}}}{2}\right)} = 0 = Y_{c-skewness}$$
(33)

## **4** Conclusion

In this paper, we have showed that when the mode of a triangular distribution is the average of the extreme values, then the resulting distribution referred as transformed triangular distribution (TTD) is a probability function that can be reasonably approximated, simulated and modeled by the continuous uniform distribution. The limit theorem takes care of the behavior of the distribution at large sample.

## Acknowledgements

We would like to express sincere thanks to the anonymous reviewer(s) and referee(s) for their constructive comments and valuable suggestions towards the improvement of the paper.

## **Competing Interests**

Authors have declared that no competing interests exist.

## References

- Mun J. Modeling risk: Applying Monte Carlo, risk simulation, strategic real options, stochastic forecasting, and portfolio optimization, 2<sup>nd</sup> Ed, Wiley & Sons; 2010. ISBN: 978-0-470-59221-2.
- Paul M. Introduction to probability and statistical applications. Addison-Wesley, Reading MA; 1970. ISBN: 9780201047103.
- [3] Johnson D. The triangular distribution as a proxy for the beta distribution in risk analysis. The Statistician. 1979;46:387-398.
- [4] David FN, Barton DE. Combinatorial chance. Lubrecht & Cramer Ltd; 1962. ISBN: 9780852640579.
- [5] John R. Combinatorial chance. The Annals of Mathematical Statistics. 1962;33(4):1477-1479.
- [6] Seal HL. The historical development of the use of generating functions in probability theory. Mitt Verein Schweiz Versich Math. 1949;49:209-228.
- [7] Kotz S, Dorp JRV. Beyond beta. World Scientific Publishing Co. Ltd, Singapore; 2004. ISBN: 981-256-115-3.
- [8] Jance M, Thomopoulos N. Min and max triangular extreme interval values and statistics. Journal of Business and Economics Research. 2010;8:139-143.

- [9] Donahue JD. Products and quotients of random variables and their applications. Aerospace Research Laboratories, Office of Aerospace Research, US Air Force, Wright-Patterson Air Force Base, Ohio; 1964.
- [10] Dorp JRV, Kotz S. A novel extension of the triangular distribution and its parameter estimation. The Statistician. 2002;51(1):63-79.
- [11] Dorp JRV, Kotz S. Generalizations of two-sided power distributions and their convolution. Comm Statist Theory Methods. 2003;32:1703-1723.
- [12] Clark CE. The PERT model for the distribution of an activity. Operations Research. 1962;10(3):405-406.
- [13] Grubbs FE. Attempts to validate certain PERT statistics or 'picking on PERT. Operations Research. 1962;10(6):912-915.
- [14] Keefer DL, Verdini WA. Better estimation of PERT activity time parameters. Management Science. 1993;39(9):1086-1091.
- [15] Moder J, Rodgers EG. Judgement estimates of the moments of PERT type distributions. Management Science. 1968;15(2):B76-B83.
- [16] Kamburowski J. New validations of PERT times. Omega. 1997;25:323-328.
- [17] Johnson N, Kotz S. Non smooth sailing or triangular distributions revisited after some 50 years. The Statistician. 1999;48:179-187.
- [18] Winston WL. Operations research: Applications and algorithm. 2<sup>nd</sup> Ed. Kent Publishing, Boston MA; 1993.
- [19] Altiok T, Melamed B. Simulation modeling and analysis with ARENA. Academic Press; 2010.
- [20] David Kelton W, Randall P. Sadowski, David T. Sturrock. Stimulation with Arena. McGraw Hill; 2008.
- [21] Sprow FB. Evaluation of research expenditures using triangular distribution functions and Monte Carlo Methods . Ind Eng Chem. 1967;59(7):35-38.
- [22] Chau KW. The validity of triangular distribution assumptions in Monte Carlo simulation of a construction costs: Empirical evidence from Hong Kong. Construction Management and Economics. 1995;13(1):15-21.
- [23] Nadarajah S. A polynomial model for bivariate extreme value distributions. Statistical Probab Letters. 1999;42:15-25.
- [24] Griffilths RC. On a Bivariate Triangular Distribution. Australian Journal of Statistics. 1978;20(2):183-185.

- [25] Karlis D, Xekalaki E. On some distributions stemming from the triangular distribution. Technical Report, Dept. of Statistics, Athens University of Economics. 2000;111.
- [26] Sarireh M. Estimation of HD drilling time using deterministic and triangular distribution functions. JETEAS. 2013;4(3):438-445.
- [27] Johnson D. Triangular approximations for continuous random variables in risk analysis. JORS. 2002;53:457-467.
- [28] Back EW, Boles W, Fry G. Defining triangular probability distributions from historical cost data. Journal of Construction Engineering and Management. 2000;126(1):29-37.
- [29] Brizzi M. A skewed model combining triangular and exponential features: The two-faced distribution and its statistical properties. Austrian Journal of Statistics. 2006;35(4):455-462.
- [30] Gary M, Choudhary S, Kalla SL. On the sum of two triangular random variables. International Journal of Optimization: Theory, Methods and Applications. 2009;1(3):279-290.
- [31] Kkonendji CC, Senga Kiesse T, Zocchi SS. Discrete triangular disributions and nonparametric estimation for probability mass function. Journal of Non-parametric Statistics. 2007;19:241-254.

© 2014 Okagbue et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

#### Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar) www.sciencedomain.org/review-history.php?iid=699&id=6&aid=6427