Varying influence of physical maturity 1

1	Title:	Influence of physical maturity status on sprinting speed among
2		youth soccer players
3		
4	Running Head:	Varying influence of physical maturity
5		
6	Authors:	Robert McCunn <sup>1</sup> , Matthew Weston <sup>2</sup> , John KA Hill <sup>3</sup> , Rich D
7		Johnston <sup>4</sup> , Neil V Gibson <sup>3</sup>
8		
9	Institutions:	<sup>1</sup> Institute of Sport and Preventive Medicine, Saarland
10		University, Saarbrücken, Germany
11		<sup>2</sup> Department of Sport and Exercise Sciences, School of Social
12		Sciences and Law, Teesside University, Middlesbrough, UK
13		<sup>3</sup> Oriam: Scotland's sports performance centre, Heriot-Watt
14		University, Edinburgh, UK
15		<sup>4</sup> School of Exercise Science, Australian Catholic University,
16		Brisbane, Australia
17		
18	Corresponding Author:	Robert McCunn
19		Institute of Sport and Preventive Medicine
20		Saarland University
21		Campus Bldg. B8.2
22		66123 Saarbrücken, Germany
23		Phone: +49 681 302 70410
24		Email: robert.mccunn@uni-saarland.de
25		Fax: +49 (0) 681 302 4296

1	Influence of physical maturity status on sprinting speed among youth
2	soccer players
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	

#### 26 Abstract

27

The relative age effect is well documented with the maturation-selection hypothesis the most 28 common explanation; however, conflicting evidence exists. We observed the birth-date 29 30 distribution within an elite junior soccer academy. The influence of physical maturity status on anthropometric variables and sprinting ability was also investigated. Annual fitness testing 31 32 was conducted over an eight-year period with a total of 306 players (age:  $12.5 \pm 1.7$  y [range: 9.7 - 16.6 y]; stature:  $156.9 \pm 12.9$  cm; mass:  $46.5 \pm 12.5$  kg) drawn from six age categories 33 (under-11s to -17s) who attended the same Scottish Premiership club academy. 34 35 Measurements included mass, stature, maturity offset and 0-15 m sprint. Odds ratios revealed 36 a clear bias towards recruitment of players born in quartile one compared to quartile four. The overall effect (all squads combined) of birth quartile was very likely small for maturity 37 38 offset (0.85 years; 90% confidence interval 0.44 years to 1.26 years) and stature (6.2 cm; 90% confidence interval 2.8 cm to 9.6 cm), and likely small for mass (5.1 kg; 90% 39 confidence interval 1.7 kg to 8.4 kg). The magnitude of the relationship between maturity 40 offset and 15 m sprinting speed ranged from *trivial* for under-11s (r = 0.01; 90% confidence 41 42 interval -0.14 to 0.16) to very likely large for under-15s (r = -0.62; -0.71 to -0.51). Making 43 decisions about which players to retain and release should not be based on sprinting ability 44 around the under-14 and under-15 age categories since any inter-individual differences may be confounded by transient inequalities in maturity offset... 45

46

47 Key words: association football, youth, talent identification, relative age effect, athletic

- 48 development
- 49
- 50

### 51 INTRODUCTION

52

Fielding teams at the professional level in soccer that include homegrown players, developed through a club's youth academy system has been described as cost effective (25). Despite long-term financial benefits apparent in the development of homegrown players a considerable outlay is required to ensure each player has access to adequate coaching and training facilities throughout their soccer education (25). Due to the scale of investment it is important that clubs make informed decisions, with appropriate foresight, when recruiting, selecting and releasing young players.

60

61 The relative age effect (RAE) is well documented within youth soccer and relates to the 62 uneven distribution of players' birth date relative to the general population (13). Youth soccer 63 is typically organised into one-year age bands with a bias toward recruitment of players born in the first quarter of the selection year (9); a finding that has been reported in many countries 64 (14). The existent research has documented the presence of the RAE in sport yet has failed to 65 explain why the phenomenon exists (8). Of the proposed theories the most commonly cited is 66 the maturation-selection hypothesis (27). It is posited that relatively older players are more 67 68 physically mature than their younger counterparts, which may be advantageous in sports, 69 which involve physical contact, for example soccer (21). Indeed, it is well understood that 70 during the transition from childhood to adulthood physical maturity influences many 71 characteristics relevant to sporting performance including stature, mass, aerobic power, strength and running speed (1, 18). However, it is less clear if advanced physical maturity 72 73 results in superior physical performance within the context of a one-year age band.

75 It is unclear whether any relationships between physical maturity and measures of physical 76 capacity are consistent throughout childhood and adolescence. Buchheit & Mendez-77 Villanueva (5) observed differences - varying in magnitude - in anthropometric and performance characteristics in relatively older and more physically mature under-15 players. 78 79 In contrast, Carling et al. (7) reported few differences between relatively older and younger 80 under-14 players. These conflicting studies illustrate that the relationships between relative 81 age, maturity and physical capacity in youth soccer players remain unclear. Furthermore, 82 studies focusing on one age category reveal only a partial view of the influence of maturity on physical qualities and the RAE, especially since many players are registered to the same 83 84 club for successive seasons. Furthermore, Figueiredo et al. (11) observed that within a wide 85 range of age categories (under-11s to -14s) the influence of physical maturity on measures of physical capacity differed depending on the category analysed. Similarly, Skorski et al. (23) 86 87 and Lovell et al. (17) reported varying influence of relative age on physical performance 88 markers across a wide range of age categories. These two studies, in addition to Buchheit & 89 Mendez-Villanueva (5) are, to our knowledge, the only instances where magnitude based inferences have been used to quantify the *degree* of influence relative age has upon physical 90 91 performance markers. The present study sought to contribute to this limited evidence base 92 and report not only if physical maturity status had an influence on sprinting speed, within 93 one-year age bands, but also the degree of the relationship. Understanding these relationships 94 has important implications for coaches and practitioners concerned with identifying players 95 for selection, retention and release at the end of each season.

96

97 The present study aimed to investigate the influence of relative age on physical maturity and 98 sprinting speed within six consecutive age categories (U11-U17). Data were collected over 99 eight seasons within a professional soccer academy. The first hypothesis was that relatively 100 older players would be more physically mature than their younger counterparts within all age 101 categories. The second hypothesis was that physical maturity would influence anthropometric 102 measurements (stature and mass) and sprinting speed but that the strength of these 103 relationships would not be consistent between all age categories.

104

### 105 **METHODS**

106

### 107 Experimental approach to the problem

108

An observational design was adopted for the present study. Anthropometric measures along
with physical performance test results from youth players belonging to a professional soccer
club academy were collected as part of routine fitness testing and analysed retrospectively.
Players were assessed over an eight-year period (season 2007/08 to 2014/15).

113

# 114 Subjects

115 A total of 306 male elite youth players (age:  $12.5 \pm 1.7$  y [range: 9.7 - 16.6 y]; stature: 156.9 116  $\pm$  12.9 cm; mass: 46.5  $\pm$  12.5 kg) who attended the same Scottish Premiership club academy 117 participated. These players were drawn from six age categories including under-11, under-12, 118 under-13, under-14, under-15 and under-17s. During the observation period some players 119 were retained year after year and progressed through the age categories resulting in multiple 120 observations in some instances (570 data points in total). All individuals joined the academy 121 via a selection process administered by scouts affiliated with the club (subjective assessment) 122 and were considered to be among the very best young players in Scotland. The benefits and 123 risks associated with the current investigation were explained to the participants before 124 signing an institutionally approved informed consent form. Written parental consent was also obtained prior to all physiological testing. The study was approved by The University of
Glasgow, College of Medical and Life Sciences research ethics board and conformed to the
recommendations of the Declaration of Helsinki.

128

129 **Procedures** 

130

131 *Relative age effect* 

132

133 To investigate the birth date distribution of the players, data were obtained from the General 134 Registrars Office for Scotland concerning the number of births within the general population 135 for the relevant years (1993-2004). This allowed a comparison between the expected and 136 observed birth date distribution in the sample population. Youth soccer in Scotland is 137 structured such that the selection year follows the calendar year (1<sup>st</sup> January to 31<sup>st</sup> 138 December). Hence, players born in quartile one possessed a birth date in January, February or 139 March and players born in quartile four possessed a birth date in October, November or 140 December.

141

142 Physiological assessments

143

During the first week of September each season, players completed a series of physical assessment protocols. Club support staff conducted all tests; all possessed a postgraduate degree in sport science in addition to nationally recognized strength and conditioning certifications. Mass along with standing and seated stretch stature was recorded to the nearest 0.1 kg and 0.1 cm respectively, using calibrated scales (Avery Weigh-Tronix, UK) and a wall-mounted stadiometer (Holtain Ltd, UK). For the anthropometric assessments players 150 removed their footwear and wore a training t-shirt and shorts. Maturity offset was calculated 151 using the equation developed by Mirwald et al. (20) and has been used in previous research 152 as an indicator of somatic maturity among youth soccer players (4, 6). Maturity offset 153 represents the amount of time (in years) until or since an individual's predicted peak height 154 velocity (PHV) and is calculated using an individual's stature, seated stature, mass, date of birth and the date of measurement (19). Maturity offset offers a logistically feasible way to 155 156 estimate physical maturity among large groups such as in the present study. Over the course 157 of the eight-year observation period a number of different tests were employed to characterise 158 the players' physical capabilities. As such, the results from season to season were not always 159 directly comparable. For example, a variety of different yoyo tests were used during the 160 observation period. The only physical performance test included in the analysis was the 0-15 161 m sprint since this test was used with all squads every season. After the players had 162 completed the anthropometric assessments they performed a standardized 15-minute warm 163 up comprising light aerobic exercise and dynamic stretches. The sprint test was always the first task to be performed in the test battery after the warm up each year. The 0-15 m sprint 164 165 test protocol allowed three attempts per player from a standing start 0.5 m behind the first 166 timing gate; the fastest time was recorded for analysis. Players had approximately three 167 minutes rest between efforts. The sprints were measured using electronic timing gates 168 (Smartspeed, Fusion Sports, Australia) and conducted on the same indoor synthetic pitch 169 each year. All participants wore soccer boots with moulded studs. The technical error of 170 measurement for the 0-15m sprinting assessment according to the club's own quality control 171 testing was 0.21 seconds.

172

173 Statistical Analysis

174 Data are presented as the mean  $\pm$  SD. Prior to all analyses plots of the residuals versus the 175 predicted values revealed no evidence of non-uniformity of error. In athletic research, it is not 176 whether there is an effect but how big the effect is that is important; use of the P value alone 177 provides no information about the direction or size of the effect or the range of feasible 178 values (2). The odds ratio, with uncertainty expressed as 90% confidence intervals, was used 179 to examine birth date distribution of our players against an expected equal distribution (e.g., 180 the general population). Here, all comparisons were made between quartile 1 and quartile 4 181 and the magnitude of the odds ratio was assessed against thresholds of trivial >1.5, small, >3.4, and moderate >9.0 (15). The effects of birth guartile (quartile 1 versus quartile 4) on 182 183 player maturity, stature and mass were analysed using a mixed linear model (SPSS v.22, 184 Armonk, NY: IBM Corp) with random intercepts. Standardised thresholds for small, 185 moderate and large changes (0.2, 0.6 and 1.2, respectively) calculated from between-player 186 standard deviations of all players in each respective squad, were used to assess the magnitude 187 of all effects (15). Inference was subsequently based on the disposition of the confidence 188 interval for the mean difference to these standardised thresholds and calculated as per the 189 magnitude-based inference approach using the following scale: 25–75%, possibly; 75–95%, 190 likely; 95–99.5%, very likely; >99.5%, most likely (15). Inference was categorised as unclear 191 if the 90% confidence limits overlapped the thresholds for the smallest worthwhile positive 192 and negative effects (15). To interpret the magnitude of the variability in maturity offset 193 within each squad, we doubled the standard deviation for each respective squad and 194 compared against a scale of 0.2 (small), 0.6 (moderate), and 1.2 (large) of the between-player 195 standard deviation across all squads (24). Finally, Pearson's correlations were used to 196 determine the relationship between player maturity and sprinting speed and the following 197 scale of magnitudes was used to interpret the magnitude of the correlation coefficients: <0.1,

trivial; 0.1-0.3, small; 0.3-0.5, moderate; 0.5-0.7, large; 0.7-0.9, very large; >0.9, nearly
perfect (15).

200

## 201 **RESULTS**

202

## 203 Age distribution

Odds ratio's revealed a clear bias in frequency, when compared to our reference population, of players born in quartile 1 versus quartile 4 within each playing squad. The magnitude of this bias was small for under-11s (Odds ratio 2.7; 90% confidence interval 1.7 to 4.3), under-12s (2.1; 1.4 to 3.2) and under-13s (3.1; 2.0 to 4.9), and moderate for under-14s (3.7; 2.3 to 6.0), under 15s (4.7; 2.6 to 8.7) and under 17s (4.3; 1.7 to 10.6).

209

# 210 Effect of birth quartile on player maturity, stature and mass

211 Descriptive anthropometry for each age category is presented in Table 1. The overall effect 212 (all squads combined) of birth quartile (quartile 1 versus quartile 4) was very likely small for 213 player maturity (0.85 years; 90% confidence interval 0.44 years to 1.26 years) and player 214 stature (6.2 cm; 90% confidence interval 2.8 cm to 9.6 cm), and likely small for player 215 weight (5.1 kg; 90% confidence interval 1.7 kg to 8.4 kg). Within-squad analyses for player 216 maturity, stature and mass are presented in Tables 2, 3, and 4, respectively; differences 217 ranged from unclear to large for player maturity and stature, and unclear to moderate for 218 player mass. After doubling the standard deviation of maturity offset within each playing 219 squad, the magnitude of variability was small for under-11s and under-12s, and moderate for 220 all remaining squads.

221

\*\*\*Insert Tables 1, 2, 3 and 4 near here\*\*\*

223

# 224 Relationship between player maturity and sprinting speed

The magnitude of the relationship between maturity offset and 15 m sprinting speed was trivial for under-11s (r = 0.01; 90% confidence interval -0.14 to 0.16) and under-12s (r = -0.04; -0.20 to 0.13), very likely small for under-13s (r = -0.26; -0.39 to -0.11), possibly large for under-14s (r = -0.53; -0.62 to -0.41), very likely large for under-15s (r = -0.62; -0.71 to -0.51), and likely small for under-17s (r = -0.26; -0.50 to 0.02).

230

### 231 **DISCUSSION**

232

233 The uneven birth date distribution observed was commensurate with that reported by many 234 others (13, 16). A widely reported explanation for the RAE phenomenon is the maturation-235 selection hypothesis, which proposes that relatively older players are more advanced in 236 physical maturity than their younger counterparts and that this confers a performance 237 advantage (27). This theory makes intuitive sense since it is well established that attributes 238 relevant to soccer performance such as sprinting speed, strength and aerobic capacity 239 improve during growth and maturation (18). However, the magnitude of the relationship 240 between physical maturity and such performance attributes within the context of one-year age 241 categories has not been widely investigated. Specifically, to our knowledge only three other 242 studies have assessed the practical relevance of the relationships between relative age, physical maturity and physical performance measures using magnitude based inferences (5, 243 244 17, 23).

245

Overall, physical maturity was related to chronological age, with older players displayinggreater maturity offset values, although the strength of the relationship differed depending on

248 the specific category considered (Table 2). This superior maturity status manifested itself as 249 both greater stature (Table 3) and mass (Table 4) up until the under-17 age category when the 250 trend was reversed, however, again the magnitude of the relationships varied depending on 251 age category. The stature and mass of the players in the present study were comparable to 252 results reported previously (17, 23). The strength of the relationships between stature, mass 253 and birth quartile increased from the under-11 ('likely small' for both stature and mass) 254 through to the under-15 age categories ('possibly moderate' for stature; 'likely moderate' for 255 mass) and then reversed among the under-17 players. This reversal should be interpreted with 256 caution since the number of under-17 players observed in the current study was small. This is 257 an interesting finding as it demonstrates that the influence of physical maturity is not 258 necessarily consistent throughout childhood and adolescence. Vaeyens et al. (26) also 259 reported that the influence of physical maturity on numerous performance parameters varied 260 depending on age category. Indeed, our analysis demonstrates that the magnitude of 261 variability in relation to maturity offset status differed between younger (under-11 and -12s) 262 and older (under-13 to -17s) players perhaps explaining some of the inconsistencies.

263

264 Similarly, the influence of physical maturity on 0-15 m sprinting speed varied depending on 265 age category. The greatest magnitudes were observed in the under-14 and -15 age categories 266 where physical maturity had a possible and very likely large positive effect respectively. 267 Combined with the fact that the older players in these two age categories were generally more 268 physically mature than their younger counterparts; the maturation-selection hypothesis 269 appears valid. It seems very plausible that scouts could associate physical precocity – in the 270 form of sprinting ability and physical dimensions - with 'talent' especially when one 271 considers how valuable a commodity speed is within the sport of soccer (10). The most 272 common action prior to scoring a goal at the professional level is straight-line sprinting,

273 highlighting the importance of this attribute (10). Adolescent boys typically pass through 274 their PHV around 14 years of age and peak weight velocity follows soon after (18, 22). The 275 greatest inter-individual discrepancies in stature and muscle mass are likely to occur around 276 the chronological age of 14 when some players will be pre- and others will be post-pubertal. 277 Beunen et al. (3) reported that differences in physical maturity between players influenced 278 physical performance to the greatest degree around the chronological ages of 14-15 years in 279 Belgian teenagers, reinforcing this theory. Maturity-associated differences between players at 280 this developmental stage are temporary and likely to diminish as less developed players 281 mature. Indeed, the present results hint at this, with minimal differences in sprinting speed 282 observed among players of differing physical maturity status within the under-17 age 283 category. The potential for players to be released from their clubs based on transient 284 maturational differences during early adolescence may result in a loss of available talent at 285 the upper echelons of the game when age categories are no longer important.

286

287 In contrast, the influence of physical maturation on sprinting speed within the younger age 288 categories (under-11 to -13s) was minimal. This suggests that relatively older and more 289 physically mature players in the earlier age categories were not selected because they were 290 faster than their younger counterparts. Within the younger age categories (under-11, -12 and -291 13s) the mean differences in stature and mass between those born in guarters one and four 292 were small to non-existent; ranging from one to four centimeters and one to two kilograms 293 respectively (see Tables 3 & 4). It is questionable whether such small differences could have 294 resulted in such a profound influence on selection. This raises the question; if differences in 295 stature, mass and sprinting ability are so small why were relatively older players 296 disproportionally chosen? At the elite youth level it may be that only the most biologically 297 advanced late-born players are considered for selection, thus, creating homogenous groups.

Gil et al. (12) reported superior sprinting ability, agility and stature among relatively older compared to relatively younger non-elite youth soccer players. The RAE may simply appear to be unrelated to physical capacity at the elite youth level because of the formation of homogenous groups.

302

303 The present results demonstrate some likeness to previous findings; however, some 304 discrepancies are apparent. Lovell et al. (17) found the greatest disparities in birthdate 305 distribution at the youngest age category observed (under-9) in addition to the age categories 306 around expected PHV (under-13 to -16s). The under-11 age category was the youngest 307 observed in the present study and so a direct comparison cannot be made, however, like 308 Lovell et al. (17) we observed the greatest RAE to be present among under-15 players. In 309 contrast to Lovell et al. (17) and Skorski et al. (23) we investigated the relationship between 310 physical maturity (rather than birth quartile directly) and sprinting ability. However, we also 311 demonstrated that physical maturity and birth quartile were likely related (Table 2). Lovell et 312 al. (17) reported superior anaerobic performance – including sprinting ability – among 313 relatively older players in the under-10 to -14 age categories. In contrast, the present results 314 indicate minimal advantages in sprinting ability related to relative age within the under-11 to 315 -13 age categories. The explanation for this discrepancy is unclear; however, it may be 316 attributable to differences in the sample populations. The data presented herewith originate 317 from a single academy whereas Lovell et al. (17) included data from 17 separate clubs. The 318 present data may be indicative of a particular selection strategy at the club in question. 319 However, since data were collected over the course of eight seasons any nuances related to 320 the club's selection strategy at least highlight a consistent approach. In addition, the academy 321 observed was attached to a Scottish top-division club whereas the club academies observed 322 by Lovell et al. (17) represented the third and fourth tier of English professional soccer.

# 324 PRACTICAL APPLICATIONS

The current results support the maturation-selection hypothesis but only at specific developmental stages (under-14 and 15s). However, questions remain especially within the earlier age categories; which are synonymous with players' initial selection into performance programmes. At the under-14 and under-15 age categories relatively older players were generally more mature and this manifested as faster sprinting speed. However, at the younger age categories while older players were generally more mature this did not translate to superior sprinting ability. Practitioners should be aware that the influence of physical maturity on sprinting speed varies throughout physical development. Crucially, it would appear that making decisions about which players to retain and release should not be based on sprinting ability around the under-14 and under-15 age categories since any inter-individual differences may be confounded by transient inequalities in physical maturity status. 

### 348 **REFERENCES**

349

Balyi I, Way R, and Higgs C. *Long-term athlete development*. Champaign, IL:
 Human Kinetics, 2013.

352

Batterham AM and Hopkins WG. Making meaningful inferences about magnitudes.
 *Int J Sports Physiol Perform* 1(1): 50-57, 2006.

- Beunen GP, Malina RM, Lefevre J, Claessens AL, Renson R, and Simons J.
   Prediction of adult stature and noninvasive assessment of biological maturation. *Med Sci Sports Exerc* 29(2): 225-230, 1997.
- 359
- 360
  4. Buchheit M, and Mendez-Villanueva A. Reliability and stability of anthropometric
  361
  and performance measures in highly-trained young soccer players: effect of age and
  362
  maturation. *J Sports Sci* 31(12): 1332-1343, 2013.
- 363
- 364 5. Buchheit M, and Mendez-Villanueva A. Effects of age, maturity and body
  365 dimensions on match running performance in highly trained under-15 soccer players.
  366 *J Sports Sci* 32(13): 1271-1278, 2014.
- 367
- Buchheit M, Mendez-Villanueva A, Mayer N, Jullien H, Marles A, Bosquet L, Maille
  P, Morin JB, Cazorla G, and Lambert P. Locomotor performance in highly-trained
  young soccer players: Does body size always matter? *Int J Sports Med* 35(6): 494504, 2014.
- 372

373	7.	Carling C, le Gall F, Reilly T, and Williams AM. Do anthropometric and fitness
374		characteristics vary according to birth date distribution in elite youth academy soccer
375		players? Scand J Med Sci Spor 19(1): 3-9, 2009.
376		
377	8.	Cobley S, Baker J, Wattie N, and McKenna J. Annual age-grouping and athlete
378		development: a meta-analytical review of relative age effects in sport. Sports Med
379		39(3): 235-256, 2009.
380		
381	9.	Deprez D, Coutts AJ, Fransen J, Deconinck F, Lenoir M, Vaeyens R, and Philippaerts
382		R. Relative age, biological maturation and anaerobic characteristics in elite youth
383		soccer players. Int J Sports Med 34(10): 897-903, 2013.
384		
385	10.	Faude O, Koch T, and Meyer T. Straight sprinting is the most frequent action in goal
386		situations in professional football. J Sports Sci 30(7):625-631, 2012.
387		
388	11.	Figueiredo AJ, Gonçalves CE, Coelho E Silva MJ, and Malina RM. Youth soccer
389		players, 11-14 years: Maturity, size, function, skill and goal orientation. Ann Hum
390		<i>Biol</i> 36(1): 60-73, 2009.
391		
392	12.	Gil SM, Badiola A, Bidaurrazaga-Letona I, Zabala-Lili J, Gravina L, Santos-
393		Concejero J, Lekue JA, and Granados C. Relationship between the relative age effect
394		and anthropometry, maturity and performance in young soccer players. J Sports Sci
395		32(5): 479-486, 2014.
396		
397	13.	Helsen WF, Baker J, Michiels S, Schorer J, Van Winckel J, and Williams AM. The

398	relative age effect in European professional soccer: did ten years of research make
399	any difference? J Sports Sci 30(15):1665-1671, 2012.
400	
401	14. Helsen WF, van Winckel J, and Williams AM. The relative age effect in youth soccer
402	across Europe. J Sports Sci 23(6): 629-636, 2005.
403	
404	15. Hopkins WG, Marshall SW, Batterham AM, and Hanin J. Progressive statistics for
405	studies in sports medicine and exercise science. Med Sci Sports Exerc 41(1): 3-13,
406	2009.
407	
408	16. Jiménez IP, and Pain MT. Relative age effect in Spanish association football: Its
409	extent and implications for wasted potential. J Sports Sci 26(10): 995-1003, 2008.
410	
411	17. Lovell R, Towlson C, Parkin G, Portas M, Vaeyens R, and Cobley S. Soccer player
412	characteristics in English lower-league development programmes: The relationships
413	between relative age, maturation, anthropometry and physical fitness. PLoS One
414	10(9), 2015. [Epub Ahead of Print]
415	
416	18. Malina RM, Bouchard C, and Bar-Or O. Growth, Maturation and Physical Activity.
417	Champaign, IL: Human Kinetics, 2004.
418	
419	19. Malina RM, Rogol AD, Cumming SP, Coelho e Silva MJ, and Figueiredo AJ.
420	Biological maturation of youth athletes: assessment and implications. Br J Sports
421	<i>Med</i> 49: 852-859, 2015.
422	

423	
424	20. Mirwald RL, Baxter-Jones AD, Bailey DA, and Beunen GP. An assessment of
425	maturity from anthropometric measurements. Med Sci Sports Exerc 34(4): 689-694,
426	2002.
427	
428	21. Musch J, and Grondin S. Unequal competition as an impediment to personal
429	development: A review of the relative age effect in sport. Dev Rev 21(2): 147-167,
430	2001.
431	
432	22. Philippaerts RM, Vaeyens R, Janssens M, van Renterghem B, Matthys D, Craen R,
433	Bourgois J, Vrijens J, Beunen G, and Malina RM. The relationship between peak
434	height velocity and physical performance in youth soccer players. J Sports Sci 24(3):
435	221-230, 2006.
436	
437	23. Skorski S, Skorski S, Faude O, Hammes D, and Meyer T. The relative age effect in
438	German elite youth soccer: Implications for a successful career. Int J Sports Physiol
439	Perform 11(3): 370-376, 2015.
440	
441	24. Smith TB, and Hopkins WG. Variability and predictability of finals times of elite
442	rowers. Med Sci Sports Exerc 43(11): 2155-2160, 2011.
443	
444	25. Stratton G, Reilly T, Williams AM, and Richardson D. Youth Soccer: From science
445	to performance. Abingdon: Routledge, 2004.
446	
447	26. Vaeyens R, Malina RM, Janssens M, van Renterghem B, Bourgois J, Vrijens J, and

448	Philippaerts RM. A multidisciplinary selection model for youth soccer: the Ghent
449	youth soccer project. Br J Sports Med 40(11): 928-934, 2006.
450	
451	27. Wattie N, Schorer J, and Baker J. The relative age effect in sport: a developmental
452	systems model. Sports Med 45(1):83-94, 2015.
453	
454	
455	
456	
457	
458	
459	
460	
461	
462	
463	
464	
465	
466	
467	
468	
469	
470	
471	
472	

# 473 ACKNOWLEDGMENTS

- 475 The authors would like to thank \*\*\*name removed for blind review\*\*\* for his long-term
- 476 support and encouragement during the data collection period.

- . -

Table 4. Within-squad comparisons	for the	effect	of birth	quartile	(quartile	1	versus
quartile 4) on player mass.							

Squad	Quartile 1	Quartile 4	Mean	Qualitative
	Mean $\pm$ SD	Mean $\pm$ SD	difference	inference
	(kg)	(kg)	(90% CI)	
Under 11's	35.1 ± 3.8	33.1 ± 3.0	2.0	Likely small
(Q1 n=47, Q4 n=16)			(-0.2 to 4.2)	
Under 12's	$36.8\pm4.6$	$37.0 \pm 4.1$	-0.2	Unclear
(Q1 n=40, Q4 n=21)			(-2.6 to 2.2)	
Under 13's	$41.9\pm7.7$	$40.7 \pm 3.5$	1.2	Unclear
(Q1 n=53, Q4 n=17)			(-1.9 to 4.3)	
Under 14's	51.3 ± 9.8	$47.2 \pm 4.8$	4.1	Likely small
(Q1 n=54, Q4 n=12)			(-0.4 to 8.6)	
Under 15's	$61.2 \pm 9.1$	$54.3 \pm 4.5$	7.0	Likely moderate
(Q1 n=37, Q4 n=9)			(2.3 to 11.6)	
Under 17's	$65.4 \pm 6.2$	74.9 ± 15.5	-9.5	Likely moderate
(Q1 n=16, Q4 n=3)			(-19.0 to -0.1)	

Table 3. Within-squad comparisons for t	ne effect of birth quartile (quartile 1 versus
quartile 4) on player stature.	

Squad	Quartile 1	Quartile 4	Mean	Qualitative
	Mean $\pm$ SD	Mean $\pm$ SD	difference	inference
	(cm)	(cm)	(90% CI)	
Under 11's	$143.7 \pm 3.4$	$139.5 \pm 3.8$	4.2	Likely small
(Q1 n=47, Q4 n=16)			(1.9 to 6.6)	
Under 12's	$146.9 \pm 5.5$	$145.9\pm4.8$	1.0	Unclear
(Q1 n=40, Q4 n=21)			(-1.6 to 3.5)	
Under 13's	$154.1 \pm 6.0$	$151.5 \pm 3.6$	2.6	Likely small
(Q1 n=53, Q4 n=17)			(-0.1 to 5.4)	
Under 14's	$164.7 \pm 7.2$	$159.9 \pm 4.5$	4.7	Possibly moderate
(Q1 n=54, Q4 n=12)			(1.2 to 8.3)	
Under 15's	$172.4 \pm 6.6$	$168.3 \pm 4.6$	4.2	Possibly moderate
(Q1 n=37, Q4 n=9)			(0.2 to 8.1)	
Under 17's	$175.2 \pm 4.8$	$181.6 \pm 7.2$	-6.4	Possibly large
(Q1 n=16, Q4 n=3)			(-11.7 to -1.0)	
CI, confidence interval	; Q, quartile			

Table 2. Within-squad comparisons for the effect of birth quartile (quartile 1 versus
quartile 4) on maturity (as measured by the maturity offset equation).

Squad	Quartile 1	Quartile 4	Mean difference	Qualitative
	$Mean \pm SD$	Mean $\pm$ SD	(90% CI)	inference
	(years)	(years)		
Under 11's	$-2.69 \pm 0.25$	$-3.11 \pm 0.33$	0.42	Possibly large
(Q1 n=47, Q4 n=16)			(0.28 to 0.57)	
Under 12's	$-2.10 \pm 0.39$	$-2.39 \pm 0.35$	0.29	Possibly moderate
(Q1 n=40, Q4 n=21)			(0.11 to 0.47)	
Under 13's	$-1.24 \pm 0.26$	$-1.64 \pm 0.36$	0.40	Likely moderate
(Q1 n=53, Q4 n=17)			(0.19 to 0.61)	
Under 14's	$-0.02 \pm 0.62$	$-0.65 \pm 0.44$	0.63	Likely moderate
(Q1 n=54, Q4 n=12)			(0.33 to 0.94)	
Under 15's	$1.16 \pm 0.61$	$0.34 \pm 0.49$	0.82	Likely large
(Q1 n=37, Q4 n=9)			(0.50 to 1.13)	
Under 17's	$2.10 \pm 0.48$	$2.35 \pm 1.10$	-0.25	Unclear
			(-0.88 to 0.39)	

Squad	Stature Mean ± SD (cm)	Seated stature* Mean ± SD (cm)	Mass Mean ± SD (kg)	Maturity offset Mean ± SD (years)					
					Under 11's	$142.7 \pm 5.1$	$115.2 \pm 2.6$	$34.9\pm4.6$	$-2.80 \pm 0.33$
					(n=120)				
Under 12's	$147.4 \pm 5.8$	$117.1 \pm 3.0$	$38.0 \pm 5.5$	$-2.16 \pm 0.41$					
(n=96)									
Under 13's	$153.6 \pm 6.0$	$119.9 \pm 3.3$	$41.9 \pm 5.9$	$-1.34 \pm 0.46$					
(n=105)									
Under 14's	$163.9 \pm 6.9$	$125.0 \pm 3.9$	51.2 ± 8.6	$-0.20 \pm 0.59$					
(n=111)									
Under 15's	$171.8 \pm 6.6$	$130.1 \pm 3.7$	$60.5 \pm 7.8$	$0.98\pm0.57$					
(n=99)									
Under 17's	$174.7 \pm 5.4$	132.1 ± 3.7	$66.0 \pm 9.4$	$1.94 \pm 0.68$					
(n=39)									

Table 1. Descriptive anthropometric data for each age category.

\*Seated stature was measured with participants sitting on a 40cm wooden box