

Poster presentation

Open Access

## Hyper-contractility and impaired cGMP signaling in the BK<sub>Ca</sub> channel deletion model of erectile dysfunction

Matthias E Werner\*<sup>1</sup>, Andrea L Meredith<sup>2</sup>, Richard W Aldrich<sup>3</sup> and Mark T Nelson<sup>4</sup>

Address: <sup>1</sup>Division of Cardiovascular and Endocrine Science, School of Medicine, University of Manchester, Manchester M13 9NT, UK, <sup>2</sup>Department of Physiology, School of Medicine, University of Maryland, Baltimore, MD 21201, USA, <sup>3</sup>Section of Neurobiology, 1 University Station C7000, The University of Texas at Austin, Austin, TX 78712, USA and <sup>4</sup>Department of Pharmacology, College of Medicine, University of Vermont, Burlington, VT 05405, USA

Email: Matthias E Werner\* - [Matthias.Werner@manchester.ac.uk](mailto:Matthias.Werner@manchester.ac.uk)

\* Corresponding author

from 3<sup>rd</sup> International Conference on cGMP Generators, Effectors and Therapeutic Implications  
Dresden, Germany. 15–17 June 2007

Published: 25 July 2007

BMC Pharmacology 2007, 7(Suppl 1):P65 doi:10.1186/1471-2210-7-S1-P65

This abstract is available from: <http://www.biomedcentral.com/1471-2210/7/S1/P65>

© 2007 Werner et al; licensee BioMed Central Ltd.

Erectile dysfunction (ED) is frequently elicited by a multiplicity of pathogenic factors, predominantly by impaired formation of and responsiveness to nitric oxide (NO) and the downstream effectors soluble guanylate cyclase (sGC) and cGMP-dependent protein kinase I (PKG1). In smooth muscle, one important target of PKGI is the large conductance, calcium-sensitive potassium (BK<sub>Ca</sub>) channel, which upon activation hyperpolarizes the smooth muscle cell membrane, causing relaxation. In our earlier report [1], we demonstrated that ablation of the gene, encoding for the pore-forming  $\alpha$  subunit of the BK<sub>Ca</sub> channel in mice (*Slo*<sup>-/-</sup>) induced an increase of corpus cavernosum smooth muscle (CCSM) force oscillations, led to reduced nerve-evoked relaxations and ED. In our current work, we used this ED model to explore the role of the BK<sub>Ca</sub> channel in the NO/cGMP pathway. Electrical field stimulation (EFS)-induced contractions of CCSM strips from *Slo*<sup>-/-</sup> mice demonstrated a 53% increase that could be reduced by sildenafil similar to levels observed in strips from wild-type (*Slo*<sup>+/+</sup>) mice. In *Slo*<sup>-/-</sup> strips precontracted with phenylephrine (PE), SNP and sildenafil induced relaxations, which were diminished by 10% and 7% over *Slo*<sup>+/+</sup>, respectively. Neither SNP nor sildenafil was able to reduce the enhanced force oscillations, which were induced by the loss of BK<sub>Ca</sub> channel function. Yet, these oscillations could be completely elim-

inated by blocking L-type voltage-dependent calcium channels (VDCCs). The latter results indicate that loss of BK<sub>Ca</sub> channel leads to ED and hyper-contractility likely due to instability of membrane potential which activates VDCCs. Moreover, since the relaxing effects of SNP and sildenafil were reduced in *Slo*<sup>-/-</sup>, the ED phenotype in our BK<sub>Ca</sub> channel deletion model could also be the consequence of an impaired NO/cGMP signaling pathway.

### References

1. Werner ME, Zvara P, Meredith AL, Aldrich RW, Nelson MT: **Erectile dysfunction in mice lacking the large conductance calcium-activated (BK) channel.** *J Physiol* 2005, **567**:545-556/Pkt.2.