

1
2
3 **Residual colonization by vaccine serotypes in rural South Africa four years**
4
5
6 **following initiation of pneumococcal conjugate vaccine immunization**
7
8

9 Shabir A. Madhi,^{1,2} Susan A. Nzenze,^{1,2} Marta C. Nunes,^{1,2} Lilian Chinyanganya,^{1,2} Nadia
10 van Niekerk,^{1,2} Kathleen Kahn,^{3,4,5} Rhine Twine,³ Linda de Gouveia,⁶ Anne von Gottberg,⁶
11
12
13 Tinevimbo Shiri ^{1,2,7}
14
15

16
17 ¹Department of Science and Technology/National Research Foundation: Vaccine Preventable
18 Diseases, University of the Witwatersrand, Johannesburg, South Africa; ²Medical Research
19 Council: Respiratory and Meningeal Pathogens Research Unit, School of Pathology, Faculty
20 of Health Sciences University of the Witwatersrand, Johannesburg, South Africa; ³MRC/Wits
21 Rural Public Health and Health Transitions Research Unit, School of Public Health, Faculty
22 of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa; ⁴Centre for
23 Global Health Research, Umeå University, Umeå, Sweden; ⁵INDEPTH Network, Accra,
24 Ghana; ⁶Centre for Respiratory Diseases and Meningitis, National Institute for
25 Communicable Diseases, National Health Laboratory Service, Johannesburg, South Africa;
26
27 ⁷Liverpool School of Tropical Medicine, Liverpool, United Kingdom
28
29
30
31
32
33
34
35
36
37
38
39

40 Corresponding author:

41
42
43 Shabir A Madhi

44
45
46 Respiratory and Meningeal Pathogens Research Unit, Chris Hani Road; Chris Hani-
47 Baragwanath Hospital, New Nurses Residence, 11th Floor West Wing
48

49
50
51 2013 Bertsham,

52
53
54 South Africa

55
56
57 E-mail: madhis@rmpru.co.za; Telephone: +2711 983 4266
58
59
60

1
2
3 **Funding:** Funding for this project was provided by the South Africa Department of Science and
4
5 Technology and the National Research Foundation (South African Research Chair Initiative in Vaccine
6
7 Preventable Diseases) and the Respiratory and Meningeal Pathogens Research Unit of the Medical
8
9 Research Council of South Africa.
10

11
12 **Declaration of interest:** S A Madhi received research funding and honoraria from Pfizer and
13
14 GlaxoSmithKline. M C Nunes received honoraria from Pfizer. A von Gottberg received research
15
16 funding from Pfizer. The authors have no other relevant affiliations or financial involvement with any
17
18 organization or entity with a financial interest in or financial conflict with the subject matter or
19
20 materials discussed in the manuscript apart from those disclosed.
21
22

23
24 **Reviewer Disclosures:** A reviewer on this manuscript has disclosed grant funding and consulting fees
25
26 from Pfizer and Merck. All other peer reviewers on this manuscript have no relevant financial or
27
28 other relationships to disclose.
29
30

31
32 **Author Contributions:** S A Madhi and S A Nzenze designed the trial and study protocol. S A Nzenze, L
33
34 Chinyanganya, K Kahn, R Twine, oversaw the clinical data collection and clinical data management.
35
36 N van Niekerk, L de Gouveia, A von Gottberg were responsible for laboratory analyses. S A Madhi, SA
37
38 Nzenze and T Shiri conducted the statistical analysis. S A Madhi wrote the first draft of the paper
39
40 with input from S A Nzenze, M C Nunes and T Shiri. All authors inputted to subsequent drafts, read
41
42 and approved the final version of the report.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Background: We evaluated pneumococcal colonization in children and adults between the time of 7-valent pneumococcal conjugate vaccine (PCV) introduction in the immunization program in April 2009 to two years after transitioning to 13-valent PCV in 2011.

Methods: Community-based pneumococcal carriage surveillance was undertaken between May-November 2013 (Period-3; n=1884), with similar surveys in 2009 (Period-1, n=2010) and 2011 (Period-2; n=3659). Households with children below two years had a similar probability of being sampled in all surveys. Nasopharyngeal swabs were processed using standard methods and serotyped by Quellung.

Results: In children >9-59 months of age, overall pneumococcal colonization prevalence declined from 81.8% in Period-1 to 65.0% in Period-3 ($p<0.001$). Reductions of 70% (95%CI: 60%-77%; 41.2% vs. 13.6%) in PCV7-serotypes colonization and 66% (95%CI:48%-78%; 15.3% vs. 4.4%) for the six additional PCV-serotypes in PCV13 (PCV13-add6VT) were observed. There was, however, high residual prevalence of colonization by PCV7-serotypes 19F (14.9% vs. 6.3%) and 23F (8.5% vs. 4.1%), despite reduction of 57% (95%CI:35%-80%) and 52% (95%CI:21%-83%), respectively. Among individuals >12 years of age, there was 61% (95%CI:18%-82%) reduction in PCV7-serotype colonization (3.1% vs. 1.3%;) and 75% (95%CI: 11%-93%) decrease for PCV13-add6VT (2.1% vs. 0.6%) between Period-1 and Period-3.

Conclusions: The residual prevalence of serotypes 19F and 23F in PCV-immunized and -unvaccinated age-groups, four years after introducing PCV in the South African public immunization program, suggests ongoing community transmission and transient vaccine effects.

1
2
3 Keywords: Streptococcus pneumoniae, colonization, carriage, pneumococcal conjugate
4
5 vaccine, serotypes
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

1. Introduction

Pneumococcal conjugate vaccine (PCV) immunization of children reduces their risk of *Streptococcus pneumoniae* (pneumococcus) vaccine-serotype nasopharyngeal acquisition and disease^{1,2}. Since young children are the major source of transmission of pneumococci in the community^{3,4}, childhood PCV immunization has also reduced the prevalence of vaccine-serotype colonization among unvaccinated age groups^{2,5-7}. Although there has been near elimination of PCV-serotype colonization and invasive disease in some high-income countries following childhood PCV immunization^{8,9}, there are limited data from low-middle income countries^{10,11}, including from settings with high prevalence of HIV infection. Monitoring the prevalence of pneumococcal vaccine-serotype colonization, a pre-requisite for developing pneumococcal disease¹², is a potential metric for assessing the indirect effect of childhood PCV-immunization against vaccine-serotype pneumococcal disease in older, non-immunized age-groups¹.

The 7-valent PCV (PCV7) was introduced in the South African public immunization program in April 2009, using a 6, 14 and 40 weeks of age dosing schedule; with no catch-up campaign of older children. In May 2011, the immunization program transitioned to 13-valent PCV (PCV13), with a limited catch-up campaign for children less than 36 months (m.) of age and in those 36-71m. with underlying medical conditions. We previously reported, in a rural setting with high HIV prevalence, within two years of PCV7 introduction and when PCV coverage in children was modest (52%), a 50-64% reduction in PCV7-serotype colonization prevalence in age-groups targeted and not targeted for PCV immunization¹³.

The aim of this study was to evaluate the prevalence of vaccine-serotype and non-vaccine serotype (NVT) pneumococcal nasopharyngeal colonization in children (>9-59m. and 60-144m. age) and household members older than 12 years (y.) of age in a rural African setting,

1
2
3 two years following transitioning from PCV7 to PCV13 in the infant public immunization
4
5 program.
6

7 8 2.0 Methods 9

10 11 2.1 Study population and study methods 12

13
14 This community-based study was undertaken in Agincourt sub-district, a health-socio
15
16 demographic surveillance site (HDSS) in Mpumalanga province, South Africa. The
17
18 pneumococcal colonization survey performed from May to November 2013 (Period-3) was
19
20 preceded by two earlier cross-sectional surveys approximating the same calendar months in
21
22 2009 (Period-1) and 2011 (Period-2) as described ¹³. Aside from children <9m. not being
23
24 enrolled during Period-1, the sampling strategy and methodology was similar across the three
25
26 surveillance periods ¹³. Briefly, households with children aged ≤ 24 m. were identified from
27
28 the HDSS database as potential participants. A random list of villages was generated and
29
30 households with children less than two years old were selected until the required sample size
31
32 was achieved. A total of 9, 26 and 19 villages were selected in 2009, 2011 and 2013,
33
34 respectively. All 9 villages included in 2009 were also included in 2011, and 16 villages
35
36 selected in 2011 were also included in sampling during to 2013. Households were included if,
37
38 in addition to the listed child, there was at least one individual >12y. who agreed to study-
39
40 participation. Although the same household could have been randomly selected in any of the
41
42 three sampling periods, the same child would generally have contributed to an older child
43
44 age-group stratum.
45
46
47
48
49

50
51 A questionnaire was administered at the time of nasopharyngeal swab collection. HIV status
52
53 was based on self-reporting in the first two study-periods. In Period-3, HIV testing following
54
55 counselling was offered to individuals older than 18y. HIV testing was not undertaken in the
56
57 younger age-group due to an anticipated low HIV prevalence in individuals aged <18y, and
58
59
60

1
2
3 consequently we relied on self-reporting by parents/care-givers for HIV-infection status for
4 these individuals. This was premised on the successful roll-out of the mother-to-child HIV
5 prevention program between 2008 and 2011, following which the mother-to-child HIV
6 transmission rates declined from 9.6% to 2.3%.¹⁴
7
8
9
10
11

12
13 The sample collection and laboratory testing methods for culture of bacteria, including
14 *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Staphylococcus aureus*, were as
15 previously reported^{13,15}. Briefly, nasopharyngeal swabs were performed in all subjects using
16 an aluminum shafted, Dacron swab (MW and E, Medical Wire and Equipment Co. Ltd.,
17 Corsham, Wiltshire, England), in addition to which an oropharyngeal swab was collected
18 from those older than 12y. Samples were processed at the Centre for Respiratory Diseases
19 and Meningitis (CRDM) at the National Institute for Communicable Diseases, South Africa,
20 by routine microbiological methodologies. Serotyping was undertaken by the Quellung
21 method using factor and serotype-specific antisera (Statens Serum Institute, Copenhagen,
22 Denmark). Presumptive pneumococcal presenting no phenotypically detectable capsule were
23 categorized as non-typeable, once pneumococcal identification was confirmed with *lytA* PCR.
24 Where >1 distinct morphological colony type was present, each colony was serotyped.
25 Serotypes 6A, 6B, 6C and 6D were distinguished only using the relevant antisera and without
26 any additional molecular typing.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 2.2 Sample size calculation

47
48 The sample size for Period-3 was based on providing 80% power to detect a 30% reduction in
49 PCV13-no-PCV7 serotypes, i.e. PCV13-additional six serotypes (PCV13-add6VT)
50 colonization among individuals >12y. age (referred to as “adults” henceforth) relative to the
51 1.1% prevalence documented in 2011. This required enrolment of 900 adults. We estimated
52 that to enroll 900 adults, we would require visiting at least 450 households with at least one
53
54
55
56
57
58
59
60

1
2
3 child aged <24m. Based on PCV13-add6VT colonization prevalence of 15.2% in 2011
4 among children aged <24m., the enrolled number of children <24m. age (n=434) would
5 provide 80% power to detect at least a 40% reduction in prevalence of PCV13-add6VT
6 colonization in this age-group.
7
8
9
10

11 12 13 2.3 Statistical analysis 14

15
16 Multiple, simultaneous colonization by different serotypes in the same individual were
17 considered as independent events when measuring the prevalence of colonization if they
18 differed in stratification into PCV7-serotypes (4, 6B, 9V, 14, 18C, 19F and 23F), PCV13-
19 add6VT (1, 3, 5, 6A, 7F and 19A) or non-vaccine serotypes (i.e. any other pneumococcal
20 serotype including non-typeable isolates). Comparison of the prevalence of serotype-specific
21 pneumococcal colonization between Period-3 and Period-1 were calculated using χ^2 or
22 Fisher's exact test where appropriate. Serotype-specific analysis was not adjusted for any
23 potential confounders. Log-binomial regression models with the log link function were used
24 to estimate the risk ratios for pneumococcal carriage in Period-3 compared to Period-1,
25 adjusting for potential confounding factors such as age, gender, fuel used for cooking in the
26 household, presence of a child attending day care and household structure. Statistical analyses
27 were conducted using SAS version 9.4 (SAS Institute, Inc., NC).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 2.4 Ethics 45

46
47 The study was approved by the Human Research Ethics Committee (Medical, M130461) at
48 the University of the Witwatersrand and the Mpumalanga Department of Health Ethics
49 Committee. Informed written consent was obtained from participants ≥ 18 y. old, and
50 parental/guardian consent was obtained for younger participants. In addition, those aged 8-
51 17y. provided verbal assent for study participation.
52
53
54
55
56
57
58

59 3.0 Results 60

1
2
3 We enrolled 2010 (from 577 households), 3659 (1079 households) and 1884 (563
4 households) individuals in 2009 (Period-1), 2011 (Period-2) and 2013 (Period-3),
5
6 respectively; as detailed in Table 1. Differences in demographic characteristics in Period-3
7
8 compared to Period-1 included fewer individuals per household (mean 5.6 vs. 8.8; $p < 0.001$),
9
10 including fewer children ≤ 5 y. age (mean 1.5 vs. 1.8; $p = 0.014$). Also, households were less
11
12 likely to use coal/wood for cooking (43.8% vs. 69.9%), and conversely more likely to use
13
14 electricity (55.8% vs. 28.2%; $p < 0.001$); Table 1. Children were less likely to ever have been
15
16 breastfed in Period-3 (82.5%) compared to Period-1 (92.0%; $p < 0.001$); albeit similar in the
17
18 12-24m. age-group (81.8% vs. 85.7%). Children were less likely to be on antibiotics at time
19
20 of sampling in Period-3 (1.5%) than Period-1 (7.7%; $p = 0.002$). The prevalence of parental-
21
22 reported HIV-positivity in their children was comparable across the study periods (3.0 to
23
24 3.2%). None of the children were immunized in Period-1. In Period-3, 65.2% of children 3-
25
26 9m. age received at least two PCV doses; and 82.2% of those > 9 -24m. received at least three
27
28 PCV doses, as did 49.8% in the 24-59m. age-group; Table 1. None of the older children (60-
29
30 144m.) were immunized with PCV in any of the three periods.

31
32
33 Among the adults (> 12 y. age), the mean age was lower in Period-3 (31.8y.) than Period-1
34
35 (33.3y.; $p = 0.040$), albeit unlikely to be of clinical significance. The prevalence of self-
36
37 reported HIV-positivity was higher in Period-3 (22.0%) than Period-1 (7.9%; $p < 0.001$),
38
39 including those > 18 y. age (8.3% vs. 23.4%). Among adults > 18 y. old who consented for HIV
40
41 testing in Period-3 ($n = 456$), the HIV-positivity prevalence was 20.2% ($n = 92$); including
42
43 being positive in 100% ($n = 50$) of those who self-reported to be HIV-infected, and positive in
44
45 a further 36 individuals who self-reported to be HIV-uninfected. Of the self-reported HIV-
46
47 infected individuals, 55.3% and 70.8% were on antiretroviral therapy in Period-1 and Period-
48
49 3, respectively. Generally, there was a low prevalence of smoking (1.7 to 2.2%), sniffing of
50
51 tobacco (2.0 to 2.7%) or alcohol use (6.4% to 10.5%) across the study-periods. A higher
52
53
54
55
56
57
58
59
60

percentage of individuals in Period-3 (19.3%) than Period-1 (13.9%; $p=0.002$) had an underlying chronic medical illness, and 0.9 to 1.0% were on treatment for tuberculosis; Table 1.

3.1 Overall prevalence of pneumococcal colonization

Across all age-groups, the prevalence of overall pneumococcal colonization was lower in Period-3 (35.8%) than Period-1 (42.4%; adjusted risk ratio [aRR]: 0.81, $p < 0.001$); Table 2 and Figure 1. This included lower prevalence in Period-3 compared to Period-1 in children aged >9m-59m. (65.0% vs. 81.8%; aRR: 0.81, $p < 0.001$) who were eligible to have been vaccinated by Period-3, as well as in the PCV-naïve 60-144m. age-group (51.8% vs. 60.0%; aRR: 0.82, $p=0.03$); Table 2.

In individuals >12y. age, overall prevalence of pneumococcal colonization did not differ between Period-3 (10.6%) and Period-1 (11.3%); although a significant reduction was observed in the >12-18y age group (22.8% vs. 15.3%; aRR: 0.58, $p=0.04$); Table 2. This was in contrast to our previous observation of reduction in overall pneumococcal colonization in the >12y. age-group between Period-1 (11.3%) and Period-2 (6.8%; aRR: 0.53); including in the >12-18y. (22.8% vs. 12.3%; aRR: 0.51) and >18-45y. (10.7% vs. 5.8%; aRR:0.44) age-groups.

3.2 Prevalence of PCV7-serotype colonization

Overall prevalence of PCV7-serotype colonization declined by 66.0% (95% CI: 56% - 73%) between Period-1 (18.3%) and Period-3 (6.8%); including further reduction from Period-2 (11.4%) to Period-3 (aRR: 0.58, $p < 0.001$). The lower prevalence of PCV7-serotype colonization in Period-3 (13.6%) compared to Period-1 (41.2%; aRR: 0.30, $p < 0.001$) among children >9-59m. age, was evident in the >9-23m. (13.4% vs. 45.1%; aRR: 0.26, $p < 0.001$) and 24-59m. (14.7% vs. 35.5%; aRR: 0.37, $p < 0.001$) age-groups. There was also a 41% (95%

1
2
3 CI: 2% - 64%) reduction of PCV7-serotype colonization in Period-3 (10%) compared
4
5 Period-1 (19.0%) in the PCV-naïve 60-144m. age-group.
6
7

8 There was 61% (95% CI: 18% - 82%) lower prevalence of PCV7-serotype colonization in the
9
10 >12y. age-group in Period-3 (1.3%) compared to Period-1 (3.1%); although no further
11
12 significant decline was observed between Period-2 (1.2%) and Period-3 (aRR: 0.86, p=0.73).
13

14
15 Reductions in PCV7-colonization from Period-1 to Period-3 were detected in the 18-45y.
16
17 (3.0% vs. 1.0%; aRR: 0.21, p=0.02) age-group, whilst the reduction in the 12-18y. age-group
18
19 (5.7% vs. 3.4%; aRR: 0.64, p=0.42) was not significant; Table 2.
20
21

22 23 4.3 Prevalence of PCV13 additional-6 serotypes colonization 24

25
26 Overall and across specific age-groups, the prevalence of PCV13-add6VT colonization was
27
28 similar in Period-1 and Period-2; Table 2. Within two-years of transitioning from PCV7 to
29
30 PCV13, the overall prevalence of PCV13-add6VT colonization declined by 66% (95% CI:
31
32 51% - 77%) (Period-1 7.9% vs. Period-3 2.8%). The reduction in PCV13-add6VT
33
34 colonization was evident across all age-groups (albeit not analyzable for >45y. age). In those
35
36 >9-59m. age, PCV13-add6VT colonization decreased by 66% (95% CI: 48%-78%), from
37
38 15.3% in Period-1 to 4.4% in Period-3; including reductions in the >9-23m. (72% (95% CI:
39
40 50% - 84%)) and 24-59m. (60% (95% CI: 24% - 79%)) age-groups. A 59% (95% CI: 19% -
41
42 80%) reduction in PCV13-add6VT colonization was also observed in Period-3 (5.6%)
43
44 compared to Period-1 (11.0%) in the PCV-naïve 60-144m. age-group.
45
46
47

48
49 Likewise, the prevalence of PCV13-add6VT colonization was 75% (95% CI: 11% - 93%)
50
51 lower in the >12y. age-group in Period-3 (0.6%) compared to Period-1 (2.1%); including
52
53 89% (95% CI: 4% - 99%) reduction in the 12-18y. (1.1% vs. 5.7%) and 73% (95% CI: 12% -
54
55 91%) decline in the 18-45y. age-group (0.7% vs. 1.9%); Figure 1 and Table 2.
56
57

58 59 3.4 Prevalence of non-PCV13 serotype colonization 60

1
2
3 Although the overall prevalence of NVT colonization was similar between Period-1 (18.4%)
4 and Period-2 (19.3%; aRR: 1.06, p=0.44), it increased to 26.8% by Period-3 (aRR: 1.49,
5 p<0.001). The increase in NVT colonization in Period-3 (47.8%) compared to Period-1
6 (29.5%, aRR: 1.84, p<0.001) was evident for children >9-59m. age, including 2.06 (95% CI:
7 1.63 – 2.61) and 1.60 (95% CI: 1.24 – 2.07) adjusted-fold increases in the >9-23m. (48.5%
8 vs. 27.1%, <0.001) and 24-59m. (49% vs. 33%, p=0.003) age-groups, respectively. There
9 was no change in NVT colonization in the 60-144m. age-group (33.2% in Period-1 vs. 37.2%
10 in Period-3, p=0.78).
11
12
13
14
15
16
17
18
19
20
21

22 Among individuals >12y. age although there was an initial decrease in NVT colonization
23 between Period-1 (6.6%) and Period-2 (4.7%; aRR: 0.57, p=0.004); there was a 113% (95%
24 CI: 47% - 212%) increase in NVT colonization by Period-3 (8.9%) compared to Period-2.
25 Although the prevalence of NVT in Period-3 was slightly higher than in Period-1, this was
26 not significant (aRR: 1.18, p=0.37); Table 2.
27
28
29
30
31
32

33 3.5 Serotype-specific colonization by age-group 34 35

36
37 Reductions in prevalence of PCV13-serotype colonization were observed for most serotypes
38 with $\geq 1.5\%$ prevalence in Period-1 (except 9V) in the >9-59m. age-group, with similar
39 observations when stratified by >9-23m. and 24-59m. age; Table 3. In the >9-59m. age-
40 group, this included reduction in colonization prevalence between Period-1 and Period-3 for
41 serotypes 6B (84.6%, p<0.001), 14 (95.4%, p<0.001), 18C (86.7%, p=0.007), 19F (57.7%,
42 p<0.001), 23F (51.8%; p=0.001), 6A (83.1%; p<0.001) and 19A (56.5%; p=0.007); Table 3.
43
44
45
46
47
48
49
50
51 There, however, remained a prominent residual colonization prevalence in Period-3 of PCV7
52 (13.6%) and PCV13-add6VT (4.4%), with the dominant vaccine-serotypes including 19F
53 (6.3%), 23F (4.1%), 19A (2.0%), 6B (1.7%) and 6A (1.5%), which were also the top-five
54
55
56
57
58
59
60 PCV13 colonizing serotypes in Period-1 (14.9%, 8.5%, 4.6%, 10.1% and 8.9%, respectively);

1
2
3 Table 3. Furthermore, significant increases in NVT colonization between Period-1 and
4
5 Period-3 in the >9-59m. age-group included serotypes 15A (0.6% vs. 2.4%; $p=0.006$), 16F
6
7 (0.7% vs. 5.6%; $p<0.001$), 21 (0.1% vs. 1.7%; $p=0.003$), 34 (2.7% vs. 6.8%; $p<0.001$) and
8
9 35B (0.6% vs. 4.7%; $p<0.001$).

10
11
12
13 In the PCV-naïve 60-144m. age-group, although significant reductions were observed for
14
15 colonization by vaccine-serotypes 4 (100%; $p=0.007$) and 6A (91.1%, $p=0.003$), there were
16
17 no reductions between Period-1 and Period-3 for serotypes 19F (4.2% vs. 4.1%), 23F (4.5%
18
19 vs. 3.2%) and 3 (3.9% vs 3.6%); Table 3. The only NVT to show a significant increase in the
20
21 60-144m. age-group in Period-3 compared to Period-1 was 35B (4.8% vs 0.3%; $p<0.001$).

22
23
24
25 In the >12y. age-group, the prevalence of PCV13-serotype colonization in Period-1 was <1%
26
27 for all serotypes (except serotype 3), and although trending to lower prevalence in Period-3,
28
29 the differences were not statistically significant. For serotype 3, the colonization prevalence
30
31 declined to 0.2% in Period-3 compared to 1.0% in Period-1 ($p=0.030$). Significant increases
32
33 in NVT colonization in Period-3 compared to Period-1 in the >12y. age-group were evident
34
35 for serotypes 8 (0.6% vs. 0%; $p=0.01$) and 35B (1.4% vs. 0%; $p<0.001$); Table 3.

39 40 3.6 Association of HIV infection and pneumococcal colonization

41
42
43 There were very few children <12y. age in Period-3 ($n=15$) that were HIV-infected based on
44
45 self-reporting, hence, limiting any comparisons on whether differences in colonization
46
47 prevalence existed by HIV-status. For individuals >18y. age with self-reported HIV-status,
48
49 there was a higher prevalence of PCV7-serotype colonization in those living with (2.5%)
50
51 compared to those without HIV (0.4%; $p=0.010$), albeit not evident when limited to
52
53 analyzing those in whom HIV testing was done; Table 4. Also, there were no differences for
54
55 overall, PCV13-add6VT or NVT colonization in adults living with and without HIV; Table 4.

56 57 58 59 4. Discussion

60

1
2
3 In this study, undertaken in a rural, low-income African setting with high prevalence of adults
4 living with HIV, immunization of children with a two dose primary series (6 and 14 weeks of
5 age) and a booster dose at 40 weeks of age, was associated with rapid decline in colonization
6 by PCV7 and PCV13-add6VT serotypes in the age-groups targeted for vaccination as well as
7 in older-children (60-144m.) and among adolescents/adults (>12y.) who were naïve for any
8 pneumococcal vaccination. Notably, the reductions in PCV7 and PCV13-add6VT
9 colonization in the vaccine-naïve groups (>60m. age) occurred within two years of
10 introduction of PCV7 and PCV13, respectively, into the public immunization program.
11
12 Additionally, although NVT colonization increased by almost 50%, complete replacement of
13 carriage by NVT was not yet realized in this community.
14
15

16
17 Sizeable vaccine indirect effects were observed, with the percentage reduction in PCV7 and
18 PCV13-add6VT colonization in Period-3 compared to Period-1 in the 60-144m. been 41%
19 and 59%, respectively; and 61% and 75%, respectively in the >12y. age-group. These point-
20 estimates (albeit overlap of the 95%CI) were lower than the reduction in PCV7 (70%) and
21 PCV13-ad6 (66%) observed in the >9-59m. age-group that were eligible for PCV-
22 immunization. Nonetheless, despite up-to-date vaccine coverage of 82.2% in the >9-23m.
23 age-group in Period-3, there was residual colonization by PCV7-serotypes of 13% and
24 PCV13-serotypes of 18% in these children and in the combined >9-59m. age-group. The
25 prevalence of PCV7-serotypes being above 10% shows that there is still significant ongoing
26 transmission of these serotypes in the community. These findings are in contrast to
27 observations in industrialized countries where within a similar period post-PCV7
28 introduction, the prevalence of vaccine-serotype colonization among children <2 years old in
29 Boston, USA, was around 2%¹⁶ and 3.6% in England⁶. The absence of an initial catch-up
30 campaign at vaccine introduction may have left a reservoir of PCV-serotypes among
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 unvaccinated older children, resulting in continued transmission of these types in the
4
5 community.
6
7

8 Similarly, high residual prevalence of vaccine-serotype colonization in age-groups targeted
9
10 for vaccination, as well as in unvaccinated age-groups have been observed from other sub-
11
12 Saharan African countries, including in the presence of using different dosing schedules with
13
14 or without catch-up campaigns. In The Gambia PCV7-serotype colonization prevalence
15
16 among children have been observed in villages where there was no catch-up campaign
17
18 (13.6% among children aged 2.5 to 5 years) ¹⁷ and with PCV13-serotypes at five years after
19
20 PCV13 introduction (11.4% among 6 to 12 months old infants) using a three dose primary
21
22 series and no booster dose (3+0 schedule)¹⁸. In rural Malawi, the reduction in PCV13-
23
24 serotype colonization in children 1-4years eligible to have been immunized with PCV13,
25
26 declined by 63% (28.2% vs. 17.9%); and was also 67% lower in women living without HIV
27
28 (6.6% vs. 2.4%) within four years of introducing PCV13 (3+0 schedule) into the childhood
29
30 immunization program ¹⁹.
31
32
33
34
35

36 The impact of catch-up campaigns in protection of at risk children against invasive
37
38 pneumococcal disease is widely accepted ²⁰, however, its effect on carriage has not been well
39
40 described. In Kilifi, Kenya, where 10-valent PCV was introduced as a three-dose primary
41
42 series coupled with a catch-up campaign through to five years of age, there was also only
43
44 partial reduction, i.e. 74% (95%CI: 66-80) in PCV10-serotype colonization from the pre-
45
46 vaccine era (33.8% vs. 8.8%)) through to five years after introduction²¹.
47
48
49

50
51 Although limited by sample size, in our study for those PCV13-serotypes with >1%
52
53 colonization prevalence among the >9-59m. age-group in Period-1, the percentage reduction
54
55 in colonization by Period-3 varied from 95% (14), 83-87% (6A, 6B and 18C), to more
56
57 modest 38-58% (3, 23F, 19A, 19F and 9V); among the latter group, serotypes 19F and 23F
58
59
60

1
2
3 were the most common colonizing serotypes in Period-1. The high residual PCV13-serotype
4
5 colonization was dominated by these two serotypes, both of which are included in PCV7 that
6
7 was introduced into the immunization program four years before sampling was undertaken in
8
9 Period-3. Moreover, there was no change in prevalence between Period-1 and Period-3 for
10
11 serotype 19F (4.2% vs. 4.4%) and 23F (4.5% vs. 3.2%) colonization in the 60-144m. age-
12
13 group. The reasons for the differential impact of PCV immunization on colonization by
14
15 different serotypes are still hypothetical.
16
17
18
19

20 In the older age-groups (>12y.), contrary to the scenario in children, declines in PCV7-
21
22 serotype colonization were detected from Period-1 (3.1%) to Period-2 (1.2%) but no further
23
24 reductions were detected by Period-3 (1.3%), suggesting that this indirect effect of childhood
25
26 vaccination might have reached a plateau state two years after PCV7 introduction despite
27
28 increase in vaccination coverage including in a wider age range. A dynamic transmission
29
30 model using data from Blantyre, Malawi, suggested that the high residual vaccine serotype
31
32 colonization levels at seven years post-PCV13 introduction was mainly due to age-related
33
34 characteristics of the local force of infection²². Further, the model at 10 years post-PCV13
35
36 introduction, estimated that there would be approximately 75% reduction in PCV13-serotype
37
38 colonization across all age-groups²². In our setting it could be that PCV coverage is still
39
40 suboptimal for complete indirect effects to materialize in adults, leading to a lag in time
41
42 between the effects observed in children. Another explanation could be that there exists a
43
44 reservoir of unvaccinated age groups contributing to the ongoing transmission of some the
45
46 PCV7-serotypes in settings such as ours, as opposed to in high-income countries where
47
48 children 1-4 years of age are considered the main source of pneumococcal transmission.²³ A
49
50 reduction in colonization by PCV13-add6VT was, however, detected in the >12y. age-groups
51
52 between Period-2 and Period-3.
53
54
55
56
57
58
59
60

1
2
3 Of the individual PCV serotypes, 3 was the only serotype with a lower colonization
4 prevalence in Period-3 compared to Period-1 in the >12y. age-group. Interesting, no
5
6 significant temporal changes were noted for serotype 3 colonization among the >9-59m. and
7
8 60-144m. age-groups, and the highest prevalence was in the 60-144m. group. These results
9
10 suggest that the dynamics of colonization and transmission might differ for serotype 3, and
11
12 that the reduction in older individuals may be due to temporal changes rather than indirect
13
14 effects of childhood vaccination. Previously, we have reported that adults living with HIV
15
16 had higher prevalence of PCV-serotype colonization, including that of serotype 3 when
17
18 compared to HIV-uninfected individuals ¹³, and this was corroborated by data from Malawi
19
20 and Uganda ^{24,25}. In the current analysis probably due to the small number of adults
21
22 confirmed to be HIV-infected no differences were detected.
23
24
25
26
27
28

29 In the >9-144m. age-group there were gradual reductions in overall pneumococcal
30
31 colonization during the three sampling Periods despite significant increases in NVT
32
33 colonization. In the >12y. age-group, however, after an initial decrease from Period-1 to
34
35 Period-2, there was a rebound in the prevalence of overall pneumococcal colonization in
36
37 Period-3 that was largely driven by the increase in prevalence of NVT colonization between
38
39 Period-3 and Period-2. This may be explained by the fact that following vaccine introduction
40
41 NVT colonization first established a niche in children before effective transmission to adults
42
43 occurred. An increase in NVT colonization following an initial decrease was also observed in
44
45 the Gambia among adults in villages vaccinated with PCV7 ¹¹. Unlike other settings such as
46
47 the England and USA ²⁶, serotype replacement by NVT is not yet complete and competition
48
49 between NVT and vaccine serotypes such 19F and 23F is ongoing.
50
51
52
53
54

55 The results of this study should be taken in the context of several limitations. First, we did not
56
57 examine the impact of carriage of multiple serotypes simultaneously. Second, since we
58
59 sampled during the same time period (season), we could not assess the impact of seasonal
60

1
2
3 trends and we cannot infer whether pneumococcal carriage patterns are similar throughout the
4 year. In addition, the improvement in socioeconomic conditions (including smaller
5 household size) over time, may have influenced the dynamics of pneumococcal colonization
6 and contributed to the overall reduction in pneumococcal carriage. Third, we relied on
7 reported HIV status, and therefore could not assess the impact of HIV on colonization during
8 the PCV13 era.
9

10 11 12 13 14 15 16 17 18 5. Conclusion

19
20 In conclusion, four years after the introduction of PCV7 and two years after transitioning to
21 PCV13 in the public childhood immunization program, PCV-serotype colonization continues
22 to decline in children <5 years of age. Nevertheless, the residual prevalence of PCV13-
23 serotype colonization in childhood age-groups eligible to have been immunized was >10%
24 four years following implementation of the vaccine program, indicating ongoing circulation
25 in the community. Our data indicate that the direct and indirect benefits of childhood PCV
26 immunization, particularly in relation to vaccine-serotypes 19F and 23F have not yet fully
27 materialized. In addition, near complete replacement by NVT is yet to happen, suggesting
28 that the impact of PCV vaccination is yet to be fully realized in this rural setting. Monitoring
29 of colonization prevalence needs to continue in this setting.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Kayhty H, Auranen K, Nohynek H, Dagan R, Makela H. Nasopharyngeal colonization: a target for pneumococcal vaccination. *Expert Rev Vaccines*. 2006;5(5):651-667.
2. O'Brien KL, Millar EV, Zell ER, et al. Effect of Pneumococcal Conjugate Vaccine on Nasopharyngeal Colonisation among Immunized and Unimmunized Children in a Community Randomized Trial. *The Journal of Infectious Diseases*. 2007;196:1211-1220.
3. Dagan R, Givon-Lavi N, Zamir O, et al. Reduction of nasopharyngeal carriage of Streptococcus pneumoniae after administration of a 9-valent Pneumococcal Conjugate Vaccine to Toddlers attending day care centers. *The Journal of Infectious Diseases*. 2002 185:927-936.
4. O'Brien KL, Dagan R, Makela HP, eds. *Nasopharyngeal carriage*. 1 ed. Washington, DC: American Society for Microbiology press; 2008. Siber GR, Klugman, K.P and Makela, P.H, ed. Pneumococcal vaccines.
5. Cheung Y-B, Zaman, S. M. A., Nsekpong, E.D. et al. . Nasopharyngeal carriage of Streptococcus pneumoniae in Gambian children who participated in a 9-valent pneumococcal conjugate vaccine trial and their younger siblings. *The Pediatric Infectious Disease Journal*. 2009;28(11):990-995.
6. *Flasche S, Jan Van Hoek A, Sheasby E, et al. Effect of pneumococcal conjugate vaccination on serotype-specific carriage and invasive disease in England: A cross-sectional study. *PLoS Medicine*. 2011;8(4):e1001017.
Reports on rapid reduction, and near elimination of PCV-serotypes following introduction of a 2+1 PCV7/13 childhood immunization program in England
7. Hammitt LL, Bruden DL, Butler JC, Baggett HC, Hurlburt DA, Reasonover A, et al. Indirect effect of conjugate vaccine on adult carriage of S pneumoniae: an explanation of trends in invasive pneumococcal disease. *J Infect Dis*. 2006;193:1487-1494.
8. Miller E, Andrews NJ, Waight PA, Slack MP, George RC. Herd immunity and serotype replacement 4 years after seven-valent pneumococcal conjugate vaccination in England and Wales: an observational cohort study. *Lancet Infect Dis*. 2011;11(10):760-768.
9. *Whitney CG, Farley MM, Hadler J, Harrison LH, Bennett NM, Lynfield R. Decline in invasive pneumococcal disease after the introduction of protein-polysaccharide conjugate vaccine. *N Engl J Med*. 2003;348:1737-1746.
Illustrates a dramatic indirect effect against vaccine serotype-invasive pneumococcal disease across all age groups in USA, following introduction of a 3+1 PCV7/13 childhood immunization program, with a catch-up campaign.
10. *Hammitt LL, Akech DO, Morpeth SC, et al. Population effect of 10-valent pneumococcal conjugate vaccine on nasopharyngeal carriage of Streptococcus pneumoniae and non-typeable Haemophilus influenzae in Kilifi, Kenya: findings from cross-sectional carriage studies. *Lancet Glob Health*. 2014;2(7):70224-70224.
Only immunization program to have introduced PCV10 (three dose primary series for infants), with a catch-up campaign up to 5-years of age in sub-Saharan Africa; however, still report high residuals prevalence of vaccine-serotype colonization
11. Roca A, Hill P, C., Townend J, et al. Effects of Community-Wide Vaccination with PCV-7 on Pneumococcal Nasopharyngeal Carriage in The Gambia: A Cluster-Randomised Trial. *PLoS medicine*. 2011;8(10).
12. Gray BM, Converse III GM, Dillon Jr HC. Epidemiologic studies of Streptococcus pneumoniae in Infants: Acquisition, Carriage and Infection during the first 24 months of life. *The Journal of Infectious Diseases*. 1980;142(6):923-933.
13. Nzenze SA, Shiri T, Nunes MC, et al. Temporal changes in pneumococcal colonization in a rural-African community with high HIV-prevalence following routine infant pneumococcal-immunization. *Pediatr Infect Dis J*. 2013;32(11):1270-1278.

14. Barron P, Pillay Y, Doherty T, et al. Eliminating mother-to-child HIV transmission in South Africa. *Bulletin of the World Health Organization*. 2013;91(1):70-74.
15. Nzenze SA, Shiri T, Nunes MC, et al. Temporal association of infant immunisation with pneumococcal conjugate vaccine on the ecology of *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Staphylococcus aureus* nasopharyngeal colonisation in a rural South African community. *Vaccine*. 2014;32(42):5520-5530.
16. Pelton S, I., Weycker D, Klein JO, Strutton D, Ciuryla V, Oster. G. 7-valent pneumococcal conjugate vaccine and lower respiratory tract infections: effectiveness of a 2-dose versus 3-dose primary series. *Vaccine*. 2010;28(6):1575-1582.
17. Roca A, Dione MM, Bojang A, et al. Nasopharyngeal carriage of pneumococci four years after community-wide vaccination with PCV-7 in The Gambia: long-term evaluation of a cluster randomized trial. *PLoS One*. 2013;8(9).
18. Usuf E, Bottomley C, Bojang E, et al. Persistence of Nasopharyngeal Pneumococcal Vaccine Serotypes and Increase of Nonvaccine Serotypes Among Vaccinated Infants and Their Mothers 5 Years After Introduction of Pneumococcal Conjugate Vaccine 13 in The Gambia. *Clin Infect Dis*. 2019;68(9):1512-1521.
19. Heinsbroek E, Tafatatha T, Phiri A, et al. Pneumococcal carriage in households in Karonga District, Malawi, before and after introduction of 13-valent pneumococcal conjugate vaccination. *Vaccine*. 2018;36(48):7369-7376.
20. Whitney CG, Goldblatt D, O'Brien KL. Dosing Schedules for Pneumococcal Conjugate Vaccine: Considerations for Policy Makers. *The Pediatric Infectious Disease Journal*. 2014;33(Suppl 2 Optimum Dosing of Pneumococcal Conjugate Vaccine For Infants 0 A Landscape Analysis of Evidence Supportin g Different Schedules):S172-S181.
21. ***Hammit LL, Etyang AO, Morpeth SC, et al. Effect of ten-valent pneumococcal conjugate vaccine on invasive pneumococcal disease and nasopharyngeal carriage in Kenya: a longitudinal surveillance study. *Lancet*. 2019;393(10186):2146-2154.
Follow-on from reference 10, which despite having introduced PCV10 with a catch-up campaign (through to 10 year of age), residual prevalence of PCV10 colonization was still 8.8% (havng shonw a 74% reduction compared to pre-PCV era. Indicates likley differences in ompact of PCV against colonization (and dynamics of transmission) in high compared to low-middle income settings
22. ***Lourenco J, Obolski U, Swarthout TD, et al. Determinants of high residual post-PCV13 pneumococcal vaccine-type carriage in Blantyre, Malawi: a modelling study. *BMC Med*. 2019;17(1):219.
Modelling study from urban low-income setting in Balntyre, Malawi, which indicates that suinga 3+0 PCV13 dosing schedule without any catch-up campaign, PCV13 serotype colonization wil only have declined by 75% across all agegroup ten years post PCV13 introduction (as opposed to near elimination within four years of introduction in high-income settings). Again, indicates likley differences in ompact of PCV against colonization (and dynamics of transmission) in high compared to low-middle income settings.
23. Choi YH, Andrews N, Miller E. Estimated impact of revising the 13-valent pneumococcal conjugate vaccine schedule from 2+1 to 1+1 in England and Wales: A modelling study. *PLoS medicine*. 2019;16(7):e1002845.
24. Blossom DB, Namayanja-Kaye G, al N-MJe. Oropharyngeal colonization by *Streptococcus pneumoniae* among HIV-infected adults in Uganda: assessing prevalence and antimicrobial susceptibility. *International Journal of Infectious Diseases*. 2006;10:458-464.
25. French N, Gordon SB, Mwalukomo T, et al. A trial of a 7-valent pneumococcal conjugate vaccine in HIV-infected adults. *N Engl J Med*. 2010;362(9):812-822.
26. Gladstone RA, Jefferies JM, Tocheva AS, et al. Five winters of pneumococcal serotype replacement in UK carriage following PCV introduction. *Vaccine*. 2015;33(17):2015-2021.

Table 1. Demographic characteristics and selected health indicators for the study population

Characteristic	Period-1 Pre-PCV era (2009)	Period-2 PCV7 era (2011)	Period-3 PCV13 era (2013)	p-value (Period-1 vs. Period-3)
Household structure				
Number of households	577	1079	563	
Total individuals in household (mean ± SD)	8.8 ± 4.1	8.8 ± 4.2	5.6 ± 2.3	<0.001
Children ≤ 5 years (mean ± SD)	1.8 ± 1.0	2.0 ± 1.1	1.5 ± 0.8	0.014
Individuals 6 – 18 years (mean ± SD)	2.4 ± 1.8	2.2 ± 1.7	1.5 ± 1.4	<0.001
Individuals >18 years (mean ± SD)	4.6 ± 2.4	4.7 ± 2.6	2.5 ± 1.3	<0.001
Rooms used for sleeping (mean ± SD)	3.2 ± 1.5	3.2 ± 1.7	2.6 ± 1.3	<0.001
Households with a child that attends day care, n/N (%)	234/569 (41.1)	351/1078 (32.5)	235/555 (42.3)	0.68
Fuel used for cooking, n/N (%)				
Coal/Wood	394/564 (69.9)	685/1066 (64.3)	235/535 (43.8)	<0.001
Paraffin/Gas	11/564 (1.9)	17/1066 (1.6)	2/535 (0.4)	0.020
Electricity	159/564 (28.2)	364/1066 (34.1)	299/535 (55.8)	<0.001
Children ≤144 months of age				
Number of participants	982	1814	944	
Female, N (%)	494 (50.3)	931 (51.3)	439 (46.5)	0.10
Overall age in months (mean ± SD)	48.4 ± 37.7)	42.0 ± 35.9)	42.5 ± 34.5)	
≤3 months (mean ± SD); n	Not applicable	1.9 ± 0.82; 9	1.8 ± 0.81; 7	
3-9 months (mean ± SD); n	Not applicable	6.8 ± 1.7; 71	6.2 ± 2.0; 23	
>9-23 months (mean ± SD); n	17.3 ± 3.9; 399	17.1 ± 4.0; 828	17.5 ± 4.1; 404	
24-59 months (mean ± SD); n	38.1 ± 11.6; 273	37.1 ± 11.4; 422	36.3 ± 11.6; 259	
60-144 months (mean ± SD); n	97.4 ± 25.0; 310	94.8 ± 24.4; 484	93.5 ± 21.8; 251	
HIV-infected (self-reported), n/N (%)	5/163 (3.1)	20/625 (3.2)	15/503 (3.0)	0.96
Child ever breastfed, overall, n/N (%)	717/779 (92.0)	1181/1398 (84.5)	769/932 (82.5)	<0.001
<12 months.	9/17 (52.9)	30/59 (50.9)	52/66 (78.8)	0.03
12-24 months	156/182 (85.7)	342/456 (75.0)	297/363 (81.8)	0.25
Child attending day-care, n/N (%)	378/981 (38.5)	648/1811 (35.8)	398/929 (42.8)	0.06
Child with tuberculosis, n/N (%)	3/977 (0.3)	2/1805 (0.1)	7/930 (0.8)	0.18
Child on previous TB treatment, n/N (%)	4/981 (0.4)	4/1808 (0.2)	9/920 (1.0)	0.13
Child on antibiotics, n/N (%)	38/981 (7.7)	96/1801 (5.3)	14/919 (1.5)	0.002

Child ever hospitalised, n/N (%)	14/974 (1.4)	9/1804 (0.5)	11/931 (1.2)	0.62
Presence of a smoker in the household, n/N (%)	214/977 (21.9)	602/1770 (34.0)	173/927 (18.7)	0.08
PCV immunization status, n/N (%)				
3-9 months				
zero doses	Not applicable	7/71 (9.9)	1/23 (4.3)	
one dose		18/71 (25.4)	6/23 (26.1)	
two doses		32/71 (45.1)	15/23 (65.2)	
three doses		2/71 (2.8)	0/23 (0.0)	
No vaccination information		12/71 (16.9)	1/23 (4.3)	
>9-23 months				
zero doses	399/399 (0)	50/828 (6.0)	13/404 (3.2)	
one dose		107/828 (12.9)	8/404 (2.0)	
two doses		191/828 (23.1)	25/404 (6.2)	
three doses		313/828 (37.8)	332/404 (82.2)	
No vaccination information		167/828 (20.2)	26/404 (6.4)	
24 to 59 months				
zero doses	273/273 (0.0)	205/422 (48.6)	48/259 (18.5)	
one dose		17/422 (4.0)	21/259 (8.1)	
two doses		30/422 (7.1)	34/259 (13.1)	
three doses		34/422 (8.1)	129/259 (49.8)	
No vaccination information		136/422 (32.2)	27/259 (10.4)	
Participants aged >12 years(y.) of age				
Number of participants	1028	1845	940	
Female, N (%)	828 (80.5)	1563 (84.7)	752 (80.0)	0.76
Overall age in years (mean \pm SD)	33.3 \pm 16.7	32.6 \pm 16.5	31.8 \pm 16.0	0.040
>12-18 years., (mean \pm SD); n	15.0 \pm 1.8; 158	14.9 \pm 1.9; 285	14.3 \pm 1.7; 177	0.001
>18-45 years, (mean \pm SD); n	28.2 \pm 7.3; 633	28.3 \pm 7.2; 1188	29.6 \pm 6.5; 615	0.001
>45 years, (mean \pm SD); n	59.1 \pm 10.3; 237	60.2 \pm 10.7; 372	62.1 \pm 11.4; 148	0.008
HIV-infected in those self-reporting, n/N (%)	40/504 (7.9)	135/1318 (10.2)	64/745 (22.0)	<0.001
>12-18 years	0/24 (0)	1/88 (1.1)	8/79 (10.1)	
\geq 18 years	40/480 (8.3)	134/1230 (10.9)	156/667 (23.4)	
HIV-infected in those tested, n/N (%)			94/463 (20.36.%)	
>12-18 years	Not done	Not done	2/7 (28.6)	
\geq 18 years	Not done	Not done	92/456 (20.2)	
On antiretroviral therapy, n/N (%)	21/38 (55.3)	75/123 (61.0)	109/154 (70.8)	0.53

Smoker, n/N (%)	17/994 (1.7)	26/1779 (1.5)	21/935 (2.2)	0.40
Sniffing tobacco, n/N (%)	27/993 (2.7)	39/1780 (2.2)	19/931 (2.0)	0.33
Drink alcohol, n/N (%)	104/994 (10.5)	114/1778 (6.4)	83/929 (8.9)	0.26
With chronic illness, n/N (%)	138/991 (13.9)	199/1775 (11.2)	179/927 (19.3)	0.002
On tuberculosis treatment, n/N (%)	7/792 (0.9)	10/1770 (0.6)	9/934 (1.0)	0.86
On cotrimoxazole, n/N (%)	4/35 (11.4)	8/101 (7.9)	0/54 (0.0)	

χ^2 – used to compare proportions between the two periods
TTEST used to compare the differences between means between two periods

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 2. Comparison of prevalence of all pneumococci, PCV7-serotypes, additional PCV13 serotypes and non-PCV13 serotypes by age-group in the three study periods

Age-group	Total per age-group N (%) ¹			Overall colonisation n (%) ²			PCV7 serotypes n (%) ²			PCV13 additional-6 serotypes n (%) ²			Non PCV13 serotypes n (%) ²		
	Period-1	Period-2	Period-3	Period-1	Period-2	Period-3	Period-1	Period-2	Period-3	Period-1	Period-2	Period-3	Period-1	Period-2	Period-3
All age-groups	2010	3659	1884	852 (42.4)	1341 (36.6)	675 (35.8)	368 (18.3)	418 (11.4)	129 (6.8)	159 (7.9)	272 (7.4)	53 (2.8)	369 (18.4)	705 (19.3)	502 (26.6)
AOR ³ (95%CI), p-value				0.80 (0.74, 0.86), <0.001			0.34 (0.27, 0.44), <0.001			0.34 (0.23, 0.49), <0.001			1.49 (1.27, 1.74), <0.001		
All age-groups	2010	3579	1854	852 (42.4)	1283 (35.8)	661 (35.7)	368 (18.3)	398 (11.1)	127 (6.9)	159 (7.9)	262 (7.3)	49 (2.6)	369 (18.4)	674 (18.8)	494 (26.6)
ARR (95%CI), p-value ⁴				0.81 (0.75, 0.88), <0.001			0.35 (0.27, 0.45), <0.001			0.33 (0.23, 0.48), <0.001			1.51 (1.29, 1.76), <0.001		
≤9 months	ND ⁵	80 (2.2)	30 (1.6)	ND	58 (72.5)	14 (46.7)	ND	20 (25.0)	2 (6.7)	ND	10 (12.5)	4 (13.3)	ND	31 (38.8)	8 (26.7)
				0.61 (0.40, 0.93), 0.02			0.27 (0.05, 0.90), 0.03			1.07 (0.36, 3.16), 0.90			0.67 (0.35, 1.28), 0.23		
>9-23 months	399 (19.9)	828 (22.6)	404 (21.4)	331 (83.0)	661 (79.8)	268 (66.3)	180 (45.1)	212 (25.6)	54 (13.4)	62 (15.5)	137 (16.5)	20 (5.0)	108 (27.1)	342 (41.3)	196 (48.5)
ARR (95%CI), p-value				0.79 (0.72, 0.88), <0.001			0.26 (0.19, 0.37), <0.001			0.28 (0.16, 0.50), <0.001			2.06 (1.63, 2.61), <0.001		
24-59 months	273 (13.6)	422 (11.5)	259 (13.7)	219 (80.2)	307 (72.7)	175 (67.6)	97 (35.5)	121 (28.7)	38 (14.7)	41 (15)	64 (15.2)	13 (5)	90 (33)	135 (32)	127 (49)
ARR (95%CI), p-value				0.84 (0.74, 0.96), 0.009			0.37 (0.24, 0.56), <0.001			0.40 (0.21, 0.76), 0.005			1.60 (1.24, 2.07), 0.003		
>9-59 months	672 (33.4)	1250 (34.2)	663 (35.2)	550 (81.8)	915 (73.2)	431 (65.0)	277 (41.2)	314 (25.1)	90 (13.6)	103 (15.3)	192 (15.4)	29 (4.4)	198 (29.5)	449 (35.9)	317 (47.8)
ARR (95%CI), p-value				0.81 (0.75, 0.89), <0.001			0.30 (0.23, 0.40), <0.001			0.34 (0.22, 0.52), <0.001			1.84 (1.54, 2.19), <0.001		
60-144 months	310 (15.4)	484 (13.2)	251 (13.3)	186 (60)	242 (50)	130 (51.8)	59 (19.0)	61 (12.6)	25 (10.0)	34 (11.0)	50 (10.3)	14 (5.6)	103 (33.2)	138 (28.5)	93 (37.1)
ARR (95%CI), p-value				0.82 (0.69, 0.98), 0.03			0.59 (0.36, 0.98), 0.04			0.41 (0.20, 0.81), 0.01			1.04 (0.77, 1.41), 0.78		
>9-144months	982 (48.9)	1734 (47.4)	914 (48.5)	736 (74.9)	1157 (66.7)	561 (61.4)	336 (34.2)	375 (21.6)	115 (12.6)	137 (14.0)	242 (14.0)	43 (4.7)	301 (30.7)	587 (33.9)	410 (44.9)
ARR (95%CI), p-value				0.81 (0.75, 0.88), <0.001			0.32 (0.25, 0.41), <0.001			0.35 (0.24, 0.51), <0.001			1.62 (1.38, 1.90), <0.001		
>12-18 years	158 (7.9)	285 (7.8)	177 (9.4)	36 (22.8)	35 (12.3)	27 (15.3)	9 (5.7)	6 (2.1)	6 (3.4)	9 (5.7)	7 (2.5)	2 (1.1)	19 (12)	22 (7.7)	20 (11.3)
ARR (95%CI), p-value				0.58 (0.35, 0.97), 0.04			0.64 (0.22, 1.88), 0.42			0.11 (0.01, 0.96), 0.04			0.75 (0.38, 1.48), 0.41		
>18-45 years	633 (31.5)	1188 (32.5)	615 (32.6)	68 (10.7)	69 (5.8)	64 (10.4)	19 (3.0)	13 (1.1)	6 (1.0)	12 (1.9)	10 (0.8)	4 (0.7)	42 (6.6)	49 (4.1)	55 (8.9)
ARR (95%CI), p-value				0.92 (0.61, 1.40), 0.70			0.21 (0.06, 0.75), 0.02			0.27 (0.09, 0.88), 0.03			1.16 (0.71, 1.88), 0.55		
>45 years	237 (11.8)	372 (10.2)	148 (7.9)	12 (5.1)	22 (5.9)	9 (6.1)	4 (1.7)	4 (1.1)	0 (0.0)	1 (0.4)	3 (0.8)	0 (0.0)	7 (3)	16 (4.3)	9 (6.1)
ARR (95%CI), p-value				1.24 (0.50, 3.05), 0.65			-			-			2.05 (0.78, 5.39), 0.14		
>12 years	1028 (51.1)	1845 (50.4)	940 (49.9)	116 (11.3)	126 (6.8)	100 (10.6)	32 (3.1)	23 (1.2)	12 (1.3)	22 (2.1)	20 (1.1)	6 (0.6)	68 (6.6)	87 (4.7)	84 (8.9)
ARR (95%CI), p-value				0.87 (0.64, 1.18), 0.38			0.39 (0.18, 0.82), 0.01			0.25 (0.07, 0.89), 0.03			1.18 (0.82, 1.71), 0.37		

¹ Value in parenthesis indicate percentage of individuals in each age-group. ²Value in parenthesis indicate percentage of positive individuals.

³ARR=adjusted risk ratio: variables adjusted for: age, gender, period, presence of a smoker in the household, fuel used for cooking and household structure. ⁴Total excludes infants ≤9 months of age. ⁵ND= Not done, infants ≤9 months of age were not sampled in Period-1, therefore comparison is for 2011 versus 2013.

Table 3. Individual serotype prevalence by age-group in children and adults over the three study periods

Serotype	Children																Adults			
	>9 – 23 months				24-59 months				>9-59 months				60-144 months				>12 years			
	Period-1	Period-2	Period-3	p-value	Period-1	Period-2	Period-3	p-value	Period-1	Period-2	Period-3	p-value	Period-1	Period-2	Period-3	p-value	Period-1	Period-2	Period-3	p-value
	N=399	N=828	N=404		N=273	N=422	N=259		N=672	N=1250	N=663		N=310	N=484	N=251		N=1028	N=1845	N=940	
	n (%)	n (%)	n (%)		n (%)	n (%)	n (%)		n (%)	n (%)	n (%)		n (%)	n (%)	n (%)		n (%)	n (%)	n (%)	
PCV7 serotypes																				
4	3 (0.8)	5 (0.6)	2 (0.5)	0.64	2 (0.7)	5 (1.2)	1 (0.4)	0.59	5 (0.7)	10 (0.8)	3 (0.5)	0.49	9 (2.9)	3 (0.6)	0 (0.0)	0.007	5 (0.5)	2 (0.1)	2 (0.2)	0.31
6B	52 (13.0)	39 (4.7)	8 (2.0)	<0.001	16 (5.9)	24 (5.7)	3 (1.2)	0.003	68 (10.1)	63 (5)	11 (1.7)	<0.001	13 (4.2)	10 (2.1)	5 (2.0)	0.14	5 (0.5)	3 (0.2)	3 (0.3)	0.56
9V	7 (1.8)	5 (0.6)	2 (0.5)	0.09	6 (2.2)	7 (1.7)	3 (1.2)	0.35	13 (1.9)	12 (1.0)	5 (0.8)	0.06	5 (1.6)	5 (1)	2 (0.8)	0.39	3 (0.3)	3 (0.2)	1 (0.1)	0.36
14	20 (5)	29 (3.5)	0 (0.0)	<0.001	9 (3.3)	11 (2.6)	1 (0.4)	0.010	29 (4.3)	40 (3.2)	1 (0.2)	<0.001	4 (1.3)	9 (1.9)	0 (0.0)	0.07	3 (0.3)	3 (0.2)	0 (0.0)	0.10
18C	5 (1.3)	1 (0.1)	1 (0.2)	0.10	5 (1.8)	3 (0.7)	0 (0.0)	0.030	10 (1.5)	4 (0.3)	1 (0.2)	0.007	1 (0.3)	2 (0.4)	1 (0.4)	0.88	4 (0.4)	1 (0.1)	0 (0.0)	0.06
19F	63 (15.8)	77 (9.3)	24 (5.9)	<0.001	37 (13.6)	47 (11.1)	18 (6.9)	0.010	100 (14.9)	124 (9.9)	42 (6.3)	<0.001	13 (4.2)	16 (3.3)	11 (4.4)	0.91	6 (0.6)	8 (0.4)	3 (0.3)	0.39
23F	34 (8.5)	39 (4.7)	15 (3.7)	0.004	23 (8.4)	28 (6.6)	12 (4.6)	0.08	57 (8.5)	67 (5.4)	27 (4.1)	0.001	14 (4.5)	16 (3.3)	8 (3.2)	0.42	6 (0.6)	3 (0.2)	3 (0.3)	0.39
Additional six serotypes in PCV13																				
1	1 (0.3)	1 (0.1)	0 (0.0)	0.31	2 (0.7)	0 (0.0)	1 (0.4)	0.59	3 (0.4)	1 (0.1)	1 (0.2)	0.32	0 (0.0)	1 (0.2)	0 (0.0)	.	0 (0.0)	0 (0.0)	0 (0.0)	.
3	1 (0.3)	6 (0.7)	2 (0.5)	0.57	8 (2.9)	7 (1.7)	3 (1.2)	0.15	9 (1.3)	13 (1.0)	5 (0.8)	0.29	12 (3.9)	14 (2.9)	9 (3.6)	0.86	10 (1.0)	11 (0.6)	2 (0.2)	0.030
5	0 (0.0)	0 (0.0)	0 (0.0)	.	2 (0.7)	0 (0.0)	0 (0.0)	0.17	2 (0.3)	0 (0.0)	0 (0.0)	0.16	0 (0.0)	0 (0.0)	0 (0.0)	.	0 (0.0)	0 (0.0)	1 (0.1)	0.30
6A	37 (9.3)	81 (9.8)	7 (1.7)	<0.001	23 (8.4)	42 (10)	3 (1.2)	<0.001	60 (8.9)	123 (9.8)	10 (1.5)	<0.001	14 (4.5)	28 (5.8)	1 (0.4)	0.003	5 (0.5)	8 (0.4)	1 (0.1)	0.13
7F	0 (0.0)	1 (0.1)	0 (0.0)	.	0 (0.0)	0 (0.0)	0 (0.0)	.	0 (0.0)	1 (0.1)	0 (0.0)	.	0 (0.0)	0 (0.0)	0 (0.0)	.	0 (0.0)	0 (0.0)	0 (0.0)	.
19A	23 (5.8)	42 (5.1)	7 (1.7)	0.003	8 (2.9)	16 (3.8)	6 (2.3)	0.66	31 (4.6)	58 (4.6)	13 (2.0)	0.007	9 (2.9)	8 (1.7)	4 (1.6)	0.31	7 (0.7)	2 (0.1)	2 (0.2)	0.12
Non-PCV13 serotypes																				
8	2 (0.5)	2 (0.2)	0 (0.0)	0.15	0 (0.0)	1 (0.2)	1 (0.4)	0.30	2 (0.3)	3 (0.2)	1 (0.2)	0.57	5 (1.6)	5 (1)	7 (2.8)	0.34	0 (0.0)	5 (0.3)	6 (0.6)	0.010
9N	3 (0.8)	7 (0.8)	3 (0.9)	0.99	1 (0.4)	3 (0.7)	2 (0.8)	0.53	4 (0.6)	10 (0.8)	5 (0.8)	0.72	2 (0.6)	3 (0.6)	4 (1.6)	0.28	3 (0.3)	2 (0.1)	2 (0.2)	0.73
11A	4 (1.0)	18 (2.2)	16 (4.0)	0.007	11 (4.0)	5 (1.2)	6 (2.3)	0.26	15 (2.2)	23 (1.8)	22 (3.3)	0.23	10 (3.2)	2 (0.4)	6 (2.4)	0.55	3 (0.3)	5 (0.3)	6 (0.6)	0.26
12F	3 (0.8)	3 (0.4)	1 (0.2)	0.31	0 (0.0)	1 (0.2)	0 (0.0)	.	3 (0.4)	4 (0.3)	1 (0.2)	0.32	0 (0.0)	4 (0.8)	1 (0.4)	0.27	1 (0.1)	1 (0.1)	1 (0.1)	0.95
13	11 (2.8)	20 (2.4)	17 (4.2)	0.26	5 (1.8)	12 (2.8)	6 (2.3)	0.69	16 (2.4)	32 (2.6)	23 (3.5)	0.24	6 (1.9)	14 (2.9)	2 (0.8)	0.26	4 (0.4)	4 (0.2)	1 (0.1)	0.21
15A	2 (0.5)	17 (2.1)	10 (2.5)	0.020	2 (0.7)	8 (1.9)	6 (2.3)	0.13	4 (0.6)	25 (2.0)	16 (2.4)	0.006	0 (0.0)	5 (1)	3 (1.2)	0.05	1 (0.1)	6 (0.3)	2 (0.2)	0.51
15B	13 (3.3)	48 (5.8)	13 (3.2)	0.97	15 (5.5)	16 (3.8)	10 (3.9)	0.37	28 (4.2)	64 (5.1)	23 (3.5)	0.51	12 (3.9)	3 (0.6)	4 (1.6)	0.11	1 (0.1)	3 (0.2)	1 (0.1)	0.95
15C	6 (1.5)	12 (1.4)	14 (3.5)	0.07	8 (2.9)	3 (0.7)	11 (4.2)	0.41	14 (2.1)	15 (1.2)	25 (3.8)	0.07	1 (0.3)	7 (1.4)	3 (1.2)	0.22	2 (0.2)	5 (0.3)	1 (0.1)	0.62
16F	4 (1.0)	40 (4.8)	21 (5.2)	0.001	1 (0.4)	13 (3.1)	16 (6.2)	<0.001	5 (0.7)	53 (4.2)	37 (5.6)	<0.001	8 (2.6)	13 (2.7)	10 (4.0)	0.35	4 (0.4)	7 (0.4)	6 (0.6)	0.44
21	0 (0.0)	8 (1.0)	5 (1.2)	0.030	1 (0.4)	3 (0.7)	6 (2.3)	0.050	1 (0.1)	11 (0.9)	11 (1.7)	0.003	2 (0.6)	2 (0.4)	6 (2.4)	0.08	3 (0.3)	5 (0.3)	3 (0.3)	0.91
23A	2 (0.5)	3 (0.4)	6 (1.5)	0.16	1 (0.4)	4 (0.9)	2 (0.8)	0.53	3 (0.4)	7 (0.6)	8 (1.2)	0.12	1 (0.3)	3 (0.6)	0 (0.0)	0.37	0 (0.0)	0 (0.0)	1 (0.1)	0.30
23B	13 (3.3)	16 (1.9)	8 (2.0)	0.26	9 (3.3)	8 (1.9)	5 (1.9)	0.33	22 (3.3)	24 (1.9)	13 (2.0)	0.13	7 (2.3)	2 (0.4)	5 (2.0)	0.83	2 (0.2)	0 (0.0)	2 (0.2)	0.93
34	7 (1.8)	26 (3.1)	22 (5.4)	0.005	11 (4.0)	15 (3.6)	23 (8.9)	0.020	18 (2.7)	41 (3.3)	45 (6.8)	<0.001	9 (2.9)	8 (1.7)	12 (4.8)	0.24	6 (0.6)	5 (0.3)	4 (0.4)	0.62
35B	3 (0.8)	15 (1.8)	23 (5.7)	<0.001	1 (0.4)	6 (1.4)	8 (3.1)	0.010	4 (0.6)	21 (1.7)	31 (4.7)	<0.001	1 (0.3)	10 (2.1)	12 (4.8)	0.001	0 (0.0)	2 (0.1)	13 (1.4)	<0.001
NT	12 (3.0)	20 (2.4)	0 (0.0)	<0.001	6 (2.2)	8 (1.9)	0 (0.0)	0.020	18 (2.7)	28 (2.2)	0 (0.0)	<0.001	2 (0.6)	4 (0.8)	0 (0.0)	0.20	1 (0.1)	11 (0.6)	0 (0.0)	0.34
OTH	28 (7.0)	70 (8.5)	34 (8.4)	0.46	23 (8.4)	37 (8.8)	31 (12)	0.18	51 (7.6)	107 (8.6)	65 (9.8)	0.15	39 (12.6)	57 (11.8)	22 (8.8)	0.15	37 (3.6)	27 (1.5)	38 (4)	0.61

p-value from chi-squared test comparing differences in prevalence between Period-1 and Period-3. Fisher’s exact test was used for counts <5

1
2
3 NT=non-typeable serotypes; OTH= all other serotypes
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For Peer Review Only

Table 4. Comparison of prevalence of all pneumococci, PCV7 serotypes, additional PCV13 serotypes and non-PCV13 serotypes in HIV-uninfected and HIV-infected children and adults in 2013

Age group	Total N	Overall n (%)	PCV7 n (%)	add PCV13 n (%)	Non PCV13 n (%)
Children ≤12 years self-reporting					
HIV-uninfected	488	290(59.4)	61 (12.5)	27 (5.5)	205 (42.0)
HIV-infected	15	10 (66.7)	2 (13.3)	2 (13.3)	6 (40.0)
p-value		0.79	0.92	0.20	0.88
Adults >12 years self-reporting					
HIV-uninfected	585	60 (10.3)	4 (0.7)	4 (0.7)	52 (8.9)
HIV-infected	156	14 (9.0)	3 (1.9)	1 (0.6)	10 (6.4)
p-value		0.64	0.16	0.95	0.32
Adults >18 years tested					
HIV-uninfected	364	28(7.7)	0	3 (0.8)	26 (7.1)
HIV-infected	92	10 (10.9)	1 (1.1)	0	9 (9.9)
p-value		0.32	-	-	0.40
Adults >18 years self-reporting					
HIV-uninfected	514	49 (9.5)	2 (0.4)	6 (1.2)	44 (8.6)
HIV-infected	157	13 (8.3)	4 (2.5)	4 (2.5)	10 (6.4)
p-value		0.64	0.01	0.21	0.38

Note that some individuals carried both vaccine and non-vaccine serotypes.




Figure 1. Prevalence of pneumococcal colonization in children and adults in a rural South African setting.

Footnote: The p-values in the panel for all ages represent the difference in prevalence in 2009 and 2013.