

**Prospects for CP violation in  $B_s^0 \rightarrow J/\psi\phi$  from first LHCb data**

OLIVIER LEROY on behalf of the LHCb COLLABORATION

CPPM, Aix-Marseille Université IN2P3 CNRS - Aix-Marseille, France

(ricevuto il 29 Settembre 2011; pubblicato online il 26 Gennaio 2012)

**Summary.** —

The determination of the  $CP$ -violating phase in  $B_s^0 \rightarrow J/\psi\phi$  decays is one of the key goals of the LHCb experiment. Its value is predicted to be very small in the Standard Model but can be significantly enhanced in many models of New Physics. The steps towards a precise determination of this phase with a flavour-tagged, time-dependent angular analysis of the decay  $B_s^0 \rightarrow J/\psi\phi$  are reviewed and first studies performed with data collected in 2010 at LHC in pp collisions at 7 TeV center-of-mass energy are presented for the first time. In particular, we report the first LHCb measurements of lifetime in the channels  $B^+ \rightarrow J/\psi K^+$ ,  $B^0 \rightarrow J/\psi K^{*0}$ ,  $B^0 \rightarrow J/\psi K_S^0$ ,  $B_s^0 \rightarrow J/\psi\phi$ ,  $\Lambda_b \rightarrow J/\psi\Lambda$ ; the polarization amplitudes in  $B^0 \rightarrow J/\psi K^{*0}$  and  $B_s^0 \rightarrow J/\psi\phi$ ; the width and mass differences of the  $B_s^0$  mass eigenstates,  $\Delta\Gamma_s$  and  $\Delta m_s$ . The data sample used corresponds to an integrated luminosity of  $36\text{pb}^{-1}$ .

PACS 12.15.Hh – Determination of Cabibbo-Kobayashi & Maskawa (CKM) matrix elements.

**1. – Introduction**

The interference between  $B_s^0$  decays to  $J/\psi\phi$  either directly or via  $B_s^0\text{-}\bar{B}_s^0$  oscillation gives rise to a  $CP$ -violating phase  $\phi_s^{J/\psi\phi}$ . In the Standard Model, this phase is predicted to be  $\simeq -2\beta_s$ , where  $\beta_s = \arg(-V_{ts}V_{tb}^*/V_{cs}V_{cb}^*)$ . The indirect determination via global fits to experimental data gives  $2\beta_s = (0.0363 \pm 0.0017)$  rad [1], within the Standard Model. The direct measurement of this phase is one of the key goals of the LHCb experiment. Indeed,  $\phi_s^{J/\psi\phi}$  is one of the  $CP$  observables with the smallest theoretical uncertainty in the Standard Model, and New Physics could significantly modify this prediction, if new particles contribute with a new phase to the  $B_s^0\text{-}\bar{B}_s^0$  box diagram. Both CDF and DØ have reported constraints on  $\phi_s^{J/\psi\phi}$  with large uncertainties [2, 3].

In this document, we present the steps towards a measurement of  $\phi_s^{J/\psi\phi}$  at LHCb and give the first preliminary results obtained with the 2010 data. The  $CP$ -violation phase will be extracted from a tagged time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi\phi$  decays. Therefore, the following steps are required:

- in sect. 2, we present the selection and lifetime of  $B_s^0 \rightarrow J/\psi(\mu\mu)\phi(KK)$  channel, together with other  $b \rightarrow J/\psi(\mu\mu)X$  control channels;




Fig. 1. –  $B^+$  mass (left) and proper time (right) projections of the two-dimensional fit to the  $B^+ \rightarrow J/\psi K^+$  candidates with  $t > 0.3$  ps. The total fit is represented by the blue solid line, the signal contribution by the green dashed line and the background contribution by the red dashed line. The mass range for the fit is  $m \in [5.15, 5.40]$   $\text{GeV}/c^2$ .

- in sect. **3**, we report on the untagged angular analysis of  $B_s^0 \rightarrow J/\psi \phi$ , together with the control channel  $B^0 \rightarrow J/\psi K^{*0}$ ;
- the tagging of the  $B_s^0$  flavour at production is discussed in sect. **4**, together with the measurement of  $\Delta m_d$  and  $\Delta m_s$ ;
- other channels can be used to measure the mixing-induced  $CP$  violation in  $B_s^0$ -decays amongst which the  $B_s^0 \rightarrow J/\psi f_0$  channel. The first observation of this decay is reported in Sect. **5**.

## 2. – Selections and lifetime measurement

The trigger and selection of  $B_s^0 \rightarrow J/\psi(\mu\mu)\phi(KK)$  and control channels are described in [4]. The measurement of  $\phi_s^{J/\psi\phi}$  requires a good understanding of detector effects such as the proper time acceptance and resolution, angular acceptance, mistag fraction and background. The strategy is to trigger and select several  $b \rightarrow J/\psi X$  decay modes in a similar way and use them as control channels to calibrate the detector and validate the analysis procedures used when studying  $B_s^0 \rightarrow J/\psi\phi$ .

The b-hadron lifetimes are extracted from a maximum-likelihood fit to the proper time distributions of the fully reconstructed candidates. In order to avoid as much as possible a proper time dependent efficiency both the trigger and the offline selection are chosen to be lifetime unbiased: the selections avoid cutting on variables that are correlated with the b-hadron proper time, such as impact parameters of final state particles with respect to the primary vertex. The only exception is a cut on proper time  $t > 0.3$  ps which allows to remove the huge prompt background dominated by combinations of tracks originating from the primary vertex.

The reconstructed mass and proper time projections of  $B^+ \rightarrow J/\psi K^+$ ,  $B^0 \rightarrow J/\psi K^{*0}$ ,  $B^0 \rightarrow J/\psi K_S^0$ ,  $B_s^0 \rightarrow J/\psi\phi$  and  $\Lambda_b \rightarrow J/\psi\Lambda$  are shown in figs. 1 to 5. The extracted lifetimes and the signal yields in the proper time range  $t \in [0.3, 14]$  ps are shown in table I. They are compatible with the PDG values [5]. The proper time resolution measured in  $B_s^0 \rightarrow J/\psi\phi$  event is 50 fs. The systematics uncertainties are given in table II and detailed in [4].




Fig. 2. –  $B^0$  mass (left) and proper time (right) projections of the two-dimensional fit to the  $B^0 \rightarrow J/\psi K^{*0}$  candidates with  $t > 0.3$  ps. The total fit is represented by the blue solid line, the signal contribution by the green dashed line and the background contribution by the red dashed line. The mass range for the fit is  $m \in [5.20, 5.36]$  GeV/c<sup>2</sup>.




Fig. 3. –  $B^0$  mass (left) and proper time (right) projections of the two-dimensional fit to the  $B^0 \rightarrow J/\psi K_S^0$  candidates with  $t > 0.3$  ps. The total fit is represented by the blue solid line, the signal contribution by the green dashed line and the background contribution by the red dashed line. The mass range for the fit is  $m \in [5.15, 5.40]$  GeV/c<sup>2</sup>.




Fig. 4. –  $B_s^0$  mass (left) and proper time (right) projections of the two-dimensional fit to the  $B_s^0 \rightarrow J/\psi\phi$  candidates with  $t > 0.3$  ps. The total fit is represented by the blue solid line, the signal contribution by the green dashed line and the background contribution by the red dashed line. The mass range for the fit is  $m \in [5.20, 5.55]$  GeV/c<sup>2</sup>.




Fig. 5. –  $\Lambda_b$  mass (left) and proper time (right) projections of the two-dimensional fit to the  $\Lambda_b \rightarrow J/\psi\Lambda$  candidates with  $t > 0.3$  ps. The total fit is represented by the blue solid line, the signal contribution by the green dashed line and the background contribution by the red dashed line. The mass range for the fit is  $m \in [5.47, 5.77]$  GeV/ $c^2$ .

TABLE I. – Signal event yields and lifetimes extracted from the likelihood fits to the candidates with proper time  $t \in [0.3, 14]$  ps. A single exponential is used to fit the proper time distribution.

Channel	Lifetime (ps)	Yield
$B^+ \rightarrow J/\psi K^+$	$1.689 \pm 0.022(\text{stat.}) \pm 0.047(\text{syst.})$	$6741 \pm 85$
$B^0 \rightarrow J/\psi K^{*0}$	$1.512 \pm 0.032(\text{stat.}) \pm 0.042(\text{syst.})$	$2668 \pm 58$
$B^0 \rightarrow J/\psi K_S^0$	$1.558 \pm 0.056(\text{stat.}) \pm 0.022(\text{syst.})$	$838 \pm 31$
$B_s^0 \rightarrow J/\psi \phi$	$1.447 \pm 0.064(\text{stat.}) \pm 0.056(\text{syst.})$	$570 \pm 24$
$\Lambda_b \rightarrow J/\psi \Lambda$	$1.353 \pm 0.108(\text{stat.}) \pm 0.035(\text{syst.})$	$187 \pm 16$

TABLE II. – Systematic uncertainties in the lifetime measurements (ps).

	$B^+ \rightarrow J/\psi K^+$	$B^0 \rightarrow J/\psi K^{*0}$	$B_s^0 \rightarrow J/\psi \phi$	$B^0 \rightarrow J/\psi K_S^0$	$\Lambda_b \rightarrow J/\psi \Lambda$
Signal mass model	0.002	0.002	0.010	0.014	0.012
Signal time model	0.043	0.038	0.040	0.015	0.022
Bkg. mass model	0.009	0.020	0.005	0.008	0.023
Bkg. time model	0.003	0.006	0.003	0.006	0.006
Time resol. model	0.005	0.005	0.005	0.005	0.005
Momentum scale	0.001	0.001	0.001	0.001	0.001
Decay length scale	0.001	0.001	0.001	0.001	0.001
Quadratic sum	0.047	0.042	0.056	0.022	0.035

### 3. – Untagged angular analysis of $B^0 \rightarrow J/\psi K^{*0}$ and $B_s^0 \rightarrow J/\psi \phi$

The decays  $B_s^0 \rightarrow J/\psi \phi$  and  $B^0 \rightarrow J/\psi K^{*0}$  are both pseudo-scalar to vector-vector transitions. Both decays are described by three time-dependent decay amplitudes corresponding to transitions in which the  $J/\psi$  and  $\phi$  (or  $K^{*0}$ ) have a relative orbital momentum




Fig. 6. – Fitted PDF with  $S$ -wave included projected on the transversity angles compared to the data distributions for the selected  $B^0 \rightarrow J/\psi K^{*0}$  candidates. Shown are the total PDF, the PDFs for signal (blue),  $S$ -wave (green), total background (red) and wrong-signal (purple).

$L$  of 0, 1, or 2. In the transversity formalism [6], the initial amplitudes at time  $t = 0$ ,  $A_0(0)$  and  $A_{\parallel}(0)$  describe the decays with  $L = 0, 2$  while  $A_{\perp}(0)$  describes the  $L = 1$  final states. The arguments of these complex amplitudes are strong phases denoted  $\delta_0$ ,  $\delta_{\parallel}$  and  $\delta_{\perp}$ . The measurement of the polarization amplitudes and strong phases using untagged events is presented in [7]. For the  $B^0 \rightarrow J/\psi K^{*0}$  channel, we find:

$$\begin{aligned} |A_{\parallel}(0)|^2 &= 0.252 \pm 0.020 \pm 0.016, \\ |A_{\perp}(0)|^2 &= 0.178 \pm 0.022 \pm 0.017, \\ \delta_{\parallel} &= -2.87 \pm 0.11 \pm 0.10, \\ \delta_{\perp} &= 3.02 \pm 0.10 \pm 0.07. \end{aligned}$$

The first error is the statistical uncertainty from the 5-dimensional fit (mass, proper time and 3 angles). The second error is the systematic uncertainty, details of which are given in [7]. The 1-dimensional projections of the 5-dimensional fit function are compared to the measured data in fig. 6.

For the  $B_s^0 \rightarrow J/\psi\phi$  channel, assuming  $\phi_s^{J/\psi\phi} = 0$ , we measure:

$$\begin{aligned} \Gamma_s &= 0.680 \pm 0.034 \pm 0.027 \text{ ps}^{-1}, \\ \Delta\Gamma_s &= 0.084 \pm 0.112 \pm 0.021 \text{ ps}^{-1}, \\ |A_{\perp}(0)|^2 &= 0.279 \pm 0.057 \pm 0.014, \\ |A_0(0)|^2 &= 0.532 \pm 0.040 \pm 0.028, \\ \cos \delta_{\parallel} &= -1.24 \pm 0.27 \pm 0.09, \end{aligned}$$

where the first error is the statistical error from the fit and the second error is the systematic uncertainty detailed in table III. The 1-dimensional projections of the 5-dimensional fit function are compared to the measured data in fig. 7.

#### 4. – Flavour tagging and measurement of $\Delta m_s$ using $B_s^0 \rightarrow D_s^-(3)\pi^+$

The tagging of the initial  $B$ -flavour in LHCb is a key step towards the measurement of  $\phi_s^{J/\psi\phi}$ . It is described in [8]. The algorithm exploits charged tracks originating from the  $b$ -hadron opposite to the signal  $B$ -meson (kaon, muon, electron and vertex charge) and also tracks close to the signal  $B$ -meson (same-side tagging). The algorithm is optimized using  $B^0 \rightarrow D^{*-}\mu^+\nu_{\mu}$  and  $B^+ \rightarrow J/\psi K^+$  events and calibrated using  $B^+ \rightarrow J/\psi K^+$ ,  $B^0 \rightarrow J/\psi K^{*0}$

TABLE III. – *Systematic uncertainties assigned to the extracted physics parameters of the decay  $B_s^0 \rightarrow J/\psi\phi$ .*

Systematic effect	$\Gamma_s [\text{ps}^{-1}]$	$\Delta\Gamma_s [\text{ps}^{-1}]$	$ A_{\perp}(0) ^2$	$ A_{\parallel}(0) ^2$	$\cos \delta_{\parallel}$
Proper time resolution	0.0001	–	–	–	–
Angular acceptance	–	–	–	0.0007	–
Acceptance parametrisation	0.0002	0.001	0.0017	0.0013	–
Proper time acceptance	0.0272	0.001	0.0003	0.0002	–
$S$ -wave treatment	0.003	0.003	0.013	0.028	0.09
Background treatment	0.0002	0.02	0.0016	0.0012	–
Mass model	0.0004	0.004	0.0032	0.0006	–
Total (quadratic sum)	0.0274	0.0206	0.0136	0.0281	0.09




Fig. 7. – Fitted PDF projected on the lifetime and the transversity angles compared to the data distributions for the selected  $B_s^0 \rightarrow J/\psi\phi$  candidates. Shown are the total PDF, the PDFs for signal, the PDFs for the CP-even and CP-odd signal components and the total background PDF.

TABLE IV. – Number of  $B_s^0$  signal candidates used in the  $\Delta m_s$  measurement.

Decay mode	# signal candidates
$B_s \rightarrow D_s^-(\phi\pi^-)\pi^+$	$515 \pm 25$
$B_s \rightarrow D_s^-(K^*K)\pi^+$	$338 \pm 27$
$B_s \rightarrow D_s^-(K^+K^-\pi^-)\pi^+$	$283 \pm 27$
$B_s \rightarrow D_s^-(K^+K^-\pi^-)3\pi$	$245 \pm 46$

events. In [9], the calibration is cross-checked using  $B^0 \rightarrow K^+\pi^-$  events and the  $B^0\text{-}\bar{B}^0$  mixing frequency is measured to be

$$\Delta m_d = 0.499 \pm 0.032 \text{ (stat.)} \pm 0.003 \text{ (syst.) ps}^{-1}.$$

An additional crucial test is performed in [10], by measuring the  $B_s^0\text{-}\bar{B}_s^0$  mixing frequency using  $B_s^0 \rightarrow D_s^-(3)\pi^+$  events. In that case, only opposite side tagging is used. The effective tagging efficiency is  $(3.8 \pm 2.1)\%$ . Using the events sample given in table IV, we measure:

$$\Delta m_s = 17.63 \pm 0.11 \text{ (stat.)} \pm 0.04 \text{ (syst.) ps}^{-1},$$

which is compatible and competitive with the world best measurement [11]. The details of the systematics uncertainties are given in [10]. The likelihood profile as a function of the mixing frequency  $\Delta m_s$  is shown in fig. 8. The statistical significance of the signal is evaluated by comparing the likelihood value at the measured  $\Delta m_s$  value of  $17.63 \text{ ps}^{-1}$  with the likelihood value obtained on the same sample in the limit of infinitely high mixing frequency. We find a significance of  $4.6 \sigma$  for the observed mixing signal. The statistical size of the sample is not large enough to illustrate the oscillation pattern of the time dependent asymmetry. However we can more clearly observe the oscillation if we plot the asymmetry as a function of the proper time modulo  $\frac{2\pi}{\Delta m_s}$  (fig. 9). Additionally we provide an amplitude scan in fig. 9, with all details given in [10].




Fig. 8. – Likelihood scan for  $\Delta m_s$  in the range from  $[0.0, 25.0] \text{ ps}^{-1}$ . The line at 20.94 indicates the likelihood value evaluated in the limit of infinite mixing frequency.




Fig. 9. – Left: Mixing asymmetry for signal  $B_s^0$  candidates as a function of proper time modulo  $\frac{2\pi}{\Delta m_s}$ . The fitted signal asymmetry is superimposed. Right: Fitted amplitude as a function of  $\Delta m_s$ . See the text for further explanation.

### 5. – First observation of $B_s^0 \rightarrow J/\psi f_0$

When LHCb will have accumulated more data, the measurement of  $\phi_s$  will not only be done in  $B_s^0 \rightarrow J/\psi\phi$ , but also in other similar channels. One of them,  $B_s^0 \rightarrow J/\psi f_0$ , has been observed for the first time in LHCb [12]. The  $J/\psi\pi^+\pi^-$  and  $\pi^+\pi^-$  invariant masses are shown in fig. 10. We measure

$$R_{f_0/\phi} \equiv \frac{\Gamma(B_s^0 \rightarrow J/\psi f_0, f_0 \rightarrow \pi^+\pi^-)}{\Gamma(B_s^0 \rightarrow J/\psi\phi, \phi \rightarrow K^+K^-)} = 0.252_{-0.032-0.033}^{+0.046+0.027}.$$

Despite a smaller branching ratio, with respect to  $B_s^0 \rightarrow J/\psi\phi$ , the fact that  $J/\psi f_0$  is a pure CP-odd final state makes the measurement of  $\phi_s$  simpler, since no angular analysis is required.




Fig. 10. – Left: The invariant mass of  $J/\psi\pi^+\pi^-$  combinations when the  $\pi^+\pi^-$  pair is required to be within  $\pm 90$  MeV of the  $f_0(980)$  mass. The data have been fit with a signal Gaussian and several background functions. The thin (red) solid curve shows the signal, the long-dashed (brown) curve the combinatorial background, the dashed (green) curve the  $B^+ \rightarrow J/\psi K^+(\pi^+)$  background, the dotted (blue) curve the  $B^0 \rightarrow J/\psi K^{*0}$  background, the dash-dot curve (purple) the  $B^0 \rightarrow J/\psi\pi^+\pi^-$  background, the barely visible dotted curve (black) the sum of  $B_s^0 \rightarrow J/\psi\eta'$  and  $J/\psi\phi$  backgrounds, and the thick-solid (black) curve the total. Right: The invariant mass of  $\pi^+\pi^-$  combinations when the  $J/\psi\pi^+\pi^-$  combination is required to be within  $\pm 30$  MeV of the  $B_s^0$  mass. The dashed curve is the like-sign background that is taken from the data both in shape and absolute normalization. The dotted curve is the result of the fit described in [12].



## 6. – Epilogue

While completing these proceedings LHCb has released its first preliminary results on  $\phi_s^{J/\psi\phi}$  [13]. The dataset was too small to calibrate the same-side tagger; the opposite side tagger has a measured effective efficiency of  $2.2 \pm 0.4\%$ . Although it was not possible to give a point estimate contours in the  $\phi_s$ - $\Delta\Gamma_s$  space could be calculated.

## 7. – Conclusions

The  $B_s^0 \rightarrow J/\psi\phi$  channel will allow LHCb to probe possible New Physics effects in the  $B_s^0$ - $\bar{B}_s^0$  box diagram. We have presented, for the first time at this conference, the preliminary results needed for a  $\phi_s^{J/\psi\phi}$  measurement, obtained with the data taken in 2010. The data sample used corresponds to an integrated luminosity of  $35 \text{ pb}^{-1}$ . We have measured the lifetime of  $B^+ \rightarrow J/\psi K^+$ ,  $B^0 \rightarrow J/\psi K^{*0}$ ,  $B^0 \rightarrow J/\psi K_s^0$ ,  $B_s^0 \rightarrow J/\psi\phi$ ,  $\Lambda_b \rightarrow J/\psi\Lambda$ , the polarization amplitudes in  $B^0 \rightarrow J/\psi K^{*0}$  and  $B_s^0 \rightarrow J/\psi\phi$ , the width and mass differences of the  $B_s^0$  mass eigenstates,  $\Delta\Gamma_s$  and  $\Delta m_s$ . In particular, we measure:  $\Delta m_s = 17.73 \pm 0.11$  (stat.)  $\pm 0.04$  (syst.)  $\text{ps}^{-1}$ . With the data currently being taken in 2011, we expect to obtain this year the world best measurement of  $\phi_s^{J/\psi\phi}$ .

\* \* \*

I wish to thank the organizers of the “Rencontres de Physique de la vallee d’Aoste”, for the very nice atmosphere during the conference in La Thuile, and all my LHCb colleagues who make possible the first presentation of these 17 new beautiful results.

## REFERENCES

- [1] CHARLES J. *et al.* (CKMFITTER GROUP), *Eur. Phys. J. C*, **41** (2005) 1-131, hep-ph/0406184, updated results and plots available at <http://ckmfitter.in2p3.fr/>.
- [2] THE CDF COLLABORATION, public note CDF/ANAL/BOTTOM/PUBLIC/10206 (2010).
- [3] THE D0 COLLABORATION, D0 Conference note 6098-CONF (2010).
- [4] THE LHCb COLLABORATION, “ $b$ -hadron lifetime measurements with exclusive  $b \rightarrow J/\psi X$  decays reconstructed in the 2010 data”, LHCb-CONF-2011-001.
- [5] NAKAMURA K. *et al.* (PARTICLE DATA GROUP), *J. Phys. G*, **37** (2010) 075021.
- [6] THE LHCb COLLABORATION, “Roadmap for selected key measurements of LHCb.”, LHCb-PUB-2009-029, arXiv:0912.4179.
- [7] THE LHCb COLLABORATION, “Untagged angular analysis of  $B^0 \rightarrow J/\psi K^{*0}$  and  $B_s^0 \rightarrow J/\psi\phi$  decays”, LHCb-CONF-2011-002.
- [8] THE LHCb COLLABORATION, “Optimization and Calibration of the Tagging performances using 2010 data”, LHCb-CONF-2011-003.
- [9] THE LHCb COLLABORATION, “Measurement of  $\Delta M_d$  in  $B^0 \rightarrow D\pi$ ”, LHCb-CONF-2011-010.
- [10] THE LHCb COLLABORATION, “Measurement of  $\Delta M_s$  and calibration and tuning of the Same-Side Tagging algorithm with  $B_s^0 \rightarrow D_s\pi$  decays using the 2010 data sample”, LHCb-CONF-2011-005.
- [11] ABULENCIA A. *et al.* (CDF COLLABORATION), *Phys. Rev. Lett.*, **97** (2006) 242003.
- [12] LHCb COLLABORATION, “First observation of  $B_s^0 \rightarrow J/\psi f^0(980)$  decays”, PLB... 2011.
- [13] THE LHCb COLLABORATION, “Tagged time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi\phi$  decays with the 2010 LHCb data”, LHCb-CONF-2011-006.