

IL NUOVO CIMENTO
DOI 10.1393/ncc/i2012-11284-7

VOL. 35 C, N. 4

Luglio-Agosto 2012

COLLOQUIA: PAVI11

The MOLLER experiment

J. MAMMEI for the MOLLER COLLABORATION

University of Massachusetts, Amherst - Massachusetts, USA

ricevuto il 14 Ottobre 2011; approvato il 5 Maggio 2012
pubblicato online il 3 Luglio 2012

Summary. — The MOLLER experiment will measure the weak charge of the electron, $Q_W^e = 1 - 4 \sin^2 \theta_W$, with a precision of 2.3% by measuring the parity-violating asymmetry in electron-electron (Møller) scattering. This measurement will provide an ultra-precise measurement of the weak mixing angle, $\sin^2 \theta_W$, which is on par with the two most precise collider measurements at the Z^0 -pole. The precision of the experiment, with a fractional accuracy in the determination of $\sin^2 \theta_W \approx 0.1\%$, makes it a probe of physics beyond the Standard Model with sensitivities to mass scales of new physics up to 7.5 TeV.

PACS 12.15.-y – Electroweak interactions.

PACS 12.15.Mm – Neutral currents.

PACS 11.30.Er – Charge conjugation, parity, time reversal, and other discrete symmetries.

1. – Introduction

The Standard Model (SM) summarizes our knowledge of fundamental particles and their interactions. Polarized electron scattering off unpolarized targets provides a clean window to study weak neutral current interactions by measuring the parity-violating asymmetry, A_{PV} . The MOLLER experiment will test the SM prediction of the weak mixing angle, $\sin^2 \theta_W$, by measuring A_{PV} in polarized electron scattering from atomic electrons (Møller scattering). The A_{PV} is defined by

$$(1) \quad A_{PV} = \frac{\sigma_R - \sigma_L}{\sigma_R + \sigma_L},$$

where σ_R (σ_L) is the scattering cross section of incident right- (left-) handed electrons. At four-momentum transfers much smaller than the mass of the Z^0 boson, $Q^2 \ll M_Z^2$, A_{PV} is dominated by the interference between photon and Z^0 boson exchange [1]. The leading order Feynman diagrams relevant for Møller scattering, which involve both direct




Fig. 1. – Feynman diagrams for Møller scattering at tree level (reproduced from ref. [3]).

and exchange diagrams that interfere with each other, are shown in fig. 1. The resulting asymmetry is given by [2]

$$(2) \quad A_{PV} = mE \frac{G_F}{\sqrt{2}\pi\alpha} \frac{4 \sin^2 \theta}{(3 + \cos^2 \theta)^2} Q_W^e = mE \frac{G_F}{\sqrt{2}\pi\alpha} \frac{2y(1-y)}{1+y^4+(1-y)^4} Q_W^e,$$

where α is the fine structure constant, G_F the Fermi constant, E the incident beam energy, m the electron mass, θ the scattering angle in the center of mass frame, and $y \equiv 1 - E'/E$, where E' is the energy of one of the scattered electrons.

Within the SM the weak charge of the electron, Q_W^e , is proportional to the product of the electron's vector and axial-vector couplings to the Z^0 boson, and the weak neutral current amplitudes are functions of the weak mixing angle $\sin^2 \theta_W$. The world average of the two most precise independent determinations of $\sin^2 \theta_W$ is consistent with other electroweak measurements and constraints on the Higgs boson mass M_H , but the measurements actually differ by over 3 standard deviations. Choosing one or the other central value ruins this consistency and implies very different new high-energy dynamics. We propose to measure $\sin^2 \theta_W$ to a sensitivity of $\delta(\sin^2 \theta_W) = \pm 0.00029$, the only method available in the next decade to directly address this issue at the same level of precision and interpretability.

MOLLER will also be *the* most sensitive probe of new flavor and CP-conserving neutral current interactions in the leptonic sector, sensitive to interaction amplitudes as small as 1.5×10^{-3} times the Fermi constant. New neutral current interactions are best parameterized model-independently at low energies by effective four-fermion interactions. Focusing on vector and axial-vector interactions between electrons and/or positrons, such an interaction Lagrangian takes the form [4]

$$(3) \quad \mathcal{L}_{e_1 e_2} = \sum_{i,j=L,R} \frac{g_{ij}^2}{2\Lambda^2} \bar{e}_i \gamma_\mu e_i \bar{e}_j \gamma^\mu e_j,$$

where $e_{L/R} = \frac{1}{2}(1 \mp \gamma_5)\psi_e$ are the usual chirality projections of the electron spinor, Λ the mass scale of the new contact interaction and $g_{ij} = g_{ij}^*$ coupling constants, with $g_{RL} = g_{LR}$. For the proposed measurement with 2.3% total uncertainty (and no additional theoretical uncertainty) the resulting sensitivity to new four-electron contact interaction amplitudes $g_{RR}^2 - g_{LL}^2$ is ~ 7.5 TeV. The proposed measurement will greatly extend the current sensitivity of four-electron contact interactions, both qualitatively and quantitatively, and is complementary to direct searches.

It is straightforward to examine the reach MOLLER will have in specific models (the mass scale depends on the size of the coupling) [5]. As a specific example, a comprehensive analysis of the MOLLER sensitivity to TeV-scale Z' bosons has recently been carried out [6] for a large class of models contained in the E_6 gauge group, where Z' bosons with the same electroweak charges as SM particles are motivated because they arise in many




Fig. 4. – CAD model of the conceptual layout of the experiment, looking upstream. The target chamber is located in the upper left, then the upstream and hybrid toroids, Roman pots for the tracking detectors, and the main detector stand, with background detectors at the far downstream end in the lower right.

2. – Experimental overview

The MOLLER experiment will run in Hall A at Jefferson Laboratory, making use of the 11 GeV longitudinally polarized ($\sim 85\%$) electron beam, generated via photoemission on a GaAs photocathode by circularly polarized laser light. The target is a 1.5 m liquid hydrogen target capable of dissipating 5 kW of beam power. Møller electrons in the full range of the azimuth (achieved by using an odd number of coils and identical particle scattering) and polar angles $5 \text{ mrad} < \theta_{lab} < 17 \text{ mrad}$, will be separated from background and focused $\sim 30 \text{ m}$ downstream of the target by a spectrometer system consisting of a pair of toroidal magnet assemblies and precision collimators (fig. 4). The upstream magnet is a traditional resistive toroidal magnet, while the downstream magnet has a novel shape designed to focus the large range of scattered electron angles and energies.

The Møller electrons will be incident on a system of quartz detectors in which the resulting Cherenkov light will provide a relative measure of the scattered flux. Rapid polarization (helicity) reversal of the electrons will be used to suppress spurious systematic effects. A_{PV} will be extracted from the fractional difference in the integrated Cherenkov light response between helicity reversals. Additional systematic suppression below the part-per-billion (ppb) level will be accomplished by periodically reversing the sign of the physics asymmetry. We plan to introduce this “slow helicity reversal” with three independent methods: the introduction of an additional half-cycle $g - 2$ rotation to the electrons in the recirculating arcs of the accelerator, the use of an insertable half-wave plate in the injector, and a full flip of the beam polarization direction with the aid of two Wien rotators and a solenoid lens (the “Double Wien”).

Simultaneously with data collection, fluctuations in the electron beam energy and trajectory and their potential systematic effects on A_{PV} will be precisely monitored, active feedback loops will minimize beam helicity correlations, and detector response to beam fluctuations will be continuously calibrated. Background fractions and their helicity-correlated asymmetries will be measured by dedicated auxiliary detectors. The absolute value of Q^2 will be calibrated using tracking detectors. The electron beam

TABLE I. – *Summary of projected fractional systematic errors on the measurement of Q_W^e . The fractional statistical error is 2.1%.*

| Error Source | Fractional Error (%) |
|---|----------------------|
| absolute value of Q^2 | 0.5 |
| beam (second order) | 0.4 |
| beam polarization | 0.4 |
| $e + p(+\gamma) \rightarrow e + X(+\gamma)$ | 0.4 |
| beam (position, angle, energy) | 0.4 |
| beam (intensity) | 0.3 |
| $e + p(+\gamma) \rightarrow e + p(+\gamma)$ | 0.3 |
| $\gamma^{(*)} + p \rightarrow \pi + X$ | 0.3 |
| transverse polarization | 0.2 |
| neutrals (soft photons, neutrons) | 0.1 |
| Total systematic | 1.1 |

polarization will be measured continuously by two independent polarimeter systems. The predicted value of A_{PV} for the proposed experimental design is ~ 35 ppb and our goal is to measure this quantity with a precision of 0.73 ppb. We tabulate our estimates of the most important systematic errors in table I.

3. – Experiment status and plans

We are proposing to take data in three separate run periods to ensure that important technical milestones are met and that each run will provide publishable results that will significantly add to our knowledge of electroweak physics to date. While the MOLLER apparatus is being designed for a beam current of $85 \mu\text{A}$ at 11 GeV, we have assumed a beam current of $75 \mu\text{A}$ and a beam polarization of 80% in formulating the run plan. If higher beam current and/or higher beam polarization are considered routine, the amount of time needed to run could correspondingly be reduced using the appropriate P^2I factor.

In order to estimate the time needed to reach the desired statistical accuracy it is necessary to take into account both the overall efficiency and how close one can approach counting statistics in an instantaneous raw asymmetry measurement. The overall efficiency depends on the efficiency of the apparatus itself and the accelerator efficiency, which we estimate as 90% and 70% respectively, for an overall efficiency of 60%. At 960 Hz, the width of the measured asymmetry per pulse pair, $\sigma(A_i)$, is 83 ppm, but this width depends on the sources of additional fluctuations. However, we are unlikely to achieve the best efficiency or asymmetry width at first, so we assumed total efficiencies of 40, 50 and 60%, respectively, and asymmetry widths of 100, 95 and 90 ppm, respectively, for the three running periods. This leads to run periods of 11, 30 and 60 weeks (including commissioning) with expected statistical errors of 11%, 4.0% and 2.4%, respectively.

MOLLER is a fourth generation parity-violation experiment at Jefferson Laboratory. Apart from the obvious challenge of measuring a raw asymmetry with a statistical error less than 1 ppb, an equally challenging task is to calibrate and monitor the absolute normalization of A_{PV} at the sub-1% level. The collaboration continues to gain extensive

experience on all aspects of such measurements as work continues on executing the third generation experiments PREX [9] and Qweak [10]. Simulation and design of the various aspects of MOLLER are ongoing, and is expected to take two to three years. Work is also being done to improve beam transport and instrumentation. In addition, upgrades to the Compton and Møller polarimeters are being planned. It is envisioned that construction and assembly will take three years, to be followed by three data collection periods with progressively improved statistical errors and systematic control over a subsequent three to four year period.

* * *

Thanks to the organizers for a wonderful meeting. This work is funded in part by US Department of Energy Grant No. DE-FG02-88R40415-A018.

REFERENCES

- [1] ZEL'DOVICH YA. B., *Sov. Phys. JETP*, **94** (1959) 262.
- [2] DERMAN E. and MARCIANO W. J., *Ann. Phys. (N.Y.)*, **121** (1979) 147.
- [3] CZARNECKI A. and MARCIANO W. J., *Int. J. Mod. Phys. A*, **15** (2000) 2365.
- [4] EICHTEEN E., LANE K. D. and PESKIN M. E., *Phys. Rev. Lett.*, **50** (1983) 811.
- [5] RAMSEY-MUSOLF M. J., *Phys. Rev. C*, **60** (1999) 015501.
- [6] ERLER J. and ROJAS E., private communication.
- [7] ERLER J., *Nucl. Phys. B*, **586** (2000) 73.
- [8] CHANG W. F., NG J. N. and WU J. M. S., *Phys. Rev. D*, **79** (2009) 055016.
- [9] KUMAR K., MICHAELS R., SOUDER P. and URCIUOLI G., *The Lead Radius Experiment PREX, E06-002*, <http://hallaweb.jlab.org/parity/prex/>.
- [10] CARLINI R., *The Qweak Experiment*, <http://www.jlab.org/Hall-C/Qweak/index.html>.