

## Gonad development of hard clam (*Meretrix xlyrata*)

\*[Hadi Hamli](#), Mohd Hanafi Idris, Abu Hena Mustafa Kamal and Amy Halimah Rajae

*Department of Animal Science and Fishery, Faculty of Agriculture and Food Sciences,  
Universiti Putra Malaysia Bintulu Sarawak Campus, Nyabau Road, P. O. Box 396, 97008  
Bintulu, Sarawak, Malaysia*

\*Corresponding author. Tel.: +6086 8555373/5382; email address: [hadihamli@gmail.com](mailto:hadihamli@gmail.com)

Hard clam are mollusc belongs to bivalvia class that dominating the benthic area from deep sea to lowland area particularly tropical area. This invertebrate is suspension feeder that feeds upon plankton, detritus and dissolved organic matter as the source of metabolic energy to keep it grow. Those foods source are abundance at the open sea which make hard clam proper for mariculture practice. Hard clam production through mariculture gradually increase each year even still behind oyster and cockle production. Knowledge regarding gonad stage development is important for hard clam culture in order to determine proper condition for spawning. Development of bivalve gonad is categorized into resting, early development, late development, mature, spawning and spawned. Gonad development and gender of *Meretrix lyrata* can be identified accurately through histological study.

**Keywords:** *Meretrix lyrata*, gametogenesis, histological