## MALAYSIA INTERNATIONAL BIOLOGY SYMPOSIUM 2016 26th – 27th OCTOBER 2016 PICC, PUTRAJAYA

## Isolation, molecular identification and biodegradation capacities of indigenous hydrocarbondegrading bacteria from tarball at Terengganu Beach, Malaysia

<u>Bruno Martins Nkem</u><sup>a</sup>, Normala Halimoon<sup>a,\*</sup>, Fatimah Md Yusoff<sup>b</sup>, Wan Lufti Wan Johari<sup>a</sup>, Mohamad Pauzi Zakaria<sup>a</sup>, Srikanth Reddy<sup>a,b</sup> and Narayanan Kannan<sup>a</sup>

<sup>a</sup>Faculty of Environmental Science, Department of Environmental Studies, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor Darul Ehsan, Malaysia.

<sup>b</sup>Laboratory of Marine Biotechnology, Institute of Bioscience, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor Darul Ehsan, Malaysia.Science.

## Abstract

In this study, we isolated four indigenous hydrocarbon-degrading bacteria from tarball found in Rhu Sepuluh beach, Terengganu, Malaysia. These bacteria were identified based on their physiological characteristic and 16S rRNA gene sequence analysis, and they showed 99% similarity with *Pseudomonas stutzeri* strain A1501, *Pseudomonas balearica* strain SP1402, *Cellulosimicrobium cellulans* DSM 43879 and *Acinetobacter baumannii* ATCC 19606 respectively. Their hydrocarbon-degrading capabilities were tested using diesel-oil as sole carbon source. Results analysed using GC-MS, showed diesel-oil alkanes were degraded an average 30.5%, 16.4%, 64.4% and 58.1% with medium optical density reaching 1.172, 1.503, 0.967 and 1.515 by *Pseudomonas stutzeri*, *Pseudomonas balearica, Cellulosimicrobium cellulans* and *Acinetobacter baumannii* respectively in minimal salt media at 32°C for 10 days. Individual diesel-oil alkanes were degraded between 19.7% - 49.2% (*P. stutzeri*), 2.5% to 27.5% (*P. balearica*), 10% - 95.4% (*C. cellulans*) and 0.2% - 95.9% (*A. baumannii*). All strains utilized diesel-oil for growth. The study suggests all strains are part of indigenous hydrocarbon-degrading bacteria in tarball with potential for bioremediation of oil-polluted marine environment.

Keywords: Bacteria, biodegradation, bioremediation, diesel, hydrocarbones, tarballs.

\*Corresponding author: mala\_upm@upm.edu.my