

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

**4,800**

Open access books available

**122,000**

International authors and editors

**135M**

Downloads

Our authors are among the

**154**

Countries delivered to

**TOP 1%**

most cited scientists

**12.2%**

Contributors from top 500 universities



**WEB OF SCIENCE™**

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



## Treatment of childhood pneumonia in developing countries

Hasan Ashraf, Mohammad Jobayer Chisti and Nur Haque Alam  
*Clinical Sciences Division, International Centre for Diarrhoeal Disease Research,  
Bangladesh (ICDDR,B)  
Bangladesh*

### Introduction

#### Definition of community acquired pneumonia (CAP)

CAP can be defined clinically as the presence of signs and symptoms of pneumonia in a previously healthy child due to an acute infection (of less than 14 days' duration) of the lower respiratory tract (usually occurs below terminal bronchioles) leading to cough or difficult breathing, tachypnoea, or lower chest-wall indrawing, which has been acquired in the community outside hospital (Zar et al., 2005; BTS, 2002). In developed countries this can be verified by the radiological finding of consolidation (BTS, 2002). In resource poor setting of the developing world, a more practical term - acute lower respiratory infection (ALRI) - is preferred, reflecting the difficulties in obtaining a chest radiograph, especially in rural areas (BTS, 2002).

#### Disease burden and epidemiology of CAP

In the developing world, pneumonia is not only more common than it is in Europe and North America (Riley et al., 1983; Berman & McIntosh, 1985; Selwyn, 1990), but also more severe and is the largest killer disease of children (Bulla & Hitze, 1978; Baqui et al., 1998). The fourth Millennium Development Goal has concentrated efforts on addressing the priority areas for improving child survival worldwide, with an aim of reducing the national child mortality rates by two-thirds by 2015 (UN, 2000). ALRI, particularly pneumonia, are currently the leading and biggest single cause of deaths among under-5 children 1 to 59 months of age in the developing countries (UNICEF & WHO, 2006; Graham et al., 2008), being responsible for at least 19% of the annual 8.8 million deaths in this age-group (Wardlaw et al., 2006). ALRI causes more than 2 million child deaths (one million in children aged 1 to 59 months and additional one million in neonates) worldwide each year, mostly from pneumonia, accounting for 20% of deaths in under-5 children (Bryce et al., 2005; Rudan et al., 2004), and 90-95% of all these deaths occur in the developing countries (Rudan et al., 2008; Murray & Lopez, 1997; Garenne et al., 1992; Mulholland, 1999; Williams et al., 2002; WHO, 1998). In terms of magnitude of the problem, there is an estimated incidence of 151 million new cases of pneumonia each year globally, and 11-20 million (7-13%) are severe enough to require hospitalization in the developing countries (Rudan et al., 2004; Rudan et

al., 2008). Recent estimates also suggest that 1.9 million (95% CI 1.6 to 2.2 million) children died from acute respiratory infection (ARI) throughout the world in 2000 and 70% of them occurred in Africa and Southeast Asia, one dying in every 7 seconds (Mulholland, 1999; Williams et al., 2002). Actually, CAP is a major cause of health care utilization, hospitalization, and death in children in the developing countries (Mulholland, 1999; Williams et al., 2002; WHO, 1998). Therefore, improvement in the case-management strategies of the major causes of child death, such as pneumonia and neonatal illnesses in developing countries, should be a priority in improving the child survival in the developing countries (Walley et al., 2008; Duke & Tamburlini, 2003). ARI is also a major cause of visits to the outpatient and emergency departments as well as admissions to the hospitals. Although bronchiolitis, tracheobronchitis and pneumonia, each accounts for one-third of ALRI cases, pneumonia is responsible for most of the ALRI deaths. In Bangladesh, ALRI account for 25% of deaths among under-5 children and constitute 40% of all infantile deaths (Baqui et al., 1998). In a study conducted at the Dhaka Hospital of ICDDR,B among 401 under-5 children with ALRI, it was observed that the most common manifestation was pneumonia and a respiratory pathogen (both bacterial and viral) was identified in 30% cases and the case fatality rates were 14% in bacterial pneumonia and 3% in viral pneumonia (Rahman et al., 1990). In another study also conducted at the Dhaka Hospital of ICDDR,B among 601 under-5 children with ALRI, it was observed that the most common manifestation was pneumonia (86.5%), and a viral pathogen was detected in 21% cases, and the overall case fatality rate was 6.8%, and that of viral pneumonia was 4.8% (Huq et al., 1990).

### **Objective**

To develop guidelines for the physicians and nurses of the developing countries for the diagnosis and management of CAP in children

### **Options**

- Clinical assessment
- Radiographic assessment
- Laboratory testing
- Empirical antimicrobial therapy

### **Outcomes**

- Increased awareness of the age-related causes of CAP including those children with severe acute malnutrition (SAM), and those with dehydrating diarrhoea
- Improved accuracy of clinical diagnosis of CAP in children
- Better utilization of the available diagnostic testing
- Rational use of empirical antimicrobial therapy
- Decreased morbidity and mortality due to CAP

### **Benefits, harms, and costs**

- Increased awareness of the causes of paediatric pneumonia
- Accurate diagnosis
- Prompt treatment
- Reduced cost associated with unnecessary investigations and complications due to inappropriate treatment

**Rationale**

As pneumonia is a major cause of morbidity and mortality in children in the developing countries (Williams et al., 2002), early and appropriate treatment of pneumonia can reduce the morbidity and mortality (Sazawal & Black, 2003), which has been the rationale for the development of guidelines for the management of CAP (Zar et al., 2005). The guidelines also aim to provide recommendations for effective therapy and to minimize the development of bacterial resistance through judicious use of antibiotics (Zar et al., 2005). This document aims to provide guidelines for the diagnosis and effective management of children with CAP so as to improve pneumonia-associated morbidity and mortality, thereby improving the case-management strategies of the major cause of child death, such as pneumonia in the developing countries which should be a priority in improving child survival globally (Walley et al., 2008; Duke & Tamburlini, 2003).

**Clinical classification and management plan of CAP**

Depending on the clinical presentation, pneumonia can be classified as very severe, severe, or non-severe according to the World Health Organization (WHO) (WHO, 1990; WHO, 1984; WHO, 2000; WHO, 1991). The specific treatment guidelines for each of them, and the diagnosis of pneumonia should primarily be based on the visible clinical parameters, including respiratory rate and lower chest-wall indrawing (WHO, 1990; WHO, 1991; Cashat et al., 2005). Pneumonia is usually caused by viruses or bacteria, but the most serious episodes are caused by bacteria. However, in the absence of clear cut demarcation for aetiological diagnosis of pneumonia on the basis of clinical and/or radiological features, empiric antibiotic therapy is needed for all cases of CAP. Hospitalization of children with severe pneumonia is recommended for giving supportive treatment including oropharyngeal or nasopharyngeal suction, oxygen therapy for hypoxaemia, fluid and nutritional management, and close monitoring (WHO, 1990; WHO, 1984; WHO, 2000; WHO, 1991). Therefore, management of severe childhood pneumonia relies on hospital-based treatment, but practical barriers often prevent children in areas with highest rates from receiving hospital care (Ashraf et al., 2008). In the developing countries, there are not enough paediatric beds in hospitals for admission of all severe cases of pneumonia (Ashraf et al. 2008). In addition, hospitalization may not be possible because of the inability of the parents to visit the hospital because of the long distances to travel or financial or other domestic reasons, such as the need to care for siblings at home and the need for the mother to work (Ashraf et al., 2008; Ashraf et al., 2007). It is, therefore, important to provide some form of institutional care for children who cannot be hospitalized, at least until stabilization of their acute condition (Ashraf et al., 2008). Radiological examination and determination of hypoxaemia by pulse oximetry, have been recently considered as the optimal methods for diagnosing pneumonia. But, they are clearly suitable only for use in the institutional settings like the day-care centres or out-patient clinics. Two prospective observational studies have shown that the day-care facility-based, modified primary care management of severe childhood pneumonia (Ashraf et al., 2008) and severe acute malnutrition (SAM) (Ashraf et al., 2007) is successful and cost-effective as an alternative to hospitalization. Provision of broad-spectrum antibiotics and appropriate supportive care during a stay at established day-care centre during their working hours, followed by the continuation of care at home at night, is an effective alternative to hospitalization of children with severe pneumonia without associated co-morbidities such as SAM (Ashraf et al., 2008). The results of this study

indicate that severe childhood pneumonia without SAM can be successfully managed on a day-care basis at established day-care clinics, if adequately trained and motivated staff and the necessary logistic support can be made available (Ashraf et al., 2008). The results of a randomized controlled clinical trial (RCT) have shown that in a select group of under-5 children with severe pneumonia, without associated co-morbidities such as SAM, can be safely and effectively managed on a day-care basis as effectively as the hospital set-up management, except those children with hypoxaemia requiring prolonged oxygen therapy for more than six hours, and that day-care based treatment option is less expensive than hospital-care (Ashraf et al., 2010). The results also indicate that severe pneumonia without hypoxaemia can be successfully managed on a day-care basis at a day-care clinic, however, identification of those severely pneumonic children having hypoxaemia requiring prolonged oxygen therapy for more than six hours are necessary, as they are at increased risk of death and therefore need to be hospitalized for support and care for a longer period of time (Ashraf et al., 2010). These results would have great impact in the treatment and care of childhood pneumonia, particularly in resource-poor countries where hospital beds are scarce. It can be easily replicated in most urban and rural out-patient clinics and day-care centres, provided that proper training and motivation of the staff as well as provision of logistic support are guaranteed. The additionally needed funds are well invested facing the lower costs of the day-care treatment model compared to those of hospital-care (US\$ 114 vs. 178) (Ashraf et al., 2010). However, policy and programme changes would be necessary to add such components to the out-patient clinics and day-care centres, and this would require additional human and financial resources, which is not an easy task. Therefore, this would be important in selecting the intervention for wider implementation in national programmes. The results of our RCT identified a way to more efficiently use scarce hospital beds in developing countries by selecting out children with severe pneumonia for the day-care management, who following existing guidelines, would have been identified as the ones requiring hospitalization. This would be a practical approach in developing countries - a smaller investment in upgrading the day-care facilities through development of trained resources and procurement of some supporting equipment, which could pay back in a much greater way.

#### **Rapid/fast breathing (WHO)**

Rapid/fast breathing is defined as when the age-specific respiratory rates become  $\geq 60$ /minute in neonates and infants aged  $< 2$  months,  $\geq 50$ /minute in infants aged 2 to  $< 12$  months, and  $\geq 40$ /minute in children aged 12-59 months, as shown below:

- Age  $< 2$  months:  $\geq 60$ /minute
- Age 2 to  $< 12$  months:  $\geq 50$ /minute
- Age 12 months-5 years:  $\geq 40$ /minute

(N. B. A child who is exactly 12 months old would have fast breathing if s/he breaths  $\geq 40$ /minute. Tachypnoea is the best single predictor of pneumonia in children of all ages).

#### **Lower chest-wall indrawing**

Indrawing of the chest wall is a manifestation of reduced lung compliance resulting from the tendency of declining intra-alveolar pressure due to pneumonic consolidation or

airways obstruction. WHO recommends the use of lower chest wall indrawing as a sign of pneumonia requiring admission to hospital, defined as the inward movement of the bony structures of the lower chest wall with inspiration. Lower chest-wall indrawing is also called “subcostal indrawing”/“subcostal retractions”

#### **Clinical types of pneumonia (WHO)**

- Very severe pneumonia (up to 5 years)
- Severe pneumonia (up to 5 years)
- Pneumonia (not severe) (only for children aged 2 months to 5 years)
- No pneumonia: cough or cold (up to 5 years)

#### **Very severe pneumonia**

If a child with cough or difficult breathing has any one or more of the following danger signs, s/he is classified as having very severe pneumonia (WHO, 2005)

- Not able to drink
- Cyanosis
- Head nodding

“No danger signs” are defined as the absence of all of the following danger signs: not able to drink/feed, central cyanosis, head nodding, stridor in calm child, abnormally sleepy, convulsion, and severe clinical malnutrition.

If a child presents with severe malnutrition with any sign of pneumonia (any of the WHO defined signs of pneumonia or severe pneumonia or very severe pneumonia or crackles or bronchial breath sound in lungs or radiological pneumonia) should be considered as very severe pneumonia.

#### **Severe pneumonia**

Severe pneumonia is defined as an young infant (< 2 months ) with cough or difficult breathing having fast breathing and/or lower chest-wall indrawing, or a child (2 months to 5 years) with cough or difficult breathing having only lower chest-wall indrawing.

#### **Pneumonia (not severe)**

Pneumonia (not severe) is defined as a child (2 months to 5 years) with cough or difficult breathing having only fast breathing, but no lower chest-wall indrawing, or no signs of very severe pneumonia.

#### **Aetiology of CAP**

Rational treatment for pneumonia depends on knowing the most likely pathogens in each community, as the relative frequency of different agents may vary from one geographical region to another and depends on the age of the patient, vaccination status, immunological status, relevant exposure and clinical setting at which pneumonia was acquired. Mixed bacterial and viral infections may occur in 30-40% of cases of CAP (Zar et al., 2005). *Streptococcus pneumoniae* is the most common bacterial cause of childhood pneumonia (BTS, 2002)), followed by *Haemophilus influenzae*, *Mycoplasma pneumoniae*, *Chlamydia pneumoniae*, and *Mycobacterium tuberculosis*. The less common bacterial causes are *Staphylococcus aureus*,

*Bordetella pertussis*, *Pneumocystis jiroveci* (previously known as *Pneumocystis carinii*), and *Nisseria meningitidis*. In older children, when a bacterial cause is found, it is most commonly *Streptococcus pneumoniae* followed by *Mycoplasma* and *Chlamydia* (BTS, 2002). Viruses are the most common causes of pneumonia in younger children (infancy, pre-school and school age children), except neonates (BTS, 2002). In neonates, the most common causes are *Group B streptococcus beta haemolyticus* (GBB), *Escherichia coli*, *Klebsiella pneumoniae*, *Chlamydia trachomatis* (3-19 weeks), *Cytomegalovirus* (CMV), and the less common causes are *Staphylococcus aureus*, *Listeria monocytogenes*, and *Pseudomonas*. The common viruses are Respiratory Syncytial Virus (RSV), influenza, and parainfluenza viruses, adenovirus, and human metapneumovirus (HMV).

The spectrum and frequency of causative agents of bacterial pneumonia in severely malnourished children often differs in pneumonic children without severe malnutrition (Chisti et al., 2009).

Table 1. Causes of CAP according to various age groups and SAM

#### All age groups

Bacteria are the major causes of CAP in children

1. *Streptococcus pneumoniae*: commonest
2. *Haemophilus influenzae* (including Hib & nontypable strains)
3. *Staphylococcus aureus*
- Atypical bacteria\*
4. *Mycoplasma pneumoniae*\* (> 5 years)
5. *Chlamydia pneumoniae*\*
6. *Chlamydia trachomatis*\* (3-19 weeks)
7. *Moraxella catarrhalis*
- Gram-negative bacteria\*\*
8. *Klebsiella pneumoniae*\*\*
9. *Escherichia coli*\*\*
10. *Pseudomonas*\*\*
- Viruses\*\*\*
11. Respiratory syncytial virus (RSV)\*\*\*
12. Influenza A or B\*\*\*
13. Parainfluenza virus types 1, 3\*\*\*
14. Adenovirus\*\*\*
15. Human metapneumovirus (HMV)\*\*\*
16. Rhinovirus\*\*\*
17. Coronavirus\*\*\*
18. Enterovirus\*\*\*
19. CMV\*\*\*
20. *Pneumocystis jiroveci* (previously known as *Pneumocystis carinii*)

**0-2 months**

1. *Group B streptococcus*
- Gram-negative bacteria\*\*
2. *Klebsiella pneumoniae*\*\*
3. *Escherichia coli*\*\*
4. *Pseudomonas*\*\*
5. *Staphylococcus aureus*
6. *Chlamydia trachomatis*\*
7. *Listeria monocytogenes*
8. Viruses\*\*\*
9. *Ureaplasma urealyticum*
10. *Bordetella pertussis*

**2 months-5 years**

1. Viruses\*\*\*
2. *Streptococcus pneumoniae*
3. *Haemophilus influenzae* (including Hib & nontypable strains) (Common in developing countries where vaccination is still not widely used)
4. *Staphylococcus aureus*
5. *Mycoplasma pneumoniae*\*

**Above 5 years**

1. *Streptococcus pneumoniae*
2. *Haemophilus influenzae* (including Hib & nontypable strains)
3. *Staphylococcus aureus*
- Atypical bacteria\*
4. *Mycoplasma pneumoniae*\*
5. *Chlamydia pneumoniae*\*
6. Viruses\*\*\*

**SAM**

1. *Klebsiella pneumoniae*\*\* (26%)
2. *Staphylococcus aureus* (25%)
3. *Streptococcus pneumoniae* (18%)
4. *Escherichia coli*\*\* (8%)
5. *Haemophilus influenzae* (including Hib) (8%)
6. *Salmonella species*\*\*
7. *Pseudomonas*\*\*
8. *Acinetobacter species*\*\*
9. Methicillin-resistant *Staphylococcus aureus* (MRSA)
10. *Pneumocystis jiroveci* (previously known *Pneumocystis carinii*)
11. CMV\*\*\*
12. *Candida*

(N.B. \* indicates atypical bacteria  
indicates viruses)

\*\* indicates gram-negative bacteria \*\*\*



### **Risk factors for CAP**

Some risk factors for CAP are shown below (Zar et al., 2005; Chisti et al., 2009)

#### *Host factors*

- Infancy (Age <1 year)
- Prematurity
- Low birth weight (including low weight for age)
- Malnutrition
- Immunosuppression

#### *Social/environmental*

- Overcrowding
- Air pollution
- Inadequate housing
- Low socioeconomic status
- Passive exposure to tobacco smoke
- Indoor fuel exposure
- Winter season
- Lack of breast feeding
- Failure to complete immunization
- Attendance at day-care centres
- Presence of coughing sibling (s) at home

### **Pneumonia in SAM**

The WHO defines malnutrition as “the cellular imbalance between the supply of nutrients and energy and the energy and the body’s demand for them to ensure growth, maintenance, and specific functions (de Onis et al., 1993). Among the four principal causes of deaths in young children worldwide, undernutrition has been ascribed to be the cause of death in 60.7% children with diarrhoeal diseases, 52.3% of those with pneumonia (Caulfield et al., 2004). More than half of all the childhood deaths are associated with malnutrition (Rice et al., 2000). Pneumonia is common in malnourished children and frequently associated with fatal outcome (Bryce et al., 2005; Rice et al., 2000; Loeb & High, 2005; Nannan et al., 2007). Of children with malnutrition requiring hospital admission, up to two-thirds are diagnosed with pneumonia (Shimeles & Lulseged, 1994; Ahmed et al., 1999). A most recent systematic review revealed those children with pneumonia and moderate or severe malnutrition are at higher risk of death (Chisti et al., 2009). For SAM, the relative risks ranged from 2.9-121.2 with odds ratios ranged from 2.5-15.1. For moderate malnutrition, the relative risks ranged from 1.2-36.5 (Chisti et al., 2009). The clinical classification of pneumonia based on the diagnostic criteria according to the WHO guidelines should be carefully evaluated in children with SAM. A Gambian study evaluated that the respiratory rate cut-off required in malnourished children should be taken approximately 5 breaths per minute less than that in well nourished children and this finding may be related to the lower body temperatures found in severely malnourished children with pneumonia (Falade et al., 1995). Similarly, intercostal indrawing was more common and lower chest wall indrawing was less common in severely malnourished children (Falade et al., 1995; Chisti et al., 2010). In addition, lower chest wall indrawing is not sufficiently sensitive as predictors of pneumonia in SAM and no

visible clinical signs are consistently reliable for the diagnosis of pneumonia in SAM. Malnourished children may not have the strength to manifest some of these physical signs in the same manner as well nourished children (Falade et al., 1995; Chisti et al., 2010). Moreover, data from a recent systematic review suggest that a reliance on simple clinical signs will underestimate the burden of the disease and potentially delay the diagnosis of pneumonia in severely malnourished children (Chisti et al., 2009). Therefore, WHO recommends that children with SAM who present with cough, fast or difficult breathing irrespective of having clinical signs of pneumonia or not, should be treated with appropriate antibiotics to save the lives of these high-risk group of children (Falade et al., 1995).

Occult pneumonia is another entity characterized by the absence of the clinical signs and may be diagnosed by performing a chest radiograph; it may occur in SAM with dehydrating diarrhoea (Hall & Simon, 1987; Murphy et al., 2007; Bachur et al., 1999). The typical clinical signs of pneumonia may be absent in SAM due to sub-optimal inflammatory responses, reduced power of the respiratory muscles, and depletion of potassium and magnesium (Suskind et al., 1990). SAM also contributes to immune deficiency and reduced host defense.

#### **Pneumonia with dehydrating diarrhoea**

The clinical classification of pneumonia based on the diagnostic criteria according to WHO should be carefully evaluated in children presenting with dehydrating diarrhoea caused by *Vibrio cholerae*, *ETEC*, as well as rotavirus. If a child presents with severe or some degree of dehydration, then s/he would likely to have acidosis, which is responsible for the development of tachypnoea and it would be very difficult to distinguish clinically whether the increased respiratory rate is due to pneumonia, or due to acidosis, or both. In this situation, it is generally recommended to fully rehydrate the child first with IV/oral fluid (according to the type of dehydration present) within 4-6 hours and then count the respiratory rate for detecting pneumonia according to the standard WHO guideline. It is also recommended to perform a chest radiograph after full hydration of the patient for confirming the diagnosis of pneumonia.

#### **Diagnosis of CAP**

The diagnosis of CAP should be considered in any child who has an acute onset of respiratory symptoms, particularly cough, fast breathing, or difficulty in breathing. Diagnosis includes clinical evaluation, radiographic evaluation and etiological investigations to: (i) establish whether pneumonia is present; (ii) assess the severity of pneumonia; (iii) determine the clinical type of pneumonia; and (iii) determine the causative organism. In general, diagnostic investigations to determine the cause of pneumonia are indicated only in children requiring hospitalization (Zar et al., 2005). Bacterial pneumonia cannot be reliably distinguished from viral pneumonia on the basis of any single parameter: clinical, laboratory, or chest radiographic findings.

- Clinical evaluation
- Radiographic evaluation
- Aetiological investigations
- Pulse oximetry

The best objective measurement of hypoxaemia is by pulse oximetry which avoids the need for arterial blood gases. Oxygen saturation ( $SaO_2$ ) measurements provide a non-invasive estimate of the arterial oxygenation (BTS, 2002). The human eye is poor in recognizing

hypoxaemia. Even under ideal conditions, skilled observers cannot consistently detect hypoxaemia until the oxygen ( $O_2$ ) saturation is below 80% (Comroe & Bothello, 1947). Pulse oximetry is probably one of the most important advances in monitoring the respiratory problems and these instruments have a reasonable degree of accuracy (Jurban, 1999). The pulse oximeter is easy to use and requires no calibration. However, it requires a pulsatile signal from the patient. When using paediatric wrap around probes, the emitting and receiving diodes need to be carefully opposed. It is also highly subject to motion artifacts. To obtain a reliable reading (i) the child should be resting, still and quiet, not crying or irritable (Zar et al., 2005; BTS, 2002); (ii) a good pulse signal (plethysmograph) should be obtained; and (iii) once a signal is obtained, the saturation reading should be watched over at least 30 seconds and a value recorded once an adequate stable trace is obtained (Zar et al., 2005; BTS, 2002). Pulse oximetry as a potentially useful diagnostic tool for the detection of hypoxaemia as an indicator of severe pneumonia; have not been sufficiently evaluated for the diagnosis of children with SAM (WHO, 2005). Widespread use of pulse oximetry is now recommended for monitoring children with severe pneumonia in the developing countries. Perhaps the major challenge facing pulse oximetry is whether this technology can be incorporated effectively into the diagnostic and management algorithms that can improve the efficiency of clinical management of CAP in the developing countries (Jurban, 1999).

### **Significance of hypoxaemia with CAP and its management**

Assessment of oxygenation is important in the evaluation of a child with pneumonia and pulse oximetry should be performed in every child admitted to a hospital with CAP (Zar et al., 2005; BTS, 2002). Hypoxaemia is defined as the arterial oxygen saturation of less than 90% in room air at sea level as recorded by the pulse oximetry, which is the most serious manifestation of childhood pneumonia (Weber et al., 1997). In white patients, an  $S_{pO_2}$  target of 92% resulted in a satisfactory level of oxygenation, whereas a higher  $S_{pO_2}$  target of 95% was required in black patients (Jurban, 1999; Jurban & Tobin, 1990). Alternatively, no hypoxaemia is defined as the arterial oxygen saturation of  $\geq 90\%$  in room air as recorded by the pulse oximetry. The median prevalence of hypoxaemia in WHO-defined pneumonia requiring hospitalization (severe and very severe pneumonia) was 13% but the prevalence varied widely (Subhi, 2009). This corresponds to at least 1.5 to 2.7 million annual cases of pneumonia with hypoxaemia presenting to the health-care facilities (Subhi, 2009). WHO recommends for children older than 2 months, the use of oxygen in severe/very severe pneumonia, as ascertained by the presence of a number of clinical indicators of hypoxaemia, including cyanosis, inability to drink, severe lower chest wall indrawing, respiratory rate greater than 70 breaths per minute, grunting respiration, or head nodding (WHO, 2005). Head nodding is a movement of the head synchronous with each breath, which is caused by increased use of auxiliary muscles of respiration and therefore indicates severe respiratory distress and an important clinical sign predicting hypoxaemia (Weber et al., 1997). Only one Gambian study showed that hypoxaemia could be predicted in only half of the children by the presence of a combination of three clinical signs, such as extreme respiratory distress, cyanosis, and severely compromised general status (Weber et al., 1997). Agitation may be an indication that the child is hypoxaemic (BTS, 2002). Hypoxaemia is also a good indicator for detecting the severity of pneumonia (Wang et al., 1995; Hall et al., 1979; Shann et al., 1989). It is a common complication of childhood infections, particularly ALRI, and case fatality rate of pneumonia is inversely related to the arterial haemoglobin oxygen saturation ( $SaO_2$ )

(Onyango et al., 1993). In pneumonia - a disease that disproportionately impacts the developing countries, hypoxaemia is an important risk factor for death (Onyango et al., 1993; Duke et al. 2000), as hypoxaemic children are five times more likely to die than non-hypoxaemic children. In patients admitted with pneumonia to a general medical service, it was found that O<sub>2</sub> saturation <90% of at least 5 minutes duration occurred in 26% of the patients (Bowton et al., 1994). On follow-up over the next 4-7 months, those patients experiencing hypoxaemia during the first 24 hour of hospitalization had more than a threefold higher mortality than patients who did not desaturate (Bowton et al., 1994). In the critical care setting especially for evaluating the progress of children suffering from severe pneumonia, pulse oximetry is also used as one of the most commonly employed monitoring modalities in the critical care setting especially for evaluating the progress of children suffering from severe pneumonia (Jurban, 1999). Moreover, hypoxaemia is also an important risk factor for failure to the day-care management as well as the need for future follow-up admissions (Ashraf et al., 2010). Therefore, measurement of oxygen saturation should be routinely done in all children with severe pneumonia with pulse oximetry, and those with hypoxaemia requiring prolonged oxygen therapy for more than six hours should be referred to a hospital for long-term oxygen therapy (Ashraf et al., 2010). Results of that study were consistent with earlier reports that hypoxaemia in pneumonic children are predictors of severe disease and is a risk factor for death (Onyango et al., 1993; Duke et al., 2000). Hypoxaemia has been overlooked in world-wide strategies for pneumonia control and reducing child mortality (Subhi, 2009). It is also often overlooked in the developing countries, mainly due to the low accuracy of clinical predictors and the limited availability of pulse oximetry, despite of its more accurate detection of hypoxaemia and oxygen therapy for treatment (Subhi, 2009). Many more people do not have access to the health care facilities. Oxygen therapy in developing countries continues to be a low priority on the child health agenda (Subhi, 2009). Oxygen was never mentioned in the recent publication by the WHO and UNICEF efforts to control pneumonia (Wardlaw et al., 2006). The accurate detection of hypoxaemia is important as delivery of oxygen to the hypoxaemic children may improve the outcome. Especially in a setting where oxygen has to be bought in cylinders, a pulse oximeter might be a cost effective purchase, as it allows identification of children who are in need of oxygen, and the amount of oxygen given can be titrated to the actual need of the patient, thus avoiding unnecessary wastage of valuable oxygen (Weber et al., 1997). There is now evidence that ensuring ample supplies of oxygen and promoting a routine and systematic approach of screening for hypoxaemia by using pulse oximetry is associated with improved quality of care and reduced mortality, and that the technology required to do so is affordable and sustainable in district level hospitals and day-care centres in the developing countries (Duke et al., 2000, Dobson et al., 1996; Duke et al., 2008; Dobson, 1991; Steinhoff & Black, 2007; Matai et al., 2008). Pulse oximetry would enable accurate identification of hypoxaemia and might increase the safety and cost effectiveness of this recommendation and this diagnostic tool should be included. This was demonstrated by our recent study reporting successful day-care case management of severe pneumonia using pulse oximetry as an important part of the treatment algorithm (Ashraf et al., 2008). Hypoxaemia is a very common and treatable complication of childhood pneumonia in developing countries and it is a recognized predictor of severe disease and a risk factor for death and correlates with disease severity. For home management to be safe and ethical, it is essential that only children without hypoxaemia are managed outside health-care facilities (Subhi et al., 2009).

Children with hypoxaemic pneumonia requiring prolonged oxygen therapy for more than six hours, need to be identified (which is often difficult using only clinical signs), admitted, and given supplemental oxygen for prolonged duration, and close monitoring (Ashraf et al. 2010). The measurement of SpO<sub>2</sub> as a regularly measured vital sign by using pulse oximetry should be incorporated as an important part of the treatment algorithm for improving the diagnosis of hypoxaemia, and categorizing the severity of pneumonia (Onyango et al., 1993; Steinhoff & Black, 2007; Lozano et al., 1994; Weber & Mulholland, 1998). For treating children with pneumonia, there is an urgent need to increase the widespread availability as well as use of pulse oximetry for monitoring patients with severe pneumonia and effective oxygen delivery systems in the developing countries.

### **Classification of hypoxaemia**

There are two ways of classifying hypoxaemia in children: (i) WHO classification and (ii) British Thoracic Society (BTS) classification as defined below:

#### **(i) WHO classification of hypoxaemia**

Experts from WHO often classifies hypoxaemia as mild, moderate and severe as defined below:

- Mild hypoxaemia: when the arterial oxygen saturation lies between 85 to 90%, the patient is known to have mild hypoxaemia.
- Moderate hypoxaemia: when the arterial oxygen saturation lies between 80 to 85%, the patient is known to have moderate hypoxaemia.
- Severe hypoxaemia: when the arterial oxygen saturation is less than 80%, the patient is known to have severe hypoxaemia.

#### **(ii) BTS classification of hypoxaemia**

Similarly, experts from paediatric respiratory medicine in developed countries including BTS often categorized hypoxaemia as mild, moderate and severe as defined below:

- Mild hypoxaemia: when the arterial oxygen saturation lies between 88 to 92%, the patient is known to have mild hypoxaemia.
- Moderate hypoxaemia: when the arterial oxygen saturation lies between 85 to 88%, the patient is known to have moderate hypoxaemia.
- Severe hypoxaemia: when the arterial oxygen saturation is less than 85%, the patient is known to have severe hypoxaemia.

### **Indications for oxygen therapy**

1. Hypoxaemia (oxygen saturation <90% in room air at sea level)
2. Central cyanosis
3. Severe lower chest-wall in-drawing
4. Grunting respiration
5. Restlessness (due to hypoxaemia)
6. Inability to drink or feed
7. Respiratory rate >70 breaths/min
8. Head Nodding

**Management of hypoxaemia**

- Oxygen should be available at any health care facility where sick children are seen regularly. Oxygen therapy reduces mortality associated with severe pneumonia. It should be given to children who are restless, who had tachypnoea with severe lower chest wall in-drawing, head nodding, cyanosis, or not tolerating oral feeds. The SpO<sub>2</sub> should be maintained above 92%.
- If oxygen is required infrequently then cylinders are the most practical source of oxygen. Cylinders also allow oxygen therapy to be used while the patient is transferred to a facility with more resources.
- If oxygen is used more frequently, then oxygen concentrators are the preferred source of oxygen.
- In hospitals with oxygen supplies, wall oxygen units should be available.
- Low flow meters must be available to give appropriate oxygen flow to children. In most hospitals these will be variable orifice units, but fixed orifice units may be more practical in some units.

**Key messages about hypoxaemia in relation to CAP (WHO, 2009)**

- Hypoxaemia is a common complication in ALRI in children, and is a strong risk factor for death.
- At least 13% of children presenting to hospitals with severe or very severe pneumonia have hypoxaemia, and the rates are much higher in some hospitals; some exceeding 50%.
- The prevalence of hypoxaemia is higher in referral hospitals than in primary care settings. Hypoxaemia is more common at higher altitude, in younger ages and in certain geographical regions.
- SpO<sub>2</sub> <90% is the most clinically useful definition of hypoxaemia and is considered by most clinicians as an appropriate indication for giving oxygen.
- If pulse oximetry is available only at the time of admission, screen all patients if time allows, or those patients with any clinical signs of hypoxaemia, including all children with emergency or priority signs.
- If oximetry is used at outpatient triage, screen all children with any emergency or priority signs.
- Any child with an SpO<sub>2</sub> <90% should receive oxygen.
- Use oximetry, at least daily, to check any patients who are already on oxygen, and screen any patient who develops any emergency signs or shows other clinical signs of deterioration.
- Explain the meaning of oximetry to parents. This will help them understand the importance of oxygen and other treatments and will involve them in their child's care.

- Children should not be discharged until their SpO<sub>2</sub> has been stable at 90% or more while breathing room air for at least 24 hours, until all emergency and priority signs have resolved, and until appropriate home treatment can be organized.

### Methods of oxygen administration

- Nasal prongs: are recommended for most children. Nasal prongs give a maximum fractional concentration of inspired oxygen (F<sub>1</sub>O<sub>2</sub>) of 28-35% except in small infants when higher concentrations may be obtained. This method does not require humidification of oxygen and ensures that the child receives oxygen during feeding. Oxygen flow rates of 0.5-1 l/minute are required in children less than 2 months old and 2-3 l/minute in infants and children aged 2 months to 5 years.
- Nasal catheters: are usually well tolerated and humidification is not required, but they can be blocked by mucous. Oxygen via nasal catheters gives a maximum F<sub>1</sub>O<sub>2</sub> of 35-40%.
- Nasopharyngeal catheters: have the advantage of requiring the lowest flow rate to achieve a given oxygen concentration in the airways. Infants under the age of 2 months can usually be treated with 0.5 l/minute and infants up to 1 year with 1 l/minute. However, humidification of oxygen is required and the catheter may be easily blocked. Further, potentially lethal complications including gastric distension, airway obstruction, apnoea, pneumo-orbitus and pneumocephalus may occur. Continuous skilled nursing is therefore necessary to prevent these complications. Consequently, oxygen administration via nasopharyngeal catheter is not recommended.
- Headbox: oxygen is well tolerated by young infants. Headbox oxygen requires no humidification but requires a high flow and a mixing device to ensure the correct F<sub>1</sub>O<sub>2</sub> is delivered. This is the *least preferred* method as there is wastage of oxygen and delivered F<sub>1</sub>O<sub>2</sub> is unpredictable.
- Facemask: oxygen is designed to deliver 28%-65% oxygen at a flow rate of 6-10 l/minutes.
- Polymask: In severely hypoxaemic infants who are not ventilated, oxygen should be administered using a polymask whereby F<sub>1</sub>O<sub>2</sub> concentrations of 60-80% may be achieved. The flow rate should be regulated to keep the bag of the mask inflated during inspiration and expiration.
- Using the prone position for infants may improve hypoxaemia and the respiratory system compliance (Chaisupamongkollarp et al., 1999) and should be attempted if hypoxaemia is difficult to treat.
- Oxygen should be discontinued when the child is improving and the transcutaneous saturation is above 90% in room air, as recorded by the pulse oximetry.

### **Radiological diagnosis**

Radiological changes may be vague or inconclusive or even absent despite the presence of clinical signs of pneumonia (Doherty, 1991; Hamid et al., 1996; Wafula et al., 1998; Chisti et al., 2009). Conversely, clinical signs of pneumonia can be absent in the presence of radiological signs of pneumonia (Murphy et al., 2007; Bachur et al., 1999; Chisti et al., 2009). There are two main clinical definitions of pneumonia based on the radiological findings:

Bronchopneumonia is defined as a febrile illness with cough, respiratory distress with evidence of localized or more than one or generalized patchy infiltrates on the chest x-ray.

Lobar pneumonia is defined as a febrile illness with cough, respiratory distress with an illness similar to that of bronchopneumonia, except that the physical findings (affected lobe reveals woody dull on percussion, rales, increased vocal resonance and/or bronchial breath sound on auscultation) and radiographic examination indicate lobar consolidation.

Occult pneumonia: It was observed that approximately 25% of the febrile children defined as the rectal temperature becoming more than 38<sup>o</sup> Celsius (Ashraf et al., 2010), with a WBC count >20,000/mm<sup>3</sup>, but without any lower respiratory tract findings on examination, had radiographic evidence of pneumonia, known as occult pneumonia, commonly found in children with SAM and children with dehydrating diarrhoea (Murphy et al., 2007; Bachur et al., 1999; Chisti et al., 2009).

### **Limitations of chest radiography**

- Less useful in discriminating the causative pathogens
- Cannot accurately discriminate viral from bacterial pneumonias (Swingler, 2000).
- Wide range of inter- and intra-observer variation in the interpretation (Swingler, 2001; Bada et al., 2007; Pauls et al., 2007; Sarria et al., 2003), not only among the paediatricians, but also among the radiologists, even paediatric radiologists
- Does not result in the improved outcome or change in the treatment of ambulatory settings (Swingler et al., 2000)

### **Indications for CXR**

- To confirm the presence of pneumonia
- To detect clinical pneumonia, unresponsive to the standard ambulatory management
- To identify suspected cavitations or military mottling in PTB
- To identify suspected foreign body aspiration
- To identify hospitalized children for the detection of local complications of pneumonia, such as pleural effusion, pneumothorax, empyema thoracis, advanced stage of bronchiectasis, and lung abscess
- High fever, leukocytosis with no obvious focus of infections (26% such cases may have radiographic pneumonia) (Bachur et al., 1999).
- Children developing secondary heart failure
- Children with congenital problems, such as Congenital Heart Disease (CHD), cystic fibrosis, Down's Syndrome
- Children with re-current attacks of pneumonia



### Indications for follow-up CXR

- Children with lobar collapse
- To document the resolution of a round pneumonia (as this may mimic the appearance of a Ghon focus)
- Children with ongoing respiratory symptoms
- Children with re-current attacks of pneumonia

### General tests of infection

They may not be useful to differentiate bacterial from viral pneumonia (Nohynek et al., 1995; Toikka et al., 2000; Korppi et al., 2003)

- WBC count, neutrophil count (PMN leukocytosis suggests bacterial pneumonia (Marks & Klein, 1995; Klein, 1992) and lymphocytic leukopenia suggests viral pneumonia) (Austrian & Gold, 1964)
- C-reactive protein (>40 mg suggests bacterial infection)
- ESR
- Procalcitonin
- Copeptin
- Blood culture: to identify the causative bacterial pathogen, that is possible in only less than 30% (10-30%) of cases of CAP (Donowitz & Mandell, 1990), and to determine its sensitivity
- Pleural fluid: if present, should be aspirated and investigated for specific infectious agents

The C-reactive protein, ESR, procalcitonin, and copeptin are the non-specific markers for the diagnosis of pneumonia.

### Indications for hospital admission

1. Hypoxaemia (oxygen saturation <90% in room air at sea level)
2. Toxic appearance
3. Respiratory rate >70/minute, or severe respiratory distress
4. Infants < 2 months
5. Impaired level of consciousness
6. Inability to drink or eat
7. Cyanosis
8. Stridor in calm child
9. Chronic lung disease
10. Systemic manifestation
11. Intermittent apnoea
12. Grunting respiration
13. Severe lower chest-wall indrawing
14. SAM
15. Family unable to provide adequate care/non-compliant parents
16. Failure to respond ambulatory care/no response to previous oral antimicrobial therapy
17. Clinical deterioration on treatment
18. Immunocompromised host/immunodeficiency
19. Recurrent pneumonia

**Indications for transfer and admission to Paediatric Intensive Care Unit (PICU)**

1. Needs ventilator support
2. Failure to maintain a saturation of >90% with oxygen therapy
3. Apnoea or slow irregular breathing
4. Severely acidotic patient
5. Exhaustion with rising respiratory rate and pulse rate
6. Patient is in shock

**Differential diagnosis of a child with cough, or difficult breathing****Diagnosis****Points in favour****1. Pneumonia**

Fast breathing  
 Lower chest wall indrawing  
 Crepitations on auscultation  
 Bronchial breathing  
 Nasal flaring  
 Grunting respiration  
 Head nodding

**2. Cardiac failure**

Tachycardia  
 Tachypnoea  
 Enlarged tender liver  
 Dependent oedema  
 Raised jugular venous pressure  
 Central cyanosis  
 Heart murmur  
 Gallop rhythm

**3. Pneumothorax**

Sudden onset  
 Sudden onset of unexplained tachypnoea and respiratory distress (disproportionate to the severity of pneumonia)  
 Hyper-resonance chest on the affected side on percussion  
 Shift of mediastinum (trachea, apex beat to opposite side)  
 Diminished or absent breath sound on auscultation on the affected side

**4. Pleural effusion, empyaema**

Stony dull on percussion (it is difficult to elicit in young child)  
 Diminished breath sound on the affected side of lesion  
 Shift of mediastinum (trachea, apex beat to opposite side)

**5. Pericardial effusion**

Oedema feet  
 Raised jugular venous pressure  
 Apex beat not visible/not palpable  
 Increased area of cardiac dullness

Heart sound (muffled or absent)  
Pulsus paradoxus  
Enlarged liver

### **Pneumonia in older children often present in many ways**

The following two classic presentations have been described for pneumonia (Jadavi et al., 1997)

#### Typical presentation (predominantly respiratory signs)

- Fever
- Chills
- Pleuritic chest pain
- Productive cough
- Fast breathing
- Lower chest wall indrawing
- Cyanosis

#### Atypical presentations (single or in combinations) gradual onset over several days to weeks:

- Nonproductive cough
- Low-grade fever
- Headache
- Malaise
- Meningism
- Acute abdominal pain
- Acute pain in chest or shoulder
- Convulsion

### **Antibiotic use in the treatment of CAP**

When treating CAP, the clinical, laboratory and radiographic findings should be considered, especially when the child is hospitalized. As it is difficult to distinguish bacterial from viral pneumonia and because of the frequency of mixed bacterial-viral infections ( $\approx 30-40\%$ ) (Zar et al., 2005), all children with CAP would require an antibiotic. The age of the child, nutritional status, immunologic status of the host, local epidemiology of respiratory pathogens, and sensitivity of these pathogens to particular antimicrobial agents and the emergence of antimicrobial resistance usually play a big role to determine the choice of antibiotic therapy. The severity of pneumonia and the drug costs also have a great impact on the selection of antimicrobial therapy, particularly in the developing countries. The management of a child with CAP involves a number of decisions regarding treatment with antibiotics (BTS, 2002).

- Whether to treat with antibiotics?
- Which antibiotic and by which route?
- When to change to oral treatment?
- Total duration of antibiotic therapy;
- When to change antibiotic and why?

### Outpatient management

In children with mild pneumonia (non-severe pneumonia), the breathing is fast, but there is no lower chest wall indrawing. Oral antibiotics at an appropriate dose for an adequate duration are effective for treatment. The mother is advised to return in two days for reassessment, or earlier if the child appears to deteriorate.

Antimicrobial agent	Recommended dosage	Route	Comment
<b>Neonate (0-4 weeks)</b>			
Ampicillin plus Gentamicin	100 mg/kg/day 12 hourly 6 mg/kg/day 12 hourly	I/V I/V	1 <sup>st</sup> line of treatment
Ceftazidime plus Flucloxacillin	75-100 mg/kg/day 8 hrly 100 mg/kg/day 6 hourly	I/V I/V	2 <sup>nd</sup> line of treatment
<b>Infants &gt; 4-8 weeks</b>			
Ampicillin plus Gentamicin	100 mg/kg/day 6 hourly 6 mg/kg/day 12 hourly	I/V I/V	1 <sup>st</sup> line of treatment
Ceftriaxone plus Gentamicin	75-100 mg/kg/day od 6 mg/kg/day 12 hourly	I/V I/V	2 <sup>nd</sup> line of treatment
<b>Infants &gt;8 weeks to children of 5 years plus above 5 years</b>			
Amoxicillin	100 mg/kg/day 8 hourly	Oral	1 <sup>st</sup> line of treatment
Azithromycin	10 mg/kg/day once daily	Oral	2 <sup>nd</sup> line of treatment
Ceftriaxone	75-100 mg/kg/day od	I/V	3 <sup>rd</sup> line of treatment
<b>Severe pneumonia with SAM</b>			
Ampicillin plus Gentamicin	100 mg/kg/day 6 hourly 6 mg/kg/day 12 hourly	I/V I/V	1 <sup>st</sup> line of treatment
Ceftriaxone plus Gentamicin	75-100 mg/kg/day od 6 mg/kg/day 12 hourly	I/V I/V	2 <sup>nd</sup> line of treatment
Ceftazidime plus Flucloxacillin	75-100 mg/kg/day 8 hrly 100 mg/kg/day 6 hourly	I/V I/V	3 <sup>rd</sup> line of treatment
<b>Suspected Staph aureus pneumonia</b>			
Amoxicillin plus Flucloxacillin (Oxacillin/ Nafcillin for MRSA)	100 mg/kg/day 8 hourly 100 mg/kg/day 6 hourly	I/V I/V	1 <sup>st</sup> line of treatment
Ceftriaxone plus Flucloxacillin (Clindamicin/Vancomycin for MRSA)	75-100 mg/kg/day od 100 mg/kg/day 6 hourly	I/V I/V	2 <sup>nd</sup> line of treatment
Vancomycin	10 mg/kg iv 8 hourly	I/V	3 <sup>rd</sup> line of treatment

Table 2. Antimicrobial treatment of CAP

<b>Suspected pseudomonas pneumonia</b>			
Ceftazidime plus Ciprofloxacin	100 mg/kg/day 8 hrly 20 mg/kg/day 12 hourly	I/V I/V	1 <sup>st</sup> line of treatment
<b>Suspected PCP pneumonia</b>			
Co-trimoxazole (SXT)	20 mg/kg loading dose followed by 10 mg/kg in 2 divided doses or, 20 mg/kg/day in 4 divided doses	Oral	1 <sup>st</sup> line of treatment. Therapy to be continued for at least 14 days (sometimes 21 days)
<b>Suspected hospital acquired pneumonia</b>			
<b>a) For early onset HAI (if HAI occurs &lt; 96 hours of admission)</b>			
Fluroquinolone (e.g. Ciprofloxacin) plus aminoglycosides (e.g. Gentamicin)	20 mg/kg/day in 2 divided doses  6 mg/kg/day 12 hourly	I/V  I/V	1 <sup>st</sup> line of treatment
3 <sup>rd</sup> generation cephalosporin e. g. Ceftriaxone Monotherapy with Fluroquinolone	75-100 mg/kg/day od  6 mg/kg/day 12 hourly	I/V  I/V	2 <sup>nd</sup> line of treatment (HAI where we don't suspect <i>Staph</i> )
3 <sup>rd</sup> generation cephalosporin e. g. Ceftriaxone plus Flucloxacillin	75-100 mg/kg/day od 100 mg/kg/day 6 hourly	I/V I/V	3 <sup>rd</sup> line of treatment (HAI where we suspect <i>Staph</i> )
<b>b) For late onset HAI (if HAI occurs &gt; 96 hours of admission)</b>			
Anti-pseudomonal penicillin (e. g. Ticarcillin/clavulanic acid) plus Anti-MRSA (e. g. Vancomycin)	50 mg/kg/dose (Ticarcillin base: max. 3.0 gm/dose) 6 hourly 10 mg/kg iv 8 hourly	I/V  I/V	1 <sup>st</sup> line of treatment
Anti-pseudomonal cephalosporin (e. g. Ceftazidime) plus Aminoglycosides with wide coverage (e.g. Amikacin), or Anti-MRSA (e. g. Vancomycin)	75-100 mg/kg/day 8 hrly  6 mg/kg/day 12 hourly 10 mg/kg iv 8 hourly	I/V  I/V I/V	2 <sup>nd</sup> line of treatment

### Duration of antibiotic treatment

Antibiotic therapy is generally recommended for 5 to 7 days for uncomplicated cases of childhood pneumonia (Zar et al., 2005; Mehta, 2003). For suspected *Staphylococcus aureus* infection, the duration of treatment may be extended for 14 to 21 days, depending on the clinical response. Gram-negative bacilli, or *Legionella* species, may also require longer courses of therapy for 10 to 21 days (BTS, 2002; Mehta, 2003). Alternatively, the newer macrolides such as azithromycin may be used for 3-5 days (Zar et al., 2005). In case of neonates, the treatment should be continued for at least 2 weeks. If pneumonia is complicated with empyema, the treatment should be continued for at least 4 weeks.

Switching from parenteral therapy to oral therapy is a key management issue for childhood pneumonia. Children receiving parenteral therapy for 2 to 4 days can usually be switched to oral therapy provided there is clinical improvement as children becoming afebrile defined as the rectal temperature becoming 38<sup>o</sup> Celsius or less and remaining so for at least 24 hours (Ashraf et al., 2010), they can tolerate medication orally, they do not have any diarrhoea and vomiting and have no relevant local complications of pneumonia (Shalit et al., 1994; Dagan et al., 1994). Switching over to oral antibiotics will help for early discharge from the hospital and subsequently prevent hospital acquired infection and vacant hospital beds for other sick children.

#### **Indications for the use of antipyretics and analgesics in CAP**

- Rectal temperature >39<sup>o</sup> Celsius
- There is a known risk of febrile convulsions
- There is central nervous system pathology that may be aggravated by high fever

Children with CAP are generally pyrexial and may also have some pain, including headache, chest pain, arthralgia (in cases of *Mycoplasma pneumoniae*), referred abdominal pain, and possibly earache from associated otitis media. Pleural pain may interfere with the depth of breathing and may impair the ability of the child to cough. Antipyretics and analgesics can be used to keep the child comfortable and to help coughing. Minimal handling helps to reduce metabolic and oxygen requirements and this should be considered when planning and carrying out procedures, investigations, and treatments. Pain associated with pneumonia may be due to pleurisy or to pathology involving the upper airways. Pain or discomfort should be treated as it may severely compromise respiratory function and adequate clearance of secretions. The most appropriate agent is paracetamol at a dose of 15 mg/kg/dose given 4-6-hourly orally or 20-40 mg/kg/dose per-rectally for two-three times daily. If this dose does not provide adequate analgesia, a mixture of paracetamol and codeine (0.5 mg/kg/dose 8-hourly) is very effective. Aspirin is contraindicated in most children because of the association with Reye's syndrome (Zar et al., 2005).

#### **Calorie requirements**

Adequate nutrition is of particular concern, especially when there are underlying factors such as malnutrition. A minimum of 50-60 kcal/kg/day should be given to a child with pneumonia with continuation of regular breast feeding for breast-fed children. A calorie intake of 80-100 kcal/kg/day should be given to a non-breast fed child with CAP. Ensuring adequate calorie intake is essential as there is an excessive demand on the energy reserves in children with pneumonia, in whom the work of breathing is increased. Children should not be starved for more than 24 hours to prevent the development of hypoglycaemia. In the presence of malnutrition, and following several days of poor nutrition, this needs to be increased considerably. In the early phase of pneumonia, ketosis should be avoided by ensuring adequate carbohydrate intake. With time, a greater proportion of intake can be lipids. The intake of calories should be adequate to meet the metabolic requirements and to promote growth.

**Enteral feeds**

Children with pneumonia should be encouraged to feed orally unless there are indications for nasogastric feeding/intravenous fluid infusions. If children are too distressed to take fluid and feeds orally, continuous enteral feeds via a nasogastric tube may be provided.

**Indications for N/G tube feeding**

- Too distressed to drink or swallow safely
- Having frequent severe coughing episodes that may be associated with vomiting and possible aspiration of gastric contents
- Hypovolaemia with associated poor peripheral perfusion (may even require I/V fluid)
- Painful oral sore/condition which interfere with feeding by mouth

**Fluid therapy**

Children with uncomplicated pneumonia should receive normal maintenance fluids. Appropriate rehydration is required in children who are dehydrated. Oral intake should cease when a child is in severe respiratory distress. In severe pneumonia, inappropriate secretion of anti-diuretic hormone (ADH) is increased (Dhawan et al., 1992), dehydration is therefore uncommon. It is important that the child should not be over hydrated. A study of 264 hospitalized children with CAP in India has shown hyponatraemia on admission in 27% cases and it was calculated that, in 68% of these children, the hyponatraemia was secondary to inappropriate ADH secretion (SIADH) (Singhi & Dhawan, 1992). Treatment is with fluid restriction.

Intravenous fluids must be used with great care and with caution, and only if adequate monitoring is available (Zar et al., 2005). Children who are vomiting or who are severely ill may require intravenous fluids. These should be given at 80% of the basal levels (once hypovolaemia has been corrected). In children with severe or complicated pneumonia, serum urea and electrolytes should be measured before instituting I/V fluids as among them (SIADH) is common. In these children, fluid intake should be restricted to 40-60% of normal requirements, i.e. 50 ml/kg/day of I/V fluids. They should be frequently monitored as severely ill children with CAP might develop SIADH as a recognized complication (Dhawan et al., 1992; Singhi & Dhawan, 1992).

**Indications for I/V fluid**

- Shock
- Inability to tolerate enteral feeds
- Sepsis
- Severe dehydration
- Gross electrolyte imbalance
- Hypoglycaemia

**Monitoring of the child with CAP**

The following parameters should be routinely monitored in every child with CAP, the frequency of which will depend on the severity of illness as well as the availability of

resources. Special attention should be given to child receiving I/V fluid therapy and those with SAM.

- Heart rate
- Respiratory rate
- Temperature
- Respiratory pattern including chest recession
- Lower chest wall indrawing
- Use of accessory muscles
- Establishment of oral feeding
- Liver size
- Oxygen saturation level
- Chest auscultation: rales, rhonchi, bilateral basal crepitation, pleural rub
- Fluid and calorie intake

Children on oxygen therapy should have at least 4-hourly observations of all the above parameters and children without getting oxygen should be observed at least 12-hourly. If a child remains pyrexial or unwell 48 hours after admission with pneumonia, re-evaluation is necessary with consideration given to exclude possible local complications of CAP (BTS, 2002).

#### **Micronutrient supplementation**

In children with CAP, as an adjuvant therapy with antibiotics, 20 mg of Zinc po daily until discharge was found to accelerate the recovery from severe pneumonia, reducing the duration of hypoxaemia (Brooks et al., 2004; Shakur et al., 2004; Mahalanabis et al., 2002). Zinc should therefore be considered for use in children hospitalized with CAP, particularly if there is co-existing SAM. Zinc also reduces the incidence of pneumonia, especially in children with SAM. But, studies from Vellore, India showed that zinc has no significant role to reduce the morbidity and mortality in children with acute severe pneumonia (Bose et al., 2006). Zinc should therefore be considered cautiously for use in children hospitalized with CAP.

#### **Non-response to therapy**

If a child remains pyrexial or unwell 48 hours after admission with CAP, re-evaluation is necessary with consideration given to the possible complications:

- Inappropriate drug (antibiotic): in relation to choice of antibiotic, adequate dosage, route of administration, and duration of antibiotic therapy
- Development of local lung complications of CAP such as pleural effusion, empyema, lung abscess, bronchieactasis, pneumothorax, liver abscess etc.
- Immunosuppression
- Coexisting disease such as cystic fibrosis, chronic suppurative bronchitis, congenital heart disease (CHD), Down's Syndrome
- Underlying SAM: slow, delayed and poor response to conventional antibiotic therapy
- Underlying TB (PTB, miliary TB, disseminated TB, TBM)



- Underlying malignancy
- Underlying HIV infection
- Development of heart failure
- Metastatic infection can rarely occur as a result of the septicaemia associated with CAP. Osteomyelitis or septic arthritis should be considered, particularly with *Staph aureus* infections.
- Incorrect diagnosis of CAP

### Complications of CAP

- Fortunately, most children with CAP recover without any complication
- Persistent effusions and empyemas are the most common serious complications of bacterial pneumonia
- Pulmonary abscess
- Respiratory distress
- Sepsis (bacteraemia may occur in 10-30% cases of pneumonia; sepsis in <10% cases especially in SAM)
- Tension pneumothorax (very rare: less than 1%)

### Patient education

- Parents should be cautioned to look for the signs of increasing respiratory distress, danger signs of very severe pneumonia, clinical signs of hypoxaemia and advised to seek medical attention immediately if any of these signs appear
- Most children with CAP treated with outpatient antibiotics will be much improved within 48 hours after the initiation of treatment. If such improvement does not occur, medical attention should be sought.

### Medicolegal Pitfalls

- Attempting to treat neonates and very young infants on an outpatient basis
- Failure to recognize and treat signs of respiratory compromise, heart failure, and sepsis
- Failure to recognize and treat associated SAM
- Failure to give parents clear discharge instructions

### Measures of no value in the treatment of CAP

- Chest physiotherapy: The function of chest physiotherapy is to assist the removal of tracheobronchial secretions resulting in an increase of gas exchange and reduction in work of breathing. However, trials have found no clinically discernible benefit or impact of chest physiotherapy on the course of illness in bronchiectasis, cystic fibrosis, pneumonia, bronchiolitis, asthma, acute atelectasis, inhaled foreign body and post extubation babies. There is no evidence to support the use of chest physiotherapy including postural drainage, percussion of the chest, or deep breathing exercises that should be routinely performed in children with uncomplicated CAP (Levine, 1978; Britton et al., 1985; Stapleton, 1985; Wallis & Prasad, 1999). There is a suggestion that physiotherapy is counterproductive, with patients who receive chest physiotherapy being at risk of having a longer duration of fever than the control group (Britton et al., 1985). In addition, there is also no

evidence to show that physiotherapy is beneficial in the resolving stage of pneumonia (Wallis & Prasad, 1999). Therefore, chest physiotherapy is not beneficial and should NOT be routinely prescribed for children with CAP. A supported sitting position may help to expand the lungs and improve the respiratory symptoms in children with respiratory distress. Chest physiotherapy only works in children with pneumonia having chronic lung disease (cystic fibrosis, suppurative chronic lung disease, primary ciliary dyskinesia).

- Mucolytic agents: Anti-tussive remedies are not recommended as they cause suppression of cough and interfere with airway clearance. Adverse effects and overdose have been reported. Therefore, they should not be advised in children with CAP.
- Postural drainage: There is no evidence for the use of a head-down position for postural drainage.
- Nebulized bronchodilators: Nebulized bronchodilators or saline do not improve the outcome of CAP.
- Corticosteroids: There is no evidence to support the use of oral or inhaled corticosteroids in CAP.

## References

- Ahmed T, Ali M, Ullah MM et al. Mortality in severely malnourished children with diarrhoea and use of a standardised management protocol. *Lancet* 1999;353:1919-22.
- Ashraf H, Ahmed T, Hossain MI, Alam NH, Mahmud R, Kamal SM, Salam MA, Fuchs GJ. Day-care management of children with severe malnutrition in an urban health clinic in Dhaka, Bangladesh. *J Trop Pediatr* 2007;53:171-8.
- Ashraf H, Jahan SA, Alam NH, Mahmud R, Kamal SM, Salam MA, Gyr N. Day-care management of severe and very severe pneumonia, without associated comorbidities such as severe malnutrition, in an urban health clinic in Dhaka, Bangladesh. *Arch Dis Child* 2008;93:490-4.
- Ashraf H, Mahmud R, Alam NH, Jahan SA, Kamal SM, Haque F, Salam MA, Gyr N. Randomized-controlled trial of day-care vs hospital-care management of severe childhood pneumonia in Bangladesh. *Pediatr* (in press).
- Austrian R, Gold J. Pneumococcal bacteraemia with especial reference to bacteremic pneumococcal pneumonia. *Ann Intern Med* 1964;60:759-76.
- Bachur R, Perry H, Harper MB. Occult pneumonias: empiric chest radiographs in febrile children with leukocytosis. *Ann Emerg Med* 1999;33:166-73.
- Bada C, Carreazo NY, Chalco JP, Huicho L. Interobserver agreement in interpreting chest X-rays on children with acute lower respiratory tract infections and concurrent wheezing. *Sao Paulo Medical Journal* 2007;125:150-4.
- Baqui AH, Black RE, Arifeen SE, Hill K, Mitra SN, al Sabir A. Causes of childhood deaths in Bangladesh: results of a nationwide verbal autopsy study. *Bull World Health Organ* 1998;76:161-71.
- Berman S, McIntosh K. Selective primary health care: strategies for control of disease in the developing world. XXI. Acute respiratory infections. *Rev Infect Dis* 1985;7:674-91.

- Bose A, Coles CL, Gunavathi, John H, Moses P, Raghupathy P, Kirubakaran C, Black RE, Brooks WA, Santosham M. Efficacy of zinc in the treatment of severe pneumonia in hospitalized children <2 y old. *Am J Clin Nutr* 2006;83:1089-96.
- Bowton DL, Scuderi PE, Haponik EF. The incidence and effect on outcome of hypoxemia in hospitalized medical patients. *Am J Med* 1994;97:38-46.
- British Thoracic Society Standards of Care Committee. BTS Guidelines for the Management of Community Acquired Pneumonia in Childhood. *Thorax* 2002;57:1-24.
- Britton S, Bejstedt M, Vedin L. Chest physiotherapy in primary pneumonia. *BMJ (Clin Res Ed)* 1985;290:1703-5.
- Brooks WA, Yunus M, Santosham M, et al. Zinc for severe pneumonia in very young children: double-blind placebo-controlled trial. *Lancet* 2004;363:1683-88.
- Bryce J, Boschi-Pinto C, Shibuya K, Black RE. WHO estimates of the causes of death in children. *Lancet* 2005;365:1147-52.
- Bulla A, Hitze KL. Acute respiratory infections: a review. *Bull World Health Organ* 1978;56:481-98.
- Cashat-Cruz M, Morales-Aguirre JJ, Mendoza-Azpiri M. Respiratory tract infections in children in developing countries. *Semin Pediatr Infect Dis* 2005;16:84-92.
- Caulfield LE, de Onis M, Blossner M, et al. Undernutrition as an underlying cause of child deaths associated with diarrhea, pneumonia, malaria, and measles. *Am J Clin Nutr* 2004;80:193-8.
- Chaisupamongkollarp T, Preuthipan A, Vaicheeta S, Chantarojanasiri T, KongvivekkajornkijW, Suwanjutha S. Prone position in spontaneously breathing infants with pneumonia. *Acta Paediatr* 1999;88:1033-4.
- Chisti MJ, Tebruegge M, Vincente SL, Graham SM, Duke T. Pneumonia in severely malnourished children in developing countries - mortality risk, aetiology and validity of WHO clinical signs: a systematic review. *Trop Med Inter Health* 2009;14:1173-89.
- Chisti MJ, Salam MA, Sharifuzzaman, Pietroni MAC. Occult pneumonia: an unusual but perilous entity presenting with severe malnutrition and dehydrating diarrhoea. *J Health Popul Nutr* 2009;27:808-12.
- Chisti MJ, Ahmed T, Faruque AS, Salam MA. Clinical and laboratory features of radiologic pneumonia in severely malnourished infants attending an urban diarrhoea treatment center in Bangladesh. *Pediatr Infect Dis J* 2010;29:174-7.
- Comroe JH, Bothello S. The unreliability of cyanosis in the recognition of arterial anoxemia. *Am J Med Sci* 1947;214:1-9.
- Dagan R, Syrogiannopoula G, Ashkenazi S, et al. Parenteral-oral switch in the management of paediatric pneumonia. *Drugs* 1994;47:43-51.
- de Onis M, Monterio C, Clugston G. The worldwide magnitude of protein energy malnutrition: an overview from the WHO global database on child growth. *Bull World Health Organ* 1993;71:703-12.
- Dhawan A, Narang A, Singhi S. Hyponatraemia and the inappropriate ADH syndrome in pneumonia. *Ann Trop Paediatr* 1992;12:455-62.
- Dobson MB. Oxygen concentrators offer cost savings for developing countries: a study based in Papua New Guinea. *Anaesthesia* 1991;146:217-9.
- Dobson M, Peel D, Khallaf N. Field trial of oxygen concentrations in upper Egypt. *Lancet* 1996;347:1597-9.

- Doherty J. WHO guidelines on detecting pneumonia in children. *Lancet* 1991;338:1454.
- Donowitz GR, Mandell GL. Acute pneumonia. In: Mandell GL, Douglas RG, Bennet JE, editors. *Principles and practice of infectious diseases*. New York: Churchill Livingstone;1990:540-54.
- Duke T, Frank D, Mgone J. Hypoxaemia in children with severe pneumonia in Papua New Guinea. *Int J TB Lung Dis* 2000;5:511-9.
- Duke T, Tamburlini G. The Paediatric Quality Care Group. Improving the quality of paediatric care in peripheral hospitals in developing countries. *Arch Dis Child* 2003;88:563-5.
- Duke T, Wandt F, Jonathan M, et al. Impact of improved oxygen systems on child deaths from pneumonia: a multi-hospital effectiveness study in Papua New Guinea. *Lancet* 2008;372:1328-33.
- Falade AG, Tschappeler H, Greenwood BM, Mulholland EK. Use of simple clinical signs to predict pneumonia in young Gambian children: the influence of malnutrition. *Bulletin of the World Health Organization* 1995;73:299-304.
- Garenne M, Ronsmans C, Campbell H. The magnitude of mortality from acute respiratory infections in children under 5 years in developing countries. *World Health Stat Q* 1992;45:180-91.
- Graham SM, English M, Hazir T, Enarson P, Duke T. Challenges to improving case management of childhood pneumonia at health facilities in resource-limited settings. *Bull World Health Organ* 2008;86:349-55.
- Hall CB, Hall WJ, Speers DM. Clinical and physiological manifestations of bronchiolitis and pneumonia. Outcome of respiratory syncytial virus. *Am J Dis Child* 1979;133:798-802.
- Hall FM, Simon M. Occult pneumonia associated with dehydration: myth or reality. *Am J Roentgenol* 1987;148:853-4.
- Hamid M, Qazi SA, Khan MA. Clinical, nutritional and radiological features of pneumonia. *Journal of the Pakistan Medical Association* 1996;46:95-9.
- Huq F, Rahman M, Nahar N, Alam A, Haque M, Sack DA, Butler T, Haider R. Acute lower respiratory tract infection due to virus among hospitalized children in Dhaka, Bangladesh. *Rev Infect Dis* 1990;12 Suppl 8:S982-7.
- Jadavi T, Law B, Lebel MH, Kennedy WA, Gold R, Wang EEL. A practical guide for the diagnosis and treatment of pediatric pneumonia. *Can Med Assoc J* 1997;156:S703-S711.
- Jurban A, Tobin MJ. Reliability of pulse oximetry in titrating supplemental oxygen therapy in ventilator-dependent patients. *Chest* 1990;97:1420-5.
- Jurban A. Pulse oximetry. *Crit Care* 1999; 3:R11-R17.
- Klein JO. Bacterial pneumonias. In: *Textbook of pediatric infectious diseases*. 3<sup>rd</sup> ed. Philadelphia: WB Saunders;1992:299-314.
- Korppi M, Remes S, Heiskanen-Kosma T. Serum, procalcitonin concentrations in bacterial pneumonia in children: a negative result in primary healthcare settings. *Pediatr pulmonol* 2003;35:56-61.
- Levine A. Chest physiotherapy for children with pneumonia. *J Am Osteopath Assoc* 1978;78:122-5.
- Loeb M, High K. The effect of malnutrition on risk and outcome of community-acquired pneumonia. *Respiratory Care Clinics of North America* 2005;11:99-108.

- Lozano JM, Steinhoff M, Ruiz JG, Mesa ML, Martinez N, Dussan B. Clinical predictors of acute radiological pneumonia and hypoxaemia at high altitude. *Arch Dis Child* 1994;71:323-27.
- Mahalanabis D, Chowdhury A, Jana S, et al. Zinc supplementation as adjunct therapy in children with measles accompanied by pneumonia: a double-blind randomized controlled trial. *A J Clin Nutr* 2002;76:604-7.
- Marks MI, Klein JO. Bacterial infections of the respiratory tract. In: Feigin RD, Cherry JD, editors. *Infectious diseases of the fetus and newborn infant*. 4<sup>th</sup> ed. Philadelphia: WB Saunders;1995:891-908.
- Matai S, Peel D, Jonathan M, Wandu F, Subhi R, Duke T. Implementing an oxygen programme in hospitals in Papua New Guinea. *Ann Trop Paediatr* 2008;28:71-8.
- Mehta PN. Choosing antibiotics for community acquired pneumonia. *Indian Pediatr* 2003;40:958-64.
- Mulholland K. Magnitude of the problem of childhood pneumonia. *Lancet* 1999;354:590-2.
- Murphy CG, van de Pol AC, Harper MB, Bachur RG. Clinical predictors of occult pneumonia in the febrile child. *Acad Emer Med* 2007;14:243-9.
- Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global Burden of Disease Study. *Lancet* 1997;349:1269-76.
- Nohynek H, Valkeila E, Leinonen M, et al. Erythrocyte sedimentation rate, white blood cell count and serum C-reactive protein in assessing etiologic diagnosis of acute lower respiratory infections in children. *Pediatr Infect Dis J* 1995;14:484-90.
- Nannan N, Norman R, Hendricks M, Dhansay MA, Bradshaw D. Estimating the burden of disease attributable to childhood and maternal undernutrition in South Africa in 2000. *South African Medical Journal* 2007;97:733-9.
- Onyango FE, Steinhoff MC, Wafula EM, Wariua S, Musia J, Kitonyi J. Hypoxaemia in young Kenyan children with acute lower respiratory infection. *BMJ* 1993;306:612-5.
- Pauls S, Kruger S, Richter K, et al. Interobserver agreement in the assessment of pulmonary infiltrates on chest radiography in community-acquired pneumonia. *Rofortschritte auf dem Gebiete der Rontgenstrahlen und der Nuklearmedizin* 2007;79:1152-8.
- Rahman M, Huq F, Sack DA, Butler T, Azad AK, Alam A, Nahar N, Islam M. Acute lower respiratory tract infections in hospitalized patients with diarrhea in Dhaka, Bangladesh. *Rev Infect Dis* 1990;12 Suppl 8:S899-906.
- Rice AL, Sacco L, Hyder A, Black RE. Malnutrition as an underlying cause of childhood deaths associated with infectious diseases in developing countries. *Bull World Health Organ* 2000;78:1207-21.
- Riley I, Carrad E, Gratten H, et al. The status of research in acute respiratory infections in children in Papua New Guinea. *Pediatr Res* 1983;17:1041-3.
- Rudan I, Tomaskovic L, Boschi-Pinto C, Campbell H, WHO Child Health Epidemiology Reference Group. Global estimate of the incidence of clinical pneumonia among children under five years of age. *Bull World Health Organ* 2004;82:895-903.
- Rudan R, Boschi-Pinto C, Biloglav Z, Mulholland K, Campbell H. Epidemiology and etiology of childhood pneumonia. *Bull World Health Organ* 2008;86:408-16.
- Sarria E, Fisher GB, Lima JA, Menna Barreto SS, Flores JA, Sukiennik R. Interobserver agreement in the radiological diagnosis of lower respiratory tract infections in children. *J Pediatr (Rio J)* 2003;79:497-503.

- Sazawal S, Black RE. Pneumonia Case Management Trials Group. Effect of pneumonia case management on mortality in neonates, infants, and preschool children: a meta-analysis of community-based trials. *Lancet Infect Dis* 2003;3:547-56.
- Selwyn BJ. The epidemiology of acute respiratory infection in young children: comparison of findings from several developing countries. *Rev Infect Dis* 1990;12:Suppl 8:S870-S888.
- Shakur MS, Malek MA, Bano N, Islam K. Zinc status in well nourished Bangladeshi children suffering from acute lower respiratory infection. *Indian J Pediatr* 2004;41:478-81.
- Shalit I, Dagan R, Engelhard D, et al. Cefuroxime efficacy in pneumonia: sequential short course i.v./oral suspension therapy. *Isr J Med Sci* 1994;30:684-9.
- Shann F, Barker J, Poore P. Clinical signs that predict death in children with severe pneumonia. *Pediatr Infect Dis J* 1989;8:852-5.
- Shimeles D, Lulseged S. Clinical profile and pattern of infection in Ethiopian children with severe protein-energy malnutrition. *East African Medical Journal* 1994;71:749-67.
- Singhi S, Dhawan A. Frequency and significance of electrolyte abnormalities in pneumonia. *Indian Pediatr* 1992;29:735-40.
- Stapleton T. Chest physiotherapy in primary pneumonia. *BMJ* 1985;291:143.
- Steinbock M, Black R. Childhood pneumonia: we must move forward. *Lancet* 2007;369:1409-10.
- Subhi R, Adamson M, Campbell H, Weber M, Smith K, Duke T, et al. The prevalence of hypoxaemia among ill children in developing countries: a systematic review. *Lancet Infect Dis* 2009;9:219-27.
- Suskind D, Murthy KK, Suskind RM. The malnourished child: an overview. *In: Suskind RM, Lewinter-Suskind L, editors. The malnourished child. New York, NY: Vevey/Raven Press, 1990:1-22.*
- Swingler GH, Hussey GD, Zwarenstein M. Randomised controlled trial of clinical outcome after chest radiograph in ambulatory acute lower respiratory infection in children. *Lancet* 1998;351:404-8.
- Swingler GH. Radiologic differentiation between bacterial and viral lower respiratory infection in children: a systematic literature review. *Clin Pediatr* 2000;39:627-33.
- Swingler GH. Observer variation in chest radiography of acute lower respiratory infections in children: a systematic review. *BMC Medical Imaging* 2001;1:1.
- Toikka P, Irjala K, Juven T, et al. Serum procalcitonin, C-reactive protein and interleukin-6 for distinguishing bacterial and viral pneumonia in children. *Pediatr Infect Dis J* 2000;19:598-602.
- UN General Assembly. United Nations Millennium Declaration. New York: United Nations, 2000. <http://www.un.org/millennium/declaration/ares552e.pdf> (accessed March 2, 2009).
- UNICEF & World Health Organization. Pneumonia: the forgotten killer of children. New York: UNICEF/WHO, 2006
- Wafula EM, Ngamau DW, Onyango FE, Mirza NM, Njeru EK. X-ray diagnosable pneumonia in children with severe malnutrition at Kenyatta National Hospital. *East African Medical Journal* 1998;75:567-71.
- Walley J, Lawn J, Tinker A, et al. Primary health care: making Alma-Ata a reality. *Lancet* 2008;372:1001-7.

- Wallis C, Prasad A. Who needs chest physiotherapy? Moving from anecdote to evidence. *Arch Dis Child* 1999;80:393-7.
- Wang EFL, Law BJ, Stephens D, PICNIC. Pediatric Investigators Collaborative Network on Infections in Canada (PICNIC) study of morbidity and mortality risk factors with RSV disease. *J Pediatr* 1995;126:212-9.
- Wardlaw TM, Johansson EW, Hodge M, WHO, UNICEF. Pneumonia: the forgotten killer of children. Geneva: World Health Organization, 2006.
- Weber MW, Usen S, Palmer A, Jaffar S, Mulholland EK. Predictors of hypoxaemia in hospital admissions with acute lower respiratory tract infection in a developing country. *Arch Dis Child* 1997;76:310-4.
- Weber MW, Mulholland EK. Pulse oximetry in developing countries. *Lancet* 1998;351:1589.
- Williams BG, Gouws E, Boschi-Pinto C, Bryce J, Dye C. Estimates of world-wide distribution of child deaths from acute respiratory infections. *Lancet Infect Dis* 2002;2:25-32.
- World Health Organization. A programme for controlling acute respiratory infections in children: memorandum from a WHO meeting. *Bull World Health Organ* 1984; 62:47-58.
- World Health Organization. Acute respiratory infections in children: case management in small hospitals in developing countries. A manual for doctors and other senior health workers. WHO/ARI/ 90.5. Geneva: WHO, 1990.
- World Health Organization. Management of the young child with an acute respiratory infection. Programme for control of acute respiratory infections. Geneva: WHO, 1991.
- World Health Organization. Technical basis for the WHO recommendations on the management of pneumonia in children at first-level health facilities (WHO/ARI/91.20). Geneva: WHO, 1991.
- World Health Organization. Life in the 21st century. A vision for all. Geneva: WHO, 1998:66.
- World Health Organization. Management of the child with a serious infection or severe malnutrition Guidelines for care at the first-referral level in developing countries (WHO/FCH/CAH/00.1). Geneva: WHO, 2000.
- World Health Organization. WHO Pocket book of hospital care for children: Guidelines for the management of common illness with limited resources. Geneva: WHO, 2005.
- World Health Organization. The clinical use of oxygen in hospitals with limited resources. Guidelines for health-care workers, hospital engineers and managers. Editor: Duke T; Geneva: WHO, 2009.
- Zar HJ, Jeena P, Argent A, Gie R, Madhi SA. Diagnosis and management of community-acquired pneumonia in childhood- South African Thoracic Society Guidelines. *S Afr Med J* 2005;95:977-90.



## **Health Management**

Edited by Krzysztof Smigorski

ISBN 978-953-307-120-6

Hard cover, 112 pages

**Publisher** Sciyo

**Published online** 27, September, 2010

**Published in print edition** September, 2010

Possibilities of medical intervention have thrived over the last decades. Our knowledge about mechanisms of the development of diseases and factors influencing it has increased. Effective treatment requires a holistic approach that takes into consideration aspects at first sight not related to a course of a specific disorder. This book contains a few chapters focusing on issues related to health management. The chapters are arranged in an order reflecting multidimensionality of issues constituting this theoretical and practical area - starting from the studies focusing on a general, administrative level, to considerations related to situations of individuals suffering from a specific illness. The discussed problems concern different age groups - children, adults and the elderly. We hope that readers professionally engaged in healthcare - both theoretically and clinically - will find it interesting, useful and inspiring.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Hasan Ashraf, Mohammud Jobayer Chisti and Nur Haque Alam (2010). Treatment of Childhood Pneumonia in Developing Countries, Health Management, Krzysztof Smigorski (Ed.), ISBN: 978-953-307-120-6, InTech, Available from: <http://www.intechopen.com/books/health-management/treatment-of-childhood-pneumonia-in-developing-countries>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821



© 2010 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen