

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Pesticide Exposure and Health Related Issues in Male and Female Reproductive System

Stavros Sifakis¹, Mihalis Mparmpas¹, Offie P. Soldin² and Aristides Tsatsakis³

¹*Department of Obstetrics & Gynecology University of Crete, Heraklion, Greece*

²*Departments of Medicine, Oncology, Physiology and Obstetrics & Gynecology,
PregnaTox, Georgetown University Medical Center, Washington D.C.,*

³*Laboratory of Toxicology, Department of Medicine, University of Crete, Heraklion,*

^{1,3}*Greece*

²*USA*

1. Introduction

During the last several decades there have been widespread uses of potent substances that, although effective in their intended use, have also been suspected of being harmful to reproductive health. This mixture of environmental contaminants that may adversely affect human fertility includes heavy metals (lead, mercury, arsenic), phthalates (plasticizers), bisphenol-A (building block of several plastics), polychlorinated biphenyls (lubricants), dioxins (byproducts of manufacture), pesticides and other agents. The impact of adverse effects on reproductive health include impaired gametogenesis, sperm maturation, decreased semen quality, miscarriages, ovulation and menstrual disturbances, infertility, stillbirths, developmental anomalies, cryptorchidism, hypospadias and cancer. The effects can be reversible, permanent or even transgenerational, take place in the offspring, as exposure can occur during pregnancy and intrauterine life, childhood or later. The route of exposure, dose, age, gender and genotype (susceptibility of the individual) are important factors that can determine the reproductive disorder. In this chapter we focus on the effects of exposure to pesticides during adulthood, on human male and female fertility, giving emphasis to semen quality and time to pregnancy.

2. Pesticides

Pesticides are mainly utilized in agriculture for crop protection, often replacing the natural processes on which agricultural production had previously depended. Pesticides are also applied in homes and gardens. More than 140.000 tones of pesticides are used annually in the European Union for agricultural purposes only (Ramazzini, 2009).

A pesticide is “any substance or mixture of substances intended for preventing, destroying or controlling any pest, including vectors of human or animal disease, unwanted species of plants or animals causing harm during or otherwise interfering with the production processing, storage, transport or marketing of food, agricultural commodities, wood and wood products or animal feedstuffs, or substances which may be administered to animals for the control of insects, arachnids or other pests in or on their bodies” (FAO, 2003). They

fall into three major classes: insecticides, fungicides, and herbicides, classification based upon the target organism. There are also rodenticides (for control of vertebrate pests), nematicides (to kill eelworms, etc), molluscicides (to kill slugs and snails), and acaricides (to kill mites) (Cox & Sorgan, 2006). Pesticides may differ according to their chemical structure, their mechanism of action and the toxicity they exhibit, but typically each pesticide consists of one (or more) active ingredient, which exerts the pesticidal activity, and an inert ingredient, which is inactive and helps in handling the active ingredient. Several studies have shown that the inert ingredient is not as inactive as it was previously believed to be (Sorgan, 2005; Cox & Sorgan, 2006). Over 700 active ingredients are in use worldwide as pesticides, each with distinct chemical and toxicological properties (Toppari et al., 1996). Another classification categorizes pesticides according their chemical structure (Table 1). Insecticides include organochlorines, organophosphates, and carbamates. Organochlorine hydrocarbons (DDT, heptachlor) operate by disrupting the sodium/potassium balance of the nerve fiber. They are persistent in human tissue with the potential to bioaccumulate. Organophosphate (parathion, malathion) and carbamates (carbaryl, carbofuran) are less toxic and largely replaced organochlorines. They are inhibitors of acetylcholinesterase, causing paralysis. Common herbicides include phenoxy and benzoic acid herbicides (2,4-D) and triazines (atrazine). Fungicides (vinclozolin, mancozeb) have sulfur as the most common active ingredient. Nicotinoids and pyrethroids are plant-derived pesticides (fenvalerate, pyrethrin). In addition to the desired effects of crop protection and pest management, pesticides have some recognized adverse impacts on human health and the environment. Humans have a great risk of exposure through several pathways in occupational, agricultural and household use. Inhalation, oral, dermal and ocular, are four possible routes for pesticide exposure. Ingestion of food and water is thought to be the main routes of pesticide exposure in the general population, while dermal absorption is suspected to be the main source of occupational exposure (Toppari et al., 1996). Over 25% of fruits, vegetables, and cereals are known to contain detectable residues of at least two pesticides and more than 300 different pesticides are known to contaminate food products sold in the EU (Ramazzini, 2009). In the majority of cases, however, human exposure is unintentional and unintended (Ribas-Fitó, 2002).

Pesticides are accused of causing short-term adverse health effects. Acute health effects include stinging eyes, rashes, blisters, blindness, nausea, dizziness, diarrhea and death (Jeyaratnam, 1990; Sanborn et al., 2007). They are suspected also for a wide range of chronic effects, which can occur months or years after the exposure, such as cancers, neurological and developmental toxicity, immunotoxicity, genotoxicity, respiratory effects and disruption of the endocrine system. Pesticides may affect not only the exposed individual but also subsequent generations (Alavanja et al., 2004; Ritter et al., 2006; McCauley et al., 2006; Bassil et al., 2007).

There has been rising concern in many developed countries about the adverse effects of pesticides on human reproduction, ranging from female and male subfertility to abortion, stillbirths, birth defects and malformations (García, 2003; Weselak et al., 2007; Peiris-John & Wickremasinghe, 2008). They may cause reproductive toxicity with direct damage to the structure of the cells or as a result of biotransformation into metabolites, or interference with processes necessary for the natural homeostasis and equilibrium. They may act like hormones in the endocrine system and disrupt the function of the natural endogenous hormones, when doing so they are often called endocrine disrupting chemicals (EDC) (Lathers, 2002; Diamanti-Kandarakis et al., 2009). This group of compounds identified as EDC is heterogeneous and includes synthetic or natural chemicals.

organochlorides	DDD, DDT, DDE, chlordane, kepone, dieldrin, endosulfan, heptachlor, lindane, mirex, methoxychlor, toxaphene
organophosphorus	chlorpyrifos, glyphosate, diazinon, dimethoate, malathion, methamidophos, parathion, terbufos, tribufos, trichlorfon
carbamates	aldicarb, carbaryl, carbofuran, fenoxycarb, propoxur
pyrethroids	cypermethrin, fenvalerate, permethrin, pyrethrin, pyrethrum, resmethrin, tetramethrin
anilides/anilines	metolachlor, pretilachlor, propachlor, trifluralin
phenoxy	2,4-D, 2,4-DB, 2,4,5-T, MCPA, MCPB, fenoprop
triazines	atrazine, cyanazine, hexazinone, prometryn, propazine, simazine, terbutryn
quaternary	diquat, MPP, paraquat
ureas	chlortoluron, DCMU, metsulfuron-methyl, monolinuron
others	acetamiprid, amitraz, chlordimeform, cyromazine, diflubenzuron, nithiazine, sulfuramid, thiachlopid, xanthone

Table 1. Classification of pesticides based on their chemical structure.

3. Pesticides as Endocrine Disrupting Chemicals (EDC)

Pesticides may act as endocrine disruptors and alter the hormonal homeostasis in both males and females and lead to subfertility. The term “endocrine disruptors” was introduced into literature with an article published in 1993 (Colborn, 1993). An endocrine disruptor was defined by the U.S. Environmental Protection Agency (EPA) as “an exogenous agent that interferes with the synthesis, secretion, transport, binding, action, or elimination of natural hormones in the body that are responsible for the maintenance of homeostasis, reproduction and developmental process” (Kavlock et al., 1996). The group of EDC includes pesticides and various synthetic substances such as polychlorinated biphenils (PCB), polybrominated biphenyls (PBB), bisphenol-A (BPA), phthalates and dioxins natural compounds such as phytoestrogens (Fourth National Report on Human Exposure to Environmental Chemicals, 2009) are used as solvents, lubricants, plasticizers, cosmetics etc. They are usually small molecules (mass 1000 Daltons) and often have a phenolic moiety that probably mimics natural steroid hormones (Diamanti-Kandarakis et al., 2009). They contain chlorine or other halogens (bromine, iodine or fluorine) with strong interaction and so they resist degradation. They usually have a long half-life and accumulate in the environment, sometimes remotely from the place they were produced. Substances banned even decades ago can still be found in the environment and in living organisms. In humans and animals EDCs are stored in fatty tissue or they may be metabolized into more toxic compounds. The predominant sources of exposure are food, water and air. The routes of exposure include ingestion, inhalation and dermal absorption.

Natural hormones act in very low concentrations, similarly, can elicit adverse effects in low doses. Occasionally, the EDCs do not follow the classic dose-response effect and low doses may result in stronger effects than high doses (vom Saal et al., 2009). Humans are exposed concomitantly to a large number of compounds. These substances may enhance and have a synergistic or antagonistic effect, and as cited above, some substances can be metabolized in more toxic products (Crews et al., 2003). One of the reasons of the difficulty studying the damage after exposure to a single agent is this mixture of compounds which is accumulated in human organisms.

An issue of critical importance is the timing of exposure, since damage is age sensitive. The same dose can have different effects in fetuses, newborn, infants or adults. Given the same doses, a developing organism (embryo, neonate) whose growth is highly controlled by the endocrine system is more vulnerable to EDCs, than an adult (Guo et al., 1995; Bigsby et al., 1999; Lilienthal et al., 2006). The damage incurred by the exposure may not be immediate and may only be manifested in adulthood or during aging. The consequences may be apparent even in subsequent generations and the classic example is the cases of vaginal carcinoma in daughters of mothers who were exposed to DES during their pregnancies (Herbst et al., 1999; Anway & Skinner, 2006; Rubin, 2007). The susceptibility of an individual may vary due to genetic polymorphism and so the results of the same exposure could be different.

Initially it was thought that EDCs act via nuclear hormone receptors (e.g. estrogen, progesterone, androgen, thyroid receptors), but now it is believed that they act also via membrane, non-steroid receptors (e.g. dopamine, serotonin, nor-epinephrine receptors). Catecholamine hormones following synthesis are stored in granular vesicles intracellularly. Steroid hormones are not stored but readily synthesized following gonadotropin stimulation of the gonads and are usually found in the circulation bound by carrier proteins (only free hormones are biologically active). Once reaching their tissue targets steroid hormones exert their action by binding to different kinds of nuclear receptors. Explicit hormones bind specific receptors and individualized mechanisms follow by intracellular signalling (e.g. protein kinase-C activation or phosphatidylinositol turnover). Hormones are mostly catabolised in the liver. Consequently, EDCs can participate in most aforementioned pathways thereby changing hormone synthesis patterns, mimicking hormone function or blocking it by occupying the receptor site, modulating the number of the receptors and their affinities for specific molecules and altering hormone clearance (Gore et al., 2006; Gore, 2007; Gore, 2008). Sex hormones synthesis is regulated by the hypothalamic-pituitary-gonadal axis. LH and FSH are synthesized by the anterior pituitary under the influence of pulsatile secretion of GnRH, released by the hypothalamus.

Several pesticides have been reported to act as estrogen agonists, e.g. methoxychlor, endosulfan, toxaphene, kepone, DDT, fenarimol, alachlor, pentachlorophenol, fenvalerate, chlordecone (Soto et al., 1995; Cummings & Gray, 1997; Garey & Wolff, 1998; Andersen et al., 2002; Kojima et al., 2004). On the other hand, other pesticides such as vinclozolin, p,p-DDE and o,p-DDT may have anti-androgenic activity, or both estrogenic and antiandrogenic activity (Kelce et al., 1994; Kelce et al., 1995; Kelce & Wilson, 1999). The fungicide methyl-2-benzimidazole carbamate decreases estradiol production in primary cultures of human ovarian granulosa (Can & Albertini, 1997). Treatment of rats with heptachlor suppresses progesterone and estradiol concentrations in blood (Oduma et al., 2006). Progesterone concentrations also decrease during early pregnancy in the rabbit following exposure to the pesticide DDT (Lindenau et al., 1994). DDT was found to be

estrogenic, but its major metabolite DDE, has considerable antiandrogenic activity (Kelce et al., 1995). Further, atrazine seems to have estrogenic and antiandrogenic properties and was suggested to reduce testicular testosterone in male rats exposed to it (Stoker et al., 2000). Lindane intercalates into the sperm membrane and may inhibit sperm responsiveness to progesterone in vitro (Silvestroni & Palleschi, 1999). Lindane also inhibits steroidogenesis by reducing StAR (steroidogenic acute regulatory) protein-mediated cholesterol transfer (Walsh & Stocco, 2000). Neonatal exposure to either DES or flutamide also inhibited steroidogenesis and StAR protein expression in the fetal rat Leydig cell (Mikkilä et al., 2006). This protein mediates cholesterol passage through mitochondrial membranes and impaired expression results in decreased testosterone production in vitro (Manna et al., 2001). Testosterone concentration is also reduced withazole fungicides (ketoconazole) due to impaired enzymatic activity of 17 α -hydroxylase and 17,20-lyase (Wang et al., 1992; Bahshwan et al., 1998; Taxvig et al., 2008). In rats, fenarimol, a different fungicide, was found to cause a dose-related decrease in fertility (Hirsch et al., 1986). In vitro studies of some pesticides such as fenarimol, prochloraz, imazalil and dicofol, indicate these pesticides inhibit the conversion of androgens to estrogens through CYP 19 aromatase inhibition (Vinggaard et al., 2000). Rats treated with mancozeb demonstrate a decrease in the number of healthy follicles and an increase in the number of atretic follicles (Mahadevaswami et al., 2000). EDCs have also been associated with breast cancer, PCO and endometriosis in women, cryptorchidism, hypospadias, testicular and prostate cancer in men, alteration in pituitary and thyroid gland functions (Crisp et al., 1998; Bretveld et al., 2006; Diamanti-Kandarakis et al., 2009).

4. Pesticides and semen quality

Approximately 6% of adult males are thought to be infertile (Purvis & Christiansen, 1992). Male factors are responsible for at least 20% of cases of infertility. Male infertility is related to impaired semen quality and may be due to a variety of causes including genetic (Klinefelter's syndrome), congenital (cryptorchidism), endocrine (hypogonadism), obstructive (vasectomy), infective (chlamydia), vascular (varicocele), neoplastic (carcinoma of the testis), lifestyle, and environmental (heat, drugs, pesticides, irradiation) factors. Others causes include sexual dysfunction related to erection and ejaculation (Purvis & Christiansen, 1992; Dohle et al., 2005). However, in many cases male infertility is regarded as idiopathic (40-75%), and no cause can be identified. Semen analysis is used to evaluate semen quality, which is taken as a surrogate measure of male infertility. The World Health Organization has provided reference values for human semen characteristics (Cooper et al., 2010).

There is evidence of regional variation in semen quality that may be an expression of gene polymorphisms, different climate, lifestyle and exposure to magnetic fields or chemical substances such as pesticides (Mallidis et al., 1991; Jørgensen et al., 2001; Swan et al., 2003; Li et al., 2010). Seasonal variation has also been detected (decrease of sperm density and total sperm count during summer) (Levine, 1999; Krause & Krause, 2002). These factors, together with the variability in techniques and methodologies used, are reasons for some controversial results in studies that analyze semen quality changes overtime. Thus, the issue of a decline in semen quality overtime is equivocal. Following the publication of the meta-analysis conducted by Carlsen et al. in 1992, demonstrating a decline in human semen quality over the last 50 years (mean sperm count from 113 millions/ml in 1940 to 66

millions/ml in 1990), numerous related studies have been published. Studies by Swan et al. had consistent results with those of Carlsen et al. (1992) and supported that historical data on sperm density, despite large random error, are reliable (Swan et al., 1997; Swan & Elkin, 1999; Swan et al., 2000). However, Olsen et al. (1995) reanalyzed the data used in a linear model to predict sperm quality deterioration in the last 50 years, advocate that the data are only robust during the last 20 years (1975-1995), in which other statistical models (quadratic, spline fit and stairstep), except the linear model, suggest constant or slightly increasing sperm counts. Studies from Italy, Denmark, Canada, Tunisia, India, Poland, Israel, Scotland, Greece and Germany suggest that there has been a decline, or sperm parameters are impaired in young populations (Adamopoulos et al., 1996; Younglai et al., 1998; Bilotta et al., 1999; Almagor et al., 2003; Vicari et al., 2003; Jørgensen et al., 2006; Sripada et al., 2007; Adiga et al., 2008; Paasch et al., 2008; Horak et al., 2008; Feki et al., 2009). Conversely, other reports from US, Japan, Korea, Sweden, Spain, Israel and Czech Republic showed no significant evidence of deterioration in sperm quality (Fisch et al., 1996; Paulsen et al., 1996; Benschushan et al., 1997; Berling & Wölner-Hanssen, 1997; Andolz et al., 1999; Seo et al., 2000; Itoh et al., 2001; Zvěřina et al., 2002). The materials and methods of the studies mentioned varied widely, as well as time period (one, two, or more decades), population sample (e.g. individuals in infertile relationship or fertile subjects who participated voluntarily) and the level of pollution in the various geographical regions. Merzenich et al. (2010) conclude that former meta-analyses of sperm count data show a global downward trend, but this conclusion should be interpreted with caution, because the included studies are of great heterogeneity. The geographic variation in semen quality may reflect different exposures to endocrine disruptors, such as pesticides (Swan, 2006). Pesticides might have the ability to interrupt male fertility at several different sites in the reproductive pathway and by one or more mechanisms, as cited previously. Thus, they can interfere with the hypothalamopituitary axis that regulates, through the production of the gonadotrophins FSH and LH, the function of Sertoli and Leydig cells, impairing spermatogenesis and steroidogenesis.

Tables 2a,b,c lists the studies published evaluating the association between exposure to pesticides and human sperm quality. Literature reviews and articles investigating chemical compounds, without including pesticides, were excluded. Studies evaluating pregnancy outcomes and no sperm quality were excluded too. Sperm quality was assessed evaluating conventional parameters (concentration, motility, morphology) or sperm DNA/chromatin integrity and aneuploidy. Sixty-three reports, satisfying these criteria were identified and included in Tables 2a,b,c. Among them six (6/63) studies evaluated the recovery of sperm quality, years after cessation of exposure to DBCP (5) and kepone (1). Thirty-six (36/63) studies examined exposure to single, specific pesticides or metabolites and included DBCP, DDT, DDE, EPB, 2,4-D, kepone (chlordecone), molinate, carbaryl, fenvalerate, ethylparathion, methamidophos, 1N (metabolite of carbaryl and naphthalene), TCPY (metabolite of chlorpyrifos and chlorpyrifos-methyl) and 3PBA, CDCCA, TDCCA (pyrethroid metabolites). The majority of the pesticides cited above are now banned or severely restricted, at least in USA or EU. Twenty-one (21/63) studies evaluated mixture of compounds such as fungicides, insecticides or herbicides with or without specifying the exact pesticides. Some of the pesticides involved in these studies are: alachlor, diazinon, acetochlor, malathion, atrazine, metolachlor, DEET (insect repellent), 2,4-D, aldicarb and cadusaphos, ethoprothos, isazophos, terbufos, pyrimiphos-ethyl (organophosphorus pesticides).

4.1 Studies with little or no evidence of an association

Eighteen studies of the 63 (18/63) found no or little evidence of association between pesticide exposure and sperm quality (Table 2a). Eight studies (8/18) involved a mixture of compounds and ten (10/18) single pesticides. Carbaryl and molinate were reported in one study each. Ten studies (10/18) were related to DDT and metabolites most of the times p,p'-DDE, the known persistent pesticide banned many years ago (not all over the world). Two of the ten studies evaluated DDT in mixtures and eight as a single pesticide. Five (5/10) of the reports that found no or little association between DDT, DDE and semen quality were carried out by the INUENDO project (including the two substudies of INUENDO by Rignell-Hydbom et al. (2004 & 2005). INUENDO (INUit-ENDOcrine) is the acronym for "Biopersistent organochlorines in diet and human fertility. Epidemiological studies in time to pregnancy and semen quality in Inuit and European populations". This EU project (2002-2005) used serum levels of CB-153 (polychlorinated biphenyl) and p,p'-DDE, the main DDT metabolite, to estimate the impact in human fertility in epidemiological studies including Inuits from Greenland and Caucasians from Poland, Sweden and Ukraine. Hauser et al. (2003) conducted two cross sectional studies and found a limited evidence of an inverse association between p,p-DDE and sperm motility as well as no strong relationships between the levels of this compound and sperm DNA damage. Furthermore, Charlier et al. (2005) estimated serum and seminal plasma concentrations of p,p-DDE in fertile and sub or infertile young men. Blood concentrations of p,p'-DDE were very low in both groups. No p,p'-DDE detected in seminal plasma of either groups. Of note, the mothers of the exposed subfertile men had serum level of p,p'-DDE significantly higher than the mothers of the control group.

Two studies did not evaluate DDT exposure separately, but it was included in a sum of other compounds. Magnusdottir et al. (2005) concluded that poor semen quality is associated with sedentary work and obesity but not with plasma levels of fourteen organochlorine pesticides including DDT and metabolites. Weiss et al. (2006) evaluating exposure, in Germany and in Tanzania, to a mixture of PCBs and pesticides, including DDT, found these pesticides had no impact on sperm quality. However high serum concentrations of DDT-DDE were associated with lower pregnancy rates in Germany.

Two out of the eighteen reports that showed little or no association between pesticide exposure and semen quality were performed by ASCLEPIOS (Larsen et al., 1998a; Härkönen et al., 1999). It was an EU project (1993-1998), that was carried out in 14 European centers and focused on occupational exposure to the fungicides styrene and inorganic lead. Questionnaire studies of time to pregnancy were combined with longitudinal and cross sectional studies of semen quality.

Tielemans et al. (1999a) conducted a case-control study and found no associations between exposure to pesticides and poor semen quality, but few subjects were exposed to pesticides (the rest of the sample was exposed to chemical pollutants different from pesticides, such as solvents and metals). Juhler et al. (1999) found similar results indicating minor association, comparing traditional and organic farmers, but for all the groups of exposure the average dietary intake of pesticides was low. Smith et al. (2004) found no significant differences in sperm aneuploidy or diploidy frequencies between men exposed to a mixture of pesticides and control groups, but the sample was rather small (n=20+20). In a cross-sectional study conducted by Multigner et al. (2008) semen characteristics were evaluated in association to exposure to a mixture of organophosphorus pesticides (cadusaphos, ethoprothos, isazophos, terbufos, pyrimiphos-ethyl and one carbamate the aldicarb). No significant difference was

found between exposed and unexposed workers (in banana plantations), but exposure was assessed by a questionnaire and not by chemical analysis. Tomenson et al. (1999) conducted a longitudinal study and found no evidence that sperm and serum hormones levels were related to molinate exposure (thiocarbamate herbicide). Whorton et al. (1979) found no apparent effects on sperm count in workers exposed to carbaryl, but three subsequent relative studies indicated contrary results.

Authors & Year Country	Pesticide	N	Study Type
Whorton et al, 1979 (USA)	carbaryl	47 workers	CS
Larsen et al, 1998a (ASCLEPIOS) (Denmark)	Mixture	248 farmers users or not users of pesticides during a spraying season	L
Härkönen et al, 1999 (ASCLEPIOS) (Finland)	fungicides	30 healthy farmers before and after exposure	CS
Tomenson et al, 1999 (UK)	molinate (thiocarbamate herbicide)	272 workers at three US plants.	L
Tielemans et al, 1999a (The Netherlands)	Occupational exposures (solvents, metals, pesticides)	Male partners of couples having their first consultation in two infertility clinics (n=899)	CC
Juhler et al, 1999 (Denmark)	mixture	171 traditional and 85 organic farmers.	CC
Hauser et al, 2003a (USA)	DDT, DDE, PCBs	212 male partners of subfertile couples	CS
Hauser et al, 2003b (USA)	DDT, DDE, PCBs	212 male partners of subfertile couples	CS
Rignell-Hydbom et al, 2004, (Sweden)	CB-153 p,p'-DDE	195 Swedish fishermen aged 24-65	CS
Smith 2004 et al, (Canada)	mixture	20 exposed, 20 non exposed	CC
Charlier 2005 et al, (Belgium)	p,p'-DDE	73 fertile young men (controls) & 23 mothers; 82 sub or infertile young men (cases) & 19 mothers	CC
Magnusdottir 2005 et al, (Iceland)	PCBs, organochlorine pesticides, (p,p'DDE)	25 poor semen quality, 20 normal semen quality & idiopathic subfertility, 27 normal semen quality & female subfertility	CS
Rignell-Hydbom et al, 2005 (Sweden)	CB-153 p,p'-DDE	176 Swedish fishermen (low & high intake of fatty fish)	CS
Spano 2005 et al, INUENDO, (Italy)	CB-153 p,p'-DDE	193 Inuits - Greenland; 178 Swedish fishermen; 141 men from Poland; 195 men from Ukraine	CS

Stronati et al, 2006, INUENDO, (Italy)	CB-153 p,p'-DDE	200 Inuits -Greenland; 166 men from Sweden; 134 men from Poland; 152 men from Ukraine	CS
Toft et al, 2006, INUENDO, (Denmark)	CB-153 p,p'-DDE	194 men from Greenland; 185 men from Sweden; 189 men from Poland; 195 men from Ukraine	CS
Weiss et al, 2006 (Germany)	mixture of PCBs & pesticides including DDT,DDE	31 women, 16 men from Tanzania, 21 couples from Germany	CS
Multinger et al, 2007 (France)	organophosphorus pesticides and one carbamate	42 exposed & 45 non exposed in banana plantations in Guadeloupe (parallel assessment in wild rats)	CS

Table 2a. Studies evaluating the association between exposure to pesticides and human sperm quality. Studies with no or little evidence of an association; 2b: Studies with evidence of an association; 2c: Studies evaluating the recovery of sperm quality (CS: cross sectional study; CC: case-control study; L: longitudinal study; R: retrospective study; P: pilot study).

4.2 Studies with evidence of an association

Thirty-nine (39/63) studies that varied widely in materials, methods, exposure and assessment of exposure concluded that there was evidence of an association between exposure to pesticides and impaired semen quality. Twenty-six (26/39) studies evaluated exposure to single pesticides or metabolites and thirteen (13/39) studies evaluated exposure to mixtures of compounds including pesticides or mixtures of pesticides. Evidence of an association was found in studies involving:

- DDT and DDE: Only in six of the 16 studies overall involved, five as a single pesticide and one in a mixture.
- DBCP: All studies showed evidence of impairment. Thrupp in his article examined a remarkable case of massive sterilization of approximately 1500 workers exposed in banana plantations in Costa Rica (Thrupp, 1991). Slutsky et al. in his report which represents the largest cohort of DBCP exposed workers, found that after a median exposure of three years, 64.3% of these men overall, and 90.1% of men studied from the Philippines, had oligospermia or azospermia (1999). Whorton et al. (1977), Potashnik et al. (1978), Lipshultz et al. (1980) and Egnatz et al. (1980) had the same results of a significant association. There are five studies more involving DBCP, but related to sperm recovery after cessation of exposure and are cited in the related paragraph.
- EPB: One cross-sectional and one longitudinal study with small sample size.
- Kepone: A cross-sectional study measuring blood levels found oligospermia and decreased sperm motility. One more study involving kepone was related to the recovery of sperm after exposure and is cited in the related paragraph.
- 2,4-D: One cross sectional study evaluated exposure using urine samples and observed asthenospermia, teratospermia and necrospermia. The second study evaluated this compound in a mixture of eight pesticides and found relation with poor semen quality.
- Fenvalerate: All studies conducted in China; three of them reporting impairment of conventional parameters of sperm quality and one increased percentage of sperm aneuploidy and altered morphology.

- Carbaryl: The first, a cross-sectional study, showed increase in abnormal morphology, the second study found altered seminal volume and sperm motility and the third study showed increased frequencies of aneuploidy.
- Ethylparathion and Methamidophos: Semen and urine samples were collected in order to estimate exposure to these organophosphate pesticides. There was a decrease in sperm concentration and motility but no significant difference was found in sperm morphology.
- 1N and TCPY: The first compound is a metabolite of carbaryl and naphthalene and the second a metabolite of chlorpyrifos and chlorpyrifos-methyl. This study used biological markers of exposure (urine analyses for metabolites) and several modeling approaches to test the robustness of the data. Statistically significant inverse dose- response relationships between 1N and sperm concentration and motility were found. There was a suggestive association between TCPY and sperm concentration and motility. Sperm morphology was not significantly associated with both 1N and TPCY. Only a single urine sample collected to estimate exposure and only a single semen sample collected to assess semen quality.
- 3PBA: It is the main metabolite of pyrethroids such as cypermethrin, deltamethrin, permethrin with high detection rate in the general population. The first study with rather large sample size showed suggestive association between increased urinary 3-PBA concentration (creatinine adjusted) and sperm concentration. There was an association between straight line velocity and curvilinear velocity (sperm progression and motion parameters) with urinary 3-PBA concentration (creatinine adjusted), while sperm volume, sperm number per ejaculum and sperm motility were weakly or not significantly associated. A second study that evaluated two more pyrethroid metabolites, CDCCA and TDCCA, was similar in design and found evidence for reduced semen quality and increased DNA damage related to the urine pyrethroid metabolites. Swan et al. (2003b) evaluated exposure to a mixture of eight pesticides in two different populations within Missouri and Minnesota. The small sample size limited statistical power. Exposure was assessed by urine analysis. Study concluded that Alachlor, atrazine, 2,4-D, metolachlor and a diazinon metabolite were associated with poor semen quality in Missouri. However, no significant associations were found for acetochlor, DEET and malathion dicarboxylic acid. Within Minnesota, the levels of pesticides were low for any of the pesticides, no significant associations were found too, but because of the overall results this study is classified in this category.

Dalvie et al. (2004) evaluated exposure to DDT in workers in South Africa, in relation to sperm quality and sexual function. Exposure was assessed by serum levels of *o'p'* and *p'p'* isomers of DDT, DDE, DDD. Sperm count and density were in the normal range. 84% of morphology scores were below the WHO criteria and *p'p'*DDT was negatively associated with sperm count (after correction for age, abstinence, physical abnormality and fever). Although no strong evidence for a DDT overall effect in sexual function and reproductive outcomes was found semen quality was impaired resulting in this study being cited in this category.

Dallinga et al. (2002) studied a group of men with poor semen quality vs. group of men with normal, based on the progressive motility of sperm. Blood samples were collected in order to determine whether differences in sperm quality were related with differences in serum concentrations of organochlorines, including DDT. No significant differences in

organochlorine levels were found initially, but after adjustment for age and sperm count, sperm progressive motility were inversely related to the concentrations of metabolites in the group of men with normal semen quality. Because of this finding the study is cited in this category.

Authors & Year Country	Pesticide	N	Study Type
Whorton et al, 1977 (USA)	DBCP	25 workers	CS
Cannon et al, 1978 (USA)	kepone	133 workers	CS
Potashnik et al, 1978 (Israel)	DBCP	6 workers	CS
Lipshultz et al, 1980 (USA)	DBCP	228 workers (exposed & non exposed cohort)	CS
Egnatz et al, 1980 (USA)	DBCP	232 workers exposed, 97 workers non exposed	CS
Wyrobek et al, 1981 (USA)	carbaryl	50 men occupationally exposed, 34 unexposed newly-hired	CS
Henderson et al, 1986 (Australia)	Mixture of chemicals including pesticides	1695 men with abnormal semen quality or fertility impairment	CS
Ratcliffe et al, 1987 (USA)	EDB	46 exposed in papaya fumigation industry in Hawaii; 43 unexposed from a sugar refinery.	CS
Schrader et al, 1988 (USA)	EDB	10 exposed forestry employees, 6 unexposed (exposing time 6 wks)	L
Thrupp et al, 1991 (USA)	DBCP	exposed workers in a banana plantation in Costa Rica	L
Lerda et al, 1991 (Argentina)	2,4-D	32 farm sprayers	CS
Strohmer et al, 1993 (Austria)	mixture	101 couples seeking artificial insemination (poor semen quality), controls couples with female infertility IVF treated	CS
Slutsky 1999 et al, (USA)	DBCP	26400 workers in banana & pineapple plantation in 12 countries	R
Abell et al, 2000 (Denmark)	mixture	122 greenhouse workers	CS
Padungtod et al, 2000 (USA)	Ethylparathion Methamidophos	32 exposed workers & 43 not exposed workers in China factories	CS
Ayotte et al, 2001 (Mexico)	DDT P,P'-DDE	24 Mexican men living in endemic malaria areas not occupationally exposed	CS
Oliva et al, 2001 (Argentina)	mixture of pesticides & solvents	225 male partners from infertility clinics	CS

Dallinga et al, 2002 (The Netherlands)	Organochlorine compounds PCBs, HCB, p,p'-DDT, p,p'-DDE	34 men with poor semen quality vs. 31 men with normal (based on progressively motile sperm concentration)	CS
Tan et al, 2002 (China)	fenvalerate	32 exposed workers, 46 administrators (internal control group), 22 administrators (external control group)	CC
Swan et al, 2003b (USA)	8 metabolites of pesticides (alachlor, diazinon, acetochlor, metolachlor, 2,4-D atrazine, DEET, malathion)	Men in whom all semen parameters were low (cases) or within normal limits (controls), in Missouri & Minnesota (50 & 36 respectively)	CC
Wong et al, 2003 (The Netherlands)	Chemicals including pesticides & other factors	73 subfertile & 92 fertile men	CC
Bian et al, 2004 (China)	fenvalerate	21 exposed workers, 23 non exposed workers (internal control group), 19 non exposed workers (external control group)	CC
Dalvie et al, 2004 (South Africa)	DDT, DDE	60 workers in South Africa	CS
Kamijima et al, 2004 (Japan)	Organophosphorus & pyrethroid insecticides	18 male sprayers, 18 age matched students or medical doctors as unexposed controls	L
Meeker et al, 2004 (USA)	1N (a metabolite of carbaryl & naphthalene), TCPY (a chlorpyrifos & chlorpyrifosmethyl metabolite)	272 men recruited through a Massachusetts Infertility clinic	CS
Pant et al, 2004 (India)	DDT, DDE, DDD, HCH	45 fertile & 45 infertile men	CS
Sánchez-Peña et al, 2004 (Mexico)	mixture of organophosphorus pesticides	33 men occupationally exposed (initially 227)	CS
Xia 2004 (China)	Fenvalerate	12 exposed workers, 12 internal control group, 18 external control group	CC
Xia et al, 2005 (China)	carbaryl	16 exposed workers, 12 internal control group, 18 external control group	CC
Tan et al, 2005 (China)	carbaryl	31 exposed workers, 46 internal control group, 22 external control group	CC

De Jager et al, 2006 (Canada)	DDT p,p'-DDE	116 men aged 27 years, non occupationally exposed living in malaria, endemic areas in Mexico	CS
Lifeng et al, 2006 (China)	fenvalerate	32 exposed workers, 46 internal control group, 22 external control group	CC
Yucra et al, 2006 (Peru)	Organophosphate pesticides	31 pesticide sprayers, 80 not exposed men	CS
Aneck-Hahn et al, 2007 (South Africa)	DDT p,p'-DDE	311 healthy men 18-40 years in an endemic malaria area in which DDT is sprayed (non occupational exposure)	CS
Perry et al, 2007 (USA)	organophosphorus & pyrethroid insecticides	18 males of reproductive age in China	P
Tuc et al, 2007 (Thailand)	Mixture	1036 rice farmers: 156 of these with abnormal semen & 314 with normal semen (as controls)	CC
Meeker et al, 2008 (USA)	Metabolites of pyrethroid insecticides: 3PBA, CDCCA, TDCCA	207 men recruited from an infertility clinic	R CC
Recio-Vega et al, 2008 (Mexico)	Organophosphorus pesticides	52 men in three occupational exposure levels	L
Xia et al, 2008 (China)	3-PBA (urinary metabolite of pyrethroids)	376 men with non obstructive infertility	R CC

Table 2b. Studies evaluating the association between exposure to pesticides and human sperm quality. Studies with evidence of an association (CS: cross sectional study; CC: case-control study; L: longitudinal study; R: retrospective study; P: pilot study.)

4.3 Studies evaluating the recovery of sperm quality

There are six (6/63) studies, evaluating the recovery of sperm quality, years after cessation of exposure to DBCP and kepone (five and one study respectively). In 1986 Eaton et al. conducted a follow-up study among 44 male agricultural workers who were exposed to DBCP years ago. These workers were found to be azoospermic or oligospermic due to DBCP exposure in a previous study and their sperm was reevaluated 5 to 8 years after exposure was terminated. Only 2 of the 8 originally azoospermic workers produced sperm during the follow up and only one had normal sperm count. There was no increase in sperm production for the oligospermic men. Authors suggested a permanent destruction of germinal epithelium. Potashnik et al. conducted two follow-up studies evaluating 15 DBCP production factory workers (Potashnik & Yanai-Inbar, 1987; Potashnik & Porath, 1995). The reports reassessed their testicular function and reproductive performance, 8 and 17 years after cessation of exposure. The first study showed recovery of spermatogenesis in 4 oligo- and 3 azoospermic men, while testosterone levels of all patients were normal at all times.

The second follow up showed a significant increase in plasma FSH and LH levels in the severely affected men and no increase in the rate of spontaneous abortions or congenital malformations among pregnancies conceived during or after exposure. Recovery was evident in three of the nine azoospermic men and in three of the six oligospermic men. Olsen et al. (1990) conducted a follow-up study among azoospermic and oligospermic workers who had a maximum of 18 months of DBCP exposure. After an 11-year period, 73% of the previously azoospermic showed recovery of spermatogenesis; 13 of the men had normospermic levels and normospermic levels were found among all of the previously oligospermic men (17/17). Another study was undertaken by Lantz et al. (1981), among 14 oligospermic workers who had a maximum of 30 months of DBCP exposure. Follow up (18-21 months) showed an increase in sperm count suggesting that there is a recovery after a short term exposure. In a cross-sectional study conducted in 1982 Guzelian treated oligospermic patients who had high serum levels of chlordecone (0.6-32.0 µg/g) with cholestyramine, an anion-exchange resin. Cholestyramine binds chlordecone and increases its fecal excretion by seven-fold, resulting in reduction of chlordecone blood levels. The author found sperm motility restoration after treatment and suggested this indicated reversibility of the results.

Authors & Year Country	Pesticide	N	Study Type
Lantz et al, 1981 (USA)	DBCP	14 oligospermic workers	L
Guzelian et al, 1982 (USA)	kepone	13 workers highly exposed	CS
Eaton et al, 1986 (USA)	DBCP	44 workers 7 years after termination of exposure	L
Potashnik et al, 1987 (Israel)	DBCP	15 workers with DBCP induced azoospermia & oligospermia, 8 years after exposure.	L
Olsen et al, 1990 (USA)	DBCP	26 azoospermic and 17 oligospermic men with a maximum of 18 months of exposure	L
Potashnik et al, 1995 (Israel)	DBCP	15 workers with DBCP induced azoospermia & oligospermia, 17 years after exposure.	L

Table 2c. Studies evaluating the association between exposure to pesticides and human sperm quality. Studies evaluating the recovery of sperm quality. (CS: cross sectional study; L: longitudinal study)

4.4 Comments

The studies evaluating pesticide exposure and sperm quality were usually cross sectional. This design is relatively inexpensive and can be conducted over a short period of time. However, inherent to the study design, it is difficult to separate cause from effect, because the measurement of exposure and the possible health impairment are conducted the same simultaneously. Thus if the effect takes place later on in life, it may be misevaluated and critical windows of exposure (intrauterine, neonatal and early adulthood life) are not considered adequately.

The studies involving sperm collection have low participation rate and this, in combination with small sample size, decreases statistical power. Men providing sperm for analysis are usually self-selected volunteers that may have concerns about their fertility and this may influence the results of semen characteristics. Some embarrassment associated with sperm collection results in men who are considered fertile to be less likely to participate in such studies (selection bias). Exposure assessment in half of the reports was performed without using specific chemical analyses. Modern studies provide more accurate estimations of exposure measuring the exact levels of contaminants in humans, although there is a certain limitation regarding non-persistent pesticides that are metabolized and excreted relatively fast. Very recent studies, using biological monitoring of exposure, evaluate the susceptibility of individuals after pesticide exposure and demonstrate that genetic polymorphism modifies exposure effects on semen quality (Pérez-Herrera et al., 2008). Many of the studies that showed no or little association of exposure to pesticides and semen quality come from the north of Europe. Maybe the cumulative exposure to pollutants in this geographic area is limited compared to others. Following the evaluation of the above epidemiological studies, we conclude that overall there is suggestive evidence of an association between pesticide exposure and semen quality.

5. Pesticides and time to pregnancy

Baird et al. (1986) proposed “time to pregnancy” (TTP) as a simple measure in epidemiological studies to investigate the effects of environmental exposures on reproduction. Time to pregnancy is defined as the time interval (number of menstrual cycles, expressed in months) between the start of unprotected intercourse and a clinically recognizable pregnancy. Fecundability is defined as the probability of conception for each menstrual cycle and varies among sexually active couples not using any contraception method. The fecundability of a couple is estimated by the inverse of time to pregnancy (time to pregnancy = $1/\text{fecundability}$). TTP does not consider the exact biological pathways (including gametogenesis, fertilisation, transport and implantation of the zygote, early survival of the conception product) or mechanisms involved in human fertility. Semen quality impairment, discussed earlier, is not a direct parameter of fecundity and altered semen quality does not necessarily entail changes in TTP (Joffe, 2000). TTP appears to be a non expensive, easy-to-determine indicator (obtained by questionnaires and interviews) and a sensitive measure of fecundability in either sex (Joffe, 1997).

The evaluation of pesticide effects on fecundability using TTP could be considerably susceptible to bias as highlighted by the following discussion. TTP studies are usually retrospective and based on questionnaires and interviews, susceptible to selection, response and recall bias. Individuals that have encountered problems of infertility or subjects occupationally exposed may not have the same response to participate in related studies, compared to fertile or unexposed population. Questionnaires may concern pregnancies that occurred several years back and require the recall of events that are not well remembered. Low participation rates and small sample size may limit the statistical power of the study. Time-trend bias may be prominent if the exposure under study has changed over time. TTP studies concern planned pregnancies and fertile couples with unwanted, mistimed or because of contraception failures pregnancies are more likely to be excluded (planning bias). Because of censoring prolonged time to pregnancy period (12 months for the majority of the

studies), subfertile couples who need periods longer than 12 months to have conceive are underestimated and infertile couples are excluded (sterility bias). On the other hand couples who are in search of assisted reproduction programmes or benefit from a therapeutic effect of a treatment, after a long waiting period, are not included. It often difficult to recognize early fetal loss, therefore, early spontaneous abortions may lead to an overestimation of TTP. Studies that are not limited to first pregnancy may be biased by the “reproductively unhealthy worker effect”: fertile women might be more likely to have children, spend their time taking care of them, or having part time jobs and as a result be less occupationally exposed. Most of the TTP studies do not assess exposure by directly measuring the concentrations of pollutants in human tissues or liquids. Usually, exposure is evaluated by the same questionnaires and interviews used to estimate TTP (self-reported data), which is simple and not expensive, but inaccurate, even if the questionnaires are detailed (specific pesticides mentioned, exact hours of spraying, quantity of pesticides applied, methods of application, protective equipment used, etc). Biological monitoring seems to be a better method to evaluate exposure, despite some limitations. The measurement of non persistent pesticides concentrations in humans cannot always reflect exposure in the time to pregnancy period accurately, as they are excreted in a relatively short period of time and many pesticides cannot be measured in biological matrices (Bradman & Whyatt, 2005). Important potential confounders that have to be taken into account include: age, ethnicity, parity, previous contraceptive use, medical conditions, and frequency of intercourse, BMI, breastfeeding, smoking, caffeine and alcohol consumption.

Table 3 shows the 29 studies that evaluate exposure to pesticides and time to pregnancy. Among these studies, three studies estimate fertilization rates and time to cleavage in women attending IVF programmes, and three studies evaluate fecundity relative to dietary intake of contaminated fish. The results are conflicting, perhaps because of the heterogeneity of study design and misclassification of exposure. The only pesticides evaluated as single pesticides and not in mixtures were glyphosate (one study), and DDT and metabolites (four studies). Some of the pesticides included in mixtures were: abamectine, imidacloprid, methiocarb, pirimicarb, deltamethrin, acephate, methomyl, cyromazine, propoxur, hexaflumuron, dichlorvos (insecticides), metalaxyl, captan, procymidon, pyrazophos, toclofosmethyl, zineb, benomyl (fungicides), acrinathrin, propargite (acaricides). DDT, DDE, HCB (hexachlorobenzene), trichlorobenzene, tetrachlorobenzene, lindane, heptachlor, aldrin, chlordane, oxychlordane, endosulfan, methoxychlor, mirex were the organochlorine pesticides most commonly used.

Sanin et al. (2009) found that reduced fecundability in some regions was not associated with the use of glyphosate. DDT and metabolites were involved in four studies as a single pesticide. Toft et al. (2005), INUENDO group, showed that fecundability is inversely related with serum concentration of p,p'-DDE. Cocco et al. (2005) concluded that the fecundity ratio among spouses of DDT applicators compare to the unexposed was decreased, but the low statistical power of the study did not allow definitive conclusions. Cohn et al. (2003) measured serum levels of DDT and DDE in mothers and evaluated TTP in their daughters. Daughters' probability of pregnancy fell by 32% per 10µg/L p,p'-DDT in maternal serum, but increased 16% per 10µg/L p,p'-DDE. Axmon et al. (2006), in a big study of INUENDO, evaluating exposure to CB-153 (biphenyl) and p,p'-DDE found no effect on TTP of either male or female exposure in Sweden, Poland and Ukraine. In Greenland there seemed to be an association, but it was not possible to determine if it was due to CB-153 or p,p'-DDE.

Four studies evaluated DDT and its metabolites in mixtures. One found decreased fecundity related to exposure (Gerhard et al., 1999) and three of the studies indicated no significant association (Law et al., 2005; Axmon et al., 2006; Harley et al., 2008). Two studies examined DDE exposure in women undergoing IVF with contrary results (Jarrell et al., 1993; Younglai et al., 2002). Three studies evaluated fecundity in relation to dietary intake of contaminated fish (Courval et al., 1999; Axmon et al., 2000; Buck et al., 2000). The first suggests a modest association for men only. The second resulted in a negative association between exposure to persistent organochlorine compounds and fertility among heavy smokers and the third concluded that maternal consumption of contaminated fish may reduce fecundability. Three studies overall, evaluated the association between pesticide exposure and fertilization rates, in women attending IVF. Tielemans et al. (1999b) found an association, but exposure was assessed by questionnaires and interviews. Jarrell et al. (1993) had contrary results, but there were trace amounts of contaminants in the follicular fluid. The study conducted by Younglai et al. (2002) did not show any significant association, except for p,p'-DDE, which was associated with failed fertilization.

Curtis et al. (1999) found no significant or consistent pattern of associations between time to pregnancy and pesticides. However 6 of 13 pesticide categories were associated with decreased fecundability, during the exposure periods in which women and most of the men participated in pesticide-related activities. In a prospective study, Sallmen et al. (2003) illustrated a significant association for pyrethroids and suggestive association for carbamates and organophosphates. Restricting analyses for the first pregnancy only, a prolonged TTP was found among male and female greenhouse workers (Bretveld et al., 2006; Bretveld et al., 2008a & 2008b). According to Harley et al. (2008) prolonged TTP was related to maternal occupational pesticide exposure, home pesticide use and residence within 200 feet of an agriculture field. There was no relation with paternal occupational exposure (assessed by maternal interviews) and DDT and DDE (levels measured in maternal serum). Cole et al. (2006) using biological monitoring of exposure found that factors associated with prolonged TTP in multivariate analysis were high caffeine consumption and maternal mercury serum level. Among several pesticides analyzed, only higher maternal benzene hexachloride levels in bivariate analysis were related with prolonged TTP. However, the sample size of this study was small.

De Cock et al. (1994), Fuortes et al. (1997), Abell et al. (2000), Petrelli et al. (2001) and Idrovo et al. (2005) showed an association between impaired fecundity and exposure to pesticides, while Heacock et al. (1998), Larsen et al. (1998b), Thonneau et al. (1999), Law et al. (2005) and Lauria et al. (2006) found insignificant effects. Overall 11 studies showed an association between pesticide exposure and prolonged TTP or decreased fecundity, 7 studies found a rather inconclusive or insignificant association and 11 studies had results that are more complex to interpret and difficult to be categorised in the two groups cited above. Twenty (20/29) studies used retrospective design and only nine (9/29) used biological monitoring of exposure. Even among the studies that used biological assessment of exposure, results are conflicting. Analyses of pollutants in human tissue and liquids give important and accurate information about the cumulative exposure of an individual but there is a lack of information about timing of exposure, which may be critical for the manifestation of a reproductive impairment, such as subfecundity. Moreover, the synergistic effect of pollutants and individual susceptibility to harmful agents are factors that are difficult to estimate.

Authors & Year Country	Pesticide	N	Stuy Type
Jarrell 1993 <i>et al</i> , (Canada)	chlordan, HCB, DDE, epoxide-oxychlordane, PCB	74 women undergoing IVF	CS
De Cock 1994 <i>et al</i> , (The Netherlands)	mixture	43 fruit growers, (91 pregnancies)	CS
Fuortes <i>et al</i> , 1997 (USA)	Agriculture related exposures including pesticides	281 infertile women (cases), 216 postpartum women (controls)	CC
Heacock <i>et al</i> , 1998 (Canada)	Chlorophenate. fungicides	26487 sawmill workers, 23829 exposed, 2658 unexposed	R
Larsen <i>et al</i> , 1998b ASCLEPIOS, (Denmark)	mixture	450 traditional farmers spraying pesticides, 72 traditional not spraying, 94 organic not exposed	R
Courval <i>et al</i> , 1999 (USA)	mixture, including pesticides (dietary intake of sport fish)	626 couples (anglers)	R
Curtis <i>et al</i> , 1999 (USA)	mixture	1048 farm occupants, 2012 pregnancies	R
Gerhard <i>et al</i> , 1999 (Germany)	chlorinated hydrocarbons (CHC) (including pesticides)	489 infertile women	CS
Thonneau <i>et al</i> , 1999 ASCLEPIOS (France)	mixture	142 rural exposed workers and 220 not exposed (France); 326 exposed and 123 not exposed farmers (Denmark); 121 greenhouse workers exposed (Denmark)	R
Tielemans <i>et al</i> , 1999b (The Netherlands)	mixture	836 couples who sought in IVF treatment; 20 men exposed - 816 reference group	CC
Abell <i>et al</i> , 2000b (Denmark)	mixture used in greenhouses	492 pregnancies of women employed when they stopped contraception to have a child (starting time)	R
Axmon <i>et al</i> , 2000 (Sweden)	Organochlorine compounds dietary intake of fatty fish	Fishermen's wives from the Swedish east (n=399) and west coasts (n=936)	R
Buck <i>et al</i> , 2000 (USA)	mixture, including pesticides (dietary intake of sport fish)	895 women after years of fish consumption from lake Ontario	R
Petrelli <i>et al</i> , 2001 (Italy)	mixture of pesticides	127 greenhouse workers 173 clerical workers	R
Younglai <i>et al</i> , 2002 (Canada)	mixture of compounds including pesticides	21 couples attending IVF programme	CS
Cohn <i>et al</i> , 2003 (USA)	p,p'DDT, p,p'DDE	preserved maternal serum (n=289)	R

Sallmen et al, 2003 (Finland)	mixture	578 couples, Finnish green house employers and employees	P
Toft et al, 2005 INUENDO (Denmark)	CB-153 p,p'DDE	Poland: 472 women, 198 male spouses; Ukraine: 640 women, 208 male spouses; Greenland: 598 women, 201 male spouses; Sweden: 559 women, 191male spouses	R CS
Cocco et al, 2005 (Italy)	DDT	spouses of 105 men exposed	R
Idrovo et al, 2005 (Colombia)	mixture of pesticides including thiocarbamates (propineb, mancozeb)	2085 women working in cut flower production	CS R
<i>Law et al, 2005 (USA)</i>	PCBs, DDE	390 pregnant women enrolled at 12 study centers in US between 1959-1965 (before PCBs or DDE was banned in US)	R
<i>Axmon et al, 2006 INUENDO (Sweden)</i>	CB 153 p,p'-DDE	778 men, 1505 women (couples from Greenland, Poland, Ukraine & a cohort of Swedish wives)	R
Bretveld et al, 2006 (The Netherlands)	mixture of pesticides	398 female greenhouse workers - 524 referents	R
<i>Cole et al, 2006 (Canada)</i>	Mixture (metals, PCBs, organochlorine pesticides)	41 couples having their first pregnancy	R
<i>Lauria et al, 2006 (Italy)</i>	mixture	713 female greenhouse workers	R
Bretveld et al, 2008a (The Netherlands)	mixture	694 greenhouse workers exposed, 613 unexposed reference group	R
Bretveld et al, 2008b (The Netherlands)	mixture	101 couples with female greenhouse workers, 957 couples with male greenhouse workers; 1408 referents	R
Harley et al, 2008 (USA)	mixture of pesticides, DDT, DDE	402 pregnant women living in a farmworker community; in 289 of them DDT and DDE serum levels measured.	CS
<i>Sanin et al, 2009 (Mexico)</i>	Glyphosate (herbicide)	2592 fertile Colombian women from 5 regions with different exposure	R

Table 3. Studies evaluating exposure to pesticides and time to pregnancy. In italics: Studies that reveal no association. (CS: cross sectional study; CC: case-control study; L: longitudinal study; R: retrospective study; P: prospective study.)

6. Conclusive remarks

It is challenging to estimate if exposure to pesticides has harmful effects on human fertility, because there are many factors that can influence, limit or bias the results of the studies related to fertility. There are two main issues that have to be considered: the methods used to estimate fertility and accuracy of exposure assessment. Related epidemiological studies use basically two methods of fertility evaluation, the semen quality analysis, which is a surrogate and is indirect, and TTP assessment, which is considerably susceptible to bias. As cited before, fecundity is the important parameter that we have to study and that is the capability of producing offspring and not the actual production (fertility). Fertility rates are increasing in developed countries but this is not a proof that fecundity is increasing too. In actuality, human fecundity seems to be decreasing. Fertility rates are influenced by many factors such as socioeconomic status, immigration, health condition and medical service, the practice of contraception and abortion, the sexual behaviour and frequency of intercourse, the methods of assisted reproduction. Given the influence of age on female fecundity, the effects of the baby boom generation, the evolution of the two-child family and the trend for marriage and childbearing later on in life, a decline in fecundity would be expected.

The decline in semen quality overtime is still equivocal, perhaps because the standard assessments (count, motility and morphology) are insufficient. It is interesting that documented changes in semen quality are not always correlated with parallel changes in TTP. In the future, consideration of additional parameters of sperm quality that provide information about the sperm-egg binding event would be helpful. Modern techniques employed to evaluate DNA damage in spermatozoa such as single cell gel electrophoresis (COMET assay), Terminal transferase dUTP Nick End Labelling (TUNEL), sperm chromatin structure assay (SCSA), in situ nick translation (ISNT) as well as and sperm aneuploidy (FISH) have to be incorporated in future studies.

Pesticides are endocrine disrupting chemicals warranting research to elucidate their mechanisms of action. Timing of exposure is critical, as the same dosage seems to have diverse effects in individuals of different age and exposure in critical windows of development (intrauterine life, neonatal period, childhood) seems to be more harmful. The classical dosage-effect rule of toxicology is contradicted by some EDCs and low doses may damage more than high doses. Individuals are often exposed to a mixture of toxicants, not a single agent, and the effects may be synergistic or antagonistic. The results of an exposure may be apparent immediately but can also become apparent on at a later stage and sometimes in subsequent generations, with varying susceptibility that may depend on genetic polymorphism. Often, studies requiring semen collection have low participation rates and thus the men providing sperm for analysis are usually self selected volunteers that may have concerns about their own fertility and may not be a true reflection of the general population. Geographical and seasonal variation of semen parameters make results difficult to compare. Most of the epidemiological studies reporting chemical exposure and semen quality damage are cross-sectional. In this design, the measurement of exposure and the possible health impairment are conducted simultaneously and, because of the factors cited above, results can be easily biased. A need for a longitudinal design in studies of semen quality is apparent.

TTP studies are usually retrospective. TTP estimation and exposure assessment are carried out using questionnaires and interviews, a method relatively inexpensive, but not precise. Unplanned pregnancies are not included in TTP studies. Therefore, fertile couples, because

of mistimed, unwanted or contraceptive failure pregnancies are excluded. A percentage of early fetal losses is not recognised and not included in these studies. Subfertile couples are underestimated because of censoring prolonged time to pregnancy to a period of around 12 months. Fertile women might be more likely to have children and be less busy in occupational activities and therefore less occupationally exposed (reproductively unhealthy worker effect). Restricting TTP studies for women attending only their first pregnancy is required. A prospective design that recruits couples early at the beginning of their attempt at a pregnancy, with more than one chemical analyses of specific pollutants per individual, during study period, might be less susceptible to bias and give more concrete results. In the majority of the studies cited above, exposure was evaluated by a simple questionnaire. There are few studies that directly measure the concentration of a toxicant in human tissue or liquids. Precise measurements of follicular fluid are more difficult to obtain. Biological monitoring is more precise, although with certain limits for non persistent pesticides and for the estimation of the primary source of exposure, since cumulative levels of pollutants are measured.

In evaluating epidemiological studies of exposure to pesticides and human semen quality, there is suggestive evidence of an association between exposure and semen quality impairment. The evaluation of the scientific literature related to human pesticide exposure and TTP results are conflicting. However, research results in animal and in vitro studies are clear and suggest that pesticides can adversely affect fertility. Unfortunately, these results cannot be taken as proof for humans strong enough to stop or modulate pesticide production. In the future, pesticides might be less toxic and it will therefore be even more difficult to estimate their adverse effects in human fertility. The evidence for impaired fecundity is strong in both in vivo and in vitro studies and investigators have to cooperate in order to optimise the research and raise public awareness and concern. Therefore, scientists need to emphasize the need for stricter rules dealing with chemical safety.

7. References

- Abell A, Ernst E, Bonde JP. Semen quality and sexual hormones in greenhouse workers. *Scand J Work Environ Health* 2000a;26:492-500.
- Abell A, Juul S, Bonde JP. Time to pregnancy among female greenhouse workers. *Scand J Work Environ Health* 2000b;26:131-6.
- Adamopoulos DA, Pappa A, Nicopoulou S, Andreou E, Karamertzanis M, Michopoulos J, et al. Seminal volume and total sperm number trends in men attending subfertility clinics in the greater Athens area during the period 1977-1993. *Hum Reprod* 1996;11:1936-41.
- Adiga SK, Jayaraman V, Kalthur G, Upadhya D, Kumar P. Declining semen quality among south Indian infertile men: A retrospective study. *J Hum Reprod Sci* 2008;1:15-8.
- Alavanja MC, Hoppin JA, Kamel F. Health effects of chronic pesticide exposure: cancer and neurotoxicity. *Ann Rev Public Health* 2004;25:155-97.
- Almagor M, Ivnitzki I, Yaffe H, Baras M. Changes in semen quality in Jerusalem between 1990 and 2000: a cross-sectional and longitudinal study. *Arch Androl* 2003;49:139-44.
- Andersen HR, Vinggaard AM, Rasmussen TH, Gjermandsen IM, Bonefeld-Jørgensen EC. Effects of currently used pesticides in assays for estrogenicity, androgenicity, and aromatase activity in vitro. *Toxicol Appl Pharmacol* 2002;179:1-12.

- Andolz P, Bielsa MA, Vila J. Evolution of semen quality in North-eastern Spain: a study in 22,759 infertile men over a 36 year period. *Hum Reprod* 1999;14:731-5.
- Aneck-Hahn NH, Schulenburg GW, Bornman MS, Farias P, de Jager C. Impaired semen quality associated with environmental DDT exposure in young men living in a malaria area in the Limpopo Province, South Africa. *J Androl* 2007;28:423-34.
- Anway MD, Skinner MK. Epigenetic transgenerational actions of endocrine disruptors. *Endocrinology* 2006;147(6 Suppl):S43-9.
- Axmon A, Rylander L, Strömberg U, Hagmar L. Time to pregnancy and infertility among women with a high intake of fish contaminated with persistent organochlorine compounds. *Scand J Work Environ Health* 2000;26:199-206.
- Axmon A, Thulstrup AM, Rignell-Hydbom A, Pedersen HS, Zvezday V, Ludwicki JK, et al. INUENDO. Time to pregnancy as a function of male and female serum concentrations of 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) and 1,1-dichloro-2,2-bis(p-chlorophenyl)-ethylene (p,p'-DDE). *Hum Reprod* 2006;21:657-65.
- Ayotte P, Giroux S, Dewailly E, Hernández Avila M, Farias P, Danis R, et al. DDT spraying for malaria control and reproductive function in Mexican men. *Epidemiology* 2001;12:366-7.
- Bahshwan SA, Owen CP, Nicholls PJ, Smith HJ, Ahmadi M. Some 1-((benzofuran-2-yl)methyl)imidazoles as inhibitors of 17 alpha-hydroxylase: 17, 20-lyase (P450 17) and their specificity patterns. *J Pharm Pharmacol* 1998;50:1109-16.
- Baird DD, Wilcox AJ, Weinberg CR. Use of time to pregnancy to study environmental exposures. *Am J Epidemiol* 1986;124:470-80.
- Bassil KL, Vakil C, Sanborn M, Cole DC, Kaur JS, Kerr KJ. Cancer health effects of pesticides: systematic review. *Can Fam Physician*. 2007;53:1704-11.
- Benshushan A, Shoshani O, Paltiel O, Schenker JG, Lewin A. Is there really a decrease in sperm parameters among healthy young men? A survey of sperm donations during 15 years. *J Assist Reprod Genet* 1997;14:347-53.
- Berling S, Wölner-Hanssen P. No evidence of deteriorating semen quality among men in infertile relationships during the last decade: a study of males from Southern Sweden. *Hum Reprod* 1997;12:1002-5.
- Bian Q, Xu LC, Wang SL, Xia YK, Tan LF, Chen JF, et al. Study on the relation between occupational fenvalerate exposure and spermatozoa DNA damage of pesticide factory workers. *Occup Environ Med* 2004;61:999-1005.
- Bigsby R, Chapin RE, Daston GP, Davis BJ, Gorski J, Gray LE, et al. Evaluating the effects of endocrine disruptors on endocrine function during development. *Environ Health Perspect* 1999;107(Suppl 4):613-8.
- Bilotta P, Guglielmo R, Steffè M. Analysis of decline in seminal fluid in the Italian population during the past 15 years. *Minerva Ginecol* 1999;51:223-31.
- Bradman A, Whyatt RM. Characterizing exposures to non persistent pesticides during pregnancy and early childhood in the National Children's Study: a review of monitoring and measurement methodologies. *Environ Health Perspect* 2005;113:1092-9.
- Bretveld RW, Thomas CM, Scheepers PT, Zielhuis GA, Roeleveld N. Pesticide exposure: the hormonal function of the female reproductive system disrupted? *Reprod Biol Endocrinol* 2006;4:30.

- Bretveld RW, Hooiveld M, Zielhuis GA, Pellegrino A, van Rooij IA, Roeleveld N. Reproductive disorders among male and female greenhouse workers. *Reprod Toxicol* 2008a;25:107-14.
- Bretveld R, Kik S, Hooiveld M, van Rooij I, Zielhuis G, Roeleveld N. Time-to-pregnancy among male greenhouse workers. *Occup Environ Med* 2008b;65:185-90.
- Buck GM, Vena JE, Schisterman EF, Dmochowski J, Mendola P, Sever LE, et al. Parental consumption of contaminated sport fish from Lake Ontario and predicted fecundability. *Epidemiology* 2000;11:388-93.
- Can A, Albertini DF. Stage specific effects of carbendazim (MBC) on meiotic cell cycle progression in mouse oocytes. *Mol Reprod Dev* 1997;46:351-62.
- Cannon SB, Veazey JM Jr, Jackson RS, Burse VW, Hayes C, Straub WE, et al. Epidemic kepone poisoning in chemical workers. *Am J Epidemiol* 1978;107:529-37.
- Carlsen E, Giwercman A, Keiding N, Skakkebaek NE. Evidence for decreasing quality of semen during past 50 years. *BMJ* 1992;305:609-13.
- Charlier CJ, Foidart JM. Comparative study of dichlorodiphenyldichloroethylene in blood and semen of two young male populations: lack of relationship to infertility, but evidence of high exposure of the mothers. *Reprod Toxicol* 2005;20:215-20.
- Cocco P, Fadda D, Ibba A, Melis M, Tocco MG, Atzeri S, et al. Reproductive outcomes in DDT applicators. *Environ Res.* 2005;98:120-6.
- Cohn BA, Cirillo PM, Wolff MS, Schwingl PJ, Cohen RD, Sholtz RI, et al. DDT and DDE exposure in mothers and time to pregnancy in daughters. *Lancet* 2003;361:2205-6.
- Colborn T, vom Saal FS, Soto AM. Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ Health Perspect* 1993;101:378-84
- Cole DC, Wainman B, Sanin LH, Weber JP, Muggah H, Ibrahim S. Environmental contaminant levels and fecundability among non-smoking couples. *Reprod Toxicol* 2006;22:13-9.
- Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, et al. World Health Organization reference values for human semen characteristics. *Hum Reprod Update* 2010;16:231-45
- Courval JM, De Hoog JV, Stein AD, Tay EM, He J, Humphrey HE, et al. Sport-caught fish consumption and conception delay in licensed Michigan anglers. *Environ Res* 1999;80:183-8.
- Cox C, Sorgan M. Unidentified inert ingredients in pesticides: implications for human and environmental health. *Environ Health Perspect* 2006;114:1803-6.
- Crews D, Putz O, Thomas P, Hayes T, Howdeshell K. Animal models for the study of the effects of mixtures, low doses, and the embryonic environment on the action of endocrine disrupting chemicals. *Pure and Applied Chemistry, SCOPE/IUPAC Project Implications of Endocrine Active Substances for Humans and Wildlife.* 2003;75:2305-20
- Crisp TM, Clegg ED, Cooper RL, Wood WP, Anderson DG, Baetcke KP, et al. Environmental endocrine disruption: an effects assessment and analysis. *Environ Health Perspect* 1998;106(Suppl 1):11-56.
- Cummings AM, Gray LE Jr. Methoxychlor affects the decidual cell response of the uterus but not other progestational parameters in female rats. *Toxicol Appl Pharmacol* 1987;90:330-6.

- Curtis KM, Savitz DA, Weinberg CR, Arbuckle TE. The effect of pesticide exposure on time to pregnancy. *Epidemiology* 1999;10:112-7.
- Dallinga JW, Moonen EJ, Dumoulin JC, Evers JL, Geraedts JP, Kleinjans JC. Decreased human semen quality and organochlorine compounds in blood. *Hum Reprod* 2002;17:1973-9.
- Dalvie MA, Myers JE, Thompson ML, Robins TG, Dyer S, Riebow J, et al. The long-term effects of DDT exposure on semen, fertility, and sexual function of malaria vector-control workers in Limpopo Province, South Africa. *Environ Res* 2004;96:1-8.
- De Cock J, Westveer K, Heederik D, te Velde E, van Kooij R. Time to pregnancy and occupational exposure to pesticides in fruit growers in The Netherlands. *Occup Environ Med* 1994;51:693-9.
- De Jager C, Farias P, Barraza-Villarreal A, Avila MH, Ayotte P, Dewailly E, et al. Reduced seminal parameters associated with environmental DDT exposure and p,p'-DDE concentrations in men in Chiapas, Mexico: a cross-sectional study. *J Androl* 2006;27:16-27.
- Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, et al. Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocr Rev* 2009;30:293-342.
- Dohle GR, Colpi GM, Hargreave TB, Papp GK, Jungwirth A, Weidner W. EAU Working Group on Male Infertility EAU guidelines on male infertility. *Eur Urol* 2005;48:703-11
- Eaton M, Schenker M, Whorton MD, Samuels S, Perkins C, Overstreet J. Seven-year follow-up of workers exposed to 1,2-dibromo-3-chloropropane. *J Occup Med* 1986;28:1145-50.
- Egnatz DG, Ott MG, Townsend JC, Olson RD, Johns DB. DBCP and testicular effects in chemical workers: an epidemiological survey in Midland, Michigan. *J Occup Med* 1980;22:727-32.
- Feki NC, Abid N, Rebai A, Sellami A, Ayed BB, Guermazi M, et al. Semen quality decline among men in infertile relationships: experience over 12 years in the South of Tunisia. *J Androl* 2009;30:541-7.
- Fisch H, Goluboff ET, Olson JH, Feldshuh J, Broder SJ, Barad DH. Semen analyses in 1,283 men from the United States over a 25-year period: no decline in quality. *Fertil Steril* 1996;65:1009-14.
- Food and Agriculture Organization of the United Nations. *FAO international code of conduct on the distribution and use of pesticides*. Rome: Food and Agriculture Organization of the United Nations; 2003.
- Fourth National Report on Human Exposure to Environmental Chemicals. Centers for Disease Control and Prevention, U.S. Department of Health and Human Services, 2009
- Fuortes L, Clark MK, Kirchner HL, Smith EM. Association between female infertility and agricultural work history. *Am J Ind Med* 1997;31:445-51.
- García AM. Pesticide exposure and women's health. *Am J Ind Med* 2003;44:584-94.
- Garey J, Wolff MS. Estrogenic and antiprogesteragenic activities of pyrethroid insecticides. *Biochem Biophys Res Commun* 1998;251:855-9.
- Gerhard I, Monga B, Krähe J, Runnebaum B. Chlorinated hydrocarbons in infertile women. *Environ Res* 1999;80:299-310.

- Gore AC, Heindel JJ, Zoeller RT. Endocrine disruption for endocrinologists (and others). *Endocrinology* 2006;147(Suppl 6):S1-3.
- Gore A. Endocrine-disrupting chemicals: from basic research to clinical practice. In: Cohn PM, ed. *Contemporary Endocrinology*. Totowa, NJ: Humana Press 2007.
- Gore AC. Neuroendocrine systems as targets for environmental endocrine-disrupting chemicals. *Fertil Steril* 2008;89(Suppl 2):e101-2.
- Guo YL, Lambert GH, Hsu CC. Growth abnormalities in the population exposed in utero and early postnatally to polychlorinated biphenyls and dibenzofurans. *Environ Health Perspect* 1995;103(Suppl 6):117-22.
- Guzelian PS. Chlordecone poisoning: a case study in approaches for detoxification of humans exposed to environmental chemicals. *Drug Metab Rev* 1982;13:663-79.
- Härkönen K, Viitanen T, Larsen SB, Bonde JP, Lähdetie J. Aneuploidy in sperm and exposure to fungicides and lifestyle factors. ASCLEPIOS. A European Concerted Action on Occupational Hazards to Male Reproductive Capability. *Environ Mol Mutagen* 1999;34:39-46.
- Harley KG, Marks AR, Bradman A, Barr DB, Eskenazi B. DDT exposure, work in agriculture, and time to pregnancy among farmworkers in California. *J Occup Environ Med* 2008;50:1335-42.
- Hauser R, Chen Z, Pothier L, Ryan L, Altshul L. The relationship between human semen parameters and environmental exposure to polychlorinated biphenyls and p,p'-DDE. *Environ Health Perspect* 2003a;111:1505-11.
- Hauser R, Singh NP, Chen Z, Pothier L, Altshul L. Lack of an association between environmental exposure to polychlorinated biphenyls and p,p'-DDE and DNA damage in human sperm measured using the neutral comet assay. *Hum Reprod* 2003b;18:2525-33
- Heacock H, Hogg R, Marion SA, Hershler R, Teschke K, Dimich-Ward H, et al. Fertility among a cohort of male sawmill workers exposed to chlorophenyl fungicides. *Epidemiology* 1998;9:56-60.
- Henderson J, Rennie GC, Baker HW. Association between occupational group and sperm concentration in infertile men. *Clin Reprod Fertil* 1986;4:275-81.
- Herbst AL, Ulfelder H, Poskanzer DC, Longo LD. Adenocarcinoma of the vagina. Association of maternal stilbestrol therapy with tumor appearance in young women. 1971. *Am J Obstet Gynecol* 1999;181:1574-5.
- Hirsch KS, Adams ER, Hoffman DG, Markham JK, Owen NV. Studies to elucidate the mechanism of fenarimol-induced infertility in male rats. *Toxicol Appl Pharmacol* 1986;86:391-9.
- Horak S, Kaminska J, Olejek A. The fertilization potential of donor semen between 1982 and 2004 in the industrial area of Upper Silesia (Poland). *Ann Agric Environ Med* 2008;15:113-8.
- Idrovo AJ, Sanin LH, Cole D, Chavarro J, Cáceres H, Narváez J, et al. Time to first pregnancy among women working in agricultural production. *Int Arch Occup Environ Health* 2005;78:493-500.
- Itoh N, Kayama F, Tatsuki TJ, Tsukamoto T. Have sperm counts deteriorated over the past 20 years in healthy, young Japanese men? Results from the Sapporo area. *J Androl* 2001;22:40-4.

- Jarrell JF, Villeneuve D, Franklin C, Bartlett S, Wrixon W, Kohut J, et al. Contamination of human ovarian follicular fluid and serum by chlorinated organic compounds in three Canadian cities. *CMAJ* 1993;148:1321-7.
- Jeyaratnam J. Acute pesticide poisoning: a major global health problem. *World Health Stat Q* 1990;43:139-44.
- Joffe M. Time to pregnancy: a measure of reproductive function in either sex. *Asclepios Project. Occup Environ Med* 1997;54:289-95.
- Joffe M. Time trends in biological fertility in Britain. *Lancet* 2000;355:1961-5.
- Jørgensen N, Andersen AG, Eustache F, Irvine DS, Suominen J, Petersen JH, et al. Regional differences in semen quality in Europe. *Hum Reprod* 2001;16:1012-9.
- Jørgensen N, Asklund C, Carlsen E, Skakkebaek NE. Coordinated European investigations of semen quality: results from studies of Scandinavian young men is a matter of concern. *Int J Androl* 2006;29:54-61.
- Juhler RK, Larsen SB, Meyer O, Jensen ND, Spanò M, Giwercman A, et al. Human semen quality in relation to dietary pesticide exposure and organic diet. *Arch Environ Contam Toxicol* 1999;37:415-23.
- Kamijima M, Hibi H, Gotoh M, Taki K, Saito I, Wang H, et al. A survey of semen indices in insecticide sprayers. *J Occup Health* 2004;46:109-18.
- Kavlock RJ, Daston GP, DeRosa C, Fenner-Crisp P, Gray LE, Kaattari S, et al. Research needs for the risk assessment of health and environmental effects of endocrine disruptors: a report of the U.S. EPA-sponsored workshop. *Environ Health Perspect* 1996;104 Suppl 4:715-40.
- Kelce WR, Monosson E, Gamcsik MP, Laws SC, Gray LE Jr. Environmental hormone disruptors: evidence that vinclozolin developmental toxicity is mediated by antiandrogenic metabolites. *Toxicol Appl Pharmacol* 1994;126:276-85.
- Kelce WR, Stone CR, Laws SC, Gray LE, Kemppainen JA, Wilson EM. Persistent DDT metabolite p,p'-DDE is a potent androgen receptor antagonist. *Nature* 1995;375:581-5.
- Kelce WR, Wilson EM. Environmental antiandrogens: developmental effects, molecular mechanisms, and clinical implications. *J Mol Med* 1997;75:198-207.
- Kojima H, Katsura E, Takeuchi S, Niiyama K, Kobayashi K. Screening for estrogen and androgen receptor activities in 200 pesticides by *in vitro* reporter gene assays using Chinese hamster ovary cells. *Environ Health Perspect* 2004;112:524-31.
- Krause A, Krause W. Seasonal variations in human seminal parameters. *Eur J Obstet Gynecol Reprod Biol* 2002;101:175-8.
- Lantz GD, Cunningham GR, Huckins C, Lipshultz LI. Recovery from severe oligospermia after exposure to dibromochloropropane. *Fertil Steril* 1981;35:46-53.
- Larsen SB, Giwercman A, Spanò M, Bonde JP. A longitudinal study of semen quality in pesticide spraying Danish farmers. The ASCLEPIOS Study Group. *Reprod Toxicol* 1998a;12:581-9.
- Larsen SB, Joffe M, Bonde JP. Time to pregnancy and exposure to pesticides in Danish farmers. ASCLEPIOS Study Group. *Occup Environ Med* 1998b;55:278-83.
- Lathers CM. Endocrine disruptors: a new scientific role for clinical pharmacologists? Impact on human health, wildlife, and the environment. *J Clin Pharmacol* 2002;42:7-23.
- Lauria L, Settini L, Spinelli A, Figà-Talamanca I. Exposure to pesticides and time to pregnancy among female greenhouse workers. *Reprod Toxicol* 2006;22:425-30.

- Law DC, Klebanoff MA, Brock JW, Dunson DB, Longnecker MP. Maternal serum levels of polychlorinated biphenyls and 1,1-dichloro-2,2-bis(p-chlorophenyl)ethylene (DDE) and time to pregnancy. *Am J Epidemiol* 2005;162:523-32.
- Lerda D, Rizzi R. Study of reproductive function in persons occupationally exposed to 2,4-dichlorophenoxyacetic acid (2,4-D). *Mutat Res* 1991;262:47-50.
- Levine RJ. Seasonal variation of semen quality and fertility. *Scand J Work Environ Health* 1999;25(Suppl 1):34-7
- Li DK, Yan B, Li Z, Gao E, Miao M, Gong D, et al. Exposure to magnetic fields and the risk of poor sperm quality. *Reprod Toxicol* 2010;29:86-92.
- Lifeng T, Shoulin W, Junmin J, Xuezhao S, Yannan L, Qianli W, et al. Effects of fenvalerate exposure on semen quality among occupational workers. *Contraception* 2006;73:92-6.
- Lilienthal H, Hack A, Roth-Härer A, Grande SW, Talsness CE. Effects of developmental exposure to 2,2,4,4,5-pentabromodiphenyl ether (PBDE-99) on sex steroids, sexual development, and sexually dimorphic behavior in rats. *Environ Health Perspect* 2006;114:194-201.
- Lindenau A, Fischer B, Seiler P, Beier HM. Effects of persistent chlorinated hydrocarbons on reproductive tissues in female rabbits. *Hum Reprod* 1994;9:772-80.
- Lipshultz LI, Ross CE, Whorton D, Milby T, Smith R, Joyner RE. Dibromochloropropane and its effect on testicular function in man. *J Urol* 1980;124:464-8.
- Magnusdottir EV, Thorsteinsson T, Thorsteinsdottir S, Heimisdottir M, Olafsdottir K. Persistent organochlorines, sedentary occupation, obesity and human male subfertility. *Hum Reprod* 2005;20:208-15.
- Mahadevaswami MP, Jadaramkunti UC, Hiremath MB, Kaliwal BB. Effect of mancozeb on ovarian compensatory hypertrophy and biochemical constituents in hemicastrated albino rat. *Reprod Toxicol* 2000;14:127-34.
- Mallidis C, Howard EJ, Baker HW. Variation of semen quality in normal men. *Int J Androl* 1991;14:99-107.
- Manna PR, Kero J, Tena-Sempere M, Pakarinen P, Stocco DM, Huhtaniemi IT. Assessment of mechanisms of thyroid hormone action in mouse Leydig cells: regulation of the steroidogenic acute regulatory protein, steroidogenesis, and luteinizing hormone receptor function. *Endocrinology* 2001;142:319-31.
- McCauley LA, Anger WK, Keifer M, Langley R, Robson MG, Rohlman D. Studying health outcomes in farmworker populations exposed to pesticides. *Environmental Health Perspectives* 2006;114:953-60.
- Meeker JD, Ryan L, Barr DB, Herrick RF, Bennett DH, Bravo R, et al. The relationship of urinary metabolites of carbaryl/naphthalene and chlorpyrifos with human semen quality. *Environ Health Perspect* 2004;112:1665-70.
- Meeker JD, Barr DB, Hauser R. Human semen quality and sperm DNA damage in relation to urinary metabolites of pyrethroid insecticides. *Hum Reprod* 2008;23:1932-40.
- Merzenich H, Zeeb H, Blettner M. Decreasing sperm quality: a global problem? *BMC Public Health* 2010;10:24
- Mikkilä TF, Toppari J, Paranko J. Effects of neonatal exposure to 4-tert-octylphenol, diethylstilbestrol, and flutamide on steroidogenesis in infantile rat testis. *Toxicol Sci* 2006;91:456-66.

- Multigner L, Kadhel P, Pascal M, Huc-Terki F, Kercret H, Massart C, et al. Parallel assessment of male reproductive function in workers and wild rats exposed to pesticides in banana plantations in Guadeloupe. *Environ Health* 2008;7:40.
- Oduma JA, Oduor Okelo D, Odongo H, Makawiti DW. The pesticide heptachlor affects steroid hormone secretion in isolated follicular and luteal cells of rat. *Comp Biochem Physiol C Toxicol Pharmacol* 2006;144:76-84.
- Oliva A, Spira A, Multigner L. Contribution of environmental factors to the risk of male infertility. *Hum Reprod* 2001;16:1768-76.
- Olsen GW, Lanham JM, Bodner KM, Hylton DB, Bond GG. Determinants of spermatogenesis recovery among workers exposed to 1,2-dibromo-3-chloropropane. *J Occup Med* 1990;32:979-84.
- Olsen GW, Bodner KM, Ramlow JM, Ross CE, Lipshultz LI. Have sperm counts been reduced 50 percent in 50 years? A statistical model revisited. *Fertil Steril* 1995;63:887-93.
- Paasch U, Salzbrunn A, Glander HJ, Plambeck K, Salzbrunn H, Grunewald S, et al. Semen quality in sub-fertile range for a significant proportion of young men from the general German population: a co-ordinated, controlled study of 791 men from Hamburg and Leipzig. *Int J Androl* 2008;31:93-102.
- Padungtod C, Savitz DA, Overstreet JW, Christiani DC, Ryan LM, Xu X. Occupational pesticide exposure and semen quality among Chinese workers. *J Occup Environ Med* 2000;42:982-92.
- Pant N, Mathur N, Banerjee AK, Srivastava SP, Saxena DK. Correlation of chlorinated pesticides concentration in semen with seminal vesicle and prostatic markers. *Reprod Toxicol* 2004;19:209-14.
- Paulsen CA, Berman NG, Wang C. Data from men in greater Seattle area reveals no downward trend in semen quality: further evidence that deterioration of semen quality is not geographically uniform. *Fertil Steril* 1996;65:1015-20.
- Peiris-John RJ, Wickremasinghe R. Impact of low-level exposure to organophosphates on human reproduction and survival. *Trans R Soc Trop Med Hyg* 2008;102:239-45.
- Pérez-Herrera N, Polanco-Minaya H, Salazar-Arredondo E, Solís-Heredia MJ, Hernández-Ochoa I, Rojas-García E et al. PON1Q192R genetic polymorphism modifies organophosphorous pesticide effects on semen quality and DNA integrity in agricultural workers from southern Mexico. *Toxicol Appl Pharmacol* 2008;230:261-8.
- Perry MJ, Venners SA, Barr DB, Xu X. Environmental pyrethroid and organophosphorus insecticide exposures and sperm concentration. *Reprod Toxicol* 2007;23:113-8.
- Petrelli G, Figà-Talamanca I. Reduction in fertility in male greenhouse workers exposed to pesticides. *Eur J Epidemiol* 2001;17:675-7.
- Potashnik G, Ben-Aderet N, Israeli R, Yanai-Inbar I, Sober I. Suppressive effect of 1,2-dibromo-3-chloropropane on human spermatogenesis. *Fertil Steril* 1978;30:444-7.
- Potashnik G, Yanai-Inbar I. Dibromochloropropane (DBCP): an 8-year reevaluation of testicular function and reproductive performance. *Fertil Steril* 1987;47:317-23.
- Potashnik G, Porath A. Dibromochloropropane (DBCP): a 17-year reassessment of testicular function and reproductive performance. *J Occup Environ Med* 1995;37:1287-92.
- Purvis K, Christiansen E. Male infertility: current concepts. *Ann Med* 1992;24:259-72.


- Ramazzini C. Collegium ramazzini statement on the control of pesticides in the European Union: A call for action to protect human health. *Am J Ind Med* 2009;52:176-7.
- Ratcliffe JM, Schrader SM, Steenland K, Clapp DE, Turner T, Hornung RW. Semen quality in papaya workers with long term exposure to ethylene dibromide. *Br J Ind Med* 1987;44:317-26.
- Recio-Vega R, Ocampo-Gómez G, Borja-Aburto VH, Moran-Martínez J, Cebrian-Garcia ME. Organophosphorus pesticide exposure decreases sperm quality: association between sperm parameters and urinary pesticide levels. *J Appl Toxicol* 2008;28:674-80.
- Ribas-Fitó N. *Silent Invaders: Pesticides, Livelihoods and Women's Health*. London: ZED Books, 2002. *Int J Epidemiol* 2006;35:4-5.
- Rignell-Hydbom A, Rylander L, Giwercman A, Jönsson BA, Nilsson-Ehle P, Hagmar L. Exposure to CB-153 and p,p'-DDE and male reproductive function. *Hum Reprod* 2004;19:2066-75.
- Rignell-Hydbom A, Rylander L, Giwercman A, Jönsson BA, Lindh C, Eleuteri P, et al. Exposure to PCBs and p,p'-DDE and human sperm chromatin integrity. *Environ Health Perspect* 2005;113:175-9.
- Ritter L, Goughleff NC, Arbuckle T, Cole D, Raizenne M. Addressing the linkage between exposure to pesticides and human health effects – research trends and priorities for research. *J Toxicol Environ Health B Crit Rev* 2006;9:441-56.
- Rubin MM. Antenatal exposure to DES: lessons learned...future concerns. *Obstet Gynecol Surv* 2007;62:548-55.
- Sallmén M, Liesivuori J, Taskinen H, Lindbohm ML, Anttila A, Aalto L, et al. Time to pregnancy among the wives of Finnish greenhouse workers. *Scand J Work Environ Health* 2003;29:85-93.
- Sanborn M, Kerr KJ, Sanin LH, Cole DC, Bassil KL, Vakil C. Non-cancer health effects of pesticides: systematic review and implications for family doctors. *Can Fam Physician* 2007; 53:1712-20.
- Sánchez-Peña LC, Reyes BE, López-Carrillo L, Recio R, Morán-Martínez J, Cebrián ME, et al. Organophosphorous pesticide exposure alters sperm chromatin structure in Mexican agricultural workers. *Toxicol Appl Pharmacol* 2004;196:108-13.
- Sanin LH, Carrasquilla G, Solomon KR, Cole DC, Marshall EJ. Regional differences in time to pregnancy among fertile women from five Colombian regions with different use of glyphosate. *J Toxicol Environ Health A* 2009;72:949-60.
- Schrader SM, Turner TW, Ratcliffe JM. The effects of ethylene dibromide on semen quality: a comparison of short-term and chronic exposure. *Reprod Toxicol* 1988;2:191-8.
- Seo JT, Rha KH, Park YS, Lee MS. Semen quality over a 10-year period in 22,249 men in Korea. *Int J Androl* 2000;23:194-8.
- Silvestroni L, Palleschi S. Effects of organochlorine xenobiotics on human spermatozoa. *Chemosphere* 1999;39:1249-52.
- Slutsky M, Levin JL, Levy BS. Azoospermia and oligospermia among a large cohort of DBCP applicators in 12 countries. *Int J Occup Environ Health* 1999;5:116-22.
- Smith JL, Garry VF, Rademaker AW, Martin RH. Human sperm aneuploidy after exposure to pesticides. *Mol Reprod Dev* 2004;67:353-9.

- Soto AM, Sonnenschein C, Chung KL, Fernandez MF, Olea N, Serrano FO. The E-SCREEN assay as a tool to identify estrogens: an update on estrogenic environmental pollutants. *Environ Health Perspect* 1995;103(Suppl 7):113-22.
- Spanò M, Toft G, Hagmar L, Eleuteri P, Rescia M, Rignell-Hydbom A, et al. Exposure to PCB and p,p'-DDE in European and Inuit populations: impact on human sperm chromatin integrity. *Hum Reprod* 2005;20:3488-99.
- Sripada S, Fonseca S, Lee A, Harrild K, Giannaris D, Mathers E, et al. Trends in semen parameters in the northeast of Scotland. *J Androl* 2007;28:313-9.
- Stoker TE, Laws SC, Guidici DL, Cooper RL. The effect of atrazine on puberty in male wistar rats: an evaluation in the protocol for the assessment of pubertal development and thyroid function. *Toxicol Sci* 2000;58:50-9.
- Strohmer H, Boldizar A, Plöckinger B, Feldner-Busztin M, Feichtinger W. Agricultural work and male infertility. *Am J Ind Med* 1993;24:587-92.
- Stronati A, Manicardi GC, Cecati M, Bordicchia M, Ferrante L, Spanò M, et al. Relationships between sperm DNA fragmentation, sperm apoptotic markers and serum levels of CB-153 and p,p'-DDE in European and Inuit populations. *Reproduction* 2006;132:949-58.
- Surgan MH. Toxicity tests: "inert" and active ingredients. *Environ Health Perspect* 2005;113:A657
- Swan SH, Elkin EP, Fenster L. Have sperm densities declined? A reanalysis of global trend data. *Environ Health Perspect* 1997;105:1228-32.
- Swan SH, Elkin EP. Declining semen quality: can the past inform the present? *Bioessays* 1999;21:614-21.
- Swan SH, Elkin EP, Fenster L. The question of declining sperm density revisited: an analysis of 101 studies published 1934-1996. *Environ Health Perspect* 2000;108:961-6.
- Swan SH, Brazil C, Drobnis EZ, Liu F, Kruse RL, Hatch M, et al. Geographic differences in semen quality of fertile U.S. males. *Environ Health Perspect* 2003a;111:414-20.
- Swan SH, Kruse RL, Liu F, Barr DB, Drobnis EZ, Redmon JB, et al. Semen quality in relation to biomarkers of pesticide exposure. *Environ Health Perspect* 2003b;111:1478-84.
- Swan SH. Semen quality in fertile US men in relation to geographical area and pesticide exposure. *Int J Androl* 2006;29:62-8
- Tan LF, Wang SL, Sun XZ, Li YN, Wang QL, Ji JM, et al. Effects of fenvalerate exposure on the semen quality of occupational workers. *Zhonghua Nan Ke Xue* 2002;8:273-6.
- Tan LF, Sun XZ, Li YN, Ji JM, Wang QL, Chen LS, et al. Effects of carbaryl production exposure on the sperm and semen quality of occupational male workers. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 2005;23:87-90.
- Taxvig C, Vinggaard AM, Hass U, Axelstad M, Metzdorff S, Nellemann C. Endocrine-disrupting properties in vivo of widely used azole fungicides. *Int J Androl* 2008;31:170-7.
- Thonneau P, Abell A, Larsen SB, Bonde JP, Joffe M, Clavert A, et al. Effects of pesticide exposure on time to pregnancy: results of a multicenter study in France and Denmark. ASCLEPIOS Study Group. *Am J Epidemiol* 1999;150:157-63.
- Thrupp LA. Sterilization of workers from pesticide exposure: the causes and consequences of DBCP-induced damage in Costa Rica and beyond. *Int J Health Serv* 1991;21:731-57.

- Tielemans E, Burdorf A, te Velde ER, Weber RF, van Kooij RJ, Veulemans H, et al. Occupationally related exposures and reduced semen quality: a case-control study. *Fertil Steril* 1999a;71:690-6.
- Tielemans E, van Kooij R, te Velde ER, Burdorf A, Heederik D. Pesticide exposure and decreased fertilisation rates in vitro. *Lancet* 1999b;354:484-5.
- Toft G, Axmon A, Giwercman A, Thulstrup AM, Rignell-Hydbom A, Pedersen HS, et al. Fertility in four regions spanning large contrasts in serum levels of widespread persistent organochlorines: a cross-sectional study. *Environ Health* 2005;4:26
- Toft G, Rignell-Hydbom A, Tyrkiel E, Shvets M, Giwercman A, Lindh CH, et al. Semen quality and exposure to persistent organochlorine pollutants. *Epidemiology* 2006;17:450-8.
- Tomenson JA, Taves DR, Cockett AT, McCusker J, Barraij L, Francis M, et al. An assessment of fertility in male workers exposed to molinate. *J Occup Environ Med* 1999;41:771-87.
- Toppari J, Larsen JC, Christiansen P, Giwercman A, Grandjean P, Guillette LJ Jr, et al. Male reproductive health and environmental xenoestrogens. *Environ Health Perspect* 1996;104(Suppl. 4):741-803
- Tuc VP, Wangsuphachart V, Tasanapradit P, Fungladda W, Van Trong P, Nhung NT. Impacts of pesticide use on semen characteristics among rice farmers in Kienxuong District, Thaibinh Province, Vietnam. *Southeast Asian J Trop Med Public Health* 2007;38:569-75.
- Vicari E, Conticello A, Battiato C, La Vignera S. Sperm characteristics in fertile men and healthy men of the south-east Sicily from year 1982 to 1999. *Arch Ital Urol Androl* 2003;75:28-34.
- Vinggaard AM, Hnida C, Breinholt V, Larsen JC. Screening of selected pesticides for inhibition of CYP19 aromatase activity in vitro. *Toxicol In Vitro* 2000;14:227-34.
- Von Saal FS, Akingbemi BT, Belcher SM, Birnbaum LS, Crain DA, Eriksen M, et al. Chapel Hill bisphenol A expert panel consensus statement: integration of mechanisms, effects in animals and potential to impact human health at current levels of exposure. *Reprod Toxicol* 2007;24:131-8.
- Walsh LP, Stocco DM. Effects of lindane on steroidogenesis and steroidogenic acute regulatory protein expression. *Biol Reprod* 2000;63:1024-33.
- Wang JM, Wu XL, You W, Ling LX, Wu J, Zhang GY. Pharmacokinetic and pharmacodynamic studies of the effect of ketoconazole on reproductive function in male rats. *Int J Androl* 1992;15:376-84.
- Weiss JM, Bauer O, Blüthgen A, Ludwig AK, Vollersen E, Kaisi M, et al. Distribution of persistent organochlorine contaminants in infertile patients from Tanzania and Germany. *J Assist Reprod Genet* 2006;23:393-9.
- Weselak M, Arbuckle TE, Foster W. Pesticide exposures and developmental outcomes: The epidemiological evidence. *J Toxicol Environ Health B Crit Rev* 2007;10:41-80.
- Whorton D, Krauss RM, Marshall S, Milby TH. Infertility in male pesticide workers. *Lancet* 1977;2:1259-61.
- Whorton MD, Milby TH, Stubbs HA, Avashia BH, Hull EQ. Testicular function among carbaryl-exposed employees. *J Toxicol Environ Health* 1979;5:929-41.

- Wong WY, Zielhuis GA, Thomas CM, Merkus HM, Steegers-Theunissen RP. New evidence of the influence of exogenous and endogenous factors on sperm count in man. *Eur J Obstet Gynecol Reprod Biol* 2003;110:49-54.
- Wyrobek AJ, Watchmaker G, Gordon L, Wong K, Moore D 2nd, Whorton D. Sperm shape abnormalities in carbaryl-exposed employees. *Environ Health Perspect* 1981;40:255-65.
- Xia Y, Bian Q, Xu L, Cheng S, Song L, Liu J, et al. Genotoxic effects on human spermatozoa among pesticide factory workers exposed to fenvalerate. *Toxicology* 2004;203:49-60.
- Xia Y, Cheng S, Bian Q, Xu L, Collins MD, Chang HC, et al. Genotoxic effects on spermatozoa of carbaryl-exposed workers. *Toxicol Sci* 2005;85:615-23.
- Xia Y, Han Y, Wu B, Wang S, Gu A, Lu N, et al. The relation between urinary metabolite of pyrethroid insecticides and semen quality in humans. *Fertil Steril* 2008;89:1743-50.
- Younglai EV, Collins JA, Foster WG. Canadian semen quality: an analysis of sperm density among eleven academic fertility centers. *Fertil Steril* 1998;70:76-80.
- Younglai EV, Foster WG, Hughes EG, Trim K, Jarrell JF. Levels of environmental contaminants in human follicular fluid, serum, and seminal plasma of couples undergoing in vitro fertilization. *Arch Environ Contam Toxicol* 2002;43:121-6.
- Yucra S, Rubio J, Gasco M, Gonzales C, Steenland K, Gonzales GF. Semen quality and reproductive sex hormone levels in Peruvian pesticide sprayers. *Int J Occup Environ Health* 2006;12:355-61.
- Zvěřina J, Urbánek V, Cirýn J. (Spermiologic parameters in Czech men from 1950 to 1984: analysis of data from the archives of the Sexology Institute of the First Medical School of Charles University and the General Medical School Hospital in Prague) *Sb Lek* 2002; 103:35-47.

IntechOpen



Pesticides - Formulations, Effects, Fate

Edited by Prof. Margarita Stoytcheva

ISBN 978-953-307-532-7

Hard cover, 808 pages

Publisher InTech

Published online 21, January, 2011

Published in print edition January, 2011

This book provides an overview on a large variety of pesticide-related topics, organized in three sections. The first part is dedicated to the "safer" pesticides derived from natural materials, the design and the optimization of pesticides formulations, and the techniques for pesticides application. The second part is intended to demonstrate the agricultural products, environmental and biota pesticides contamination and the impacts of the pesticides presence on the ecosystems. The third part presents current investigations of the naturally occurring pesticides degradation phenomena, the environmental effects of the break down products, and different approaches to pesticides residues treatment. Written by leading experts in their respective areas, the book is highly recommended to the professionals, interested in pesticides issues.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Stavros Sifakis, Mihalīs Mparmpas, Offie P. Soldin and Aristides Tsatsakis (2011). Pesticide Exposure and Health Related Issues in Male and Female Reproductive System, Pesticides - Formulations, Effects, Fate, Prof. Margarita Stoytcheva (Ed.), ISBN: 978-953-307-532-7, InTech, Available from:
<http://www.intechopen.com/books/pesticides-formulations-effects-fate/pesticide-exposure-and-health-related-issues-in-male-and-female-reproductive-system>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen