

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Therapeutic Approach to Focal and Segmental Glomerulosclerosis (FSGS) Recurrence in Kidney Transplant Recipients

Guillaume Canaud^{1,2,3}, Frank Martinez^{1,2} and Christophe Legendre^{1,2}

¹Université Paris Descartes, Rue de l'École de Médecine, Paris,

²Service de Transplantation et Unité de soins intensifs, Hôpital Necker, Paris,

³INSERM U845, Hôpital Necker

⁴Service de Néphrologie Pédiatrique, Hôpital des Enfants Malades, Paris, France

1. Introduction

Primary focal and segmental glomerulosclerosis (FSGS) is the most frequently acquired disease leading to end stage renal disease (ESRD) in children (NAPRTCS 2004, annual report)¹⁻⁶. The recurrence of FSGS after kidney transplantation is frequent (20-40%) and is associated with poor graft survival⁷⁻¹³. The pathophysiology of primary FSGS remains uncertain, but secretion of a circulating factor is suspected to play a key role in excessive glomerular permeability. Excepted for the recurrence of FSGS with a previous allograft, this event is almost unpredictable and characterized by early nephrotic range proteinuria. Some studies have identified risk factors for recurrent FSGS in children, including rapid progression to ESRD, young age of onset of FSGS and, of course loss of a previous graft from recurrent disease but none of these studies can clearly separate the patients who will or will not be affected by recurrence. The treatment of recurrence remains controversial, and most reports relate to use of plasma exchange (PE) in uncontrolled trials with relatively small groups of patients and conflicting results. PE and protein immunoadsorption can markedly reduce urinary protein excretion and induce complete remission in some cases, but they usually fail to achieve sustained remission¹⁴⁻¹⁶. Steroids and cyclosporine are also associated with inconstant remission and act through different ways. More recent therapeutic approach seems promising and need to be evaluated in large trials. This review summarizes the various therapeutic approaches to FSGS recurrence.

2. Pathogenesis of FSGS recurrence

To better understand the therapeutic approaches to FSGS recurrence, we might briefly comment on its pathogenesis. The pathophysiology of this disease remains largely unclear and is thought to involve at least three types of cells (T-cells, B-cells and podocyte) and a circulating factor. In 1974, to explain what would later be called idiopathic nephrotic syndrome (INS), Shalhoub proposed the hypothesis (Shalhoub hypothesis) of T-cell dysfunction resulting in the secretion of "*circulating chemical mediator toxic to an*

*immunologically innocent glomerular basement membrane*¹⁷. Although knowledge of the pathogenesis of this disease has improved in recent years, novel therapeutics have not ameliorated the rate of its recurrence. T-cell disorder is still suspected based on much evidence for Th2 cytokine bias in INS. Sahali et al. were the first to report that T-cells from INS patients are driven toward a Th2 phenotype¹⁸. By screening a cDNA library from T-cells of patients with relapsing INS they found that overexpression of truncated c-maf-inducing protein (Tc-mip) induced c-maf, transactivated the IL-4 gene, and downregulated IFN γ expression, characteristic of Th2 commitment¹⁹. These data were also supported by the observation of increased expression of cytoplasmic IL-13 mRNA in T-cells from patients with relapsing INS as compared to those in remission and control patients²⁰. Furthermore, overexpression of IL-13 in Wistar rats led to minimal change disease confirming the potential role for Th2 predominance in pathophysiology of INS²¹. More recently, the expression level of soluble ST2 protein (sST2), a marker of Th2 cells that is predicted to be regulated by c-maf, was investigated in a population of patient with FSGS recurrence after transplantation²². The level of sST2 were increased after transplantation in patients with recurrence compared to the control group; however this protein was unable to induce podocyte injury *in vitro* suggesting that it could be a marker of recurrence but might not be implicated in the pathogenesis of recurrence. Using a humanized mouse model, Sellier-Leclerc et al showed that CD34⁺ stem cells, unlike CD34⁻ peripheral blood mononuclear cells, from patients with INS induced albuminuria, suggesting the involvement of immature differentiating cells rather than mature peripheral T-cells in the pathogenesis of the disease²³. Thus, there is evidence for T-cell dysfunction in INS, but treatments targeting T-cells (calcineurin inhibitors, anti-CD3 or anti-CD52) are not completely effective in preventing or treating FSGS recurrence. Furthermore, there is recent evidence for B-cell participation from a report of remission obtained with rituximab (anti-CD20)²⁴. Indeed, it is accepted that primary FSGS is the consequence of a complex interaction between T- and B-cells leading to the secretion of a circulating factor targeting podocyte. The presence of a circulating vascular permeability factor implicated in the physiopathology of this disease is highlighted by (i) the early recurrence of nephrotic syndrome after transplantation²⁵, (ii) the fact that serum from patients with recurrent FSGS infused in rats can induce albuminuria²⁶, (iii) the occurrence of a transient nephrotic syndrome in newborn infants of women with FSGS²⁷, and (iv) the efficiency of plasma exchange and/or immunoadsorption at inducing remission¹⁵. The biochemical characteristics of this factor, however, are still unknown. Its molecular weight is suspected to be between 30-100 kDa, and Dantal et al. suggested that it could be a part of a complex with immunoglobulins²⁸. Recently, Savin et al. found that this circulating factor had a high affinity for galactose and that all its activity was eliminated by galactose affinity columns²⁹. The third player in this complex disease is the podocyte. Podocytes are a post-mitotic cell, arrested in G2/M phase of the cell cycle, and do not proliferate. It has been postulated that the circulating factor and/or immune cells directly interact with podocytes, and could induce the redistribution of the protein of the slit diaphragm, the loss of nephrin and/or podocin and the effacement of the foot processes, a hallmark of podocyte injury³⁰. It has been suggested that foot processes effacement, if reversed, can lead to the restoration of glomerular architecture, which is typically observed in steroid-sensitive minimal change disease. The failure of repair mechanisms promotes podocyte detachment, apoptosis, podocyte depletion and FSGS. A special situation is collapsing glomerulopathy, a specific variant of FSGS. Findings in human suggest that

podocytes may undergo a proliferative state and phenotypic changes in collapsing glomerulopathy^{31, 32}. Importantly, evidence suggests that, in addition to podocytes, parietal epithelial cells also participate in the collapsing glomerulopathy phenotype³³. Recently, Reiser et al. described the expression of the co-stimulatory molecule B7.1 on podocytes³⁴. The significance of the presence of this molecule is not clearly understood and remains speculative.

3. Treatment of FSGS recurrence

Despite the introduction of new immunosuppressive regimens, discovery of cyclosporine, and use of induction therapies, the incidence of FSGS recurrence has remained unchanged³⁵⁻³⁷. In the case of recurrence treatments are not standardized, and show inconstant results. Most reports concern children and involve a small number of patients^{15, 38-46}. Cyclosporine (CsA) has shown some degree of efficacy in pediatric subjects^{43, 44}. Indeed, intravenous CsA after FSGS recurrence was associated with a drastic decrease of proteinuria in 82% of patients, although PE was added in some resistant cases⁴³. A few reports have studied the use of tacrolimus to prevent or to treat recurrence. The results of these studies are almost inconstant^{47, 48}. Due to the presence of a circulating permeability factor, most transplant teams test the efficacy of PE, which substitutes the plasma of the patient with either plasma from healthy pooled donors, albumin or colloidal substance. This treatment usually induces a reduction in proteinuria and, in some cases complete remission. Determining when to begin PEs, their frequency, duration and optimal stop time are challenging to determine, and most remissions are PE-dependant. Others supporting approaches have also been used, such as anti-human immunoglobulin affinity immunoadsorption and tryptophan immunoadsorption^{28, 49}.

We recently conducted a non-randomized pilot trial of intensive and prolonged multiple treatment of FSGS recurrence in adult kidney transplant recipients⁵⁰. As this complex disease involves systemic immune dysregulation targeting podocyte, we used a strategy of concomitant high dose steroids and intravenous CsA not only for their immunosuppressive properties, but also for their podocyte cytoskeleton stabilization properties, and with PE sessions. Glucocorticoid receptors are present on podocytes, have anti apoptotic properties *in vitro*⁵¹ and could stabilize the actin cytoskeleton⁵². CsA acts directly on podocytes by blocking calcineurin-mediated dephosphorylation of synaptopodin and stabilizes actin cytoskeleton⁵³. We decided to add PE sessions based on the presence of the circulating factor and an increased five-year allograft survival rate in our own experience. Indeed, from January 1994 to December 2004, we observed that the five-year allograft survival rate was 55% in cases of FSGS recurrence (R group), compared to 93% in cases without recurrence (NR group) ($p < 0.01$). In the R group, patients who benefited from PE had an increased five-year allograft survival rate (91%) similar to the NR group, whereas patients not treated with PE had a five-year allograft survival rate of only 40% ($p < 0.01$).

The details of our therapeutic strategy consisted of high dose of oral steroids (1 mg/kg/d) for the first 4 weeks followed by tapering according to the following sequence: 0.75 mg/kg/d for 2 weeks, 0.5 mg/kg/d for two weeks, 0.25 mg/kg/d for two weeks and 0.2 mg/kg/d or a maximal daily dose of 10 mg thereafter. We used 14 days of intravenous CsA (2 mg/kg, targeting a blood level between 200 and 400 ng/ml) followed by oral treatment, targeting C2 levels between 1,200 and 1,400 ng/ml. PEs were performed with 5% albumin replacement (three sessions per week for three weeks, followed by two sessions per week

for three weeks, one session per week until month 3, two sessions per month until month 5, and once per month until month 9). Once remission was obtained, an angiotensin-converting enzyme inhibitor and/or angiotensin receptor blocker were introduced. Rescue with rituximab therapy was anticipated in case of failure. The primary endpoint was the induction of a complete (<0.3 g per 24H) and sustained (>12 months) remission of proteinuria. The treatment was started after 3 days of nephrotic-range proteinuria without spontaneous improvement. Of 286 kidney transplant procedures performed between September 2005 and March 2007 at our institution, 18 were performed in patients with primary FSGS. After transplantation these individuals were carefully monitored for proteinuria and ten (56%, eight men, two women) exhibited recurrence. The mean age at the onset of primary FSGS in native kidneys was 19.1 ± 14.8 years, leading to ESRD 8.1 ± 8.6 years later. The mean duration of hemodialysis was 2.8 ± 2.4 years. For nine of the patients, this transplant was their first; the remaining patient had received two previous kidney transplants, both of which had been quickly lost due to recurrence. After kidney transplantation, all patients received an immunosuppressive regimen including prednisone, mycophenolate mofetil, oral CsA ($n=10$) or tacrolimus ($n=8$), along with an induction of either basiliximab ($n=12$) or antithymocyte globulin (Thymoglobulin®, $n=6$). Proteinuria occurred immediately after transplantation in six patients (60%) and early post-transplant in the remaining four (range 4-55 days). Recurrence was associated with massive proteinuria (mean 12 ± 11 g/day). We began PE within the first ten days of diagnosis in all patients, while high-dose steroids and intravenous CsA were initiated at the time of diagnosis. Complete proteinuria remission was achieved in all patients at a mean of 22.9 ± 6.6 days post-diagnosis. Three months after diagnosis, all patients were still in complete remission (mean proteinuria 0.16 ± 0.09 g/day) and all but one patient remained in remission at one year (mean proteinuria 0.19 ± 0.29 g/d). In the nine patients with complete remission, PE was tapered gradually until month 9 and then stopped. During early follow-up (mean 15.8 ± 3.3 months), none of these nine patients relapsed. At one year post-transplant, mean serum creatinine was $121 \pm 29 \mu\text{mol/l}$ and the measured iohexol glomerular filtration rate was $68.5 \pm 18 \text{ml/min}$. The patient who failed to obtain sustained remission of proteinuria was the individual who had undergone a third kidney transplant. In this patient, proteinuria recurred when the frequency of PE was tapered to less than once per week. Rituximab infusion was attempted, and after two doses circulating B-cells were completely depleted ($\text{CD}19 < 1/\text{mm}^3$). Proteinuria was maintained in the range 1-2 g/day with PE administered twice per month.

Long term follow-up (34.2 ± 6.7 months) of this cohort revealed that all these patients were still in complete remission with a mean proteinuria level of 0.11 ± 0.07 g/day and a mean serum creatinine of $122.6 \pm 18 \mu\text{mol/l}$. The patient in partial remission went into complete remission by maintaining PE (two times per month). During the follow-up, none of these patients developed diabetes, cancer or severe infectious disease. Indeed, long-term follow-up of our cohort revealed that intensive treatment of FSGS induces not only an excellent short term response but also a complete and sustained remission of proteinuria without significant immunosuppressive regimen-associated adverse effects.

We then considered whether the treatment administered in case of FSGS recurrence impacted the pathological features of FSGS during the post-transplant course. To address this question, we retrospectively studied 77 patients from January 1984 to December 2007, including both children and adults with primary FSGS who underwent renal transplantation⁵⁴. Of these, 42 patients experienced a recurrence of nephrotic-range

proteinuria. After kidney transplantation, recurrence of proteinuria occurred immediately in 32 of the 42 cases (children, $n = 28$; adults, $n = 4$), early in 9 of the 42 cases (children, $n = 4$; adults, $n = 5$) and late in 1 of the 42 case (adult, $n = 1$). Briefly, at time of recurrence, day 6 to day 60 days after transplantation, no glomerular lesion was observed in 32 of 33 biopsies and we considered them as minimal change disease. Only one patient had already developed a FSGS lesion. At month 3, FSGS lesions were observed in 11 of 39 cases, and at month 12, they were observed in 14 of 37 cases. Interestingly, 17 of the 42 patients went into complete and sustained remission, and none of them developed FSGS. On the other hand, patients who never achieved complete and sustained remission developed FSGS lesions. These data suggest continuing intensive treatments if no FSGS lesion are observed on transplant biopsies.

4. Other available treatments

4.1 Preemptive PE

A few studies of preemptive PE have been reported that show inconstant efficacy and lack a control group. In 2005, Gohh et al conducted a prospective study to test whether pre-transplant PE could prevent recurrence of primary FSGS in high risk patients⁵⁵. A high risk of FSGS recurrence was defined by a rapid evolution toward ESRD ($n = 4$) or prior allograft loss due to recurrence ($n = 6$). Patients were subjected to a course of eight PE sessions over 2 weeks in the immediate peri-operative period. Recipients of living donor kidneys initiated PE treatments 1 week before transplantation and completed their course at the end of the first post-operative week. Recipients of cadaver kidneys underwent an initial PE within 24 h of implantation. FSGS recurred in 3 of 10 patients, each of whom had lost prior transplants to recurrent FSGS. Two of these progressed to ESRD and the third had significant renal dysfunction. The authors conclude that PE may decrease incidence of FSGS recurrence in this particular population (rate of recurrence expected without PE: 60%). This therapeutic approach is difficult to organize in case of deceased donors, which is the major group of donors in many countries including our own. Furthermore, PE session before or soon after transplantation, may increase the risk of major bleeding despite the finding that no adverse event were reported in the previous study. This approach would also lead to excessive treatment in 50% of patients.

4.2 Rituximab

In 2006, Pescovitz et al reported complete remission after infusion of rituximab in a young transplant recipient²⁴. After kidney transplantation this child rapidly developed FSGS recurrence resistant to PE and CsA and at month 5 developed a post-transplantation lymphoproliferative disease (PTLD). After six infusions of rituximab to treat the PTLD, proteinuria disappeared suggesting a possible interaction between B- and T-cells that leads to the secretion of permeability factor. Since that report, many transplant teams have tested the ability of rituximab to treat FSGS recurrence with inconstant results⁵⁶⁻⁵⁹. In fact, rituximab seems to induce remission in about 50% of cases, but some questions remain unsolved. When should the infusion begin: as an induction therapy or at time of recurrence? How many infusions should be administered, given that depletion of circulating B-cells does not always correlate with lymphoid organ depletion⁶⁰? What are the long term side effects of this treatment? To date, no consensus has emerged, and double-blind studies are needed to determine the therapeutic potential of rituximab.

4.3 Anti-CTLA4

The co-stimulation molecule B7.1 is normally expressed on antigen presenting cells and B-cells. Recently, Reiser et al found that B7.1 is also expressed on podocyte and could be upregulated in many proteinuric states³⁴. Again, the significance of the presence of this molecule is not clearly understood and remains speculative. To date, no published studies have evaluated blockade of this co-stimulation pathway for the treatment of FSGS recurrence.

4.4 Anti-TNF α

TNF α mRNA was found to be upregulated in macrophages preceding the development of nephrotic syndrome in Buffalo/Mna rats⁶¹. Furthermore, a high level of TNF α mRNA was detected in mononuclear cells from patients with FSGS⁶². Anti-TNF α therapy was recently tested in a child with resistant FSGS recurrence⁶³ and induced transient complete remission, but every relapse was sensitive to anti-TNF α infusion.

4.5 Retinoic acid, Roscovitine and cyclin dependant kinase inhibitor

Collapsing glomerulopathy recurrence is a situation in which podocytes can proliferate. Using *in vitro* and *in vivo* mouse models, retinoic acids were found to be efficient at reducing podocyte proliferation and proteinuria⁶⁴. An on going study is evaluating treatment of collapsing glomerulopathy in native kidneys using retinoic acid (NCT00098020).

4.6 Galactose

Recently, Savin et al. found that the circulating factor has a high affinity for galactose, and that its activity can be removed by use of galactose affinity columns²⁹. This group and one another reported a significant reduction in proteinuria following administration of galactose along with others therapeutics^{29, 65}. A clinical trial (NCT00098020) is recruiting patients to treat primary FSGS in native kidney with galactose. Galactose has an excellent safety profile and could be an interesting therapeutic candidate.

5. Conclusion

Primary FSGS remains mysterious with a poorly understood pathogenesis. Recurrence is still frequent and associated with a poor allograft prognosis. We clearly must increase our knowledge of the pathogenesis of this disease to better identify specific risk factors for recurrence and design more specific therapeutic strategies. Our pilot study, although limited by a small population size, provides very encouraging results. Combined and intensive therapy showed a markedly beneficial effect on early proteinuria recurrence in this cohort of adult kidney transplant recipients. These preliminary results require confirmation on a larger scale with extensive follow-up. We have also described patients who went into complete and sustained remission and did not develop FSGS; patients who never achieved complete and sustained remission developed FSGS lesions. New therapeutics such as rituximab, anti-TNF α , galactose and retinoic acid should be evaluated in randomized double-blind studies. A better understanding of the molecular function of podocytes give hope that new therapeutics will be available in the next future.

6. References


- [1] Habib R, Levy M, Gubler MC. Clinicopathologic correlations in the nephrotic syndrome. *Paediatrician* 1979;8(5-6):325-48.

- [2] Cameron JS. The enigma of focal segmental glomerulosclerosis. *Kidney Int Suppl* 1996;57:S119-31.
- [3] Haas M, Meehan SM, Karrison TG, Spargo BH. Changing etiologies of unexplained adult nephrotic syndrome: a comparison of renal biopsy findings from 1976-1979 and 1995-1997. *Am J Kidney Dis* 1997;30(5):621-31.
- [4] Braden GL, Mulhern JG, O'Shea MH, Nash SV, Ucci AA, Jr., Germain MJ. Changing incidence of glomerular diseases in adults. *Am J Kidney Dis* 2000;35(5):878-83.
- [5] Kitiyakara C, Eggers P, Kopp JB. Twenty-one-year trend in ESRD due to focal segmental glomerulosclerosis in the United States. *Am J Kidney Dis* 2004;44(5):815-25.
- [6] Filler G, Young E, Geier P, Carpenter B, Drukker A, Feber J. Is there really an increase in non-minimal change nephrotic syndrome in children? *Am J Kidney Dis* 2003;42(6):1107-13.
- [7] Cameron JS, Senguttuvan P, Hartley B, et al. Focal segmental glomerulosclerosis in fifty-nine renal allografts from a single centre; analysis of risk factors for recurrence. *Transplant Proc* 1989;21(1 Pt 2):2117-8.
- [8] Dantal J, Baatard R, Hourmant M, Cantarovich D, Buzelin F, Souillou JP. Recurrent nephrotic syndrome following renal transplantation in patients with focal glomerulosclerosis. A one-center study of plasma exchange effects. *Transplantation* 1991;52(5):827-31.
- [9] Ingulli E, Tejani A. Incidence, treatment, and outcome of recurrent focal segmental glomerulosclerosis posttransplantation in 42 allografts in children--a single-center experience. *Transplantation* 1991;51(2):401-5.
- [10] Dantal J, Souillou JP. Relapse of focal segmental glomerulosclerosis after kidney transplantation. *Adv Nephrol Necker Hosp* 1996;25:91-106.
- [11] Tejani A, Stablein DH. Recurrence of focal segmental glomerulosclerosis posttransplantation: a special report of the North American Pediatric Renal Transplant Cooperative Study. *J Am Soc Nephrol* 1992;2(12 Suppl):S258-63.
- [12] Artero ML, Sharma R, Savin VJ, Vincenti F. Plasmapheresis reduces proteinuria and serum capacity to injure glomeruli in patients with recurrent focal glomerulosclerosis. *Am J Kidney Dis* 1994;23(4):574-81.
- [13] Pardon A, Audard V, Caillard S, et al. Risk factors and outcome of focal and segmental glomerulosclerosis recurrence in adult renal transplant recipients. *Nephrol Dial Transplant* 2006;21(4):1053-9.
- [14] Savin VJ, Sharma R, Sharma M, et al. Circulating factor associated with increased glomerular permeability to albumin in recurrent focal segmental glomerulosclerosis. *The New England journal of medicine* 1996;334(14):878-83.
- [15] Dantal J, Bigot E, Bogers W, et al. Effect of plasma protein adsorption on protein excretion in kidney-transplant recipients with recurrent nephrotic syndrome. *The New England journal of medicine* 1994;330(1):7-14.
- [16] Artero M, Biava C, Amend W, Tomlanovich S, Vincenti F. Recurrent focal glomerulosclerosis: natural history and response to therapy. *Am J Med* 1992;92(4):375-83.
- [17] Shalhoub RJ. Pathogenesis of lipoid nephrosis: a disorder of T-cell function. *Lancet* 1974;2(7880):556-60.
- [18] Sahali D, Pawlak A, Valanciute A, et al. A novel approach to investigation of the pathogenesis of active minimal-change nephrotic syndrome using subtracted cDNA library screening. *J Am Soc Nephrol* 2002;13(5):1238-47.

- [19] Grimbert P, Valanciute A, Audard V, et al. Truncation of C-mip (Tc-mip), a new proximal signaling protein, induces c-maf Th2 transcription factor and cytoskeleton reorganization. *The Journal of experimental medicine* 2003;198(5):797-807.
- [20] Yap HK, Cheung W, Murugasu B, Sim SK, Seah CC, Jordan SC. Th1 and Th2 cytokine mRNA profiles in childhood nephrotic syndrome: evidence for increased IL-13 mRNA expression in relapse. *J Am Soc Nephrol* 1999;10(3):529-37.
- [21] Lai KW, Wei CL, Tan LK, et al. Overexpression of interleukin-13 induces minimal-change-like nephropathy in rats. *J Am Soc Nephrol* 2007;18(5):1476-85.
- [22] Bruneau S, Le Berre L, Herve C, et al. Potential role of soluble ST2 protein in idiopathic nephrotic syndrome recurrence following kidney transplantation. *Am J Kidney Dis* 2009;54(3):522-32.
- [23] Sellier-Leclerc AL, Duval A, Riveron S, et al. A humanized mouse model of idiopathic nephrotic syndrome suggests a pathogenic role for immature cells. *J Am Soc Nephrol* 2007;18(10):2732-9.
- [24] Pescovitz MD, Book BK, Sidner RA. Resolution of recurrent focal segmental glomerulosclerosis proteinuria after rituximab treatment. *The New England journal of medicine* 2006;354(18):1961-3.
- [25] Hoyer JR, Vernier RL, Najarian JS, Raij L, Simmons RL, Michael AF. Recurrence of idiopathic nephrotic syndrome after renal transplantation. *Lancet* 1972;2(7773):343-8.
- [26] Zimmerman SW. Increased urinary protein excretion in the rat produced by serum from a patient with recurrent focal glomerular sclerosis after renal transplantation. *Clinical nephrology* 1984;22(1):32-8.
- [27] Lagrue G, Branellec A, Niaudet P, Heslan JM, Guillot F, Lang P. [Transmission of nephrotic syndrome to two neonates. Spontaneous regression]. *Presse Med* 1991;20(6):255-7.
- [28] Dantal J, Godfrin Y, Koll R, et al. Antihuman immunoglobulin affinity immunoadsorption strongly decreases proteinuria in patients with relapsing nephrotic syndrome. *J Am Soc Nephrol* 1998;9(9):1709-15.
- [29] Savin VJ, McCarthy ET, Sharma R, Charba D, Sharma M. Galactose binds to focal segmental glomerulosclerosis permeability factor and inhibits its activity. *Transl Res* 2008;151(6):288-92.
- [30] Kim BK, Hong HK, Kim JH, Lee HS. Differential expression of nephrin in acquired human proteinuric diseases. *Am J Kidney Dis* 2002;40(5):964-73.
- [31] Canaud G, Bruneval P, Noel LH, et al. Glomerular Collapse Associated With Subtotal Renal Infarction in Kidney Transplant Recipients With Multiple Renal Arteries. *Am J Kidney Dis* 2009.
- [32] Bariety J, Nochy D, Mandet C, Jacquot C, Glotz D, Meyrier A. Podocytes undergo phenotypic changes and express macrophagic-associated markers in idiopathic collapsing glomerulopathy. *Kidney international* 1998;53(4):918-25.
- [33] Dijkman HB, Weening JJ, Smeets B, et al. Proliferating cells in HIV and pamidronate-associated collapsing focal segmental glomerulosclerosis are parietal epithelial cells. *Kidney international* 2006;70(2):338-44.
- [34] Reiser J, von Gersdorff G, Loos M, et al. Induction of B7-1 in podocytes is associated with nephrotic syndrome. *The Journal of clinical investigation* 2004;113(10):1390-7.
- [35] Raafat R, Travis LB, Kalia A, Diven S. Role of transplant induction therapy on recurrence rate of focal segmental glomerulosclerosis. *Pediatric nephrology (Berlin, Germany)* 2000;14(3):189-94.

- [36] Gagnadoux MF. Ask the expert. Does antibody induction therapy with daclizumab or basiliximab increase the risk of recurrence of post-transplant focal segmental glomerulosclerosis? *Pediatric nephrology (Berlin, Germany)* 2002;17(4):305.
- [37] Hubsch H, Montane B, Abitbol C, et al. Recurrent focal glomerulosclerosis in pediatric renal allografts: the Miami experience. *Pediatric nephrology (Berlin, Germany)* 2005;20(2):210-6.
- [38] Laufer J, Ettenger RB, Ho WG, Cohen AH, Marik JL, Fine RN. Plasma exchange for recurrent nephrotic syndrome following renal transplantation. *Transplantation* 1988;46(4):540-2.
- [39] Ingulli E, Tejani A, Butt KM, et al. High-dose cyclosporine therapy in recurrent nephrotic syndrome following renal transplantation. *Transplantation* 1990;49(1):219-21.
- [40] Cochat P, Kassir A, Colon S, et al. Recurrent nephrotic syndrome after transplantation: early treatment with plasmapheresis and cyclophosphamide. *Pediatric nephrology (Berlin, Germany)* 1993;7(1):50-4.
- [41] Dall'Amico R, Ghiggeri G, Carraro M, et al. Prediction and treatment of recurrent focal segmental glomerulosclerosis after renal transplantation in children. *Am J Kidney Dis* 1999;34(6):1048-55.
- [42] Cheong HI, Han HW, Park HW, et al. Early recurrent nephrotic syndrome after renal transplantation in children with focal segmental glomerulosclerosis. *Nephrol Dial Transplant* 2000;15(1):78-81.
- [43] Salomon R, Gagnadoux MF, Niaudet P. Intravenous cyclosporine therapy in recurrent nephrotic syndrome after renal transplantation in children. *Transplantation* 2003;75(6):810-4.
- [44] Raafat RH, Kalia A, Travis LB, Diven SC. High-dose oral cyclosporin therapy for recurrent focal segmental glomerulosclerosis in children. *Am J Kidney Dis* 2004;44(1):50-6.
- [45] Deegens JK, Andresdottir MB, Croockewit S, Wetzels JF. Plasma exchange improves graft survival in patients with recurrent focal glomerulosclerosis after renal transplant. *Transpl Int* 2004;17(3):151-7.
- [46] Valdivia P, Gonzalez Roncero F, Gentil MA, et al. Plasmapheresis for the prophylaxis and treatment of recurrent focal segmental glomerulosclerosis following renal transplant. *Transplant Proc* 2005;37(3):1473-4.
- [47] McCauley J, Shapiro R, Ellis D, Igdal H, Tzakis A, Starzl TE. Pilot trial of FK 506 in the management of steroid-resistant nephrotic syndrome. *Nephrol Dial Transplant* 1993;8(11):1286-90.
- [48] Kessler M, Champigneulle J, Hestin D, Frimat L, Renoult E. A renal allograft recipient with late recurrence of focal and segmental glomerulosclerosis after switching from cyclosporine to tacrolimus. *Transplantation* 1999;67(4):641-3.
- [49] Bussemaker E, Passauer J, Franz T, Gross P. Tryptophan immunoadsorption strongly reduces proteinuria in recurrent nephrotic syndrome. *Nephrol Dial Transplant* 2001;16(6):1270-2.
- [50] Canaud G, Zuber J, Sberro R, et al. Intensive and prolonged treatment of focal and segmental glomerulosclerosis recurrence in adult kidney transplant recipients: a pilot study. *Am J Transplant* 2009;9(5):1081-6.

- [51] Wada T, Pippin JW, Marshall CB, Griffin SV, Shankland SJ. Dexamethasone prevents podocyte apoptosis induced by puromycin aminonucleoside: role of p53 and Bcl-2-related family proteins. *J Am Soc Nephrol* 2005;16(9):2615-25.
- [52] Ransom RF, Lam NG, Hallett MA, Atkinson SJ, Smoyer WE. Glucocorticoids protect and enhance recovery of cultured murine podocytes via actin filament stabilization. *Kidney international* 2005;68(6):2473-83.
- [53] Faul C, Donnelly M, Merscher-Gomez S, et al. The actin cytoskeleton of kidney podocytes is a direct target of the antiproteinuric effect of cyclosporine A. *Nature medicine* 2008.
- [54] Canaud G, Dion D, Zuber J, et al. Recurrence of nephrotic syndrome after transplantation in a mixed population of children and adults: course of glomerular lesions and value of the Columbia classification of histological variants of focal and segmental glomerulosclerosis (FSGS). *Nephrol Dial Transplant* 2009.
- [55] Gohh RY, Yango AF, Morrissey PE, et al. Preemptive plasmapheresis and recurrence of FSGS in high-risk renal transplant recipients. *Am J Transplant* 2005;5(12):2907-12.
- [56] Gossmann J, Scheuermann EH, Porubsky S, Kachel HG, Geiger H, Hauser IA. Abrogation of nephrotic proteinuria by rituximab treatment in a renal transplant patient with relapsed focal segmental glomerulosclerosis. *Transpl Int* 2007;20(6):558-62.
- [57] Hristea D, Hadaya K, Marangon N, et al. Successful treatment of recurrent focal segmental glomerulosclerosis after kidney transplantation by plasmapheresis and rituximab. *Transpl Int* 2007;20(1):102-5.
- [58] Kamar N, Faguer S, Esposito L, et al. Treatment of focal segmental glomerular sclerosis with rituximab: 2 case reports. *Clinical nephrology* 2007;67(4):250-4.
- [59] Yabu JM, Ho B, Scandling JD, Vincenti F. Rituximab failed to improve nephrotic syndrome in renal transplant patients with recurrent focal segmental glomerulosclerosis. *Am J Transplant* 2008;8(1):222-7.
- [60] Leandro MJ, Cambridge G, Edwards JC, Ehrenstein MR, Isenberg DA. B-cell depletion in the treatment of patients with systemic lupus erythematosus: a longitudinal analysis of 24 patients. *Rheumatology (Oxford, England)* 2005;44(12):1542-5.
- [61] Le Berre L, Herve C, Buzelin F, Usal C, Soulillou JP, Dantal J. Renal macrophage activation and Th2 polarization precedes the development of nephrotic syndrome in Buffalo/Mna rats. *Kidney international* 2005;68(5):2079-90.
- [62] Bakr A, Shokeir M, El-Chenawi F, El-Husseni F, Abdel-Rahman A, El-Ashry R. Tumor necrosis factor-alpha production from mononuclear cells in nephrotic syndrome. *Pediatric nephrology (Berlin, Germany)* 2003;18(6):516-20.
- [63] Leroy S, Guignon V, Bruckner D, et al. Successful anti-TNFalpha treatment in a child with posttransplant recurrent focal segmental glomerulosclerosis. *Am J Transplant* 2009;9(4):858-61.
- [64] He JC, Lu TC, Fleet M, et al. Retinoic acid inhibits HIV-1-induced podocyte proliferation through the cAMP pathway. *J Am Soc Nephrol* 2007;18(1):93-102.
- [65] De Smet E, Rioux JP, Ammann H, Deziel C, Querin S. FSGS permeability factor-associated nephrotic syndrome: remission after oral galactose therapy. *Nephrol Dial Transplant* 2009;24(9):2938-40.



After the Kidney Transplant - The Patients and Their Allograft

Edited by Prof. Jorge Ortiz

ISBN 978-953-307-807-6

Hard cover, 386 pages

Publisher InTech

Published online 17, August, 2011

Published in print edition August, 2011

There are many obstacles in kidney transplantation. For the transplant team, there is the balance between immunosuppression to aid in the recipient's tolerance of the allograft and the infection risk of a suppressed immune system. These potential long term complications of kidney transplantation are relatively well known, but there are many other complications that patients and families do not consider when preparing themselves for a kidney transplant. Although the benefits of attempting a kidney transplant far outweigh downfalls of the long term sequelae, kidney transplantation is by no means a benign procedure. It is the hope of these authors that the reader will leave with a sense of understanding towards the kidney recipients.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Guillaume Canaud, Frank Martinez and Christophe Legendre (2011). Therapeutic Approach to Focal and Segmental Glomerulosclerosis (FSGS) Recurrence in Kidney Transplant Recipients, *After the Kidney Transplant - The Patients and Their Allograft*, Prof. Jorge Ortiz (Ed.), ISBN: 978-953-307-807-6, InTech, Available from: <http://www.intechopen.com/books/after-the-kidney-transplant-the-patients-and-their-allograft/therapeutic-approach-to-focal-and-segmental-glomerulosclerosis-fsgs-recurrence-in-kidney-transplant->

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen