

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Dithiocarbamate Toxicity - An Appraisal

Narayan C. Rath, Komal S. Rasaputra, Rohana Liyanage,
Gerry R. Huff and William E. Huff
*United States Department of Agriculture/Agricultural Research Service,
Department of Poultry Science,
Department of Chemistry and Biochemistry,
University of Arkansas,
Fayetteville,
USA*

1. Introduction

Dithiocarbamates (DTC) are organosulfur compounds represented by a general structure $(R_1R_2)N-(C=S)-SX$, where R can be substituted by an alkyl, alkylene, aryl, or similar other group, and X usually by a metal ion (Edwards, 1991; Kamrin, 1997; US EPA, 2001). Discovered in the 1930s, the DTC were first introduced as fungicides for commercial applications during World War II (Ware & Whitacre, 2004). Besides their wide use as fungicides for treatment of crops, vegetables, seeds, and ornamental plants, they are also used as accelerators in the rubber industry, animal repellants, and biocides in many household products (Edwards et al., 1991; Kamrin, 1997). Figure 1 shows examples of some common DTC pesticides. Thiram, disulfiram, ziram, and ferbam are analogous dialkyl DTC with differences in their R groups and the later two containing different metal ions between their S atoms. Pyrrolidinedithiocarbamate (PDTC) is a monomeric DTC which contains a five member ring attached to its N atom. It is a metabolic inhibitor used in cell physiological studies (Schreck et al., 1992; Cvek and Dvorak, 2007). In ethylene-bis-dithiocarbamates (EBDTC), the R groups of two DTC molecules form an ethylene bridge. The EBDTC are regarded as polymeric DTC because their metal ions can bind several molecules to form polymeric complexes. Some examples of EBDTC are zineb, maneb, and mancozeb which are used in preharvest agricultural applications. The DTC anions are highly reactive which can conjugate with other molecules containing SH groups and form metal chelates. The multisite interactions of DTC give them advantage to influence the biological activities of different proteins, enzymes, and exert toxic effects. Some of those modes of action of DTC compounds have been exploited for their use in clinical applications (Morrison et al., 2010). However, the extensive use of these chemicals in agriculture has raised concern for their effects as occupational and ecotoxicological hazards. Several reviews on the biological and toxicological effects of DTC summarize many studies in the field (Edwards et al., 1991; US EPA, 2001; Cvek and Dvorak, 2007). The objective of this review is to highlight some of the recent findings on the effects of dialkyl DTC and EBDTC with emphasis on studies of the avian system which has not been a focus of earlier literature.




Fig. 1. Structures of some representative dithiocarbamates, R1, R2= alkyl (CH₃, C₂H₅)
X= metal ion (Na⁺, Mn⁺², Zn⁺², Fe⁺³)

2. Metabolism and the toxic effects

The toxicological effects of DTC can occur from their absorption through skin exposure, ingestion, and inhalation. The lipophilic nature of DTC makes them suitable for their passage across the cell membrane. Metabolic studies with representative DTC (ex. thiram, disulfiram, mancozeb) have shown that these chemicals undergo detoxification through S-glucuronidation or biodegrade to different metabolites such as carbon disulfide (CS₂), thiourea, alkylamines, ethyleneamines, and several other biotransformation products (Edwards et al., 1991; US EPA, 2001). CS₂ is a general neuropathic agent and ethylenethiourea (ETU) which is a metabolite of EBDTC, has antithyroid and carcinogenic effects (Edwards et al., 1991; DeCaprio et al., 1992; Houeto et al., 1995; US EPA, 2001). Under physiological conditions most dialkyl DTC can reoxidize to form thiuram disulfide (Burkitt et al., 1998). Thus, the toxic effects of DTC can be due to the whole molecule and their decomposition products such as CS₂ and ETU. The intact DTC molecules exhibit both pro oxidant and antioxidant activities (Nobel et al., 1995; Liu et al., 1996; Orrenius et al., 1996; Burkitt et al., 1998; Wild & Mulcahy, 1999; Cereser et al., 2001). Whereas the disulfide bridges and the metal complexes contribute to their prooxidant effects, the SH contributes to their antioxidant effects (Orrenius et al., 1996; Elskens & Penninckx, 1997). Tissue or organ specific toxic effects of these chemicals may be due to the differential competency of their intracellular passage and binding to crucial structural and functional entities of the cells eventually leading to the metabolic disruptions, pathological changes, and cell death.

3. Molecular and cellular effects

The DTC compounds can form mixed disulfides with other molecules containing SH functions such as proteins, peptides and enzymes modulating their biological activities. The covalent modification of cysteine residues in the active sites can affect enzyme activities. As antioxidants, they react with hydroxyl radicals, peroxides, and superoxide ions, and inhibit their oxidative potential (Nobel et al., 1995; Liu et al., 1996). As prooxidants, DTC increase Cu catalyzed reactive oxygen species (ROS) formation and change the balance of reduced glutathione (GSH) to its oxidized form (GSSG) in favor of the later (Burkitt et al., 1998). GSH is a sulfhydryl containing tripeptide critical for protecting cells against oxidative stress. It is a major antioxidant in the body and an important regulator of cell proliferation, gene transcription, and apoptosis (Rana et al., 2002; Biswas & Rahman, 2009). GSH is necessary for detoxification of xenobiotics, carcinogens, and maintenance of immunity. Accumulation of oxidized form of glutathione (GSSG) leads to the activation of transcription factor nuclear factor kappa B (NF- κ B) stimulating stress and inflammatory response, and cell survival (Dellhale et al., 2004). The conversion of GSSG to GSH, catalyzed by glutathione reductase, is inhibited by DTC which also inactivate several different transcription factors principally, the NF- κ B and hypoxia inducible factor (Haddad, 2002 & 2003; Biswas & Rahman, 2009). PDTC is a popular inhibitor of transcription factor NF- κ B which modulates the expression of many enzymes and proteins including nitric oxide synthase, heat shock protein 70 (HSP70), and induces endoplasmic reticulum stress (Schreck et al., 1992; Cvek & Dvorak, 2007; Chen et al., 2010; Cotogni et al., 2010). Prevention of binding of NF- κ B to DNA induces apoptosis. DTC inhibit proteasome dependent protein degradation (Wang et al, 2006, 2011; Lovborg et al, 2006; Daniel et al, 2007; Chou et al, 2008) and promote peptide amidation (Mains et al., 1986). Thiram increases oxidative stress and induces formation of lipid peroxides, protein carbonyls, and stimulates changes in membrane potential of cells leading to ion influx inducing cell death (Erl et al., 2000; Sook Han et al., 2003; Grosicka et al., 2005). Similarly, disulfiram also induces oxidative stress that changes mitochondrial permeability leading to mitochondrial injury (Balakirev and Zimmer, 2001). A number of enzymes are inhibited by DTC which include cyclooxygenase, (Lee et al., 2002), heme oxygenase (Kushida et al, 2002), cytochrome P450, superoxide dismutase, glutathione reductase, and caspase (Dalvi et al., 2002; Cvek & Dvorak, 2007; Seefeldt et al., 2009). The superoxide dismutase inhibitory activity of thiram and disulfiram is implicated in their anti-angiogenic effects (Marikovskiy et al., 2002; Shian et al., 2003). The aldehyde dehydrogenase inhibitory activity of disulfiram is the basis of its therapeutic efficacy against alcoholism (Edwards et al., 1991; Cvek & Dvorak, 2007). Disulfiram also suppresses matrix metalloproteinase (MMP) expression in osteosarcoma cells through modulation of NF- κ B and activator protein-1, and possibly its metal chelating properties (Cho et al., 2007).

3.1 Neuropathic effects

Peripheral neuropathy induced by DTC is a major toxic effect which has been reported in humans and animals (Frisoni & Di Monda, 1989). Many DTC pesticides including several dialkyl dithiocarbamates and EBDTC are implicated in inducing Parkinson's-like neuropathy. The ability of DTC to inhibit acetylcholine esterase, an enzyme responsible for degradation of the neurotransmitter acetylcholine, was considered to cause neuropathy (Edwards et al., 1991), but later studies did not substantiate this mode of action. However, Viviani et al., (2008) using adrenomedullary PC12 cells showed propineb, an EBTC, to induce acetylcholine release which is mediated through depolymerization of cytoskeletal

actins. Since most DTC compounds can metabolize to CS₂, their neuropathic effects were thought to be mediated by this metabolite alone. Johnson et al. (1998) showed cross linking of neurofilament proteins induced by CS₂ as a mechanism for its axonopathic and neurotoxic effects. However, metabolic studies with different DTC have not supported the role of CS₂ as the sole mechanism for their neurotoxic effects (SAP report, 2001). Stimulation of non selective cation channels by thiram, ziram, and maneb cause the influx of Ca⁺⁺ and Cu⁺⁺ into mitochondria increasing oxidative stress which induce apoptosis of PC12 cells and dopaminergic neuronal damage (Sook Han et al., 2003; Barlow et al., 2005). DTC metal complexes induce dopamine oxidation and produce intraneuronal oxidative stress leading to neuronal damage (Fitsanakis et al., 2002). Since DTC chelate heavy metals such as Cu, Zn, and Fe, leading to their intraneuronal accumulations, these metals have been implicated in promoting lipid peroxidation, oxidative stress, and enzyme inhibitions causing neurotoxic effects (Nobel et al., 1995; Valentine et al., 2009; Viquez et al., 2009; Viola-Rhenals et al., 2007). Increased production of reactive oxygen species by the actions of mancozeb and zineb is also implicated in their neuronal toxicities (Domico et al., 2007). Maneb, an EBDTC containing Mn⁺², was found to induce nitric oxide production, lipid peroxidation, and cause Parkinson's like disease syndrome in mice (Gupta et al., 2010). Mancozeb, thiram, and disulfiram cause membrane potential changes and impair ATP dependent glutamate uptake into the synaptic vesicles and prevent binding of glutamate to its receptors resulting in excitotoxic effects in the brain (Nagendra et al., 1997; Vaccari et al., 1999). Ubiquitin proteasome pathway maintains the balance of cellular proteins through their degradation since abnormal accumulation of protein can interfere with cell functions (Myung et al., 2001). Both disulfiram and ziram inhibit ubiquitin proteasomal pathways causing dopaminergic cell damage (Lovborg et al., 2006; Chou et al., 2008). Disulfiram also reduces the activity of brain enzyme peptidoglycine-5 hydroxylating monooxygenase and alpha-melanocyte stimulating hormones affecting behavioral changes in rats (Rahman et al., 1997).

3.2 Reproductive and endocrine disruptive effects

Chemicals which interfere with endocrine functions altering the synthesis, metabolism, and secretion of hormones, or their target organ effects, are called endocrine disruptors (Diamanti-Kandarakis et al., 2009). There are several reports suggesting the endocrine disruptive actions of DTC. Studies by Stoker et al (1993; 2003) showed that thiram induces ovulatory delay and affects fecundity in rats. Some of these effects of DTC are related to the interference of enzymes involved in the synthesis of catecholamines which regulate neuroendocrine functions (Stoker et al., 1993; Goldman et al., 1994). Thiram inhibits spermatogenesis in rats (Mishra et al., 1998). Mancozeb affects ovarian function and disrupts the estrous cycle, inducing infertility in rats (Cooper et al., 1999; Cecconi et al., 2007). The hypothyroid and antithyroid effects of zineb and mancozeb are associated with their metabolite ethinylthiourea (ETU) (Houeto et al., 1995; US EPA, 2001; Panganiban et al., 2004; Axelstad et al., 2011). Both thiram and disulfiram inhibit 11 β hydroxyl steroid dehydrogenase 2, an enzyme that catalyzes conversion of hormonally active glucocorticoids, cortisol and corticosterone, to their inactive metabolites, and interfere with binding to their receptors (Atansov et al., 2003; Garbrecht et al., 2006).

3.3 Immunomodulatory effects

The immunomodulatory effects of DTC can be largely related to their ability to prevent activation of transcription factors and other signaling mechanisms. Lipopolysaccharide

induced tumor necrosis factor alpha production by promyelocytic THP-1 cells is inhibited by mancozeb (Corsini et al., 2006). Ziram interferes with the lytic function of natural killer cells through modification of their cell surface proteins such as CD16 which is necessary for their binding to target cells (Taylor & Whalen, 2009) and potentiates Concanavalin A induced interferon- γ and interleukin-6 production by the vascular lymph node cells (De Jong et al., 2002). In U937 lymphoma cells, ziram produces its toxic effects by activating intracellular caspase-3 enzyme and mitochondrial cytochrome c release which lead to their apoptosis (Li et al., 2010). However, with respect to cellular immunity, studies have shown DTC induce activation of T cells, natural killer (NK) cells, and increase immunoglobulin secretion by B cells (Corsini et al., 2006, 2008). Thiram induces lymphocyte sensitization, hypersensitivity, and allergic dermatitis (Saunders & Watkins, 2001). Although the mechanism of allergic dermatitis induced by thiram is not well understood, the involvement of T cells is likely. DTC can act as haptens which on conjugating to proteins may induce allergic hypersensitivity. Cytofluorometric study by Lombardi et al. (1991) showed increased splenic population of T cytotoxic/suppressor cells induced by dimethyl and diethyl DTC. The effects of different DTC on immunity needs better understanding.

3.4 Carcinogenic and teratogenic effects

The EBDTC in general, are considered to be carcinogenic because of their metabolite ETU that produces thyroid and pituitary tumors (Houeto et al., 1995). Steenland et al. (1997) showed the genotoxic effects of mancozeb indicated by increased chromosomal translocations and sister chromatid exchange in the blood cells of workers exposed to it. In vitro studies with zineb on human lymphocytes and CHO cells showed it to induce DNA strand breaks suggesting its carcinogenic potential in the event that the affected cells survive and propagate (Soloneski et al., 2002; 2003; Gonzalez et al., 2003). Calviello et al. (2006) showed DNA single strand breaks in rat fibroblasts exposed to mancozeb. DNA breaks and chromosomal aberration induced by thiram in CHO cells was reported by Mosseso et al. (1994), but in vivo tests employing different doses of ferbam, which is similar to thiram, showed no significant induction of aneuploidy (Shanthi & Krishnamoorthy, 2002). Although the recovery of DTC damaged cells and their survival is important for carcinogenicity, there is meagre evidence in its favor (Hasegawa et al., 1988). Studies on the effects of DTC on developing rat embryos show that these agents induce cleft palate, wavy rib formation, and long bone distortions (Roll, 1971). Several recent studies have shown sodium metam, thiram, and disulfiram caused notochord distortions, and craniofacial abnormalities in zebra fish embryos (Haendel et al., 2004; Tilton, et al., 2006; Teraoka et al., 2006; van Boxtel et al., 2010). Some effects of these chemicals on craniofacial malformation are attributed to their down-regulating effects on genes related to transforming growth factor beta-1 (TGF- β 1) which plays an important role in skeletal morphogenesis. Inhibition of lysyl oxidase, a Cu⁺⁺ dependent enzyme essential for collagen cross linking, by the chelating actions of DTC is also suggested as another possible mechanism in the induction of craniofacial abnormalities (van Boxtel et al., 2010).

4. DTC effects on avian systems

The major bulk of research on DTC has been carried out using mammalian models or cells. But their effect on avian growth plate cartilage is noteworthy because in relatively small doses and short exposure time, certain DTC can induce cartilage defects in growing birds

which render them lame (Vargas et al., 1983). The teratogenic and embryo toxic effects of DTC on avian system was recognized as early as 1955 when researchers noticed exposure to thiram caused leg problems in poultry (Waibel et al., 1955). These effects of thiram also were observed in later years (Page, 1975; Guitart et al., 1996). With the identification of tibial dyschondroplasia (TD), a defect of endochondral bone formation in young poultry by Leach & Nesheim (1965), correlations showed that DTC caused poultry leg problems (Vargas et al., 1983). Subsequent studies by different investigators showed that both dimethyl and diethyl DTC caused TD in post hatch poultry (Veltmann et al., 1985; Edwards, 1987; Orth & Cook, 1994; Rath et al., 2004). With TD, the proximal growth plates of the tibia and tibio-tarsal bones fail to ossify leading to the retention of unresolved cartilage. Feeding post hatch chickens diets containing thiram 50-100 mg /kg feed for a day or two is sufficient to induce TD (Rath et al., 2004; 2005; 2007b) (Figure 2). The incidence and severity of the disease is related to the age of the chicks; during the early phase of growth when the bones are fast growing, the effects are more severe. Subsequent studies showed that the dimeric and trimeric analogs such as thiram, disulfiram, ferbam, and ziram would induce this defect, where as the monomeric DTC such as potassium dimethyl dithiocarbamate, sodium metam, or PDTC were ineffective or less potent in similar concentrations (Rath et al., 2004; 2007b). The induction of TD by thiram is dose dependent. Feeding a thiram containing diet was more effective in inducing tibial dyschondroplasia than their subcutaneous administration (unpublished observation). Thiram reduces feed intake resulting in body weight loss but it does not stop longitudinal growth of bones. Whether, feed intake is affected through the influence of thiram on hypothalamic mechanisms is not known but based on its demonstrated neuroendocrine effects (Stoker et al., 1993), it may be a possibility. Thiram causes an elevation in serum corticosterone level (Rath et al., 2004) which can be related to its inhibitory effect on 11β -hydroxysteroid dehydrogenase-2 that mediates the conversion of corticosterone to its inactive metabolite 11-hydroxycorticosterone (Atanasov et al., 2003). Endochondral bone formation is a complex process which involves an orderly transition of cartilage from proliferative to hypertrophic state when they undergo chondrolytic degeneration and replaced by osteoblast (Reddi & Anderson, 1976; Burdan et al., 2009). Angiogenesis and neovascularization of growth plate is essential for bone formation. Thiram exerts a high level of toxicity on endothelial cells inducing death of capillary vessels in the growth plate and interferes with the hypertrophic process resulting in premature death of chondrocytes. Apoptosis of growth plate chondrocytes and blood vessels are evident by histochemical staining and the assessment of DNA fragmentation (Rath et al., 2005) (Figure 3). Treatment with thiram reduces the concentrations of enzymes and proteins associated with bone development which may be related to the cell death in growth plate (Rath et al., 2005). Both in chickens and turkeys, thiram interferes with growth plate modeling and angiogenesis by interfering with matrix metalloproteinases (MMP) production (Hasky-Negev et al., 2008; Dan et al., 2009). Vascular endothelial growth factor (VEGF) is a regulator of angiogenesis that acts through its receptors. Thiram down-regulates the expression of genes for VEGF receptor and Bcl-2, an antiapoptotic protein, in the growth plate (Rath et al., 2007a). Tian et al. (2009) showed the down-regulation of matrilin, and MMP-13 genes in growth plates of chickens that were treated with thiram. Expressions of these genes are important in growth plate maturation. Consistent with literature on the action of DTC in different systems, there was also a decrease in glutathione levels in growth plate cartilage of thiram-treated chickens (Rath et al., 2005). Comparative proteomics of growth plate tissue extracts showed decreases in several proteins in thiram-fed chickens most of which were

associated with energy metabolism, signal transduction, and secretory functions (Rasaputra et al., 2010). Down-regulation of those proteins may be responsible for chondrocyte death. The differential effect of thiram on hypertrophic chondrocytes may be related to the developmental transition of cells when they become prone to the toxic effects of DTC. Hypertrophy of chondrocytes is necessary for lengthening of bone which results in a significant change in cell volume (Farnum et al., 2002). Increases in cell volume occur from increased protein synthesis and influx of inorganic solutes, and osmolytes. The latter processes are affected by the changes in membrane permeability and increased ion channel activities. Recently, Bush




Fig. 2. Proximal tibial growth plates (arrow) of 14 day-old chickens fed either (a) a control diet or (b) a diet containing 100 mg thiram/kg feed for 48 hours between days 8 and 9 showing tibial dyschondroplasia evident by an irregular broadening of growth plate.




Fig. 3. Histology and histochemistry of hypertrophic zone chondrocytes of (a) normal tibial growth plate (b) thiram-induced dyschondroplastic growth plate with diminished chondrocyte volumes, pyknotic nuclei, and matrix rarefaction. (c&d) Dyschondroplastic zone chondrocytes showing terminal deoxynucleotidyl transferase mediated fluorescein dUTP nick end labeling (TUNEL) of apoptotic cells with yellow to green fluorescence and healthy chondrocytes with red fluorescence due to propidium iodide staining. (d) A dead capillary vessel surrounded by both healthy and apoptotic chondrocytes (adapted from Rath et al., 2005).

et al. (2010) showed an increased expression of Na⁺ K⁺ Cl⁻ cotransporter protein (NKCC) in hypertrophic chondrocytes. Pucci et al. (2007) also observed changes in mitochondrial membrane potentials of hypertrophic chondrocytes that permeate influx of cationic molecules. It is possible that changes in chondrocyte membrane permeability during hypertrophy facilitate higher influx of thiram into the cells inducing metabolic inhibitions, oxidative stress, and apoptosis. Thiram also can inhibit other molecular changes associated with the ossification process. Using microarray analysis of chicken growth plate, Horvat-Gordon et al. (2010) showed high expression of several genes associated with angiogenesis and oxido-reductive metabolism in hypertrophic chondrocytes. The proteins encoded by these genes such as the transferrin, matrix metalloproteinases, aldehyde dehydrogenase, lysyl oxidase, and superoxide dismutase contain metal ions that are prone to chelation by DTC which can modulate their activities and cause metabolic dysregulations. Marikovsky et al. (2002) have shown that both thiram and disulfiram interfere with angiogenesis through inhibition of superoxide dismutase.

4.1 Effects on chondrocyte culture

Proteomics is a powerful tool to identify biomarkers and understand the mechanisms of action of toxicants (Kennedy, 2002). To find whether thiram induces peptide and protein changes, the growth plate chondrocytes in culture were treated with sub lethal concentrations of thiram for 48 h. The viability of the cells were determined by monitoring the release of lactate dehydrogenase (LDH) into the culture medium as an indicator of cell damage (Rath et al., 1995) which showed no significant change at 48 h. The peptide profiles of these chondrocyte extracts were examined by means of matrix assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF-MS) in the *m/z* range of 1,000 to 7,000, and compared between control and thiram treated cells. Differential expression of the peptides was determined using statistical algorithms and principal component analysis by the use of ClinproTool™ software (Bruker Daltonics, Germany). Comparing approximately 50 spectral peaks, 4 showed quantitative differences in thiram treated chondrocytes with 2 peptides corresponding to *m/z* 3004.5 and 3310, elevated, and 2 corresponding to *m/z* 1778.9, 2556.3, decreased (Figures 4 & 5) (Rasaputra et al., unpublished). Although the functional significance of the changes in these peptides is currently unknown such information can be useful to identify toxicity associated peptide biomarkers. Similarly, comparing the protein profiles of control and thiram treated chondrocytes by two dimensional gel electrophoresis, several proteins were found to be decreased by thiram treatment, particularly a heat shock protein HSP70 was significantly down-regulated (Rasaputra et al., unpublished). HSP70 is necessary for protein folding and protects the cells from oxidative stress and apoptosis (Beere et al., 2000; Mosser et al., 2000; Guzhova & Margulis, 2006). Its chondroprotective effect has been shown in mammalian models (Otsuka et al., 1996; Etienne et al., 2008). It is possible that the decrement in the levels of HSP70 contributes to the loss of chondrocyte viability. In conclusion, the effect of DTC on growth plate development provides a good experimental model to study the toxicology of these compounds in skeletal system.

5. Conclusion

From the preceding discussion, it is evident that the DTC modify cellular metabolism by their direct interactions with different molecules such as signaling proteins, peptides, and




Fig. 4. The MALDI-TOF mass spectral profiles of control and thiram treated chondrocyte extracts showing peptide peaks in the m/z 1500-4000.




Fig. 5. (a) Principal component analysis of mass spectrum showing similarities and differences in peptide profiles of control (red) and thiram treated chondrocytes (green), and (b) profiles of the differentially expressed peptides ($P \leq 0.001$).

enzymes, and influence the oxido-reductive metabolism of the cells. Their metal chelating properties additionally, contribute to their prooxidative effects. The cells exposed to DTC experience increased oxidative stress and metabolic dysregulations leading to tissue damage, and apoptosis. The disparate vulnerability of tissues to the toxic effects of different DTC may be due to the differences in their membrane permeability and cellular constituents

interacting with these chemicals. Dividing and differentiating cells may be more susceptible to the toxic effects of DTC. Although some of their metabolites such as carbon disulfide and ethynylurea contribute to certain organ specific pathologies, it is most likely that the whole molecules are responsible for their acute toxicities. There is little evidence of the ecotoxicological hazards of these chemicals. High propensity of dithiocarbamates to modulate signal transduction mechanisms, provide the promise for their usefulness in various pharmaceutical applications.

6. References

- Atanasov, A. G., Tam, S., Rocken, J. M., Baker, M. E., & Odermatt, A. (2003). Inhibition of 11 beta-hydroxysteroid dehydrogenase type 2 by dithiocarbamates. *Biochemical and Biophysical Research Communications*. Vol. 308, pp. 257-262
- Axelstad, M., Boberg, J., Nellemann, C., Kiersgaard, M., Jacobsen, P. R., Christiansen, S., Hougaard, K. S., & Hass, U. (2011). Exposure to the widely used fungicide Mancozeb causes thyroid hormone disruption in rat dams but no behavioral effects in the offspring. *Toxicological Sciences*. PMID: 21266532
- Balakirev, M. Y., & Zimmer, G. (2001). Mitochondrial injury by disulfiram: two different mechanisms of the mitochondrial permeability transition. *Chemico-Biological Interactions*. Vol. 138, pp. 299-311
- Barlow, B. K., Lee, D. W., Cory-Slechta, D. A., & Opanashuk, L. A. (2005). Modulation of antioxidant defense systems by the environmental pesticide maneb in dopaminergic cells. *Neurotoxicology*. Vol. 26, pp. 63-75
- Beere, H. M., Wolf, B. B., Cain, K., Mosser, D. D., Mahboubi, A., Kuwana, T., Tailor, P., Morimoto, R. I., Cohen, G. M., & Green, D. R. (2000). Heat-shock protein 70 inhibits apoptosis by preventing recruitment of procaspase-9 to the Apaf-1 apoptosome. *Nature Cell Biology*. Vol. 2, pp. 469-475
- Biswas, S. K., & Rahman, I. (2009). Environmental toxicity, redox signaling and lung inflammation: the role of glutathione. *Molecular Aspects of Medicine*. Vol. 30, pp. 60-76
- Burdan, F., Szumilo, J., Korobowicz, A., Farooque, R., Patel, S., Patel, A., Dave, A., Szumilo, M., Solecki, M., Klepacz, R., & Dudka, J. (2009). Morphology and physiology of the epiphyseal growth plate. *Folia Histochemica Et Cytobiologica*. Vol. 47, pp. 5-16
- Burkitt, M. J., Bishop, H. S., Milne, L., Tsang, S. Y., Provan, G. J., Nobel, C. S., Orrenius, S., & Slater, A. F. (1998). Dithiocarbamate toxicity toward thymocytes involves their copper-catalyzed conversion to thiuram disulfides, which oxidize glutathione in a redox cycle without the release of reactive oxygen species. *Archives of Biochemistry and Biophysics*. Vol. 353, pp. 73-84
- Bush, P. G., Pritchard, M., Loqman, M. Y., Damron, T. A., & Hall, A. C. (2010). A key role for membrane transporter NKCC1 in mediating chondrocyte volume increase in the mammalian growth plate. *Journal of Bone and Mineral Research*. Vol. 25, pp. 1594-1603
- Calviello, G., Piccioni, E., Boninsegna, A., Tedesco, B., Maggiano, N., Serini, S., Wolf, F. I., & Palozza, P. (2006). DNA damage and apoptosis induction by the pesticide Mancozeb in rat cells: involvement of the oxidative mechanism. *Toxicology and Applied Pharmacology*. Vol. 211, pp. 87-96

- Cecconi, S., Paro, R., Rossi, G., & Macchiarelli, G. (2007). The effects of the endocrine disruptors dithiocarbamates on the mammalian ovary with particular regard to mancozeb. *Current Pharmaceutical Design*. Vol. 13, pp. 2989-3004
- Cereser, C., Boget, S., Parvaz, P., & Revol, A. (2001). Thiram-induced cytotoxicity is accompanied by a rapid and drastic oxidation of reduced glutathione with consecutive lipid peroxidation and cell death. *Toxicology*. Vol. 163, pp. 153-162
- Chen, Y. W., Chen, K. L., Chen, C. H., Wu, H. C., Su, C. C., Wu, C. C., Way, T. D., Hung, D. Z., Yen, C. C., Yang, Y. T., & Lu, T. H. (2010). Pyrrolidine dithiocarbamate (PDTC)/Cu complex induces lung epithelial cell apoptosis through mitochondria and ER-stress pathways. *Toxicology Letters*. Vol. 199, pp. 333-340
- Cho, H. J., Lee, T. S., Park, J. B., Park, K. K., Choe, J. Y., Sin, D. I., Park, Y. Y., Moon, Y. S., Lee, K. G., Yeo, J. H., Han, S. M., Cho, Y. S., Choi, M. R., Park, N. G., Lee, Y. S., & Chang, Y. C. (2007). Disulfiram suppresses invasive ability of osteosarcoma cells via the inhibition of MMP-2 and MMP-9 expression. *Journal of Biochemistry and Molecular Biology*. Vol. 40, pp. 1069-1076
- Chou, A. P., Maidment, N., Klintonberg, R., Casida, J. E., Li, S., Fitzmaurice, A. G., Fernagut, P. O., Mortazavi, F., Chesselet, M. F., & Bronstein, J. M. (2008). Ziram causes dopaminergic cell damage by inhibiting E1 ligase of the proteasome. *The Journal of Biological Chemistry*. Vol. 283, pp. 34696-34703
- Cooper, R. L., Goldman, J. M., & Stoker, T. E. (1999). Neuroendocrine and reproductive effects of contemporary-use pesticides. *Toxicology and Industrial Health*. Vol. 15, pp. 26-36
- Corsini, E., Liesivuori, J., Vergieva, T., Van Loveren, H., & Colosio, C. (2008). Effects of pesticide exposure on the human immune system. *Human & Experimental Toxicology*. Vol. 27, pp. 671-680
- Corsini, E., Viviani, B., Birindelli, S., Gilardi, F., Torri, A., Codeca, I., Lucchi, L., Bartesaghi, S., Galli, C. L., Marinovich, M., & Colosio, C. (2006). Molecular mechanisms underlying mancozeb-induced inhibition of TNF-alpha production. *Toxicology and Applied Pharmacology*. Vol. 212, pp. 89-98
- Cotogni, P., Bini, R., Trombetta, A., & Olivero, G. (2010). Pyrrolidine Dithiocarbamate Modulates HSP70, iNOS, and Apoptosis during Hemorrhagic Shock Resuscitation in Rats. *Journal of Investigative Surgery*. Vol. 23, pp. 295-302
- Cvek, B., & Dvorak, Z. (2007). Targeting of nuclear factor-kappaB and proteasome by dithiocarbamate complexes with metals. *Current Pharmaceutical Design*. Vol. 13, pp. 3155-3167
- Dalvi, P. S., Wilder-Ofie, T., Mares, B., Lane, C., Dalvi, R. R., & Billups, L. H. (2002). Effect of cytochrome P450 inducers on the metabolism and toxicity of thiram in rats. *Veterinary and Human Toxicology*. Vol. 44, pp. 331-333
- Dan, H., Simsa-Maziel, S., Hisdai, A., Sela-Donenfeld, D., & Monsonego Ornan, E. (2009). Expression of matrix metalloproteinases during impairment and recovery of the avian growth plate. *Journal of Animal Science*. Vol. 87, pp. 3544-3555
- Daniel, K. G., Chen, D., Yan, B., & Dou, Q. P. (2007). Copper-binding compounds as proteasome inhibitors and apoptosis inducers in human cancer. *Frontiers in Bioscience*. Vol. 12, pp. 135-144
- De Jong, W. H., Tentij, M., Spiekstra, S. W., Vandebriel, R. J., & Van Loveren, H. (2002). Determination of the sensitising activity of the rubber contact sensitizers TMTD,

- ZDMC, MBT and DEA in a modified local lymph node assay and the effect of sodium dodecyl sulfate pretreatment on local lymph node responses. *Toxicology*. Vol. 176, pp. 123-134
- DeCaprio, A. P., Spink, D. C., Chen, X., Fowke, J. H., Zhu, M., & Bank, S. (1992). Characterization of isothiocyanates, thioureas, and other lysine adduction products in carbon disulfide-treated peptides and protein. *Chemical Research in Toxicology*. Vol. 5, pp. 496-504
- Delhalle, S., Blasius, R., Dicato, M., & Diederich, M. (2004). A beginner's guide to NF-kappaB signaling pathways. *Annals of the New York Academy of Sciences*. Vol. 1030, pp. 1-13
- Diamanti-Kandarakis, E., Bourguignon, J. P., Giudice, L. C., Hauser, R., Prins, G. S., Soto, A. M., Zoeller, R. T., & Gore, A. C. (2009). Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocrine Reviews*. Vol. 30, pp. 293-342
- Domico, L. M., Cooper, K. R., Bernard, L. P., & Zeevalk, G. D. (2007). Reactive oxygen species generation by the ethylene-bis-dithiocarbamate (EBDC) fungicide mancozeb and its contribution to neuronal toxicity in mesencephalic cells. *Neurotoxicology*. Vol. 28, pp. 1079-1091
- Edwards, I.R., D.H. Ferry, and W.A. Temple. 1991. Fungicides and related compounds. In: *Handbook of Pesticide Toxicology, v3, W.J. Hayes, Jr. and E. R. Laws, Jr. editors Academic Press, San Diego, CA. Pp 1409-1470*
- Edwards, H. M., Jr. (1987). Effects of thiuram, disulfiram and a trace element mixture on the incidence of tibial dyschondroplasia in chickens. *The Journal of Nutrition*. Vol. 117, pp. 964-969
- Elskens, M. T., & Penninckx, M. J. (1997). Thiram and dimethyldithiocarbamic acid interconversion in *Saccharomyces cerevisiae*: a possible metabolic pathway under the control of the glutathione redox cycle. *Applied and Environmental Microbiology*. Vol. 63, pp. 2857-2862
- Erl, W., Weber, C., & Hansson, G. K. (2000). Pyrrolidine dithiocarbamate-induced apoptosis depends on cell type, density, and the presence of Cu(2+) and Zn(2+). *American Journal of Physiology. Cell Physiology*. Vol. 278, pp. C1116-25
- Etienne, S., Gaborit, N., Henrionnet, C., Pinzano, A., Galois, L., Netter, P., Gillet, P., & Grossin, L. (2008). Local induction of heat shock protein 70 (Hsp70) by proteasome inhibition confers chondroprotection during surgically induced osteoarthritis in the rat knee. *Bio-Medical Materials and Engineering*. Vol. 18, pp. 253-260
- Farnum, C. E., Lee, R., O'Hara, K., & Urban, J. P. (2002). Volume increase in growth plate chondrocytes during hypertrophy: the contribution of organic osmolytes. *Bone*. Vol. 30, pp. 574-581
- Fitsanakis, V. A., Amarnath, V., Moore, J. T., Montine, K. S., Zhang, J., & Montine, T. J. (2002). Catalysis of catechol oxidation by metal-dithiocarbamate complexes in pesticides. *Free Radical Biology & Medicine*. Vol. 33, pp. 1714-1723
- Frisoni, G. B., & Di Monda, V. (1989). Disulfiram neuropathy: a review (1971-1988) and report of a case. *Alcohol and Alcoholism*. Vol. 24, pp. 429-437
- Garbrecht, M. R., Krozowski, Z. S., Snyder, J. M., & Schmidt, T. J. (2006). Reduction of glucocorticoid receptor ligand binding by the 11-beta hydroxysteroid dehydrogenase type 2 inhibitor, Thiram. *Steroids*. Vol. 71, pp. 895-901
- Goldman, J. M., Stoker, T. E., Cooper, R. L., McElroy, W. K., & Hein, J. F. (1994). Blockade of ovulation in the rat by the fungicide sodium N-methyldithiocarbamate:

- relationship between effects on the luteinizing hormone surge and alterations in hypothalamic catecholamines. *Neurotoxicology and Teratology*. Vol. 16, pp. 257-268
- Gonzalez, M., Soloneski, S., Reigosa, M. A., & Larramendy, M. L. (2003). Effect of dithiocarbamate pesticide zineb and its commercial formulation, azzurro. IV. DNA damage and repair kinetics assessed by single cell gel electrophoresis (SCGE) assay on Chinese hamster ovary (CHO) cells. *Mutation Research*. Vol. 534, pp. 145-154
- Grosicka, E., Sadurska, B., Szumilo, M., Grzela, T., Lazarczyk, P., Niderla-Bielinska, J., & Rahden-Staron, I. (2005). Effect of glutathione depletion on apoptosis induced by thiram in Chinese hamster fibroblasts. *International Immunopharmacology*. Vol. 5, pp. 1945-1956
- Guitart, R., Mateo, R., Gutierrez, J. M., & To-Figureueras, J. (1996). An outbreak of thiram poisoning on Spanish poultry farms. *Veterinary and Human Toxicology*. Vol. 38, pp. 287-288
- Gupta, S. P., Patel, S., Yadav, S., Singh, A. K., Singh, S., & Singh, M. P. (2010). Involvement of nitric oxide in maneb- and paraquat-induced Parkinson's disease phenotype in mouse: is there any link with lipid peroxidation? *Neurochemical Research*. Vol. 35, pp. 1206-1213
- Guzhova, I., & Margulis, B. (2006). Hsp70 chaperone as a survival factor in cell pathology. *International Review of Cytology*. Vol. 254, pp. 101-149
- Haddad, J. J., (2002). Antioxidant and prooxidant mechanisms in the regulation of redox(y)-sensitive transcription factors. *Cellular Signalling*. Vol. 14, pp. 879-897
- Haddad, J. J., (2003). Science review: redox and oxygen-sensitive transcription factors in the regulation of oxidant-mediated lung injury: role for hypoxia-inducible factor-1 α . *Critical Care*. Vol. 7, pp. 47-54
- Haendel, M. A., Tilton, F., Bailey, G. S., & Tanguay, R. L. (2004). Developmental toxicity of the dithiocarbamate pesticide sodium metam in zebrafish. *Toxicological Sciences*. Vol. 81, pp. 390-400
- Hasegawa, R., Takahashi, M., Furukawa, F., Toyoda, K., Sato, H., Jang, J. J., & Hayashi, Y. (1988). Carcinogenicity study of tetramethylthiuram disulfide (thiram) in F344 rats. *Toxicology*. Vol. 51, pp. 155-165
- Hasky-Negev, M., Simsa, S., Tong, A., Genina, O., & Monsonego Ornan, E. (2008). Expression of matrix metalloproteinases during vascularization and ossification of normal and impaired avian growth plate. *Journal of Animal Science*. Vol. 86, pp. 1306-1315
- Horvat-Gordon, M., Praul, C. A., Ramachandran, R., Bartell, P. A., & Leach, R. M., Jr. (2010). Use of microarray analysis to study gene expression in the avian epiphyseal growth plate. *Comparative Biochemistry and Physiology. Part D, Genomics & Proteomics*. Vol. 5, pp. 12-23
- Houeto, P., Bindoula, G., & Hoffman, J. R. (1995). Ethylenebisdithiocarbamates and ethylenethiourea: possible human health hazards. *Environmental Health Perspectives*. Vol. 103, pp. 568-573
- Johnson, D. J., Graham, D. G., Amarnath, V., Amarnath, K., & Valentine, W. M. (1998). Release of carbon disulfide is a contributing mechanism in the axonopathy produced by N,N-diethyldithiocarbamate. *Toxicology and Applied Pharmacology*. Vol. 148, pp. 288-296

- Kamrin M. A., (1997). Pesticide Profile, Toxicity, environmental impact and fate. CRC Press, Boca Raton, FL.
- Kennedy, S., (2002). The role of proteomics in toxicology: identification of biomarkers of toxicity by protein expression analysis. *Biomarkers : Biochemical Indicators of Exposure, Response, and Susceptibility to Chemicals*. Vol. 7, pp. 269-290
- Kushida, T., Quan, S., Yang, L., Ikehara, S., Kappas, A., & Abraham, N. G. (2002). A significant role for the heme oxygenase-1 gene in endothelial cell cycle progression. *Biochemical and Biophysical Research Communications*. Vol. 291, pp. 68-75
- Leach, R. M., Jr & Nesheim, M. C. (1965). Nutritional genetic and morphological studies of an abnormal cartilage formation in young chicks. *The Journal of Nutrition*. Vol. 86, pp. 236-244
- Lee, J. E., Kim, K. M., Cho, J. W., Suh, S. I., Suh, M. H., Kwon, T. K., Park, J. W., Bae, J. H., Song, D. K., Cho, C. H., Bae, I., & Baek, W. K. (2002). Pyrrolidine dithiocarbamate induces cyclooxygenase-2 expression in NIH 3T3 fibroblast cells. *Biochemical and Biophysical Research Communications*. Vol. 298, pp. 230-234
- Li, Q., Kobayashi, M., & Kawada, T. (2010). Ziram induces apoptosis and necrosis in human immune cells. *Archives of Toxicology*. DOI 10.1007/s00204-010-0586-9
- Liu, J., Shigenaga, M. K., Yan, L. J., Mori, A., & Ames, B. N. (1996). Antioxidant activity of diethyldithiocarbamate. *Free Radical Research*. Vol. 24, pp. 461-472
- Lombardi, P., Fournier, M., Bernier, J., Mansour, S., Neveu, P., & Krzystyniak, K. (1991). Evaluation of the immunomodulatory potential of diethyl dithiocarbamate derivatives. *International Journal of Immunopharmacology*. Vol. 13, pp. 1073-1084
- Lovborg, H., Oberg, F., Rickardson, L., Gullbo, J., Nygren, P., & Larsson, R. (2006). Inhibition of proteasome activity, nuclear factor-KappaB translocation and cell survival by the antialcoholism drug disulfiram. *International Journal of Cancer*. Vol. 118, pp. 1577-1580
- Mains, R. E., Park, L. P., & Eipper, B. A. (1986). Inhibition of peptide amidation by disulfiram and diethyldithiocarbamate. *The Journal of Biological Chemistry*. Vol. 261, pp. 11938-11941
- Marikovsky, M., Nevo, N., Vadai, E., & Harris-Cerruti, C. (2002). Cu/Zn superoxide dismutase plays a role in angiogenesis. *International Journal of Cancer*. Vol. 97, pp. 34-41
- Mishra, V. K., Srivastava, M. K., & Raizada, R. B. (1998). Testicular toxicity in rat to repeated oral administration of tetramethylthiuram disulfide (Thiram). *Indian Journal of Experimental Biology*. Vol. 36, pp. 390-394
- Morrison, B. W., Doudican, N. A., Patel, K. R., & Orlow, S. J. (2010). Disulfiram induces copper-dependent stimulation of reactive oxygen species and activation of the extrinsic apoptotic pathway in melanoma. *Melanoma Research*. Vol. 20, pp. 11-20
- Mosesso, P., Turchi, G., Cinelli, S., Di Chiara, D., Fiore, M., & Palitti, F. (1994). Clastogenic effects of the dithiocarbamate fungicides thiram and ziram in Chinese hamster cell lines cultured in vitro. *Teratogenesis, Carcinogenesis, and Mutagenesis*. Vol. 14, pp. 145-155
- Mosser, D. D., Caron, A. W., Bourget, L., Meriin, A. B., Sherman, M. Y., Morimoto, R. I., & Massie, B. (2000). The chaperone function of hsp70 is required for protection against stress-induced apoptosis. *Molecular and Cellular Biology*. Vol. 20, pp. 7146-7159


- Myung, J., Kim, K. B., & Crews, C. M. (2001). The ubiquitin-proteasome pathway and proteasome inhibitors. *Medicinal Research Reviews*. Vol. 21, pp. 245-273
- Nagendra, S. N., Faiman, M. D., Davis, K., Wu, J. Y., Newby, X., & Schloss, J. V. (1997). Carbamoylation of brain glutamate receptors by a disulfiram metabolite. *The Journal of Biological Chemistry*. Vol. 272, pp. 24247-24251 .
- Nobel, C. I., Kimland, M., Lind, B., Orrenius, S., & Slater, A. F. (1995). Dithiocarbamates induce apoptosis in thymocytes by raising the intracellular level of redox-active copper. *The Journal of Biological Chemistry*. Vol. 270, pp. 26202-26208
- Orrenius, S., Nobel, C. S., van den Dobbelsteen, D. J., Burkitt, M. J., & Slater, A. F. (1996). Dithiocarbamates and the redox regulation of cell death. *Biochemical Society Transactions*. Vol. 24, pp. 1032-1038
- Orth, M. W., & Cook, M. E. (1994). Avian tibial dyschondroplasia: a morphological and biochemical review of the growth plate lesion and its causes. *Veterinary Pathology*. Vol. 31, pp. 403-404
- Otsuka, G., Kubo, T., Imanishi, J., & Hirasawa, Y. (1996). Expression of heat-shock-proteins in the differentiation process of chondrocytes. *Nippon Geka Hokan*. Vol. 65, pp. 39-48
- Page, R. K., (1975). Teratogenic activity of Arasan fed to broiler breeder hens. *Avian Diseases*. Vol. 19, pp. 463-472
- Panganiban, L., Cortes-Maramba, N., Dioquino, C., Suplido, M. L., Ho, H., Francisco-Rivera, A., & Manglicmot-Yabes, A. (2004). Correlation between blood ethylenethiourea and thyroid gland disorders among banana plantation workers in the Philippines. *Environmental Health Perspectives*. Vol. 112, pp. 42-45
- Pucci, B., Adams, C. S., Fertala, J., Snyder, B. C., Mansfield, K. D., Tafani, M., Freeman, T., & Shapiro, I. M. (2007). Development of the terminally differentiated state sensitizes epiphyseal chondrocytes to apoptosis through caspase-3 activation. *Journal of Cellular Physiology*. Vol. 210, pp. 609-615
- Rahman, M. A., Grunberg, N. E., & Mueller, G. P. (1997). Disulfiram causes sustained behavioral and biochemical effects in rats. *Pharmacology, Biochemistry, and Behavior*. Vol. 56, pp. 409-415
- Rana, S. V., Allen, T., & Singh, R. (2002). Inevitable glutathione, then and now. *Indian Journal of Experimental Biology*. Vol. 40, pp. 706-716
- Rasaputra K. S., Liyanage R., Lay, J. O. Jr, McCarthy, F. M., & Rath, N. C., (2010). Tibial dyschondroplasia associated proteomic changes in chicken growth plate cartilage. *Avian Diseases*. Vol. 54, pp. 1166-1171
- Rath, N. C., Huff, W. E., Balog, J. M., & Huff, G. R. (2004). Comparative efficacy of different dithiocarbamates to induce tibial dyschondroplasia in poultry. *Poultry Science*. Vol. 83, pp. 266-274
- Rath, N. C., Huff, W. E., Bayyari, G. R., & Balog, J. M. (1995). Effect of thiram on chick chondrocytes in culture. *Journal of Toxicology and Environmental Health*. Vol. 44, pp. 369-376
- Rath, N. C., Huff, W. E., & Huff, G. R. (2007a). Thiram-induced changes in the expression of genes relating to vascularization and tibial dyschondroplasia. *Poultry Science*. Vol. 86, pp. 2390-2395
- Rath, N. C., Huff, W. E., Huff, G. R., & Kannan, L. (2007b). Induction of tibial dyschondroplasia by carbamate and thiocarbamate pesticides. *Avian Diseases*. Vol. 51, pp. 590-593

- Rath, N. C., Richards, M. P., Huff, W. E., Huff, G. R., & Balog, J. M. (2005). Changes in the tibial growth plates of chickens with thiram-induced dyschondroplasia. *Journal of Comparative Pathology*. Vol. 133, pp. 41-52
- Reddi, A. H., & Anderson, W. A. (1976). Collagenous bone matrix-induced endochondral ossification hemopoiesis. *The Journal of Cell Biology*. Vol. 69, pp. 557-572
- Roll, R., (1971). Teratologic studies with Thiram (TMTD) on two strains of mice]. *Archiv Fur Toxikologie*. Vol. 27, pp. 173-186
- SAP Report. Common mechanism of action of dithiocarbamates and thiocarbamates (2001). <http://www.epa.gov/scipoly/sap/meetings/2001/september7/September2001finalsapreport.pdf>. Accessed February 2011.
- Saunders, H., & Watkins, F. (2001). Allergic contact dermatitis due to thiuram exposure from a fungicide. *The Australasian Journal of Dermatology*. Vol. 42, pp. 217-218
- Schreck, R., Meier, B., Mannel, D. N., Droge, W., & Baeuerle, P. A. (1992). Dithiocarbamates as potent inhibitors of nuclear factor kappa B activation in intact cells. *The Journal of Experimental Medicine*. Vol. 175, pp. 1181-1194
- Seefeldt, T., Zhao, Y., Chen, W., Raza, A. S., Carlson, L., Herman, J., Stoebner, A., Hanson, S., Foll, R., & Guan, X. (2009). Characterization of a novel dithiocarbamate glutathione reductase inhibitor and its use as a tool to modulate intracellular glutathione. *The Journal of Biological Chemistry*. Vol. 284, pp. 2729-2737
- Shanthi, R., & Krishnamoorthy, M. (2002). Evaluation of the aneugenic potential of the fungicide Ferbam in mice. *Teratogenesis, Carcinogenesis, and Mutagenesis*. Vol. 22, pp. 451-459
- Shian, S. G., Kao, Y. R., Wu, F. Y., & Wu, C. W. (2003). Inhibition of invasion and angiogenesis by zinc-chelating agent disulfiram. *Molecular Pharmacology*. Vol. 64, pp. 1076-1084
- Soloneski, S., Reigosa, M. A., & Larramendy, M. L. (2003). Effect of the dithiocarbamate pesticide zineb and its commercial formulation, the azzurro. V. Abnormalities induced in the spindle apparatus of transformed and non-transformed mammalian cell lines. *Mutation Research*. Vol. 536, pp. 121-129
- Soloneski, S., Reigosa, M. A., & Larramendy, M. L. (2002). Effect of dithiocarbamate pesticide zineb and its commercial formulation, azzurro. II. micronucleus induction in immunophenotyped human lymphocytes. *Environmental and Molecular Mutagenesis*. Vol. 40, pp. 57-62
- Sook Han, M., Shin, K. J., Kim, Y. H., Kim, S. H., Lee, T., Kim, E., Ho Ryu, S., & Suh, P. G. (2003). Thiram and ziram stimulate non-selective cation channel and induce apoptosis in PC12 cells. *Neurotoxicology*. Vol. 24, pp. 425-434
- Steenland, K., Cedillo, L., Tucker, J., Hines, C., Sorensen, K., Deddens, J., & Cruz, V. (1997). Thyroid hormones and cytogenetic outcomes in backpack sprayers using ethylenebis(dithiocarbamate) (EBDC) fungicides in Mexico. *Environmental Health Perspectives*. Vol. 105, pp. 1126-1130
- Stoker, T. E., Goldman, J. M., & Cooper, R. L. (1993). The dithiocarbamate fungicide thiram disrupts the hormonal control of ovulation in the female rat. *Reproductive Toxicology*. Vol. 7, pp. 211-218
- Stoker, T. E., Jeffay, S. C., Zucker, R. M., Cooper, R. L., & Perreault, S. D. (2003). Abnormal fertilization is responsible for reduced fecundity following thiram-induced ovulatory delay in the rat. *Biology of Reproduction*. Vol. 68, pp. 2142-2149

- Taylor, T. R., & Whalen, M. M. (2009). Effects of ziram on tumor-cell-binding capacity, cell-surface marker expression, and ATP levels of human natural killer cells. *Cell Biology and Toxicology*. Vol. 25, pp. 447-455
- Teraoka, H., Urakawa, S., Nanba, S., Nagai, Y., Dong, W., Imagawa, T., Tanguay, R. L., Svoboda, K., Handley-Goldstone, H. M., Stegeman, J. J., & Hiraga, T. (2006). Muscular contractions in the zebrafish embryo are necessary to reveal thiuram-induced notochord distortions. *Toxicology and Applied Pharmacology*. Vol. 212, pp. 24-34
- Tian, W. X., Zhang, W. P., Li, J. K., Bi, D. R., Guo, D. Z., Pan, S. Y., Zhang, Y. H., & Qin, P. (2009). Identification of differentially expressed genes in the growth plate of broiler chickens with thiram-induced tibial dyschondroplasia. *Avian Pathology*. Vol. 38, pp. 161-166
- Tilton, F., La Du, J. K., Vue, M., Alzarban, N., & Tanguay, R. L. (2006). Dithiocarbamates have a common toxic effect on zebrafish body axis formation. *Toxicology and Applied Pharmacology*. Vol. 216, pp. 55-68
- USEPA.(2001).http://www.epa.gov/scipoly/sap/meetings/2001/september7/dithiofinal_aug17.pdf. Assessed February 2011
- Vaccari, A., Saba, P., Mocci, I., & Ruiu, S. (1999). Dithiocarbamate pesticides affect glutamate transport in brain synaptic vesicles. *The Journal of Pharmacology and Experimental Therapeutics*. Vol. 288, pp. 1-5
- Valentine, H. L., Viquez, O. M., Amarnath, K., Amarnath, V., Zyskowski, J., Kassa, E. N., & Valentine, W. M. (2009). Nitrogen substituent polarity influences dithiocarbamate-mediated lipid oxidation, nerve copper accumulation, and myelin injury. *Chemical Research in Toxicology*. Vol. 22, pp. 218-226
- van Boxtel, A. L., Pieterse, B., Cenijn, P., Kamstra, J. H., Brouwer, A., van Wieringen, W., de Boer, J., & Legler, J. (2010). Dithiocarbamates induce craniofacial abnormalities and down regulate sox9a during zebra fish development. *Toxicological Sciences*. Vol. 117, pp. 209-217
- Vargas, M. I., Lamas, J. M., & Alvarenga, V. (1983). Tibial dyschondroplasia in growing chickens experimentally intoxicated with tetramethylthiuram disulfide. *Poultry Science*. Vol. 62, pp. 1195-1200
- Veltmann, J. R., Jr, Rowland, G. N., & Linton, S. S. (1985). Tibial dyschondroplasia in single-comb White Leghorn chicks fed tetramethylthiuram disulfide (a fungicide). *Avian Diseases*. Vol. 29, pp. 1269-1272
- Viola-Rhenals, M., Rieber, M. S., & Rieber, M. (2007). Role of peroxidases, thiols and Bak/Bax in tumor cell susceptibility to Cu[DEDTC]₂. *Biochemical Pharmacology*. Vol. 74, pp. 841-850
- Viquez, O. M., Lai, B., Ahn, J. H., Does, M. D., Valentine, H. L., & Valentine, W. M. (2009). N,N-diethyldithiocarbamate promotes oxidative stress prior to myelin structural changes and increases myelin copper content. *Toxicology and Applied Pharmacology*. Vol. 239, pp. 71-79
- Viviani, B., Bartesaghi, S., Binaglia, M., Corsini, E., Boraso, M., Grazi, E., Galli, C. L., & Marinovich, M. (2008). Dithiocarbamate propineb induces acetylcholine release through cytoskeletal actin depolymerization in PC12 cells. *Toxicology Letters*. Vol. 182, pp. 63-68

- Waibel, P. E., Pomeroy, B. S., & Johnson, E. L. (1955). Effect of Arasan-Treated Corn on Laying Hens. *Science*. Vol. 121, pp. 401-402
- Wang, X. F., Li, S., Chou, A. P., & Bronstein, J. M. (2006). Inhibitory effects of pesticides on proteasome activity: implication in Parkinson's disease. *Neurobiology of Disease*. Vol. 23, pp. 198-205
- Wang, F., Zhai, S., Liu, X., Li, L., Wu, S., Dou, Q. P., & Yan, B. (2011). A novel dithiocarbamate analogue with potentially decreased ALDH inhibition has copper-dependent proteasome-inhibitory and apoptosis-inducing activity in human breast cancer cells. *Cancer Letters*. Vol. 300, pp. 87-95
- Ware, G.W., & D. M. Whitcare, (2004). *The Pesticide Book*. 6th Edition. Meister Pro Information Resources, Willoughby, OH.
- Wild, A. C., & Mulcahy, R. T. (1999). Pyrrolidine dithiocarbamate up-regulates the expression of the genes encoding the catalytic and regulatory subunits of gamma-glutamylcysteine synthetase and increases intracellular glutathione levels. *The Biochemical Journal*. Vol. 338 (Pt 3), pp. 659-665

IntechOpen



Pesticides in the Modern World - Effects of Pesticides Exposure

Edited by Dr. Margarita Stoytcheva

ISBN 978-953-307-454-2

Hard cover, 376 pages

Publisher InTech

Published online 12, September, 2011

Published in print edition September, 2011

The introduction of the synthetic organochlorine, organophosphate, carbamate and pyrethroid pesticides by 1950s marked the beginning of the modern pesticides era and a new stage in the agriculture development. Evolved from the chemicals designed originally as warfare agents, the synthetic pesticides demonstrated a high effectiveness in preventing, destroying or controlling any pest. Therefore, their application in the agriculture practices made it possible enhancing crops and livestock yields and obtaining higher-quality products, to satisfy the food demand of the continuously rising world population. Nevertheless, the increase of the pesticide use estimated to 2.5 million tons annually worldwide since 1950., created a number of public and environment concerns. This book, organized in two sections, addresses the various aspects of the pesticides exposure and the related health effects. It offers a large amount of practical information to the professionals interested in pesticides issues.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Narayan C. Rath, Komal S. Rasaputra, Rohana Liyanage, Gerry R. Huff and William E. Huff (2011).
Dithiocarbamate Toxicity - An Appraisal, Pesticides in the Modern World - Effects of Pesticides Exposure, Dr.
Margarita Stoytcheva (Ed.), ISBN: 978-953-307-454-2, InTech, Available from:
<http://www.intechopen.com/books/pesticides-in-the-modern-world-effects-of-pesticides-exposure/dithiocarbamate-toxicity-an-appraisal>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen