

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

**4,800**

Open access books available

**122,000**

International authors and editors

**135M**

Downloads

Our authors are among the

**154**

Countries delivered to

**TOP 1%**

most cited scientists

**12.2%**

Contributors from top 500 universities



**WEB OF SCIENCE™**

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



# Dissociative Tendency, Anger Expression, and Frontal Activation During a Verbal Fluency Task

Toru Uehara<sup>1</sup>, Yoko Ishige<sup>2</sup> and Masashi Suda<sup>2</sup>

<sup>1</sup>General Health Support Centre, Gunma University,

<sup>2</sup>Department of Neuropsychiatry, Gunma University Graduate School of Medicine,  
Japan

## 1. Introduction

Dissociation in adolescence has been described as a part of severe psychopathology with a complex comorbidity or serious trauma including sexual abuse (e.g., Chu and Dill, 1990; Sar et al., 2006). In contrast, it is also possible for healthy subjects to experience nonpathological (nonclinical) dissociation including daydreams or defense from strong stressors (de Ruiters et al., 2006). It is important to clarify the etiology of dissociative experiences. For example, a lack of integration of consciousness, amnesia, derealization, or depersonalization, and some neural mechanisms underlying dissociation could be reasonably hypothesized as core features of dissociative disorders (DD). Particularly, self-reference related regions such as the medial prefrontal or ventral frontal area might be involved in identity diffusion or multiple identity disorder (Reinders et al., 2003). Actually, recent advances in neuroimaging technology enable us to explore the brain functioning of dissociation. Although it is ideal to examine a patient with DD who is experiencing dissociation in vivo, it is simultaneously intrusive to induce pathological dissociation using reminders or stressors. According to the spectrum of dissociative experiences, another option would be to examine nonclinical dissociation or dissociative tendencies in a typical setting. Dissociative experiences usually begin in childhood, and less than 8% of the DD diagnoses are made in adolescents between the ages of 12 and 19 (Kluft, 1984). Some of the experiences are not subjectively distressful; therefore, dissociation might be underestimated as a clinical phenomenon (Foote et al., 2006). Furthermore, recent campus mental health professionals in Japan have often indicated that a trend for defence mechanism is “to dissociate from repression” in young adults (e.g., Takaishi, 2000). Based on these assumptions, the present study focused on nonclinical dissociative tendencies and investigated frontal functioning during a cognitive task in healthy university students.

Aggression or impulsivity is another important mental health topic in youth including attention-deficit hyperactivity disorder (ADHD) or childhood bipolar spectrum including temper dysregulation disorder. Impulsivity is related to the loss of response inhibition, particularly to the lack of frontal functioning (e.g., Chamberlain and Sakhian, 2007). A recent study concluded that slower cortical thinning in the prefrontal cortex during adolescence is characteristic of ADHD, providing neurobiological evidence for the

dimensionality of this disorder (Shaw et al., 2010). Anger is the most relevant emotion closely associated with impulsivity or aggression. Activity in the dorsal anterior cingulate cortex is positively related to self-reporting of anger and individual differences in general aggression (Denson et al., 2009). Another study (Hewig et al., 2004) reported that subjects with relatively greater left than right frontal cortical activity showed higher anger-out scores and lower anger-control scores. Furthermore, anger induction was uniquely associated with increased regional cerebral blood flow to the right temporal pole and thalamus. Based on these studies, it appears reasonable to investigate functional correlates of anger expression in youth.

Interestingly, one descriptive study reported a relationship between dissociative experiences and anger proneness in late adolescent females (Calamari and Pini, 2003). The authors stated that significant correlations were obtained between the Dissociative Experience Scale (DES) and the State-Trait Anger Expression Inventory (STAXI), confirming a connection between anger proneness and dissociation described in patients with dissociative disorders in a nonclinical sample. We also speculate that dissociation may be one of the defence mechanisms to cope with anger, impulsivity, or aggression. Consequently, frontal functioning may determine individual styles including expression, control, and suppression.

Near-infrared spectroscopy (NIRS) and the newly developed optic brain functional imaging are promising techniques because of their non-invasiveness and convenience. NIRS employs near-infrared light emitted and detected on the skull skin (Boas et al., 2004). It allows the monitoring of hemodynamic changes, which include both cerebral blood volume changes and oxygenation state, using a small apparatus with a high time resolution of about 0.1 s. It also allows the monitoring of changes in both oxygenated haemoglobin concentration [o-Hb] and deoxygenated haemoglobin concentration [d-Hb]. NIRS is suitable for studies of higher brain function because it enables measurements in a natural setting compared with other brain imaging techniques. For example, subjects can undergo an NIRS examination in the sitting position, with their eyes open, or while speaking (Suda et al., 2009). Taking advantage of these characteristics, several NIRS studies on psychiatric disorders, such as schizophrenia (Grignon et al., 2008; Takizawa et al., 2008), depression (Matsuo et al., 2005), eating disorders (Suda et al., 2010a; Uehara et al., 2007), and ADHD disorder (Ehlis et al., 2008), have been conducted. These characteristics of NIRS have also enabled the investigation of subjective experiences in healthy subjects such as conversation, subjective sleepiness, and psychological fatigue (Suda et al., 2008, 2009, 2010b).

The present study aimed to examine cortical activation using a verbal fluency task, which has been applied widely in clinical and nonclinical samples as a standard and specific paradigm to activate the frontal lobe. We used a newly developed multi-channel NIRS machine specified for frontal regions to explore differences in frontal activation according to the dissociative tendency. In addition, we examined correlations among frontal functioning, dissociation scores, and anger expression styles measured by self-report.

## 2. Material and methods

### 2.1 Participants

The study participants were 44 healthy university students (29 females), with a mean age of 20.5 years (SD, 2.0). They were all Japanese, and two students were left-handed. None of the

participants had any significant medical/psychiatric history. The participants were voluntarily recruited as subjects for this scientific study, and were paid 1,600 yen for a 2 hours exam as a co-operator as per the official provision. All subjects gave written informed consent prior to their participation in the study, which conformed with the provisions of the Declaration of Helsinki revised in Edinburgh in 2000. Privacy and anonymity of all participants were carefully preserved. The data was collected from August to December in 2009 and 2010.

## 2.2 NIRS (Fig. 1-3)

NIRS allows the calculation of changes in [Hb] parameters, including [o-Hb] and [d-Hb], by measuring the attenuation of near-infrared light at an approximate 800 nm wavelength. Neural activation induces regional hemodynamic changes in brain tissue, almost identical in pattern to spontaneous cerebral neural activity. Cortical activation is typically detected as an [o-Hb] increase or an [d-Hb] decrease; however, the direction of change in [d-Hb] can be ambiguous in the frontal lobe (Sato et al., 2007). Mainly changes in [o-Hb] at a depth of 2–3 cm from the scalp, that is, the surface of the cerebral cortex, are correlated with positron emission tomography (PET) hemodynamic changes (Ohmae et al., 2006) and blood-oxygenation-level-dependent signal changes in functional magnetic resonance imaging (fMRI) (Toronov et al., 2001; Mehagnoul-Schipper et al., 2002). NIRS does not measure cerebral luminescence but measures the attenuation of irradiated light intensity. Therefore, the combination of optical irradiation and photon detection determines the resolution. It characteristically measures not the 1:1 combination of irradiation and detection, but the light from one light source with 2 or more detectors arranged geometrically in the measurement system of NIRS. Thus, information on which detector measures the signal of which portion becomes important. Some methods are available for judging this channel separation. The first method, time division multiple access, makes a light source turn on in order, and separates the signal on a time axis. The second method, frequency division multiple access, is for modulating and irradiating two or more light sources with different frequencies and separating a signal based on frequency information after detection. The third method is code division multiple access (CDMA), using spectrum diffusion attenuation, which is applied in such applications as global positioning system or mobile phone. A new machine, OEG-16 (Spectratech, Inc, Yokohama, Japan), uses CDMA and is very convenient and portable. It can generate NIRS data under natural conditions noninvasively, and artifacts induced by hair can be avoided because of the adjustments only on the front of the head. The OEG-16 measures 16 channels on the frontal lobe (according to Broadman's map, provides data on 10, 11, 12, 44, 45, and 46). Its time resolution is 0.5 s, and space resolution is 2 cm. A headset was placed on the participant's head according to the 10/20 system, by which a central hole was coordinated with Fz. The measurement points for channels 1 to 8 were placed from the right lateral to the central pole. For channels 9 to 16, the measurement points were placed from the ventral/rostral to the left lateral (refer to the video content). These placements provided for relative changes in [Hb] concentration, and the values obtained were in arbitrary units (concentration  $\times$  path length).

Details of NIRS methodology have already been described in major publications in Japan (Fukuda, 2009; 2011).




Fig. 1. Estimated areas for measuring the blood volume changes of cerebral cortex. (© Hitachi Medical Corporation, <http://www.hitachi-medical.co.jp/info/opt/qa.html#q1>)



Fig. 2. OEG-16 (18×17×4 cm, Spectratech, Inc.)



Fig. 3. Measuring points and headset. (Spectratech, Inc.)

### 2.3 Verbal fluency task

This standardized activation task is employed internationally for NIRS measurements, and it has been confirmed that this method provides widespread frontal activation reliably (Kono et al., 2007; Schecklmann et al., 2008; Kakimoto et al., 2009). The frontal activation task was a modified version of the verbal fluency task. A subject sat on a comfortable chair in a quiet room with their eyes open throughout the measurement. The activation task consisted of a 15-s pre-task baseline, a 30-s verbal fluency period, and a 15-s post-task baseline. During

the verbal fluency period, the subjects were instructed to verbally generate as many words as they could whose initial Japanese syllable (mora) was either /a/, /ki/, or /ha/. These three initial syllables were used in the above-mentioned order and changed every 10 s during the 30-s verbal fluency period to reduce the time during which the subjects remained silent. The number of words generated during the verbal fluency period was determined as a measure of task performance. During the pre-task and post-task baseline periods, the subjects were instructed to repeat the syllables /a/, /i/, /u/, /e/, and /o/ as the Japanese counterparts of A, B, and C in English. Two sets of this task were performed, and the respective data were superimposed and averaged.



Fig. 4. One epoch of the verbal fluency test (VFT).

#### 2.4 ADES and STAXI

The ADES was developed solely for adolescents by Armstrong et al. (1997) to detect dissociative behaviour in children between 11 and 17 years of age. It is a 30-item self-reporting scale composed of an 11-point scale ranging from 0 representing “never” to 10 representing “always”. The total ADES score is equal to the mean of all item scores. A subject circles the number that best describes how often a given experience happens. On the title page, respondents were instructed not to count experiences that occur under the influence of any drugs. Several versions of the ADES are available in Japanese (e.g., Tanabe, 2004). We used the one developed by Matsumoto et al. (2004) because of its excellent psychometric properties.

The STAXI tool evaluates anger isolated from hostility and aggression, covering anger experience and expression (Spielberger, et al., 1998). The STAXI is a self-reported 44 items, and individuals answered on a four-point Likert scale (score range: 0–132) to assess either the intensity of their angry feelings or the frequency as state, trait, anger-in, anger-out, and anger-control subscores (expression of anger toward other persons or objects in the environment is anger expression-out; holding in or suppressing angry feelings is anger expression-in; and controlling angry feelings by preventing the expression of anger toward other persons or objects in the environment, or controlling suppressed angry feelings by calming down or cooling off is anger-control). The Japanese version was developed by Mine et al. (1997), and the first STAXI was used in the present study.

#### 2.5 Data analysis

The continuous waveforms of the [Hb] changes on all 16 channels were acquired from all subjects during the paradigm. The individually averaged [Hb] waveforms were obtained as the average sum of two trials; a baseline realignment for 5 s before and after the task periods, and a task segment averaging two sets of 15-s image viewing periods. Thereafter, the grand average values of the baseline and task segments for each channel were calculated for all data. Figure 5 is an example of a grand average waveform for three parameters; the

red polygonal line indicates the relative changes in [o-Hb], the blue indicates those of [d-Hb], and the green indicates the changes in the total-Hb (sum of o- and d-Hb).

IntechOpen




Fig. 5. The red polygonal line indicates the relative changes in [o-Hb], the blue indicates those of [d-Hb], and the green indicates the changes in the total-Hb (sum of o- and d-Hb).

We used only [o-Hb] values as cerebral blood volume changes for statistics, based on previous reports (Suto et al., 2004; Kameyama et al., 2006). Topography (video content) was presented on the frontal portion according to the time course. In this grand average data, channels that carried significant activation were analyzed between the pre-task and task periods using the t-test (<http://www.brsystems.jp>). Differences in the mean values were tested by combined variance, as the number of samples for the two periods was not equal.

Combined variance:  $Ue2 = \{ (na - 1) ua2 + (nb - 1)ub2 \} / \{ na + nb - 2 \}$

Variance and numbers of sample a:  $ua2$  and  $na$

Variance and numbers of sample b:  $ub2$  and  $nb$

T-value:  $t0 = | mXa - mXb | / \text{root}\{ue2(1/na + 1/nb)\}$

Average of each sample:  $mXa, mXb$




Fig. 6. An presentation using the Data Viewer (BR Systems, Inc.).

In the next step, differences were investigated according to dissociative tendencies followed by the t-test for each channel during the task period between the sample grand average data divided by the ADES mean scores (“strong” and “weak”). In the last step, the relationships between the [o-Hb] changes, the STAXI subscales, and the ADES scores were investigated. The channels, with significantly correlated changes, were analyzed by a nonparametric Spearman’s correlation coefficient (two-tailed). Imaging software was used to analyze the NIRS parameters, and figure 6 shows an example of demonstration by this software (Data Viewer ver.1.1a, BR Systems Inc., Tokyo, Japan). The other statistical analyses were conducted using SPSS version 17.0 (SPSS, Inc., Tokyo, Japan).

### 3. Results

#### 3.1 Waveform and dissociation

Figure 7 shows the grand average waveforms for the [Hb] changes for all participants, and figure 8 indicates the comparisons of waveforms according to the ADES average scores (mean = 13.9, S.D. = 12.8). 27 students scored under the mean as “weak” on the left, and the others as “strong” on the right. The red polygonal line indicates the relative changes in [o-Hb], the blue indicates those of [d-Hb], and the green indicates the changes in the total-Hb (sum of o- and d-Hb). The video content depicts the topography of the [o-Hb] changes (“weak” on the left and “strong” on the right side), and redder areas represent greater



activation (grading to yellow, green, and opposite in deeper blue). Overall, gradual increases and fluctuations were generally observed in widespread channels during the task. The video file can be viewed on this website.

[http://www.youtube.com/watch?v=g7IG5HG8q\\_U](http://www.youtube.com/watch?v=g7IG5HG8q_U)




Fig. 7. Grand average waveforms

This demonstrated grand average waveforms of [Hb] changes for all participants' data. The red polygonal line indicates [o-Hb], blue indicates [d-Hb], and green indicates the total [Hb] (sum of o-Hb and d-Hb) relative changes. The vertical axis represents the relative changes in [Hb] (mMmm), and the vertical grey lines represent the start and end of task periods. The numbers on the abscissa indicate measurement time points (seconds). The data of the respective channels was placed according to each number from the top to the bottom.




Fig. 8. Comparisons of the grand average waveforms according to dissociative tendency

This demonstrated grand average waveforms of [Hb] changes for all participants according to ADES, in which two samples were divided by the mean scores (the data over the mean is in the left and the data below the mean is on the right side). The red polygonal line indicates [o-Hb], blue indicates [d-Hb], and green indicates the total [Hb] (sum of o-HB and d-HB) relative changes. The vertical axis represents the relative changes in [Hb] (mMmm), and the vertical grey lines represent the start and end of task periods. The numbers on the abscissa indicate measurement time points (seconds). The data of the respective channels was placed according to each number from the top to the bottom.

Comparisons between the baseline and task periods are summarized in Table 1. The respective arbitrary units and t-values for each channel are presented for all, “weak” and “strong” students; significant activations (two-tailed, DF = 32) were obtained on channels 3–15 for all students; on channels 2–7, 10, 12, and 15 for “weak” samples; and on channels 3–8 and 10–15 for the “strong” samples. Comparisons of the activation for the task periods between the “strong” and “weak” samples were calculated; t-values for every channel indicated that they were significantly lower on channels 1\*\*, 2\*\*, and 7\* (dorsolateral to the ventral portion) in the right PFC, and higher in channel 13\*\* and lower in channel 16\*\* in the left PFC for “strong” than for “weak” samples (\*p < .05, \*\*p < .01, DF = 52) The total ADES scores were negatively correlated with activation in the center prefrontal channel 8 (rostral ventral PFC; r = -.30, p < .05).

channel	WEAK (n = 27)			STRONG (n = 17)			All 44 Students			Comparisons for task periods between WEAK and STRONG	
	Rest	Task	t	Rest	Task	t	Rest	Task	t	t	p (DF = 52)
1	0.003	0.012	1.37	0	-0.002	-0.56	0.002	0.006	0.88	4.77	0.0002
2	0	0.015	2.36*	0.002	0	-0.61	0.001	0.008	1.44	5.48	0.00001
3	0	0.014	3.13**	0	0.012	4.00**	0	0.013	3.56**	0.51	0.61
4	0	0.018	2.91**	0.002	0.025	3.82**	0	0.022	3.79**	-1.91	0.06
5	0	0.013	4.53**	-0.002	0.014	4.31**	-0.001	0.013	4.90**	-0.40	0.70
6	0	0.016	3.53**	0	0.019	4.38**	0	0.018	4.21**	-0.76	0.45
7	0	0.012	3.42**	-0.002	0.008	2.58*	-0.002	0.010	3.53**	2.45	0.02
8	0	0.011	1.09	-0.002	0.006	3.35**	0	0.008	2.80**	1.08	0.29
9	-0.002	0.005	1.89	-0.001	0.006	1.80	-0.003	0.005	2.16*	-1.09	0.28
10	0	0.015	2.96**	-0.001	0.012	2.44*	0	0.013	3.00**	0.91	0.37
11	0	0.008	2.00	-0.004	0.009	4.18**	-0.003	0.008	3.30**	-0.65	0.52
12	0	0.014	2.76*	0	0.017	3.01**	0	0.015	3.35**	-1.40	0.30
13	0.003	0.014	1.73	-0.005	0.023	4.78**	-0.002	0.018	3.48**	-3.08	0.0031
14	0	0.009	1.30	-0.005	0.013	2.22*	-0.002	0.011	2.20*	-1.40	0.17
15	9	0.018	2.92**	0.005	0.016	2.25*	0.002	0.017	3.20**	0.83	0.41
16	0.005	0.011	1.08	0.010	-0.006	-1.66	0.007	0.003	-0.75	4.00	0.0003

arbitrary unit: mMmmol \*; p < .05, \*\*; p < .01 (DF = 32)

WEAK/STRONG; divided by the mean scores '13.9' of the adolescent dissociative experiences scale (ADES)

Table 1. Comparisons of activations according to dissociative tendency

### 3.2 Correlation with anger expression

Table 2 shows the descriptive values of the STAXI subscale scores for all participants. Spearman's correlation coefficients were calculated for the relative changes in [o-Hb] for the 16 channels and the STAXI scores. The anger-in STAXI scores were negatively correlated with channel 8 activation (frontal pole of PFC; r = -.31, p < .05). The other scale scores were not significantly correlated with any other channel activation.

	state	trait	AX/In	AX/Out	AX/Con	AX/EX
mean	12.3	20.3	19.2	14.3	22.0	27.5
S.D.	3.5	7.1	4.2	3.7	4.1	7.2

STAXI: State Trait Anger Expression Inventory

AX/In: anger expression in

AX/Out: anger expression out

AX/Con: anger expression control

AX/EX: expression, (AX/In+AX/Out)-Ax/Con

Table 2. Descriptive values of the STAXI

#### 4. Discussion

Some interesting relationships were found between frontal activation and psycho-behavioural variables that can be summarized as follows: 1. Strong dissociative tendencies were generally related to lower PFC activation. In particular, the activation was relatively dominant on the lateral and right sides. 2. Only channel 13 (left inferior frontal gyrus, close to BA46) activation could be directly associated with dissociative reactions with regard to contrast of activations; significant increases in channel 13 and significant decreases in channel 7 (right BA46). 3. Analyses of the correlations revealed that frontal pole (channel 8, close to BA10) dysfunction may be linked to dissociative tendencies. This relationship was paradoxical to anger suppression; the "ANGER-IN" style was correlated with lower activation in this portion (rostral PFC, channel 8).

Similar to the normal control data of many previous NIRS studies, the verbal fluency task provided excellent activation in widespread areas of PFC both in the "strong and "weak" dissociative groups, and this was reasonable, as our sample consisted of healthy students. However, according to ADES, a dissociative tendency might be associated with differences in prefrontal deactivation. In particular, lateral or right side deactivation may influence the dissociative tendency. The findings of the present study also suggest that suppression may be disturbed by over-function in the frontal pole (rostral PFC), and a contradiction could be seen in dissociative experiences. Considering dissociation and repression as coping mechanism to distresses, frontal pole deactivations may induce dissociative defence and the repression of anger expression.

Reviewing related studies, Bell et al. (2010) commented that dissociation is associated with increases in PFC activity and suggested that intervention by the executive system for both automatic and voluntary cognitive processing was common to both hysteria and hypnosis. We found only one study (Amrhein et al., 2008) that has investigated high/low dissociators in subjects without any psychiatric or neurological disorders and without prior trauma experiences. High dissociators show cognitive deficiencies, and a hippocampal and PFC dysfunction was assumed to be the core factor. Although the subjects had clinically disordered post-traumatic stress disorder (PTSD), Lanius et al. (2010) proposed that a dissociative subtype of PTSD is a form of emotion dysregulation that involves emotional overmodulation mediated by midline prefrontal inhibition of the limbic regions. Based on these studies, the results from the present study strongly support involvement of PFC with dissociation. A PET study stated that anger conditions are

associated with increased regional cerebral blood flow (rCBF) in the left inferior PFC and decreased rCBF in the right superior frontal cortex (Kimbrell et al., 1999). An fMRI study indicated that medial PFC activity (mostly identical to BA10 or 9) is related to self-reported rumination and individual differences in displaced aggression (Denson et al., 2009). As “ANGER-IN” means the frequency with which angry feelings are suppressed, rumination appears to be linked to this type of anger expression. Based on these findings and our results, the relationships between anger and bilateral ventral or rostral PFC involvement (dominantly left) can be assumed.

Some limitations of this study should be noted. The data could not be interpreted directly for DD because clinical subjects were strictly excluded. A future study should administer paradigms directly to induce dissociation or present specific stimulation. Similarly, tasks provoking anger such as visual stimuli should be considered. We did not reveal a correlation between frontal activation and state/trait anger, anger-out expression, or anger-control. We should still be conservative when discussing the association between anger suppression and brain function. This study used a frontal-specific methodology that should be complemented with a whole-brain measurement. Considering the importance of the clinical application, relative changes in cerebral [Hb] data as absolute values should be cautiously interpreted. Furthermore, anatomically identifying the measuring points is a major methodological challenge in multichannel NIRS, but the space resolution in NIRS ranges from 2 to 3 cm; therefore, it would not be problematic to discuss functional dynamism within broader frontal connectivity. Rather than being a weakness, high time resolution can contribute to clarifying the network or circuit based on detailed analyses of the time-course and activation patterns. Actually, the OEG-16 has already been applied in a psychological study, and its availability could be introduced (Uehara et al., 2011).

Differences in frontal activation suggest that dissociation might be related to PFC. Activation in the left inferior frontal gyrus could be associated with dissociative reactions in contrast to right side activation. We also suggest that a dissociative tendency and anger suppression could be associated with rostral PFC deactivations. Further study needs to include a clinical application and improve the task paradigm.

## 5. Acknowledgment

This study was supported, in part, by the Grants-in Aid for Scientific Research (C-21591504) from the Japan Society of Promotion of Science and by the Intramural Research Grant (20-1) for Neurological and Psychiatric Disorders of NCNP, Japan.

A summary of this study was presented at the 16th ESCAP in Helsinki on June 2011.

## 6. References

- Amrhein, C., Hengmith, S., Maragkos, M., & Hennig-Fast, K. (2008). Neuropsychological characteristics of highly dissociative healthy individuals. *Journal of Trauma and Dissociation*, Vol. 9, pp.525–542. ISSN :1529-9732
- Armstrong, J.G., Putnam, F.W., Carlson, E.B., Libero, D.Z., & Smith, S.R. (1997). Development and validation of a measure of adolescent dissociation: the Adolescent Dissociative Experiences Scale. *Journal of Nervous and Mental Diseases*, Vol. 185, pp.491–497. ISSN :0022-3018

- Bell, V., Oakley, D.A., Halligan, P.W., & Deeley, Q. (2011). Dissociation in hysteria and hypnosis: evidence from cognitive neuroscience. *Journal of Neurology Neurosurgery and Psychiatry*, Vol. 82, pp.332-339, ISSN : 0002-3050.
- Boas, D.A., Dale, A.M., & Franceschini, M.A. (2004). Diffuse optical imaging of brain activation: approaches to optimizing image sensitivity, resolution, and accuracy. *Neuroimage*, Vol. 23, pp.275-288. ISSN : 1053-8119
- Calamari, E., & Pini, M. (2003). Dissociative experiences and anger proneness in late adolescent females with different attachment styles. *Adolescence*, Vol. 38, pp.287-303. ISSN : 0001-8449
- Chamberlain, S.R., & Sahakian, B.J. (2007). The neuropsychiatry of impulsivity. *Current Opinion of Psychiatry*, Vol.20, pp.255-261. ISSN : 0951-7367
- Chu, J.A. & Dill, D.L. (1990). Dissociative symptoms in relation to childhood physical and sexual abuse. *American Journal of Psychiatry*, Vol. 147, pp87-92. ISSN : 0002-953X
- Denson, T.F., Pedersen, W.C., Ronquillo, J., & Nandy, A.S. (2009). The angry brain: neural correlates of anger, angry rumination, and aggressive personality. *Journal of Cognitive Neuroscience*, Vol. 21, pp.734-744. ISSN : 0898-929X
- Ehlis, A.C., Bähne, C.G., Jacob, C.P., Herrmann, M.J., & Fallgatter, A.J. (2008). Reduced lateral prefrontal activation in adult patients with attention-deficit/hyperactivity disorder (ADHD) during a working memory task: a functional near-infrared spectroscopy (fNIRS) study. *Journal of Psychiatry Research*, Vol. 42, pp.1060-1067. ISSN : 0022-3956
- Foot, B., Smolin, Y., Kaplan, M., Legatt, M.E., & Lipschitz, D. (2006). Prevalence of dissociative disorders in psychiatric outpatients. *American Journal of Psychiatry*, Vol. 163, pp.623-629. ISSN : 0002-953X
- Fukuda, M. (2011). *Clinical evaluation of NIRS data*, Nakayama Shoten, ISBN 978-4-521-73368-5, Tokyo, Japan (in Japanese)
- Fukuda, M. (2009). *NIRS in Psychiatry*, Nakayama Shoten, ISBN 978-4-521-73118-6, Tokyo, Japan (in Japanese)
- Grignon, S., Forget, K., Durand, M., & Huppert, T. (2008). Increased left prefrontal activation during staring/mutism episodes in a patient with resistant catatonic schizophrenia: a near infrared spectroscopy study. *Cognitive and Behavioral Neurology*, Vol.21, pp. 41-45. ISSN : 1543-3633
- Hewig, J., Hagemann, D., Seifert, J., Naumann, E., & Bartusek, D. (2004). On the selective relation of frontal cortical asymmetry and anger-out versus anger-control. *Journal of Personality and Social Psychology*, Vol.87, pp.926-939. ISSN : 0022-3514
- Kakimoto, Y., Nishimura, Y., Hara, N., Okada, M., Tanii, H., & Okazaki, Y. (2009). Intrasubject reproducibility of prefrontal cortex activities during a verbal fluency task over two repeated sessions using multi-channel near-infrared spectroscopy. *Psychiatry and Clinical Neurosciences*, Vol. 63, pp.491-499. ISSN : 1323-1316
- Kameyama, M., Fukuda, M., Yamagishi, Y., Sato, T., Uehara, T., Ito, M., & Mikuni, M. (2006). Frontal lobe function in bipolar disorder: a multichannel near-infrared spectroscopy study. *Neuroimage*, Vol.29, pp.172-184. ISSN : 1053-8119
- Kimbrell, T.A., George, M.S., Parekh, P.I., Ketter, T.A., Podell, D.M., Danielson, A.L., Repella, J.D., Benson, B.E., Willis, M.W., Herscovitch, P., & Post, R.M. (1999).

- Regional brain activity during transient self-induced anxiety and anger in healthy adults. *Biological Psychiatry*, Vol.46, pp.454–465, ISSN : 0006-3223
- Kluft, R.P. (1984) Treatment of multiple personality disorder. A study of 33 cases. *Psychiatric Clinics of North America*, Vol. 7, pp. 9–29, ISSN : 0193-953X
- Kono, T., Matsuo, K., Tsunashima, K., Kasai, K., Takizawa, R., Rogers, M.A., Yamasue, H., Yano, T., Taketani, Y., & Kato, N. (2007). Multiple-time replicability of near-infrared spectroscopy recording during prefrontal activation task in healthy men. *Neuroscience Research*, Vol. 57, pp.504–512, ISSN: 0168-0102
- Lanius, R.A., Vermetten, E., Loewenstein, R.J., Brand, B., Schmahl, C., Bremner, J.D., & Spiegel, D. (2010). Emotion modulation in PTSD: Clinical and neurobiological evidence for a dissociative subtype. *American Journal of Psychiatry*, Vol. 167, pp.640–647. ISSN: 0002-953X
- Matsumoto, T., Azekawa, T., Yamaguchi, A., Asami, T., & Iseki, E. (2004). Habitual self-mutilation in Japan. *Psychiatry and Clinical Neurosciences*, Vol. 58, pp.191–198. ISSN : 1323-1316
- Matsuo, K., Onodera, Y., Hamamoto, T., Muraki, K., Kato, N., & Kato, T. (2005). Hypofrontality and microvascular dysregulation in remitted late-onset depression assessed by functional near-infrared spectroscopy. *Neuroimage*, Vol. 26, pp. 234–242. ISSN : 1053-8119
- Mehagnoul-Schipper, D.J., van der Kallen, B.F., Colier, W.N., van der Sluijs, M.C., van Erning, L.J., Thijssen, H.O., Oeseburg, B., Hoefnagels, W.H., & Jansen, R.W. (2002). Simultaneous measurements of cerebral oxygenation changes during brain activation by near-infrared spectroscopy and functional magnetic resonance imaging in healthy young and elderly subjects. *Human Brain Mapping*, Vol. 16, pp. 14–23. ISSN :1065-9471
- Mine, H., Hama, H., & Okubo, J. (1997). An attempt for the standardization anger expression scale for Japanese. *Japanese Journal of Research on Emotion*, Vol. 4, pp.14–21. ISSN: 1882-8817 (in Japanese)
- Ohmae, E., Ouchi, Y., Oda, M., Suzuki, T., Nobesawa, S., Kanno, T., Yoshikawa, E., Futatsubashi, M., Ueda, Y., Okada, H., & Yamashita, Y. (2006). Cerebral hemodynamics evaluation by near-infrared time-resolved spectroscopy: correlation with simultaneous positron emission tomography measurements. *Neuroimage*, Vol. 29, pp.697–705. ISSN : 1053-8119
- Reinders, A.A., Nijenhuis, E.R., Paans, A.M., Korf, J., Willemsen, A.T., & den Boer, J.A. (2003) One brain, two selves. *Neuroimage*, Vol. 20, pp.2119–2125. ISSN : 1053-8119
- de Ruiter, M.B., Elzinga, B.M., & Phaf, R.H. (2006) Dissociation: cognitive capacity or dysfunction? *Journal of Trauma and Dissociation*, Vol. 7, pp. 115–134. ISSN :1529-9732
- Sar, V., Akyuz, G., Kugu, N., Ozturk, E., & Ertem-Vehid, H. (2006). Axis I dissociative disorder comorbidity in borderline personality disorder and reports of childhood trauma. *Journal of Clinical Psychiatry*, Vol. 67, pp.1583–1590. ISSN : 0160-6689
- Sato, T., Ito, M., Suto, T., Kameyama, M., Suda, M., Yamagishi, Y., Ohshima, A., Uehara, T., Fukuda, M., & Mikuni, M. (2007). Time courses of brain activation and their implications for function: a multichannel near-infrared spectroscopy study during finger tapping. *Neuroscience Research*, Vol.58, pp.297–304. ISSN : 0168-0102


- Schecklmann, M., Ehlis, A.C. Plichta, M.M., & Fallgatter, A.J. (2008). Functional near-infrared spectroscopy: a long-term reliable tool for measuring brain activity during verbal fluency. *Neuroimage*, Vol. 43, pp.147–155, ISSN : 1053-8119
- Shaw, P., Gilliam, M., Liverpool, M., Weddle, C., Malek, M., Sharp, W., Greenstein, D., Evans, A., Rapoport, J., & Giedd, J. (2010) Cortical development in typically developing children with symptoms of hyperactivity and impulsivity: support for a dimensional view of attention deficit hyperactivity disorder. *American Journal of Psychiatry*, Vol.168, pp. 143-151, ISSN : 0002-953X
- Spielberger, C.D., Krasner, S.S., & Solomon, E.P. (1988). The experience, expression and control of anger. In: *Health psychology: individual differences and stress*. Janisse M.P. (Eds.), pp. 89–108.; Springer-Verlag. ISBN : 0387966692, New York, U.S.A.
- Suda, M., Uehara, T., Fukuda, M., Sato, T., Kameyama, M., & Mikuni, M. (2010a). Dieting tendency and eating behavior problems in eating disorder correlate with right frontotemporal and left orbitofrontal cortex: a near-infrared spectroscopy study. *Journal of Psychiatry Research*, Vol. 44, pp.547–555, ISSN : 0022-3956
- Suda, M., Takei, Y., Aoyama, Y., Narita, K., Sato, T., Fukuda, M., & Mikuni, M. (2010b). Frontopolar activation during face-to-face conversation: an in situ study using near-infrared spectroscopy. *Neuropsychologia*, Vol. 48, pp. 441–447. ISSN : 0028-3932
- Suda, M., Fukuda, M., Sato, T., Iwata, S., Song, M., Kameyama, M., & Mikuni, M. (2009). Subjective feeling of psychological fatigue is related to decreased reactivity in ventrolateral prefrontal cortex. *Brain Research*, Vol. 1252, pp.152–160. ISSN : 0006-8993
- Suda, M., Sato, T., Kameyama, M., Ito, M., Suto, T., Yamagishi, Y., Uehara, T., Fukuda, M., & Mikuni, M. (2008). Decreased cortical reactivity underlies subjective daytime light sleepiness in healthy subjects: a multichannel near-infrared spectroscopy study. *Neuroscience Research*, Vol. 60, pp.319–326. ISSN : 0168-0102
- Suto, T., Fukuda, M., Ito, M., Uehara, T., & Mikuni, M. (2005). Multichannel near-infrared spectroscopy in depression and schizophrenia: cognitive brain activation study. *Biological Psychiatry*, Vol. 55, pp.501–511. ISSN : 0006-3223
- Takaishi, K. (2000). I-2 .Recent youth culture. In: *When university students seek counseling*. Sugihara, Y., Takaishi, K., and Kobayashi T. (Eds), pp18–37, Minervashobo, ISBN : 4-623-03133-0, Kyoto, Japan ( in Japanese)
- Takizawa, R., Kasai, K., Kawakubo, Y., Marumo, K., Kawasaki, S., Yamasue, H., & Fukuda, M. (2008). Reduced frontopolar activation during verbal fluency task in schizophrenia: a multichannel near-infrared spectroscopy study. *Schizophrenia Research*, Vol. 99, pp.250–262. ISSN : 0920-9964
- Tanabe, H. (2010). Measuring pathological dissociation with Dissociative Experiences Scale. IN *Manual for clinical evaluation and laboratory in Psychiatry*. *Japanese Journal of Clinical Psychiatry*, Vol.39 supplement, pp.303–320, ISSN :09342-12 (in Japanese).
- Toronov, V., Webb, A., Choi, J.H., Wolf, M., Michalos, A., Gratton, E., & Hueber, D. (2001). Investigation of human brain hemodynamics by simultaneous near-infrared spectroscopy and functional magnetic resonance imaging. *Medical Physics*, Vol.28, pp.521–527. ISSN :



- Uehara, T., Ishige, Y., Suda, M., & Sachdev, P. (2011). Gender differences in frontal activation while perceiving pathologically thin female body forms. *Journal of Behavioral and Brain Science*, in press, ISSN : 2160-5874
- Uehara, T., Fukuda, M., Suda, M., Kameyama, M., Yamagishi, Y., & Mikuni, M. (2007). Cerebral blood volume changes inpatients with eating disorders during word fluency using multi-channel near infrared spectroscopy. *Eating and Weight Disorders*, Vol. 12, pp.183-190, ISSN : 1124-4909

IntechOpen

IntechOpen



## **Psychiatric Disorders - Worldwide Advances**

Edited by Dr. Toru Uehara

ISBN 978-953-307-833-5

Hard cover, 336 pages

**Publisher** InTech

**Published online** 03, October, 2011

**Published in print edition** October, 2011

A psychiatric disorder is defined as any complex condition that involves the impairment of cognitive, emotional, or behavioral functioning. Aside from knowing the physical organic factors, its causal pathology has remained a mystery. Regarding recent advances in psychiatry and neurosciences, psychiatric disorders have been closely associated with socio-cultural, psychological, biochemical, epigenetic or neural-networking factors. A need for diverse approaches or support strategies is present, which should serve as common knowledge, empathetic views or useful skills for specialists in the field. This book contains multifarious and powerful papers from all over the world, addressing themes such as the neurosciences, psychosocial interventions, medical factors, possible vulnerability and traumatic events. Doubtlessly, this book will be fruitful for future development and collaboration in "world psychiatry".

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Toru Uehara, Yoko Ishige and Masashi Suda (2011). Dissociative Tendency, Anger Expression, and Frontal Activation During a Verbal Fluency Task, *Psychiatric Disorders - Worldwide Advances*, Dr. Toru Uehara (Ed.), ISBN: 978-953-307-833-5, InTech, Available from: <http://www.intechopen.com/books/psychiatric-disorders-worldwide-advances/dissociative-tendency-anger-expression-and-frontal-activation-during-a-verbal-fluency-task>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen