

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Entropy Generation in Viscoelastic Fluid Over a Stretching Surface

Saouli Salah and Aïboud Soraya
*University Kasdi Merbah, Ouargla,
Algeria*

1. Introduction

Due to the increasing importance in processing industries and elsewhere when materials whose flow behavior cannot be characterized by Newtonian relationships, a new stage in the evolution of fluid dynamics theory is in progress. An intensive effort, both theoretical and experimental, has been devoted to problems of non-Newtonian fluids. The study of MHD flow of viscoelastic fluids over a continuously moving surface has wide range of applications in technological and manufacturing processes in industries. This concerns the production of synthetic sheets, aerodynamic extrusion of plastic sheets, cooling of metallic plates, etc.

(Crane, 1970) considered the laminar boundary layer flow of a Newtonian fluid caused by a flat elastic sheet whose velocity varies linearly with the distance from the fixed point of the sheet. (Chang, 1989; Rajagopal et al., 1984) presented an analysis on flow of viscoelastic fluid over stretching sheet. Heat transfer cases of these studies have been considered by (Dandapat & Gupta, 1989, Vajravelu & Rollins, 1991), while flow of viscoelastic fluid over a stretching surface under the influence of uniform magnetic field has been investigated by (Andersson, 1992).

Thereafter, a series of studies on heat transfer effects on viscoelastic fluid have been made by many authors under different physical situations including (Abel et al., 2002, Bhattacharya et al., 1998, Datti et al., 2004, Idrees & Abel, 1996, Lawrence & Rao, 1992, Prasad et al., 2000, 2002). (Khan & Sanjayanand, 2005) have derived similarity solution of viscoelastic boundary layer flow and heat transfer over an exponential stretching surface.

(Cortell, 2006) have studied flow and heat transfer of a viscoelastic fluid over stretching surface considering both constant sheet temperature and prescribed sheet temperature. (Abel et al., 2007) carried out a study of viscoelastic boundary layer flow and heat transfer over a stretching surface in the presence of non-uniform heat source and viscous dissipation considering prescribed surface temperature and prescribed surface heat flux.

(Khan, 2006) studied the case of the boundary layer problem on heat transfer in a viscoelastic boundary layer fluid flow over a non-isothermal porous sheet, taking into account the effect a continuous suction/blowing of the fluid, through the porous boundary. The effects of a transverse magnetic field and electric field on momentum and heat transfer characteristics in viscoelastic fluid over a stretching sheet taking into account viscous dissipation and ohmic dissipation is presented by (Abel et al., 2008). (Hsiao, 2007) studied

the conjugate heat transfer of mixed convection in the presence of radiative and viscous dissipation in viscoelastic fluid past a stretching sheet. The case of unsteady magnetohydrodynamic was carried out by (Abbas et al., 2008). Using Kummer's functions, (Singh, 2008) carried out the study of heat source and radiation effects on magnetohydrodynamics flow of a viscoelastic fluid past a stretching sheet with prescribed power law surface heat flux. The effects of non-uniform heat source, viscous dissipation and thermal radiation on the flow and heat transfer in a viscoelastic fluid over a stretching surface was considered in (Prasad et al., 2010). The case of the heat transfer in magnetohydrodynamics flow of viscoelastic fluids over stretching sheet in the case of variable thermal conductivity and in the presence of non-uniform heat source and radiation is reported in (Abel & Mahesha, 2008). Using the homotopy analysis, (Hayat et al., 2008) looked at the hydrodynamic of three dimensional flow of viscoelastic fluid over a stretching surface. The investigation of biomagnetic flow of a non-Newtonian viscoelastic fluid over a stretching sheet under the influence of an applied magnetic field is done by (Misra & Shit, 2009). (Subhas et al., 2009) analysed the momentum and heat transfer characteristics in a hydromagnetic flow of viscoelastic liquid over a stretching sheet with non-uniform heat source. (Nandeppanavar et al., 2010) analysed the flow and heat transfer characteristics in a viscoelastic fluid flow in porous medium over a stretching surface with surface prescribed temperature and surface prescribed heat flux and including the effects of viscous dissipation. (Chen, 2010) studied the magneto-hydrodynamic flow and heat transfer characteristics viscoelastic fluid past a stretching surface, taking into account the effects of Joule and viscous dissipation, internal heat generation/absorption, work done due to deformation and thermal radiation. (Nandeppanavar et al., 2011) considered the heat transfer in viscoelastic boundary layer flow over a stretching sheet with thermal radiation and non-uniform heat source/sink in the presence of a magnetic field

Although the forgoing research works have covered a wide range of problems involving the flow and heat transfer of viscoelastic fluid over stretching surface they have been restricted, from thermodynamic point of view, to only the first law analysis. The contemporary trend in the field of heat transfer and thermal design is the second law of thermodynamics analysis and its related concept of entropy generation minimization.

Entropy generation is closely associated with thermodynamic irreversibility, which is encountered in all heat transfer processes. Different sources are responsible for generation of entropy such as heat transfer and viscous dissipation (Bejan, 1979, 1982). The analysis of entropy generation rate in a circular duct with imposed heat flux at the wall and its extension to determine the optimum Reynolds number as function of the Prandtl number and the duty parameter were presented by (Bejan, 1979, 1996). (Sahin, 1998) introduced the second law analysis to a viscous fluid in circular duct with isothermal boundary conditions.

In another paper, (Sahin, 1999) presented the effect of variable viscosity on entropy generation rate for heated circular duct. A comparative study of entropy generation rate inside duct of different shapes and the determination of the optimum duct shape subjected to isothermal boundary condition were done by (Sahin, 1998). (Narusawa, 1998) gave an analytical and numerical analysis of the second law for flow and heat transfer inside a rectangular duct. In a more recent paper, (Mahmud & Fraser, 2002a, 2002b, 2003) applied the second law analysis to fundamental convective heat transfer problems and to non-Newtonian fluid flow through channel made of two parallel plates. The study of entropy generation in a falling liquid film along an inclined heated plate was carried out by (Saouli & Aïboud-Saouli, 2004). As far as the effect of a magnetic field on the entropy generation is

concerned, (Mahmud et al., 2003) studied the case of mixed convection in a channel. The effects of magnetic field and viscous dissipation on entropy generation in a falling film and channel were studied by (Aïboud-saouli et al., 2006, 2007). The application of the second law analysis of thermodynamics to viscoelastic magnetohydrodynamic flow over a stretching surface was carried out by (Aïboud & Saouli 2010a, 2010b).

The objective of this paper is to study the entropy generation in viscoelastic fluid over a stretching sheet with prescribed surface temperature in the presence of uniform transverse magnetic field.

2. Formulation of the problem

In two-dimensional Cartesian coordinate system (x, y) we consider magneto-convection, steady, laminar, electrically conduction, boundary layer flow of a viscoelastic fluid caused by a stretching surface in the presence of a uniform transverse magnetic field and a heat source. The x -axis is taken in the direction of the main flow along the plate and the y -axis is normal to the plate with velocity components u, v in these directions.

Under the usual boundary layer approximations, the flow is governed by the following equations:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} = 0 \quad (1)$$

$$u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} = \nu \frac{\partial^2 u}{\partial y^2} - k_0 \left(u \frac{\partial^3 u}{\partial x \partial y^2} + v \frac{\partial^3 u}{\partial y^3} - \frac{\partial u}{\partial y} \frac{\partial^2 u}{\partial x \partial y} + \frac{\partial u}{\partial y} \frac{\partial^2 u}{\partial y^2} \right) - \frac{\sigma B_0^2}{\rho} u \quad (2)$$

The constant $k_0 = -\frac{\alpha_1}{\rho}$ is the viscoelastic parameter.

The boundary conditions are given by

$$y = 0, u = u_p = \lambda x, v = 0 \quad (3a)$$

$$y = \infty, u = 0, \frac{\partial u}{\partial y} = 0 \quad (3b)$$

The heat transfer governing boundary layer equation with temperature-dependent heat generation (absorption) is

$$\rho C_p \left(u \frac{\partial T}{\partial x} + v \frac{\partial T}{\partial y} \right) = k \frac{\partial^2 T}{\partial y^2} + Q(T - T_\infty) \quad (4)$$

The relevant boundary conditions are

$$y = 0, T = T_p = A \left(\frac{x}{l} \right)^2 + T_\infty \quad (5a)$$

$$y = \infty, T = T_\infty \quad (5b)$$

3. Analytical solution

The equation of continuity is satisfied if we choose a dimensionless stream function $\Psi(x, y)$ such that

$$u = \frac{\partial \Psi}{\partial y}, v = -\frac{\partial \Psi}{\partial x} \quad (6)$$

Introducing the similarity transformations

$$\eta = y\sqrt{\frac{\lambda}{\nu}}, \Psi(x, y) = x\sqrt{\nu\lambda}f(\eta) \quad (7)$$

Momentum equation (2) becomes

$$f'^2(\eta) - f(\eta)f''(\eta) = f'''(\eta) - \frac{k_0\lambda}{\nu} [2f'(\eta)f'''(\eta) - f(\eta)f^{IV}(\eta) - f''^2(\eta)] - Mn f'(\eta) \quad (8)$$

where $Mn = \frac{\sigma B_0^2}{\rho\lambda}$

Now let us seek a solution of Eq. (6) in the form

$$f'(\eta) = e^{-\alpha\eta} \quad (\alpha > 0) \quad (9)$$

which is satisfied by the following boundary conditions:

$$\eta = 0, f(0) = 0, f'(0) = 1 \quad (10a)$$

$$\eta = \infty, f'(\infty) = 0, f''(\infty) = 0 \quad (10b)$$

On substituting (7) into (6) and using boundary conditions (10a) and (10b) the velocity components take the form

$$u = \lambda x f'(\eta) \quad (11)$$

$$v = -\sqrt{\nu\lambda} f(\eta) \quad (12)$$

Where $k_1 = \frac{k_0\lambda}{\nu}$ is the viscoelastic parameter, and

$$\alpha = \sqrt{\frac{1 + Mn}{1 - k_1}} \quad (13)$$

Defining the dimensionless temperature

$$\Theta(\eta) = \frac{T - T_\infty}{T_p - T_\infty} \quad (14)$$

and using (9), (11), (12), Eq. (14) and the boundary conditions (5a) and (5b) can be written as

$$\Theta''(\eta) + \frac{\text{Pr}}{\alpha}(1 - e^{-a\eta})\Theta'(\eta) - (2\text{Pr}e^{-a\eta} - \beta)\Theta(\eta) = 0 \quad (15)$$

$$\eta = 0, \Theta(0) = 1 \quad (16a)$$

$$\eta = \infty, \Theta(\infty) = 0 \quad (16b)$$

Where $\text{Pr} = \frac{\mu C_P}{k}$ and $\beta = \frac{Q_V}{\lambda k}$ are respectively the Prandtl number and the heat/sink parameter.

Introducing the variable

$$\xi = \frac{\text{Pr}}{\alpha^2} e^{-a\eta} \quad (17)$$

And inserting (17) in (15) we obtain

$$\xi \Theta''(\xi) + \frac{\text{Pr}}{\alpha} \left(1 - \frac{\text{Pr}}{\alpha^2} + \xi\right) \Theta'(\xi) - \left(2 - \frac{\beta}{\alpha^2 \xi}\right) \Theta(\xi) = 0 \quad (18)$$

And (16a) and (16b) transform to

$$\xi = \frac{\text{Pr}}{\alpha^2}, \Theta\left(\frac{\text{Pr}}{\alpha^2}\right) = 1 \quad (19a)$$

$$\xi = 0, \Theta(0) = 0 \quad (19b)$$

The solution of Eq. (18) satisfying (19a) and (19b) is given by

$$\Theta(\xi) = \left(\frac{\alpha^2}{\text{Pr}} \xi\right)^{a+b} \frac{M(a+b-2, 2b+1, -\xi)}{M\left(a+b-2, 2b+1, -\frac{\text{Pr}}{\alpha^2}\right)} \quad (20)$$

The solution of (20) in terms of η is written as

$$\Theta(\xi) = e^{-\alpha(a+b)\eta} \frac{M\left(a+b-2, 2b+1, -\frac{\text{Pr}}{\alpha^2} e^{-a\eta}\right)}{M\left(a+b-2, 2b+1, -\frac{\text{Pr}}{\alpha^2}\right)} \quad (21)$$

where $a = \frac{\text{Pr}}{2\alpha^2}$, $b = \frac{\sqrt{\text{Pr}^2 - 4\alpha^2\beta}}{2\alpha^2}$ and $M\left(a+b-2, 2b+1, -\frac{\text{Pr}}{\alpha^2} e^{-a\eta}\right)$ is the Kummer's function.

4. Second law analysis

According to (Woods, 1975), the local volumetric rate of entropy generation in the presence of a magnetic field is given by

$$S_G = \frac{k}{T_\infty^2} \left[\left(\frac{\partial T}{\partial x} \right)^2 + \left(\frac{\partial T}{\partial y} \right)^2 \right] + \frac{\mu}{T_\infty} \left(\frac{\partial u}{\partial y} \right)^2 + \frac{\sigma B_0^2}{T_\infty} u^2 \quad (22)$$

Eq. (22) clearly shows contributions of three sources of entropy generation. The first term on the right-hand side of Eq. (22) is the entropy generation due to heat transfer across a finite temperature difference; the second term is the local entropy generation due to viscous dissipation, whereas the third term is the local entropy generation due to the effect of the magnetic field. It is appropriate to define dimensionless number for entropy generation rate N_s . This number is defined by dividing the local volumetric entropy generation rate S_G to a characteristic entropy generation rate S_{G0} . For prescribed boundary condition, the characteristic entropy generation rate is

$$S_{G0} = \frac{k(\Delta T)^2}{l^2 T_\infty^2} \quad (23)$$

therefore, the entropy generation number is

$$N_s = \frac{S_G}{S_{G0}} \quad (24)$$

using Eq. (9), (21) and (22), the entropy generation number is given by

$$N_s = \frac{4}{X^2} \Theta^2(\eta) + \text{Re}_l \Theta'^2(\eta) + \text{Re}_l \frac{Br}{\Omega} f''^2(\eta) + \frac{BrHa^2}{\Omega} f'^2(\eta) \quad (25)$$

where Re_l and Br are respectively the Reynolds number and the Brinkman number. Ω and Ha , are respectively the dimensionless temperature difference and the Hartman number. These number are given by the following relationships

$$\text{Re}_l = \frac{u_l l}{\nu}, \quad Br = \frac{\mu u_p^2}{k \Delta T}, \quad \Omega = \frac{\Delta T}{T_\infty}, \quad Ha = B_0 l \sqrt{\frac{\sigma}{\mu}} \quad (26)$$

5. Results and discussion

The flow and heat transfer in a viscoelastic fluid under the influence of a transverse uniform magnetic field has been solved analytically using Kummer's functions and analytic expressions of the velocity and temperature have been used to compute the entropy generation. Figs. 1 and 2 show the variations of the longitudinal velocity $f'(\eta)$ and the transverse velocity $f(\eta)$ as function of η for several values of magnetic parameter Mn . It can be observed that $f'(\eta)$ decreases with η and $f(\eta)$ increases with η asymptotically for Mn keeping constant. For a fixed position η , both $f'(\eta)$ and $f(\eta)$ decreases with Mn , thus the presence of the magnetic field decreases the momentum boundary layer thickness and increase the power needed to stretch the sheet.

The effects of the viscoelastic parameter k_1 on the longitudinal velocity $f'(\eta)$ and the transverse velocity $f(\eta)$ are illustrated on figs. 3 and 4. As it can be seen, for a fixed value of η , both $f'(\eta)$ and $f(\eta)$ decrease as viscoelastic parameter rises. This can be explained by

the fact that, as the viscoelastic parameter increases, the hydrodynamic boundary layer adheres strongly to the surface, which in turn retards the flow in the longitudinal and the transverse directions.




Fig. 1. Effect of the magnetic parameter on the longitudinal velocity.




Fig. 2. Effect of the magnetic parameter on the transverse velocity.




Fig. 3. Effect of the viscoelastic parameter on the longitudinal velocity.




Fig. 4. Effect of the viscoelastic parameter on the transverse velocity.

Fig. 5 depicts the temperature profiles $\Theta(\eta)$ as function of η for different values of the Prandtl number Pr . As it can be noticed, $\Theta(\eta)$ decreases with η whatever is the value of the Prandtl number, for a fixed value of η , the temperature $\Theta(\eta)$ decreases with an increase in Prandtl number which means that the thermal boundary layer is thinner for large Prandtl number.




Fig. 5. Effect of the Prandtl number on the temperature.

The temperature profiles $\Theta(\eta)$ as function of η for different values of the magnetic Mn are plotted in fig. 6. An increase in the magnetic parameter Mn results in an increase of the temperature; this is due to the fact that the thermal boundary layer increases with the magnetic parameter. Fig. 7 represents graphs of temperature profiles $\Theta(\eta)$ as function of η for various values of the heat source/sink parameter β . For fixed value of η , the temperature $\Theta(\eta)$ augments with the heat source/ sink parameter β . This is due to the fact that the increase of the heat source/sink parameter means an increase of the heat generated inside the boundary layer leading to higher temperature profile.

The influence of the magnetic parameter Mn on the entropy generation number N_s is shown on fig. 8. The entropy generation number N_s decreases with η for Mn keeping constant. For




Fig. 6. Effect of the magnetic parameter on the temperature.




Fig. 7. Effect of the heat source/sink parameter on the temperature.




Fig. 8. Effect of the magnetic parameter on the entropy generation number.

fixed value of η , the entropy generation number increases with the magnetic parameter, because the presence of the magnetic field creates more entropy in the fluid. Moreover, the stretching surface acts as a strong source of irreversibility.




Fig. 9. Effect of the Prandtl number on the entropy generation number.

Fig. 9 illustrates the effect of the Prandtl number Pr on the entropy generation number N_s . The entropy generation number is higher for higher Prandtl number near the surface, but as η increases, the entropy generation number shows different variation. This is due to the fact that according to fig. 6, the temperature profiles decrease sharply with the increase of the Prandtl number.




Fig. 10. Effect of the Reynolds number on the entropy generation number.

The influence of the Reynolds number Re_L on the entropy generation number is plotted on fig. 10. For a given value of η , the entropy generation number increases as the Reynolds number increases. The augmentation of the Reynolds number increases the contribution of the entropy generation number due to fluid friction and heat transfer in the boundary layer.




Fig. 11. Effect of the dimensionless group on the entropy generation number.

The effect of the dimensionless group parameter $Br\Omega^{-1}$ on the entropy generation number N_s is depicted in fig. 11. The dimensionless group determines the relative importance of viscous effect. For a given η , the entropy generation number is higher for higher dimensionless group. This is due to the fact that for higher dimensionless group, the entropy generation numbers due to the fluid friction increase.




Fig. 12. Effect of the Hartman number on the entropy generation number.

The effect of the Hartman number Ha on the entropy generation number N_s is plotted in fig. 12. For a given η , as the Hartman number increases, the entropy generation number increases. The entropy generation number is proportional to the square of Hartman number which is proportional to the magnetic field. The presence of the magnetic field creates additional entropy.

6. Conclusion

The velocity and temperature profiles are obtained analytically and used to compute the entropy generation number in viscoelastic magnetohydrodynamic flow over a stretching surface

The effects of the magnetic parameter and the viscoelastic parameter on the longitudinal and transverse velocities are discussed. The influences of the Prandtl number, the magnetic parameter and the heat source/sink parameter on the temperature profiles are presented. As far as the entropy generation number is concerned, its dependence on the magnetic parameter, the Prandtl, the Reynolds, the Hartmann numbers and the dimensionless group are illustrated and analyzed.

From the results the following conclusions could be drawn:

- The velocities depend strongly on the magnetic and the viscoelastic parameters.
- The temperature varies significantly with the Prandtl number, the magnetic parameter and the heat source/sink parameter.
- The entropy generation increases with the increase of the Prandtl, the Reynolds, the Hartmann numbers and also with the magnetic parameter and the dimensionless group.
- The surface acts as a strong source of irreversibility.

7. Nomenclature

A	constant, K
\vec{B}_0	uniform magnetic field strength, Wb.m ⁻²
Br	Brinkman number, $Br = \frac{\mu u_0^2}{k\Delta T}$
C_p	specific heat of the fluid, J.kg ⁻¹ .K ⁻¹
f	dimensionless function
Ha	Hartman number $Ha = B_0 l \sqrt{\frac{\sigma}{\mu}}$
k	thermal conductivity of the fluid, W.m ⁻¹ .K ⁻¹
k_1	viscoelastic parameter, $k_1 = \frac{k_0 \lambda}{\nu}$
k_0	viscoelastic parameter, m ²
l	characteristic length, m
M	Kummer's function
Mn	magnetic parameter, $Mn = \frac{\sigma B_0^2}{\rho \lambda}$
N_s	entropy generation number, $N_s = \frac{S_g}{S_{g0}}$
Pr	Prandtl number, $Pr = \frac{\mu C_p}{k}$
Q	rate of internal heat generation or absorption, W.m ⁻³ .K ⁻¹

Re_l	Reynolds number based on the characteristic length, $Re_l = \frac{u_l l}{\nu}$
S_G	local volumetric rate of entropy generation, $W.m^{-3}.K^{-1}$
S_{G0}	characteristic volumetric rate of entropy generation, $W.m^{-3}.K^{-1}$
T	temperature, K
u	axial velocity, $m.s^{-1}$
u_l	plate velocity based on the characteristic length, $m.s^{-1}$
u_p	plate velocity, $m.s^{-1}$
ν	transverse velocity, $m.s^{-1}$
x	axial distance, m
X	dimensionless axial distance, $X = \frac{x}{l}$
y	transverse distance, m
α	positive constant
β	heat source/sink parameter, $\beta = \frac{Q\nu}{\lambda k}$
λ	proportional constant, s^{-1}
η	dimensionless variable, $\eta = y\sqrt{\frac{\lambda}{\nu}}$
ξ	dimensionless variable, $\xi = \frac{Pr}{\alpha^2} e^{-\alpha\eta}$
μ	dynamic viscosity of the fluid, $kg.m^{-1}.s^{-1}$
ν	kinematic viscosity of the fluid, $m^2.s^{-1}$
ΔT	temperature difference, $\Delta T = T_p - T_\infty$
Ω	dimensionless temperature difference, $\Omega = \frac{\Delta T}{T_\infty}$
Θ	dimensionless temperature, $\Theta = \frac{T - T_\infty}{T_p - T_\infty}$
ρ	density of the fluid, $kg.m^{-3}$
σ	electric conductivity, $\Omega^{-1}.m^{-1}$
<i>subscripts</i>	
P	plate
∞	far from the sheet

8. References


- Abbas, Z.; Wang, Y.; Hayat, T. & Oberlack M. (2008). Hydromagnetic flow of a viscoelastic fluid due to the oscillatory stretching surface, *Int. J. Non-Linear Mech.*, 43, pp. 783-793, ISSN 0020-7462.
- Abel, M. S.; Khan, S. K. & Prasad, K. V. (2002). Study of viscoelastic fluid flow and heat transfer over stretching sheet with variable viscosity, *Int. J. Non-Linear Mech.* 37, pp. 81-88, ISSN 0020-7462.

- Abel, M. S.; Siddheshwar, P. G. & Nandeppanavar, M. M. (2007). Heat transfer in a viscoelastic boundary layer flow over a stretching sheet with viscous dissipation and non-uniform heat source, *Int. J. Heat Mass Transfer*, 50, pp. 960-966, ISSN 0017-9310.
- Abel, M. S. & Mahesha, N. (2008). Heat transfer in MHD viscoelastic fluid over a stretching sheet with variable thermal conductivity, non-uniform heat source and radiation, *Appl. Math. Modelling*, 32, pp. 1965-1983, ISSN 0307-904X.
- Abel, M. S.; Sanjayanand, E. & Nandeppanavar, M. M. (2008). Viscoelastic MHD flow and heat transfer over a stretching sheet with viscous and ohmic dissipation, *Comm. Nonlinear. Sci. and Num. Simulation*, 13, pp. 1808-1821, ISSN 1007-5704.
- Abel, M. S. & Nandeppanavar, M. M. (2009). Heat transfer in MHD viscoelastic boundary layer flow over a stretching sheet with non-uniform heat source/sink, *Comm. NonLinear Sci. and Num. Simu.*, 14, pp. 2120-2131, ISSN 1007-5704.
- Aïboud-Saouli, S.; Saouli, S.; Settou, N. & Meza, N. (2006). Thermodynamic analysis of gravity-driven liquid film along an inclined heated plate with hydromagnetic and viscous dissipation effects, *Entropy*, 8, pp. 188-199, ISSN 1099-4300.
- Aïboud-Saouli, S.; Settou, N.; Saouli, S. & Meza, N. (2007). Second-law analysis of laminar fluid flow in a heated channel with hydromagnetic and viscous dissipation effects, *Applied Energy*, 84, pp. 279-289, ISSN 0306-2619.
- Aïboud, S., & Saouli, S. (2010). Second law analysis of viscoelastic fluid over a stretching sheet subject to a transverse magnetic field with heat and mass transfer, *Entropy*, 12, pp. 1867-1884, ISSN 1099-4300.
- Aïboud, S., & Saouli, S. (2010). Entropy analysis for viscoelastic magnetohydrodynamics flow over a stretching surface, *Int. J. Non-Linear Mech.* 45, pp. 482-489, ISSN 0020-7462.
- Andersson, H. D. (1992). MHD flows of a viscoelastic fluid past a stretching surface, *Acta Mech.*, 95, pp. 227-230, ISSN 0001-5970.
- Bejan, A. (1982). Second-law analysis in heat transfer and thermal design, *Adv. Heat Transfer*, 15, pp. 1-58, ISBN 10-0-12-020021.
- Bejan, A. (1996). Entropy generation minimization. *CRC Press, Boca Raton*, ISBN-10:9780849396519, New York, USA.
- Bejan, A. (1979). A study of entropy generation in fundamental convective heat transfer, *J. Heat Transfer*, 101, pp. 718-725, ISSN 0022-1481.
- Bhattacharya, S.; Pal, A. & Gupta, A. S. (1998). Heat transfer in the flow of a viscoelastic fluid over a stretching surface, *Heat mass Transfer*, 34, pp. 41-45, ISSN 0947-7411.
- Chang, W. D. (1989). The non-uniqueness of the flow of viscoelastic fluid over a stretching sheet, *Q. Appl. Math.*, 47, pp. 365-366, ISSN 0033-569X.
- Chen, C. H. (2010). On the analytic solution of MHD flow and heat transfer for two types of viscoelastic fluid over a stretching sheet with energy dissipation internal heat source and thermal radiation, *Int. J. Heat mass Transfer*, 53, pp. 4264-4273, ISSN 0017-9310.
- Cortell, R. (2006). A note on flow and heat transfer of a viscoelastic fluid over a stretching sheet, *Int. J. Non-Linear Mech.*, 41, pp. 78-85, ISSN 0020-7462.
- Crane, L. J. (1970). Flow past a stretching sheet, *Z. Angew. Math. Phys.*, 21, pp. 645-647, ISSN 0044-2275.

- Dandapat, B. S. & Gupta, A. S. (1998). Flow and heat transfer in a viscoelastic fluid over a stretching sheet, *Int. J. Non-Linear Mech.*, 24, pp. 215-219, ISSN 0020-7462 .
- Datti, P. S.; Prasad, K. V.; Abel, M. S. & Joshi, A. (2004). MHD viscoelastic fluid flow over a non-isothermal stretching sheet, *Int. J. Eng. Sci.*, 42, pp. 935-946, ISSN 0020-7225.
- Hayat, T.; Sajid, M. & Pop, I. (2008). Three-dimensional flow over a stretching sheet in a viscoelastic fluid, *Nonlinear. Ana. Real World Appl.*, 9, pp. 1811-1822, ISSN 1468-1218.
- Hsiao, K. L. (2007). Conjugate heat transfer of magnetic mixed convection with viscous dissipation effects for second-grade viscoelastic fluid past a stretching sheet, *Appl. Therm. Eng.*, 27, pp. 1895-1903, ISSN 1359-4311.
- Idrees, M.K. & Abel, M. S. (1996). Viscoelastic flow past a stretching sheet in porous media and heat transfer with internal heat source, *Indian J. Theor. Phys.*, 44, pp. 233-244, ISSN 0019-5693.
- Khan, S. K. & Sanjayanand, E. (2005). Viscoelastic boundary layer flow and heat transfer over an exponential stretching sheet, *Int. J. Heat Mass Transfer*, 48, pp. 1534-1542, ISSN 0017-9310.
- Khan, S. K. (2006). Heat transfer in a viscoelastic fluid over a stretching surface with source/sink, suction/blowing and radiation, *Int. J. Heat Mass Transfer*, 49, pp. 628-639, ISSN 0017-9310.
- Lawrence, P.S. & Rao, B. N. (1992). Heat transfer in the flow of viscoelastic fluid over stretching sheet, *Acta Mech.*, 93, pp. 53-61, ISSN 0001-5970.
- Mahmud, S. & Fraser, R. A. (2003). The second law analysis in fundamental convective heat transfer problems, *Int. J. Therm. Sci.*, 42, pp. 177-186, ISSN 1290-0729.
- Mahmud, S. & Fraser, R. A. (2002). Thermodynamic analysis of flow and heat transfer inside channel with two parallel plates, *Exergy* 2, pp. 140-146, ISSN 1164-0235.
- Mahmud, S. & Fraser, R. A. (2002). Inherent irreversibility of channel and pipe flows for non-Newtonian fluids, *Int. Comm. Heat Mass Transfer*, 29, pp. 577-587, ISSN 0947-7411.
- Mahmud, S.; Tasnim, S. H. & Mamun, H. A. A. (2003). Thermodynamic analysis of mixed convection in a channel with transverse hydromagnetic effect, *In. J. Therm. Sci.*, 42, pp. 731-740, ISSN 1290-0729.
- Misra, J. C., & Shit, G. C. (2009). Biomagnetic viscoelastic fluid flow over a stretching sheet, *Appl. Math. And Compu*, 210, pp. 350-361, ISSN 0096-3003.
- Nandeppanavar, M. M.; Abel, M. S. & Vajravelu, K. (2010). Flow and heat transfer characteristics of a viscoelastic fluid in a porous medium over an impermeable stretching sheet with viscous dissipation, *Int. J. Heat Mass Transfer*, 53, pp. 4707-4713, ISSN 0017-9310.
- Nandeppanavar, M. M.; Vajravelu, K. & Abel, M. S. (2011). Heat transfer in MHD viscoelastic boundary layer flow over a stretching sheet with thermal radiation and non-uniform heat source/sink, *Comm. Nonlinear. Sci. and Num. Simulation*, 16, pp. 3578-3590, ISSN 1007-5704.
- Narusawa, U. (1998). The second-law analysis of mixed convection in rectangular ducts, *Heat Mass Transfer*, 37, pp. 197-203, ISSN 0947-7411.
- Prasad, K.V.; Abel, M. S.; Khan, S.K. & Datti, P. S. (2002). Non-Darcy forced convective heat transfer in a viscoelastic fluid flow over a non-Isothermal stretching sheet, *J. Porous Media*, 5, pp. 41-47, ISSN 1091-028X.

- Prasad, K.V.; Abel, M. S. & Khan, S. K. (2000). Momentum and heat transfer in viscoelastic fluid flow in a porous medium over a non-isothermal stretching sheet, *Int. J. Numer. Method Heat flow*, 10, pp. 786-801, ISSN 0961-5539.
- Prasad, K.V.; Pal, D.; Umesh, V. & Prasanna Rao, N. S. (2010). The effect of variable viscosity on MHD viscoelastic fluid flow and heat transfer over a stretching sheet, *Comm. Nonlinear Sci. and Num. Simulation*, 15, pp. 331-344, ISSN 1007-5704.
- Rajagopal, K. R. ; Na, T. Y. & Gupta, A. S. (1984). Flow of a viscoelastic sheet over a stretching sheet, *Rheo. Acta*, 23, pp. 213-221, ISSN 0035-4511.
- Sahin, A. Z. (1998). Second law analysis of laminar viscous flow through a duct subjected to constant wall temperature, *J. Heat Transfer*, 120, pp. 76-83, ISSN 0022-1481.
- Sahin, A. Z. (1999). Effect of variable viscosity on the entropy generation and pumping power in a laminar fluid flow through a duct subjected to constant heat flux, *Heat Mass Transfer*, 35, pp. 499-506, ISSN 0947-7411.
- Sahin, A. Z. (1998). A second law comparison for optimum shape of duct subjected to constant wall temperature and laminar flow, *Heat Mass Transfer*, 33, pp. 425-430, ISSN 0947-7411.
- Saouli, S. & Aïboud-Saouli, S. (2004). Second law analysis of laminar falling liquid film along an inclined heated plate, *Int. Comm. Heat Mass Transfer*, 31, pp. 879-886, ISSN 0947-7411.
- Singh, A. K. (2008). Heat source and radiation effects on magneto-convection flow of a viscoelastic fluid past a stretching sheet: Analysis with Kummer's functions, *Int. Comm. Heat Mass Transfer*, 35, pp. 637-642, ISSN 0947-7411.
- Vajravelu, K. & Rollins, D. (1991). Heat transfer in a viscoelastic fluid over a stretching sheet, *J. Math. Anal. Appl.* 158, pp. 241-255, ISSN 0022-247X.
- Woods, L. C. (1975). Thermodynamics of fluid systems, *Oxford University Press*, ISBN-10: 0198561806, Oxford, UK.

IntechOpen



Thermodynamics - Systems in Equilibrium and Non-Equilibrium

Edited by Dr. Juan Carlos Moreno Piraján

ISBN 978-953-307-283-8

Hard cover, 306 pages

Publisher InTech

Published online 10, October, 2011

Published in print edition October, 2011

Thermodynamics is one of the most exciting branches of physical chemistry which has greatly contributed to the modern science. Being concentrated on a wide range of applications of thermodynamics, this book gathers a series of contributions by the finest scientists in the world, gathered in an orderly manner. It can be used in post-graduate courses for students and as a reference book, as it is written in a language pleasing to the reader. It can also serve as a reference material for researchers to whom the thermodynamics is one of the area of interest.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Saouli Salah and Aïboud Soraya (2011). Entropy Generation in Viscoelastic Fluid Over a Stretching Surface, Thermodynamics - Systems in Equilibrium and Non-Equilibrium, Dr. Juan Carlos Moreno Piraján (Ed.), ISBN: 978-953-307-283-8, InTech, Available from: <http://www.intechopen.com/books/thermodynamics-systems-in-equilibrium-and-non-equilibrium/entropy-generation-in-viscoelastic-fluid-over-a-stretching-surface>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen