

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Getting Lipids for Biodiesel Production from Oleaginous Fungi

Maddalena Rossi, Alberto Amaretti, Stefano Raimondi and Alan Leonardi
*University of Modena and Reggio Emilia,
 Italy*

1. Introduction

Biomass-based biofuel production represents a pivotal approach to face high energy prices and potential depletion of crude oils reservoirs, to reduce greenhouse gas emissions, and to enhance a sustainable economy (Zinoviev et al., 2010). Microbial lipids can represent a valuable alternative feedstock for biodiesel production, and a potential solution for a bio-based economy.

Nowadays, the production of biodiesel is based mostly on plant oils, even though animal fats, and algal oils can also be used. In particular, soybean, rapeseed, and palm oils are adopted as the major feedstock for biodiesel production. They are produced on agricultural land, opening the debate on the impact of the expansion of bioenergy crop cultures, which displace land from food production. Furthermore, their price restricts the large-scale development of biodiesel to some extent.

In order to meet the increasing demand of biodiesel production, other oil sources have been explored. Recently, the development of processes to produce single cell oil (SCO) by using heterotrophic oleaginous microorganisms has triggered significant attention (Azocar et al., 2010). These organisms accumulate lipids, mostly consisting of triacylglycerols (TAG), that form the storage fraction of the cell. The occurrence of TAG as reserve compounds is widespread among all eukaryotic organisms such as fungi, plants and animals, whereas it has only rarely been described in bacteria (Meng et al., 2009). In fact, bacteria generally accumulate polyhydroxyalkanoates as storage compound and only few bacterial species, belonging to the actinobacterial genera *Mycobacterium*, *Streptomyces*, *Rhodococcus* and *Nocardia* produce relevant amounts of lipids (Alvarez & Steinbuchel, 2002).

Among heterotrophic microorganisms, oleaginous fungi, including both molds and yeasts, are increasingly been reported as good TAG producers. This chapter will focus on current knowledge advances in their metabolism, physiology, and in the result achieved in strain improvement, process engineering and raw material exploitation.

2. Ecology of oleaginous fungi

Oleaginous microorganisms are able to accumulate lipids above the 20% of their biomass, on dry basis. Several species of yeasts and filamentous fungi are regarded as oleaginous, since they have the capability to synthesize and accumulate high amounts of TAG within their cells, up to 70% of the biomass weight. These lipids have similar composition and energy

value to plant and animal oils, but their production do not compete for food resources, in particular if it is based on inexpensive carbon sources, such as raw materials, by-products, and surplus. Furthermore, fungal SCO have a short process cycle, and their production is not subjected to seasonal and cyclical weather variations.

The study of oleaginous yeasts has a long history: their ability to accumulate lipids has been known from the 70s, but only in the last years the attention has been focused on exploitation of SCO for biodiesel production. The yeasts represent a part of the microbiota in all natural ecosystems, such as soils, freshwaters and marine waters, from the ocean surface to the deep sea. Widely distributed in the natural environment, they colonize also more extreme environments, such as low temperatures, low oxygen availabilities, and oceanic waters (Butinar et al., 2007). Approximately 1500 species of yeasts belonging to over 100 genera have been described so far (Satyanarayana & Kunze, 2010). Although the vast majority of yeasts are beneficial to human life, only a few are opportunistic human pathogens. As a whole, they play a pivotal role in the food chain, and in the carbon, nitrogen and sulphur cycles. Among the huge number of species that have been described, only 30 are able to accumulate more than 25% of their dry weight as lipids (Beopoulos et al., 2009b).

Basidiomycetous yeasts strongly prevail among oleaginous yeasts, representing most of all the strains identified as lipid producers, even though some important oleaginous species have been identified among Ascomycota as well (e.g. *Yarrowia lipolytica*).

The most deeply investigated oleaginous yeasts belong to the genera *Yarrowia*, *Candida*, *Rhodotorula*, *Rhodospiridium*, *Cryptococcus*, and *Lipomyces* (Ageitos et al., 2011; Li et al., 2008; Rossi et al., 2009). *Yarrowia lipolytica*, previously referred to as *Candida lipolytica*, is a good candidate for single-cell oil production (Beopoulos et al., 2009a; Beopoulos et al., 2009b). *Yarrowia* are hemiascomycetous dimorphic fungi that belong to the order *Saccharomycetales*. They are able to degrade hydrophobic substrates such as n-paraffins and oils very efficiently and this physiological feature prompted the scientific community to explore several biotechnological applications (Bankar et al., 2009). The common habitats of these fungi are oil-polluted environments and foods such as cheese, yogurt, kefir, shoyu, meat, and poultry products. Despite *Y. Lipolytica* is distantly related to the conventional yeast *Saccharomyces cerevisiae*, the genome displays an expansion of protein families and genes involved in hydrophobic substrate (such as alkanes and lipids) utilization. Wild-type strains accumulate up to 38% of dry weight (DW) as lipids. Albeit the levels are lower than those of other oleaginous yeasts, it became a model organism because it can be subjected to genetic and metabolic engineering, having been developed a reliable and versatile system for disruption, cloning and expression of target genes.

Within the *Candida* genus, *Candida curvata* (Holdsworth & Ratledge, 1991) also referred as *Apiotrichum curvatum* and *Candida freyschussii* (Amaretti et al., 2011) synthesize and store significant amount of lipids. *Candida* comprises an extremely heterogeneous group of Ascomycota that can all grow with yeast morphology, classified in 150 heterogeneous species, among which only a minority have been implicated in human diseases, since approximately 65% of *Candida* species are unable to grow at 37°C, then they can not be successful pathogens or commensals of humans (Calderone, 2002). Therefore, most of the species can be exploited for biotechnological applications, despite of unwarranted negative public perceptions.

Lipomyces spp. present a great propensity to accumulate triacylglycerols. This genus belongs to the *Saccharomycetales* order and represents a unique branch in the evolution of the

ascomycetes (van der Walt, 1992). *Lipomyces* are true soil inhabitants and have a worldwide distribution. The oleaginous species *Lipomyces starkeyi* has the capability to accumulate over 70% of its cell biomass as lipid under defined culture conditions, and can produce lipid on xylose, ethanol, and L-arabinose, or using a mixture of glucose and xylose (Zhao et al., 2008), as well as other wastes (Angerbauer et al., 2008).

Cryptococcus curvatus is a yeast with industrial potential as single-cell oil because it can grow and accumulate lipid on a very broad range of substrates. It requires minimal nutrients for growth, accumulating up to 60% of its cellular dry weight (DW) as intracellular lipid (Meesters et al., 1996; Zhang et al., 2011). Yeasts of the *Cryptococcus* genus are widely distributed in nature and may be isolated from various substrates such as air, soil, bird excreta, water, animal surfaces and mucosae, leaves, flowers, and decomposing wood. Most species are considered as free-living (non-symbiotic) and only a few have medical importance being responsible for disease in man and animals (*C. neoformans* and *C. gattii*). *C. curvatus* is recognized as an opportunistic pathogen of animals, including humans (Findley et al., 2009).

Species belonging to the genus *Rhodospiridium*, and to its asexual counterpart *Rhodotorula*, have been claimed as oleaginous yeasts. They belong to one of the three main lineages of the Basidiomycota, the Pucciniomycotina. *Rhodotorula* is a common environmental inhabitant. The synthesis of different commercially important natural carotenoids by yeast species belonging to the genus *Rhodotorula* has led to consider these microorganisms as a potential pigment sources. Within this genus, the mesophilic red yeast *Rhodotorula glutinis* is able to synthesize and store lipids also growing on glycerol, whereas the psychrophilic species *Rhodotorula glacialis*, that are not red yeasts, accumulates lipids in a range of temperature between -3 and 20°C (Amaretti et al., 2010). The red yeast *Rhodospiridium toruloides* is an oleaginous mesophilic species. *Rhodospiridium* are able to carry out a number of diverse biochemical reactions such as biodegradation of epoxides, biphenyls and oxiranes (Smit, 2004), biosynthesis of carotenoids (de Miguel et al., 1997) and other types of biotransformations, but a major biotechnological exploitation is associated to their ability to convert glycerol and lignocellulosic biowastes into lipids (Hu et al., 2009; Yu et al., 2011). Among the oily yeasts, two novel species of the anamorphic basidiomycetous genus *Trichosporon* have been recently identified (*T. cacaoliposimilis* and *T. oleaginosus*) (Gujjari et al., 2011), despite lipid accumulation has not yet explored in the perspective of biodiesel production. *Trichosporon* are basidiomycetous yeasts widely distributed in nature, consisting of soil- and water-associated species, predominantly found in environmental substrates, such as decomposing wood. They present distinct morphological characteristics of budding yeast cells and true mycelia that disarticulate to form arthroconidia. Some species are causative agents of diseases in man and cattle. They can occasionally belong to the gastrointestinal microbiota of humans as well as transiently colonize the skin and respiratory tract.

Exploitation of oleaginous filamentous fungi for biodiesel production has a more recent history, which, with few exceptions, derives from studies focused to poly-unsaturated fatty acid production (PUFA), such as arachidonic acid and γ -linolenic acid. The most relevant example of this biotechnological application is represented by exploitation of *Mortierella alpina* to produce oils containing n-1, n-3, n-4, n-6, n-7, and n-9 PUFAs (Sakuradani et al., 2009). Among the major lipid producers there is *Mucor circinelloides*, a zygomycete fungus, which is emerging as opportunistic pathogen in immunocompromised patients (Li et al., 2011). *M. circinelloides* has been used for the first

commercial production of microbial lipids (Ratledge, 2004). Lipid accumulation in *M. circinelloides* has been extensively studied (Wynn et al., 2001), and its TAG have been proposed as feedstock for producing biodiesel by direct transformation of its lipids (Vicente et al., 2009). *M. circinelloides* represents an outstanding model within the *Zygomycota* phylum, based on the availability of an efficient transformation procedure (Gutierrez et al., 2011) and on the whole sequence of genome (<http://genome.jgi-psf.org/Mucci2/Mucci2.home.html>). Also the phylogenetically related *Umbelopsis isabellina* has emerging as a promising species to convert biomass residues to biodiesel precursors (Meeuwse et al., 2011a). To the best of our knowledge, limited are the attempts to get lipids with *Aspergillus oryzae* that, conversely, is extensively studied as lipase producer to carry out transesterification of TAG (Adachi et al., 2011).

3. Biochemistry of lipid accumulation

Lipid accumulation in oleaginous yeasts and molds has been demonstrated to occur when a nutrient in the medium (e.g. the nitrogen or the phosphorus source) becomes limited and the carbon source is present in excess. Nitrogen limitation is the most efficient condition for inducing lipogenesis. During the growth phase, nitrogen is necessary for the synthesis of proteins and nucleic acids, while the carbon flux is distributed among energetic and anabolic processes yielding carbohydrates, lipids, nucleic acids and proteins. When nitrogen gets limited, the growth rate slows down and the synthesis of proteins and nucleic acids tends to cease. In non-oleaginous species, the carbon excess remains unutilized or is converted into storage polysaccharides, while, in oleaginous species, it is preferentially channeled toward lipid synthesis, leading to the accumulation of TAG within intracellular lipid bodies (Ratledge & Wynn, 2002; Granger et al., 1993).

The biochemical pathway of lipid biosynthesis is not very different among eukaryotic organisms and does not differ in oleaginous and non-oleaginous fungi. The ability to accumulate high amounts of lipid depends mostly on the regulation the biosynthetic pathway and the supply of the precursors (i.e. acetyl-CoA, malonyl-CoA, and glycerol-3-phosphate) and the cofactor NADPH.

Most information were obtained from the model yeast *Saccharomyces cerevisiae* (Kohlwein, 2010), that does not accumulate lipids, and *Yarrowia lipolytica*, that represent a model for bio-oil production and is suitable for genetic manipulation (Beopoulos et al., 2009b).

3.1 Fatty acids biosynthesis and modifications

De novo synthesis of fatty acids (FA), the first step of lipid accumulation, is carried out in the cytosol by fatty acids synthetase (FAS) complex. In yeasts, FAS bears phosphopantetheine transferase activity to activate its acyl carrier protein (ACP) by loading the coenzyme pantothenate. FAS is a multimer of 6 α and 6 β subunits encoded by *fas2* and *fas1*, respectively, each subunit containing four functional domains. Therefore, FAS consists in a $\alpha_6\beta_6$ molecular complex of 2.6 MDa with 48 functional centers that catalyze all reactions required for synthesis of fatty acids through cycles of multistep reactions. FAS firstly loads acetyl-CoA on its β -ketoacyl-ACP synthase (KS), then it exerts β -ketoacyl-ACP reductase (KR), β -hydroxyacyl-ACP dehydratase (DH), and enoyl-ACP reductase (EAR) activities. This set of reactions is repeated cyclically seven times to yield palmitoyl-ACP (Fig. 1) (Tehlivets et al., 2007).





Fig. 1. Reactions occurring sequentially in fatty acid synthetase: condensation of acyl-ACP and malonyl-ACP mediated by KS, NADPH-dependent reduction of the keto group to a hydroxyl group by means of KR, dehydration to create a double bond with DH and reduction of the double bond by means of EAR. R = H, CH₃(CH₂)_{2n}; n_{max}=7.

The biosynthesis of FA requires the constant supply of acetyl-CoA as initial biosynthetic unit and of malonyl-CoA as the elongation unit, supplying two carbons at each step. Non-oleaginous yeasts receive acetyl-CoA mostly from glycolysis. In oleaginous yeasts, acetyl-CoA is mostly provided by the cleavage in the cytosol of citrate, which accumulated as a consequence of nitrogen limitation (Ratledge, 2002) (Fig. 2). In fact, lipid accumulation by oleaginous fungi does not occur under balanced nutrient conditions.

In oleaginous yeasts, nitrogen limitation activates AMP-deaminase (Ratledge & Wynn, 2002), which supply ammonium to the nitrogen-starved cell. As a consequence, mitochondrial AMP concentration decreases, causing isocitrate dehydrogenase activity to drop. The TCA cycle is then blocked at the level of isocitrate, which accumulates and equilibrates with citrate through aconitase. Excess of citrate from TCA cycle is exported out of the mitochondrion via the malate/citrate antiport. Cytosolic ATP-citrate lyase (ACL) cleaves citrate to give oxaloacetate and acetyl-CoA (Fig. 2).

ACL represents one of the key enzymes that contribute to the oleaginous trait of yeasts, whereas its activity is negligible in non-oleaginous species. ACL is composed of two subunits, encoded by *ACL1* and *ACL2* and is negatively regulated by exogenous FA.

Malonyl-CoA is produced from acetyl-CoA by acetyl-CoA carboxylase (ACC) that condensate an acetyl-CoA unit with bicarbonate:



ACC is also a key enzyme in *de novo* FA synthesis, since *ACC1* mutants became FA auxotrophs or maintain low levels of ACC activity (Tehlivets et al., 2007). *ACC1* undergoes allosteric activation by citrate. Furthermore the transcription of *FAS1*, *FAS2*, and *ACC1* is coordinately regulated, being negatively regulated by FA.






Fig. 2. Lipid biosynthesis from excess of citrate as a consequence of nitrogen limitation. Adapted from Ratledge, 2004.

Cytosolic NADPH is required for KR and EAR functions of FAS. For each elongation step of the acyl chain, two molecules of NADPH are required. One of the major sources of cytosolic NADPH are the pentose phosphate pathway and the transhydrogenase cycle, which transforms NADH into NADPH through the activity of pyruvate carboxylase (PC), malate dehydrogenase (MDH), and malic enzyme (ME), catalyzing the following reactions:



ME has been found in several oleaginous fungi and it has been regarded as a key enzyme involved in lipid accumulation (Ratledge, 2002). In *Mortierella circinelloides*, overexpression of ME enhanced lipid accumulation (Zhang et al., 2011), whereas overexpression of the ME homologous in *Yarrowia lipolytica* did not result in yield improvement.




Fig. 3. Biosynthesis of poly-unsaturated fatty acid. Δ^9d , $\Delta^{12}d$ and $\Delta^{15}d$ are the most common desaturases which are present in the endoplasmic reticulum (Ratledge 2004).

The final products of FAS are myristic or palmitic acids, depending on the yeast species. Reactions resulting in further elongation or desaturation occur in the endoplasmic reticulum (ER). Elongation reactions are catalyzed by elongases (such as malonyl-palmitoyl transacylase, MPT) organized in a complex that requires malonyl-CoA provided by ACC. Desaturations are introduced by ER desaturases, hydrophobic membrane-bound proteins. The most common desaturases are Δ^9 , which inserts the first double bond onto palmitic and/or stearic acids, and Δ^{12} , which catalyzes the insertion of the second unsaturation into oleic acid to produce linoleic acid. Δ^6 and Δ^{15} desaturase activities have been recently described in psychrophilic oleaginous yeasts, based on production of γ and α -linolenic acids, respectively (Fig. 3).

3.2 Biosynthesis of triacyl-glycerol

The fatty acyl-CoA produced by *de novo* synthesis are esterified with glycerol or sterols to produce triacyl-glycerol (TAG) and sterol-esters (SE), respectively. In oleaginous fungi, the neutral lipids SE and TAG are stored inside the lipid bodies (LB). TAG are mostly formed by consecutive acylation of glycerol-3-phosphate (G3P), carried out by diverse acyl transferases. G3P is formed from glycerol by glycerol kinase or can be synthesized from dihydroxyacetone phosphate (DHAP) by G3P dehydrogenase, in a reversible reaction. *S. cerevisiae* can use both G3P and DHAP as acyl-group acceptor. The addition of the first acyl group leads to 1-acyl G3P, also named lysophosphatidic acid (LPA). LPA can also be formed by reduction of acyl-DHAP, carried out by a NADPH dependent reductase. A second acyltransferase loads another acyl group, producing 1,2-diacyl G3P (phosphatidic acid, PA). Phosphate is removed from PA by phosphatidate phosphatase isoenzymes, generating diacylglycerol (DAG). DAG can be the direct precursor of TAG, or can be channeled toward phospholipids biosynthesis (Fig. 4).




Fig. 4. *de novo* synthesis of TAG (adapted from Czabany et al., 2007)

The last step of *de novo* synthesis of TAG can be carried either by using diverse acyl donors, such as acyl-CoA or with phospholipids. In the former case, DAG acyl transferases (DGAT), which are integral proteins of the ER, can directly load the third Acyl-CoA. A DGAT enzyme is present in *S. cerevisiae* and *Y. lipolytica* and is mostly active during the stationary phase, although it is expressed also during the exponential phase. A second DGAT, more active during the exponential growth phase, has been identified in *Y. lipolytica*. In *S. cerevisiae* the phospholipid:DAG acyltransferase (PDAT) is localized in the ER, whereas in *Y. lipolytica* it is present both in the ER and in the surface of LB (Fig. 4).

3.3 Biogenesis of lipid bodies

In eukaryotes, neutral lipids are stored in specialized compartments known as lipid bodies (LB). They are assembled at a specialized subdomain of the ER where most biosynthetic enzymes and structural proteins are located (Waltermann et al., 2005). The neutral lipids do not fit among phospholipids and are thus deposited between the two leaflets of the membrane bilayer. However, substantial amounts of neutral lipids cannot be incorporated into membrane bilayer of ER and ongoing neutral lipid synthesis leads to the formation of a bud which buds off of the ER as a mature LB after reaching the critical size (Fig. 5).




Fig. 5. Model of lipid bodies biogenesis from the membrane of the ER. TAG and SE accumulate between the two leaflets of the phospholipid bilayer (i to iii). The micro-droplet generated (iii, iv) evolve to lipid bodies (v) (Figure adapted from Czabany et al., 2007).

In most oleaginous yeasts, the neutral lipids of LB consist mostly of TAG (up to 90% or more) whereas a small fraction is represented by sterol esters. The presence of significant quantity of free fatty acids (FFA) within LP has been reported only for *Y. lipolytica*. In *S. cerevisiae*, which accumulates less than 15% lipids of its biomass, LB contain similar amounts of TAG and SE.

The core of LB, consisting of neutral lipids is surrounded by a phospholipid monolayer where several proteins are embedded. These proteins exert a key role in lipid metabolism, biosynthesis, and substrate trafficking. Upon requirement, storage lipids are mobilized from this compartment by triacylglycerol lipases and sterol ester hydrolases. The respective degradation products serve as energy sources and/or building blocks for membrane formation. In fact, FA hydrolyzed from TAG or SE are either channeled to the peroxisome, where β -oxidation takes place, or to phospholipid biosynthesis.

4. Metabolic engineering of oleaginous yeasts

The availability of genome data and genetic tools, such as the possibility to integrate homologous or heterologous genes, opened up the possibility to use metabolic engineering to understand the molecular mechanisms involved in lipid accumulation or to increase the yield of stored lipids in *S. cerevisiae* and *Y. lipolytica*. Whereas *S. cerevisiae*

has been used mostly as a model to investigate and understand the lipid metabolism, in *Y. lipolytica* several attempts have been done in order to address the carbon flux toward TAG production and accumulation. Similar approaches are precluded to other oleaginous fungi since they lack genome information and the necessary tools for gene manipulation and strain improvement.

In *Y. lipolytica*, the role of glycerol-3-phosphate (G3P) in triacylglycerol (TAG) biosynthesis and accumulation has been investigated (Beopoulos et al., 2008). In this yeast G3P is formed from glycerol by the glycerol kinase encoded by *GUT1*, or it is synthesized from dihydroxyacetone phosphate (DHAP) by the G3P dehydrogenase (*GPD1*). The antagonist reaction, which produces DHAP from G3P, is carried out in competition by a second isoform of the G3P dehydrogenase, encoded by *GUT2*. In order to force the conversion of DHAP into G3P, the gene *GPD1* was over-expressed and the gene *GUT2* was deleted.

A diverse strategy to increase lipid accumulation was based on the disruption of the β -oxidative metabolism, through the deletion of the 6 *POX* genes (*POX1* to *POX6*) that encode the peroxisomal acyl-coenzyme oxidases (Mlickowa et al., 2004a; Mlickowa et al., 2004b; Beopoulos et al., 2008). As a whole, the best results in terms of percentage of lipids per dry biomass, were reached coupling the increased level of G3P with the deactivation of the β -oxidation pathway (65%) (Dulermo et al., 2011).

Metabolic engineering strategies have been recently exploited to expand the range of substrates used by oleaginous fungi, also through functional expression of heterologous genes. Recently, it has been found that inulin is a good material for bio-productions (Chi et al., 2009). In order to make the oleaginous yeast *Y. lipolytica* able to accumulate lipids on inulin containing materials, the *Kluyveromyces marxianus* exo-inulinase gene (*INU1*) was heterologously expressed on a high copy plasmid (Zhao et al., 2010). The inulinase was efficiently secreted by *Y. lipolytica*, and inulin was hydrolyzed, assimilated and converted into TAG.

5. Cultivation condition of oleaginous yeasts

Lipid accumulation by oleaginous yeasts depends mostly on nutrient limitation conditions when excess carbon is present in the medium. Nutrient limitation prevents cells from being generated, while the carbon excess is converted into storage TAG. Published studies reports that phosphorus, magnesium, zinc, or iron limitation lead to lipid accumulation in model oleaginous yeasts (Hall & Ratledge, 1977; Beopoulos et al., 2009; Wu et al., 2010). However, nitrogen limitation is the most efficient form of nutrient limitation for lipogenesis induction, leading to the highest values of substrate/lipid conversion yield and lipid content within biomass (Hall & Ratledge, 1977; Wynn et al., 2001). Thus, nitrogen limitation is commonly used to induce lipogenesis in oleaginous fungi and the utilization of cultural media with appropriate C/N ratio is crucial to maximize lipid production.

Several studies focused on determining the optimal composition of cultural media for oleaginous fungi with the aim to optimize the performance of lipid-producing bioprocesses. The effect of the C/N ratio on lipid metabolism has been investigated for a number of oleaginous yeasts and molds, such as *Y. lipolytica* and many oleaginous species of *Rhodotorula*, *Candida*, *Apiotrichum*/*Cryptococcus*, *Mortierella* (Hall & Ratledge, 1977; Papanikolau et al., 2003; Granger et al., 1992; Wu et al., 2010; Park et al., 1990; Jang et al., 2005; Amaretti et al., 2010), and has been mathematically modeled for some of these

organisms (Ykema et al., 1986; Granger et al., 1993; Economou et al., 2011). *Y. lipolytica* is the oleaginous microorganism for which information about the metabolic response to different C/N ratios is most abundant (Beopoulos et al., 2009a), particularly due to the availability of molecular tools for genetic engineering of this organism. Therefore, *Y. lipolytica* is regarded as a model organism for microbial oil production and the main traits of its metabolism can be used to give a general description of the metabolic response to different C/N ratios in the majority of oleaginous yeasts. With the increase of the C/N ratio, different metabolic behaviors were observed in *Y. lipolytica*: i) growth with mobilization of storage lipids, ii) growth of fat-free biomass, iii) growth with accumulation of lipids, and iv) growth with lipid accumulation and production of organic acids (Fig. 6).




Fig. 6. Metabolic activity of oleaginous fungi (e.g. *Y. lipolytica*) as a function of carbon flow rate for a fixed nitrogen flow rate. Arrows indicate the consumption of nitrogen and carbon sources by the cells; squares indicate production rate. The dimension of arrows and squares is proportional to flow. (adapted from Beopoulos et al., 2009a)

If the medium is carbon limited or when the extracellular carbon supply gets exhausted, previously stored intracellular lipid can be mobilized and utilized by oleaginous microorganisms to sustain cells generation and production of lipid-free biomass (Park et al., 1990) (Fig. 6, i). If the medium is balanced and/or furnishes just the right amount of carbon flow to satisfy the growth need, balanced growth occurs without any accumulation of storage lipids (Fig. 6, ii). In conditions of carbon excess, a part of the carbon flow, which is proportional to nitrogen availability (Granger et al., 1993), is directed toward cells generation, whereas the carbon exceeding growth needs is channeled to the production of storage lipids (Fig. 6, iii). In some oleaginous fungi, the presence of a large carbon excess leads to the production of great amounts of organic acids, such as pyruvic acid and diverse TCA-cycle intermediates, at the expenses of lipid accumulation (Fig. 6, iv). In these latter conditions, *Y. lipolytica* produces citric acid (Levinson et al., 2007) but other oleaginous yeasts have never been reported to behave this way.

6. Batch, fed-batch and fermentation processes

Batch, fed batch, and continuous modes of culture have been developed to culture oleaginous microorganisms. Lipid production in batch cultures is carried out in a cultural medium with a high initial C/N ratio, the carbon source being present in an adequate excess with respect to the nitrogen source. In fact, in this condition, the flow of carbon utilization is limited only by the substrate uptake system of the cell, while the changes in nitrogen concentration determine the passage from a phase of balanced growth to a phase of lipid accumulation, causing the process to proceed through two phases. As nitrogen is consumed from the culture the C/N ratio tends to increase, but growth remains exponential and balanced until nitrogen is not the limiting substrate. During the growth phase, the carbon flow is mostly channeled to satisfy the growth need, therefore growth is balanced and lipid-free biomass is mostly produced (Fig. 6 ii). As nitrogen concentration becomes limiting, the growth rate and the carbon flow toward biomass generation decrease, while lipid production is triggered, resulting in a shift of microbial metabolism into the lipogenic phase (Fig. 6 iii, Fig. 7).




Fig. 7. Modeling and prediction of the timecourse of a batch fermentation (left) and the steady-state values of a continuous process (right) for microbial production of lipids. Axis are in arbitrary scales.

In batch cultures the initial C/N ratio of the cultural medium has a pivotal role in determining the bioprocess performance. In fact, both the rate and the yield of lipid production depend by the C/N ratio, which affects the duration of the exponential phase and the amount of biomass produced during growth. With a fixed carbon concentration, higher amounts of lipid-free biomass produced during the growth phase correspond to higher lipid production rates during the lipogenic phase, but to lower amounts of lipid content within cells and lipid/substrate conversion yields. Therefore, the initial C/N ratio needs to be optimized to maximize lipid productivity in batch cultures. The optimal C/N value is always high (e.g. in the range between 80 and 350 mol/mol) and strongly depends on the microorganism, the medium composition, the carbon source (e.g. glucose, glycerol, etc.), and the nitrogen source (e.g. diverse organic or inorganic sources). The minimal C/N ratio suitable for lipid accumulation can be estimated as $(Y_{X/S} \cdot q)^{-1}$, where $Y_{X/S}$ is the biomass/carbon source yield coefficient under conditions of carbon limitations (C-mol/C-

mol) and q is the nitrogen/carbon content of biomass (N-mol/C-mol) (Ykema et al., 1986). However, it should be considered that extremely high C/N ratios may cause the production of unwanted byproducts, such as organic acids (Fig. 6 iv), or may lead to severe nitrogen deficiency, causing a rapid decrease in cells viability.

Unlike batch processes, in fed-batch mode, nutrients are fed into the bioreactor in a controlled manner, with the purpose to monitor and control the specific growth rate and the flows of nitrogen and carbon utilization. Through the judicious management of the feeding rate and composition, it is possible to control the C/N ratio within the culture and maintain the oleaginous microorganism in the optimal metabolic status, as appropriate, first for the growth phase, and later for the lipogenic phase. The lipogenic phase is the most extensive, corresponding to lipid production under nitrogen limitation, with constant C/N ratio, preventing loss of viability and acids production (Beopoulos et al., 2009a).

In continuous cultures, at the steady state, the assimilation of C and N sources and the microbial growth occur at constant rates, which ultimately depend by the dilution rate (D). The concentration of the substrates within the bioreactor is steady and depends by the dilution rate as well, the actual C/N ratio of the culture remaining constant unlike in batch cultures. Likewise in batch cultures, in continuous cultures the C/N ratio of the fresh medium needs to be higher than $(Y_{X/S} \cdot q)^{-1}$ to obtain some lipid accumulation (Ykema et al., 1986). However, at the steady-state with this medium, the C/N ratio within the culture is higher than in the fresh medium, due to nitrogen consumption. The extent of substrates utilization, and also the biomass and lipid concentrations are the highest at low D values and decrease with the increase of D (Fig. 7). While low D values promote lipid production and a more complete substrate utilization, on the contrary, the volumetric productivity of continuous processes is positively affected by the increase of the dilution rate (Ykema et al., 1986; Meeuwse et al., 2011b). Therefore, both the C/N content of the medium and the dilution rate need to be thoroughly tuned to maximize lipid productivity of continuous processes.

7. Substrates and raw material

The demand for the inexpensive production of biofuels has intensified due to increasing concerns of climate change, depletion of petroleum-based fuels, and environmental problems. In a market economy, corporations aim to maximize profit, seeking the most competitive feedstock. To produce single-cell oils for biodiesel production, the carbon source has necessarily to be cheap and available in large quantities. Therefore, while the first investigations on oleaginous fungi most commonly employed glucose as carbon source, nowadays the production of single-cell oils is predominantly addressed to transformation of raw materials, by-products and surplus.

Glucose is the carbon source most commonly employed for growth of oleaginous fungi and lipid production (Boulton & Ratledge, 1984; Hansson & Dostalek, 1986; Hassan et al., 1993; Heredia & Ratledge, 1988; Jacob, 1991; Jacob, 1992; Johnson et al., 1992; Li et al., 2007; Pan et al., 1986; Ratledge, 2004; Rau et al., 2005; Saxena et al., 2008; Zhao et al., 2008). High glucose concentrations enhance the carbon flow that is directed toward TAG production, thus improving lipid production in several yeasts. However, growth of some yeasts (e.g. *R. toruloides*) is inhibited by high concentration of glucose, (Li et al., 2007). Furthermore, in batch cultures, initial glucose concentration also affects the fatty acids composition of the lipids (Amaretti et al., 2010).

Carbon sources other than glucose, such as xylose (Christopher et al., 1983; Heredia & Ratledge, 1988;), lactose (Christopher et al., 1983; Daniel et al., 1999;), arabinose, mannose (Hansson & Dostalek, 1986), mannitol (Hansson & Dostalek, 1986), ethanol (Christopher et al., 1983; Eroshin & Krylova, 1983), have been also investigated in the 80s and 90s for the production of microbial lipids.

Albeit glucose is a very good carbon source for lipid production with oleaginous fungi, molasses, which carbohydrate fraction is mainly composed of sucrose, glucose, and fructose, do not represent a promising raw material for lipid production, since they are characterized by a high nitrogen content which delays the unbalanced growth, where number of cells can not augment anymore and lipids are accumulated (Johnson et al., 1995).

Carbons sources obtained from ligno-cellulosic biomasses represent one of the most important potential to produce biodiesel. In fact, several waste biomasses containing forest residues, agricultural residues, food wastes, municipal wastes, and animal wastes can be utilized for the production of lignocellulosic based microbial lipids. Microbial oil production from sulphuric acid treated rice straw hydrolysate (SARSH) by the yeast *Trichosporon fermentans* pointed out the difficulty to perform the process of lipid accumulation in presence of the inhibitory compounds released during hydrolysis, such as acetic acid, furfural, 5-hydroxymethylfurfural, and water soluble lignin (Huang et al., 2009). Selected strains were able to grow on xylose and glucose (Zhu et al., 2008), but the crude hydrolyzate did not result an optimal substrate for a high yield process of lipid production. Cellulose and hemicellulose are generally hardly hydrolyzed and assimilated by yeasts, while they can be degraded and used as carbon source by filamentous fungi. A screening of endophytic fungi from the oleaginous plants was the selection of strains belonging to the genera *Microsphaeropsis*, *Phomopsis*, *Cephalosporium*, *Sclerocystis* and *Nigrospora* that simultaneously accumulated lipids (21.3 to 35.0% of dry weight) and produced cellulase (Peng & Chen, 2007). Albeit these strains could be exploited as microbial oil producers by utilising straw as substrate, they have never been claimed again as a SCO producers on lingo-cellulosic biomass. Attempts to carry out lipid production in Solid State Fermentation (SSF) on wheat straw have been performed exploiting a cellulolytic strain of *Aspergillus oryzae* (Lin et al., 2010). This strain is able to use cellulose as substrate and accumulate lipids in a low cost fermentation system on this abundant cellulosic by-product.

Other complex matrices have been used, such as solids from wheat bran fermentation (Jacob 1991), sewage sludge (Angerbauer et al., 2008), wastewaters of animal fat treatment (Papanikolaou et al., 2002), whey derivatives (Ykema et al., 1989; Vamvakaki et al., 2010), olive oil mill wastewaters (Yousuf et al., 2010), and tomato waste hydrolysate (Fakas et al., 2008).

Nowadays, lipid production with oleaginous yeasts is focused on selection and development of yeasts as converters of glycerol into lipid for biodiesel production, since it is the major side-product of the biodiesel production process. The biotransformation of glycerol into TAG is therefore regarded as a promising way to decrease the cost of biodiesel process through simultaneous reutilization of its major byproduct. In general, for every 100 kg of biodiesel produced, approximately 10 kg of crude glycerol are created. Crude glycerol is a mixture of glycerol (65–85%, w/w), methanol, and soap, and contains macro elements such as calcium, potassium, magnesium, sulfur and sodium. In order to minimize unknown variables introduced through the use of crude glycerol, several studies to determine whether or not glycerol could be used as substrate or co-substrate for growth have been conducted using purified glycerol.

A deep characterization of lipid accumulation on glycerol has been carried out with *Yarrowia lipolytica*, that is able to metabolize several important industrial and agro-industrial

by-products such as raw glycerol, producing large amounts of SCO and organic acids (Papanikolaou et al., 2003; Papanikolaou & Aggelis, 2002; Rymowicz et al., 2010; Rywinska et al., 2009). Biochemistry of lipid production on glycerol has been investigated in this organism: glycerol passes into the microbial cell by facilitated diffusion and the conversion is carried out via phosphorylation pathway, with direct phosphorylation to G3P and subsequent dehydrogenation. Recently, *Y. lipolytica* has been subjected to targeted and purposeful alteration of G3P shuttle pathway to better utilize glycerol for lipid production. In the genetically manipulated strains, lipid accumulation resulted from a complex interrelation between different processes in diverse cell compartments, such as lipid synthesis in the cytosol, location and storage in ER and LB, mobilization and degradation processes (Dulermo & Nicaud, 2011).

Pure glycerol supported growth and lipid accumulation of *Rhodotorula glutinis* and *Candida freyschussii* (Easterling et al., 2009; Amaretti et al., 2011), being used as sole carbon and energy source or in addition to xylose or glucose. The diverse composition of the medium affected not only the lipid/biomass yield, but also the TAG composition, in terms of ration of saturated, monounsaturated, and polyunsaturated fatty acids (Easterling et al., 2009).

Attempts to convert crude glycerol into lipids have been successfully performed exploiting the oleaginous yeast *Cryptococcus curvatus* (Liang et al., 2010). Different processes have been developed with very efficient yields and productivities. In a 12 days two-stage fed-batch where raw glycerol was fed, the biomass density and the lipid content reached 32.9 g/l and 52%, respectively. Methanol of crude glycerol did not pose a significant inhibitory effect even though it was existent in the bioreactor. Lipid accumulated by *C. curvatus* on glycerol presented high amount of monounsaturated fatty acid, turning out as excellent substrate for transformation into biodiesel.

8. Conclusions and perspectives

Oleaginous fungi, and particularly yeasts, are very efficient in the accumulation of intracellular TAG and it is expected that they will be exploited by the biofuel industry in the future. Nonetheless, the costs of microbial lipids are still too high in order to compete with plant oils for biodiesel manufacturing. Cheap carbon sources have necessarily to be used as carbon sources for the cultivation of these microorganisms and the performance of the bioprocess has to be further improved in terms of both the yield and the productivity. The exploration of the natural biodiversity is a promising strategy to identify novel oleaginous species that assimilate and get fat on agro-industrial residues, particularly the lingo-cellulosic biomass and crude glycerol from biodiesel industry. Further approaches combining genomic, transcriptomic, metabolomics, and lipidomic techniques will undoubtedly provide deeper information of lipid production by oleaginous fungi. A metabolic engineering approach is very promising, but it is still precluded for the most oleaginous species, for which genome disclosure has not been accomplished and genetic tools are not available yet.

9. References

- Adachi, D., Hama, S., Numata, T., Nakashima, K., Ogino, C., Fukuda, H., & Kondo, A., (2011) Development of an *Aspergillus oryzae* whole-cell biocatalyst coexpressing triglyceride and partial glyceride lipases for biodiesel production. *Bioresource Technology*. Vol. 102, No. 12, pp. 6723-6729, ISSN 0960-8524.

- Ageitos, J. M., Vallejo, J. A., Veiga-Crespo, P., & Villa TG. (2011) Oily yeasts as oleaginous cell factories. *Applied Microbiology and Biotechnology*. Vol. 90, No. 4, pp. 1219-1227, ISSN 0175-7598.
- Alvarez, H. M. & Steinbuchel, A. (2002) Triacylglycerols in prokaryotic microorganisms. *Applied Microbiology and Biotechnology*. Vol. 60, No. 4, pp. 367-376, ISSN 0175-7598.
- Amaretti, A., Raimondi, S., Sala, M., Roncaglia, L., De Lucia, M., Leonardi, A., & Rossi, M. (2010) Single cell oils of the cold-adapted oleaginous yeast *Rhodotorula glacialis* DBVPG 4785. *Microbial Cell Factories*. Vol. 23, No. 9, pp. 73, ISSN 1475-2859.
- Amaretti, A., Raimondi, S., Leonardi, A., & Rossi, M. (2011) Lipid production from glycerol by *Candida Freyschusii*. *Proceedings of FEMS 2011 14th Congress of European Microbiologists*, pp. 126, Geneva, Switzerland, June 26-30, 2011
- Angerbauer C, Siebenhofer M, Mittelbach M, Guebitz GM (2008) Conversion of sewage sludge into lipids by *Lipomyces starkeyi* for biodiesel production. *Bioresource Technology*. Vol. 99, No. 8, pp. 3051–3056, ISSN 0960-8524.
- Azócar, L., Ciudad, G., Heipieper, H. J., & Navia, R. (2010) Biotechnological processes for biodiesel production using alternative oils. *Applied Microbiology and Biotechnology*. Vol. 88, No. 3, pp. 621-636, ISSN 0175-7598.
- Bankar, A. V., Kumar, A. R., & Zinjarde, S. S. (2009) Environmental and industrial applications of *Yarrowia lipolytica*. *Applied Microbiology and Biotechnology*. Vol. 84, No. 5, pp. 847-865, ISSN 0175-7598.
- Beopoulos, A., Mrozova, Z., Thevenieau, F., Le Dall, M. T., Hapala, I., Papanikolaou, S., Chardot, T., & Nicaud, J. M. (2008) Control of lipid accumulation in the yeast *Yarrowia lipolytica*. *Applied and Environmental Microbiology*. Vol. 74, No. 24, pp. 7779-7789, ISSN 0099-2240.
- Beopoulos, A., Cescut, J., Haddouche, R., Uribe-larrea, J. L., Molina-Jouve, C., & Nicaud, J. M. (2009a) *Yarrowia lipolytica* as a model for bio-oil production. *Progress in Lipid Research*. Vol. 48, No. 6, pp. 375-387, ISSN 0163-7827.
- Beopoulos, A., Chardot, T., & Nicaud, J.M. (2009b) *Yarrowia lipolytica*: a model and a tool to understand the mechanisms implicated in lipid accumulation. *Biochimie*. Vol. 91, No. 6, pp. 692-696, ISSN 0300-9084.
- Boulton, C. A. & Ratledge, C. (1984) *Cryptococcus terricolus*, an oleaginous yeast re-appraised. *Applied Microbiology and Biotechnology*. Vol. 20, No. 1, pp. 72-76, ISSN 0175-7598.
- Butinar, L., Spencer-Martins, I., & Gunde-Cimerman, N. (2007) Yeasts in high Arctic glaciers: the discovery of a new habitat for eukaryotic microorganisms. *Antonie Van Leeuwenhoek*. Vol. 91, No. 3, pp. 277-289, ISSN 1572-9699.
- Calderone, R. A. (2002) *Candida and Candidiasis*. Edited by R. A. Calderone, *American Society of Microbiology Press*. ISSN 1058-4838, Washington, DC, USA.
- Chi, Z. M., Chi, Z., Zhang, T., Liu, G. L., & Yue, L. X. (2009) Inulinase expressing microorganisms and applications of inulinases. *Applied Microbiology and Biotechnology*. Vol. 82, No. 2, pp. 211-220, ISSN 0175-7598.
- Christopher, T., Scragg, A. H., & Ratledge, C. (1983) A comparative study of citrate efflux from mitochondria of oleaginous and non oleaginous yeasts. *European Journal of Biochemistry* Vol. 130, No. 1, pp. 195–204, ISSN 0014-2956
- Czabany, T., Athenstaedt, K., & Daum, G. (2007) Synthesis, storage and degradation of neutral lipids in yeast. *Biochimica et Biophysica Acta*. Vol. 1771, pp. 299–309. ISSN 0006-3002

- Dahlqvist, A., Stahl, U., Lenman, M., Banas, A., Lee, M., Sandager, L., Ronne, H., & Stymne, S. (2009) Phospholipid:diacylglycerol acyltransferase: an enzyme that catalyzes the acyl-CoA-independent formation of triacylglycerol in yeast and plants. *Proceeding of the National Accademy of Science USA*. Vol. 6, No. 12, pp. 6487-6492, ISSN 0027-8424
- Daniel, H. J., Otto, R. T., Binder, M., Reuss, M., & Syldatk, C. (1999) Production of sophorolipids from whey: development of a two-stage process with *Cryptococcus curvatus* ATCC 20509 and *Candida bombicola* ATCC 22214 using deproteinized whey concentrates as substrate. *Applied Microbiology and Biotechnology*. Vol. 51, No. 1, pp. 40-45, ISSN 0175-7598
- de Miguel, T., Calo, P., Díaz, A., & Villa, T. G. (1997) The genus *Rhodospiridium*: a potential source of beta-carotene. *Microbiologia*. Vol. 13, No. 1, pp. 67-70, ISSN 0213-4101
- Dulermo, T. & Nicaud, J. M. (2011) Involvement of the G3P shuttle and β -oxidation pathway in the control of TAG synthesis and lipid accumulation in *Yarrowia lipolytica*. *Metabolic engineering*. Vol. 40, No. 4, pp. 483-488, ISSN 1096-7176
- Easterling, E. R., French, W. T., Hernandez, R., & Licha, M. (2009) The effect of glycerol as a sole and secondary substrate on the growth and fatty acid composition of *Rhodotorula glutinis*. *Bioresource Technology*. Vol. 100, No. 1, pp. 356-361. ISSN 0960-8524
- Economou, C.N., Aggelis, G., Pavlou, S., & Vayenas, D.V. (2011) Modeling of single-cell oil production under nitrogen-limited and substrate inhibition conditions. *Biotechnology and Bioengineering*. Vol. 108, No. 5, pp. 1049-1055, ISSN 0006-3592.
- Fakas, S., Certik, M., Papanikolaou, S., Aggelis, G., Komaitis, M., & Galiotou-Panayotou, M. (2008) Gamma-linolenic acid production by *Cunninghamella echinulata* growing on complex organic nitrogen sources. *Bioresource Technology*. Vol. 99, No. 13, pp. 5986-5990, ISSN 0960-8524
- Findley, K., Rodriguez-Carres, M., Metin, B., Kroiss, J., Fonseca, A., Vilgalys, R., & Heitman, J. (2009) Phylogeny and phenotypic characterization of pathogenic *Cryptococcus* species and closely related saprobic taxa in the Tremellales. *Eukaryotic Cell*. Vol. 8, No. 3, pp. 353-361, ISSN 1535-9786
- Gujjari, P., Suh, S. O., Coumes, K., Zhou, J. J. (2011) Characterization of oleaginous yeasts revealed two novel species: *Trichosporon cacaoliposimilis* sp. nov. and *Trichosporon oleaginosus* sp. nov. *Mycologia*. in press, ISSN 1557-2536.
- Granger, L.M., Perlot, P., Goma, G., & Pareilleux, A. (1992) Kinetics of growth and fatty acid production of *Rhodotorula glutinis*. *Applied Microbiology and Biotechnology*. Vol. 37, No. 1, pp. 13-17, ISSN 0175-7598.
- Granger, L.M., Perlot, P., Goma, G., & Pareilleux, A. (1993) Efficiency of fatty acid synthesis by oleaginous yeasts: Prediction of yield and fatty acid cell content from consumed C/N ratio by a simple method. *Journal of biochemical and microbiological technology and engineering*. Vol. 42, No. 10, pp. 1151-1156, ISSN 0006-3592.
- Gutiérrez, A., López-García, S., & Garre, V. (2011) High reliability transformation of the basal fungus *Mucor circinelloides* by electroporation. *Journal of Microbiological Methods*. Vol. 84, No. 3, pp. 442-446, ISSN 0167-7012
- Hall, M. J., & Ratledge, C. (1977) Lipid accumulation in an oleaginous yeast (*Candida* 107) growing on glucose under various conditions in a one- and two-stage continuous

- culture. *Applied and Environmental Microbiology*. Vol. 33, No. 3, pp. 577-584, ISSN 0099-2240.
- Hansson, L. & Dostalek, M. (1986) Influence of cultivation conditions on lipid production by *Cryptococcus albidus*. *Applied Microbiology and Biotechnology*. Vol. 24, No. 1, pp. 12-18, ISSN 0175-7598.
- Hassan, M., Blanc, P. J., Granger, L. M., Pareilleux, A., & Goma, G. (1993) Lipid production by an unsaturated fatty acid auxotroph of the oleaginous yeast *Apiotrichum curvatum* grown in single-stated continuous culture. *Applied Microbiology and Biotechnology*. Vol. 40, No. 4, pp. 483-488, ISSN 0175-7598.
- Heredia, L. & Ratledge, C. (1988) Simultaneous utilization of glucose and xylose by *Candida curvata* D in continuous culture. *Biotechnology Letters*. Vol. 10, No. 1, pp. 25-30, ISSN 0141-5492.
- Holdsworth, J. E. & Ratledge, C. (1991) Triacylglycerol synthesis in the oleaginous yeast *Candida curvata* D. *Lipids*. Vol. 26, No. 2, pp. 111-118, ISSN 0024-4201.
- Hu, C., Zhao, X., Zhao, J., Wu, S., & Zhao, Z. K. (2009) Effects of biomass hydrolysis by-products on oleaginous yeast *Rhodospiridium toruloides*. *Bioresource Technology*. Vol. 100, No. 20, pp. 4843-4847, ISSN 0960-8524.
- Huang, C., Zong, M. H., Wu, H., & Liu, Q. P. (2009). Microbial oil production from rice straw hydrolysate by *Trichosporon fermentans*. *Bioresource Technology*. Vol. 100, No. 19, pp. 4535-4538, ISSN 0960-8524
- Jacob Z. (1991) Enrichment of wheat bran by *Rhodotorula gracilis* through solid-state fermentation. *Folia Microbiologica*. Vol. 36, No. 1, pp. 86-91, ISSN 1874-9356.
- Jacob, Z. (1992) Linear growth and lipid synthesis in the oleaginous yeast *Rhodotorula gracilis*. *Folia Microbiologica*. Vol. 37, No. 2, pp. 117-121, ISSN 1874-9356.
- Jang, H.D., Lin, Y.Y., & Yang, S.S. (2005) Effect of culture media and conditions on polyunsaturated fatty acids production by *Mortierella alpina*. *Bioresource Technology*. Vol. 96, No. 15, pp. 1633-1644, ISSN 0960-8524.
- Johnson, V. W., Singh, M., Saini, V. S., Adhikari, D. K., Sista, V. R., & Yadav, N. K. (1992) Effect of pH on lipid accumulation by an oleaginous yeast: *Rhodotorula glutinis* IIP-30. *World Journal of Microbiology and Biotechnology* Vol. 8, pp. 382-384, ISSN 0959-3993.
- Johnson, V. W., Sigh, M., Saini V. S., Adhikari, D. K., Sista, V., & Yadav, N. K. (1995) Utilization of molasses for the production of fat by an oleaginous yeast, *Rhodotorula glutinis* IIP-30. *Journal of Industrial Microbiology and Biotechnology* Vol. 14, No. 1 pp. 1-4, ISSN 1476-5535
- Kohlwein, S. D. (2010) Triacylglycerol homeostasis: insights from yeast. *Journal of Biological Chemistry*. Vol. 285, No. 21, pp. 15663-15667, ISSN 0021-9258.
- Levinson, W.E., Kurtzman, C.P., & Min, T. (2007) Characterization of *Yarrowia lipolytica* and related species for citric acid production from glycerol. *Enzyme and Microbial Technology*. Vol. 41, No. 3, pp. 292-295, ISSN 0141-0229.
- Liang, Y., Cui, Y., Trushenski, J., & Blackburn, J. W. (2010) Converting crude glycerol derived from yellow grease to lipids through yeast fermentation. *Bioresource Technology*. Vol. 101, No. 19, pp. 7581-7586, ISSN 0960-8524
- Li, C. H., Cervantes, M., Springer, D. J., Boekhout, T., Ruiz-Vazquez, R. M., Torres-Martinez, S. R., Heitman, J., & Lee, S. C. (2011) Sporangiospore Size Dimorphism Is Linked to

- Virulence of *Mucor circinelloides*. *PLoS Pathogen*. Vol. 7, No. 6, e1002086, ISSN 1553-7366
- Li, Q., Du, W., & Liu D. (2008) Perspectives of microbial oils for biodiesel production. *Applied Microbiology and Biotechnology*. Vol. 80, No. 5, pp. 749-756, ISSN 0175-7598.
- Li, Y., Zhao, Z., & Bai, F. (2007) High-density cultivation of oleaginous yeast *Rhodospiridium toruloides* Y4 in fed-batch culture. *Enzyme and Microbial Technology*. Vol. 41, No. 3, pp. 312-317, ISSN 0141-0229.
- Lin, H., Cheng, W., Ding, H. T., Chen, X. Zhou, Q. F., Zhao, Y. H. (2010) Direct microbial conversion of wheat straw into lipid by a cellulolytic fungus of *Aspergillus oryzae* A-4 in solid-state fermentation. *Bioresource technology*. Vol. 101, No. 19, pp. 7556-7562, ISSN 0960-8524
- Meesters, P. A. E. P., Huijberts, G. N. M., & Eggink, G. (1996) High-cell-density cultivation of the lipid accumulating yeast *Cryptococcus curvatus* using glycerol as a carbon source. *Applied Microbiology and Biotechnology*. Vol. 45, No. 5, pp. 749-756, ISSN 0175-7598.
- Meeuwse, P., Tramper, J., & Rinzema, A. (2011a) Modeling lipid accumulation in oleaginous fungi in chemostat cultures: I. Development and validation of a chemostat model for *Umbelopsis isabellina*. *Bioprocess and Biosystems Engineering*. Epub ahead of print, ISSN 1615-7591.
- Meeuwse, P., Tramper, J., & Rinzema A. (2011b) Modeling lipid accumulation in oleaginous fungi in chemostat cultures. II: Validation of the chemostat model using yeast culture data from literature. *Bioprocess and biosystems Engineering*. Epub ahead of print, ISSN 1615-7591.
- Meng, X., , Yang, J., Xu, X., Zhang, L., Nie, Q., & Xian, M. (2009) Biodiesel production from oleaginous microorganisms. *Renewable Energy*. Vol. 34, No. 1, pp. 1-5, ISSN 0960-1481.
- Mlickova, K., Luo, Y., d'Andrea, S., Pec, P., Chardot, T., & Nicaud, J.-M. (2004a). Acyl-CoA oxidase, a key step for lipid accumulation in the yeast *Yarrowia lipolytica*. *Journal of molecular catalysis. B, Enzymatic*. Vol. 28, No. 2-3, pp. 81-85 ISSN 1381-1177
- Mlíckova, K., Roux, E., Athenstaedt, K., d'Andrea, S., Daum, G., Chardot, T., & Nicaud, J-M. (2004b). Lipid accumulation, lipid body formation, and acyl coenzyme A oxidases of the yeast *Yarrowia lipolytica*. *Applied and Environmental Microbiology*. Vol. 70, No. 7, pp. 3918-3924 ISSN 0099-2240
- Pan, J. G., Kwak, N. Y., & Rhee, J. S. (1986) High density cell culture of *Rhodotorula glutinis* using oxygen-enriched air. *Biotechnology Letters*. Vol. 8, No. 10, pp. 715-718, ISSN 0141-5492
- Papanikolaou, S., & Aggelis, G. (2002) Lipid production by *Yarrowia lipolytica* growing on industrial glycerol in a single-stage continuous culture. *Bioresource technology*. Vol 82, No. 1, pp. 43-49, ISSN 0960-8524
- Papanikolaou, S., Chevalot, I., Komaitis, M., Marc, I., & Aggelis, G. (2002) Single cell oil production by *Yarrowia lipolytica* growing on an industrial derivative of animal fat in batch cultures. *Applied Microbiology and Biotechnology* Vol. 58, No. 3, pp. 308-312, ISSN 0175-7598
- Papanikolaou, S., Muniglia, L., Chevalot, I., Aggelis, G., & Marc, I. (2003) Accumulation of a cocoa-butter-like lipid by *Yarrowia lipolytica* cultivated on agro-industrial residues. *Current Microbiology*. Vol. 46, No. 2, pp.124-130, ISSN 0343-8651.

- Park, W.S., Murphy, P.A., & Glatz, B.A. (1990) Lipid metabolism and cell composition of the oleaginous yeast *Apiotrichum curvatum* grown at different carbon to nitrogen ratios. *Canadian Journal of Microbiology*. Vol. 36, No. 5, pp. 318-326, ISSN 0008-4166.
- Peng, X. W. & Chen, H. Z. (2007) Microbial oil accumulation and cellulase secretion of the endophytic fungi from oleaginous plants. *Annals of Microbiology*. 57 Sakaiu, 239-242, ISSN 1590-4261
- Rajakumari, S., Grillitsch, K., & Daum, G. (2008) Synthesis and turnover of non-polar lipids in yeast. *Progress in Lipid Research*. Vol. 47, No. 3, pp. 157-171, ISSN 0163-7827
- Ratledge, C. (2002) Regulation of lipid accumulation in oleaginous micro-organisms. *Biochemical Society Transactions*. Vol. 30, No. 6, pp. 1047-1050, ISSN 0300-5127
- Ratledge, C., (2004) Fatty acid biosynthesis in microorganisms being used for single cell oil production. *Biochimie*. Vol. 86, No. 11, pp. 807-815, ISSN 0300-9084.
- Ratledge, C. & Wynn, J. P. (2002) The biochemistry and molecular biology of lipid accumulation in oleaginous microorganisms. *Advances in Applied Microbiology*. Vol. 51, No. 1, pp. 1-44, ISSN 0065-2164.
- Rymowicz, W., Fatykhova, A. R., Kamzolova, S. V., Rywińska, A., & Morgunov, I.G. (2010) Citric acid production from glycerol-containing waste of biodiesel industry by *Yarrowia lipolytica* in batch, repeated batch, and cell recycle regimes. *Applied Microbiology and Biotechnology*. Vol 87, No. 3, pp. 971-979. ISSN 0175-7598
- Rywinska, A., Rymowicz, W., & Marcinkiewicz, M. (2010) Valorization of raw glycerol for citric acid production by *Yarrowia lipolytica* yeast. *Electronic Journal of Biotechnology*. Vol. 13, No. 4, ISSN 0717-3458
- Rau, U., Nguyen, L. A., Roeper, H., Koch, H., & Lang, S. (2005) Fed-batch bioreactor production of mannosylerythritol lipids secreted by *Pseudozyma aphidis*. *Applied Microbiology and Biotechnology*. Vol. 80, No. :607-613, ISSN 0175-7598
- Rossi, M., Buzzini, P., Cordisco, L., Amaretti, A., Sala, M., Raimondi, S., Ponzoni, C., Pagnoni, U. M., & Matteuzzi, D. (2009) Growth, lipid accumulation, and fatty acid composition in obligate psychrophilic, facultative psychrophilic, and mesophilic yeasts. *FEMS microbiology ecology*. Vol. 69, No. 3, pp. 363-372, ISSN 0168-6496
- Sakuradani, E., Ando, A., Ogawa, J., & Shimizu, S. (2009) Improved production of various polyunsaturated fatty acids through filamentous fungus *Mortierella alpina* breeding. *Applied Microbiology and Biotechnology*. Vol. 84, No. 1, pp. 1-10. ISSN 0175-7598
- Satyanarayana, T., & Kunze, G. (2010) Gatersleben, Germany (Eds.) *Yeast Biotechnology: Diversity and Applications* 746 pages ISBN: 1402082916
- Saxena, V., Sharma, C. D., Bhagat, S. D., Saini, V. S., & Adhikari, D. K. (2008) Lipid and fatty acid biosynthesis by *Rhodotorula minuta*. *Journal of the American Oil Chemists Society*. Vol. 75, No. 4, pp. 501-505, ISSN 0003-021X.
- Smit, M. S. (2004) Fungal epoxide hydrolases: new landmarks in sequence-activity space. *Trends in biotechnology*. Vol. 22, No. 3, pp. 123-129, ISSN 0167-7799
- Tehlivets, O., Scheuringer, K., & Kohlwein, S. D. (2007) Fatty acid synthesis and elongation in yeast. *Biochimica et Biophysica Acta*. Vol. 1171, No. 3, pp. 255-270, ISSN 0006-3002
- van der Walt, J. P. (1992) The Lipomycetaceae, a model family for phylogenetic studies. *Antonie Van Leeuwenhoek*. Vol. 62, No. 4, pp. 247-250, ISSN 0003-6072.
- Vamvakaki, A. N., Kandarakis, I., Kaminarides, S., Komaitis, M. & Papanikolaou, S. (2010) Cheese whey as a renewable substrate for microbial lipid and biomass production

- by Zygomycetes *Engineering in Life Sciences* Vol. 10, No. 4, pp. 348–360, ISSN 1618-0240
- Vicente, G., Bautista, L. F., Rodríguez, R., Gutiérrez, F. J., Sádaba, I., Ruiz-Vázquez, R. M., Torres-Martínez, S., & Garre, V. (2009) Biodiesel production from biomass of an oleaginous fungus. *Biochemical Engineering Journal*. Vol. 48, No. 1, pp. 22–27, ISSN 1369-703X
- Wang, S. L., Sun, J. S., Han, B. Z., & Wu, X. Z. (2007) Optimization of beta-carotene production by *Rhodotorula glutinis* using high hydrostatic pressure and response surface methodology. *Journal of Food Science*. Vol. 72, No. 8, pp. M325–329, 1911, ISSN 0022-1147
- Wu, S., Hu, C., Jin, G., Zhao, X., & Zhao, Z.K. (2010) Phosphate-limitation mediated lipid production by *Rhodospiridium toruloides*. *Bioresource Technology*. Vol. 101, No. 15, pp. 6124–6129, ISSN 0960-8524
- Wynn, J.P., Hamid, A. A., Li, Y., & Ratledge, C. (2001) Biochemical events leading to the diversion of carbon into storage lipids in the oleaginous fungi *Mucor circinelloides* and *Mortierella alpina*. *Microbiology*. Vol. 147, No. 10, pp. 2857–2864, ISSN 1350-0872.
- Ykema, A., Verbree, E.C., Verseveld, H.W., & Smit, H. (1986) Mathematical modelling of lipid production by oleaginous yeasts in continuous cultures. *Antonie van Leeuwenhoek*. Vol. 52, No. 6, pp. 491–506, ISSN:1572-9699
- Ykema, A., Kater, M. M. & Smit H. (1989) Lipid production in whey permeate by an unsaturated fatty acid mutant of the oleaginous yeast *Apiotrichum curvatum*. *Biotechnology Letters*. Vol.11, No. 7, pp. 477–482, ISSN 0141-5492
- Yousuf, A., Sannino, F., Addorisio, V., Pirozzi, D. (2010) Microbial conversion of olive oil mill wastewaters into lipids suitable for biodiesel production. *Journal of agricultural and food chemistry*. Vol. 11 No. 58(15), pp. 8630–8635, ISSN 0021-8561
- Yu, X., Zheng, Y., Dorgan, K. M., Chen, S. (2011) Oil production by oleaginous yeasts using the hydrolysate from pretreatment of wheat straw with dilute sulfuric acid. *Bioresource Technology*. Vol. 102, No. 10, pp. 6134–6140, ISSN 0960-8524
- Zhang, Y., Adams, I. P., & Ratledge, C. (2007) Malic enzyme: the controlling activity for lipid production? Overexpression of malic enzyme in *Mucor circinelloides* leads to a 2.5-fold increase in lipid accumulation. *Microbiology*. Vol. 153, No. 7, pp. 2013–2025, ISSN 1350-0872
- Zhang, J., Fang, X., Zhu, X. L., Li, Y., Xu, H. P., Zhao, B. F., Chen, L. & Zhang, X. D. (2011) Microbial lipid production by the oleaginous yeast *Cryptococcus curvatus* O3 grown in fed-batch culture. *Biomass and Bioenergy*. Vol. 35, No. 5, pp. 1906–1911, ISSN 0961-9534
- Zhao, C. H., Cuim W., Lium X. Y., Chi, Z. M., & Madzak, C. (2010) Expression of inulinase gene in the oleaginous yeast *Yarrowia lipolytica* and single cell oil production from inulin-containing materials. *Metabolic engineering*. Vol. 12, No. 6, pp. 510–517, ISSN 1096-7176
- Zhao, X., Kong, X., Hua, Y., Feng, B., Zhao, Z. K. (2008) Medium optimization for lipid production through co-fermentation of glucose and xylose by the oleaginous yeast *Lipomyces starkeyi*. *European Journal of Lipid Science and Technology*. Vol. 110, No. 5, pp. 405–412, ISSN 1438-9312

- Zhu, L. Y., Zong, M. H., & Wu, H. (2008) Efficient lipid production with *Trichosporon fermentans* and its use for biodiesel preparation. *Bioresource Technology*. Vol. 99, No. 16, pp. 7881-7885, ISSN 0960-8524
- Zinoviev, S., Müller-Langer, F., Das, P., Bertero, N., Fornasiero, P., Kaltschmitt, M., Centi, G., & Miertus, S. (2010) Next-generation biofuels: Survey of emerging technologies and sustainability issues. *Chemistry & sustainability, energy & materials*. Vol. 3, No. 10, pp. 1106-1133, ISSN 1864-564X



Biodiesel - Feedstocks and Processing Technologies

Edited by Dr. Margarita Stoytcheva

ISBN 978-953-307-713-0

Hard cover, 458 pages

Publisher InTech

Published online 09, November, 2011

Published in print edition November, 2011

The book "Biodiesel: Feedstocks and Processing Technologies" is intended to provide a professional look on the recent achievements and emerging trends in biodiesel production. It includes 22 chapters, organized in two sections. The first book section: "Feedstocks for Biodiesel Production" covers issues associated with the utilization of cost effective non-edible raw materials and wastes, and the development of biomass feedstock with physical and chemical properties that facilitate its processing to biodiesel. These include Brassicaceae spp., cooking oils, animal fat wastes, oleaginous fungi, and algae. The second book section: "Biodiesel Production Methods" is devoted to the advanced techniques for biodiesel synthesis: supercritical transesterification, microwaves, radio frequency and ultrasound techniques, reactive distillation, and optimized transesterification processes making use of solid catalysts and immobilized enzymes. The adequate and up-to-date information provided in this book should be of interest for research scientist, students, and technologists, involved in biodiesel production.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Maddalena Rossi, Alberto Amaretti, Stefano Raimondi and Alan Leonardi (2011). Getting Lipids for Biodiesel Production from Oleaginous Fungi, *Biodiesel - Feedstocks and Processing Technologies*, Dr. Margarita Stoytcheva (Ed.), ISBN: 978-953-307-713-0, InTech, Available from:
<http://www.intechopen.com/books/biodiesel-feedstocks-and-processing-technologies/getting-lipids-for-biodiesel-production-from-oleaginous-fungi>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen