

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



p53 as a Therapeutic Target in T-ALL

Irene Riz¹, Wenjing Yang², Weiqun Peng² and Robert G. Hawley¹

¹*Department of Anatomy and Regenerative Biology*

²*Department of Physics, The George Washington University, Washington, DC
USA*

1. Introduction

TP53 is a central hub integrating stress signals from oncogenic genetic lesions and cytotoxic anti-cancer agents. The function of p53 as a regulator of transcription is well documented. More recently it was shown to directly interact with BCL2 family members and induce mitochondrial cell death. Stress-induced activation of p53 leads to cell cycle arrest that allows metabolic adjustments and repair mechanisms to take place prior to the next cycle; p53 may also induce senescence or apoptosis depending on the strength and nature of stress stimuli and/or cell type.

In 50% of human cancers, the TP53 gene is deleted or mutated with a high proportion of gain of oncogenic function mutations. It is noteworthy therefore that TP53 is rarely mutated in T-ALL. However, its tumor suppressor activity is circumvented by genetic lesions. We will discuss here the most frequent T-ALL genetic abnormalities of INK4A/ARF, NOTCH1 and PTEN genes and how they affect TP53 expression and function. Current understanding of the signaling pathways governed by these oncogenes is advanced enough to find points of intersection with p53 downstream targets and attempt to translate the accumulated knowledge to the clinic.

In addition, we will discuss the results of our analysis of publicly available expression profiling data indicating the existence of a TP53-anchored transcriptional program targeted by T-ALL oncogenes such as NOTCH, MYC and TLX1 in primary leukemic cells and how it can be exploited for cancer intervention.

Overall, the goal of this chapter is to describe how T-ALL pathobiology affects the p53-interacting network, highlighting some new potential therapeutic targets as well as some still unresolved questions.

2. INK4A-ARF inactivation circumvents oncogene-induced p53

Inactivation of INK4A-ARF occurs in 70% of T-ALL cases by mutations, biallelic deletions or hypermethylation (Hebert et al., 1994; Gardie et al., 1998; Sulong et al., 2009; Van Vlierberghe et al., 2008; Ferrando et al., 2002; Omura-Minamisawa et al., 2000). The locus encodes two stress-induced proteins with distinct tumor suppressing functions (Quelle et al., 1995): one, INK4A, targets cell cycle entry and another, ARF, inhibits cell cycle progression and cell survival. The two completely unrelated protein sequences derive from the same locus via expression of two alternative reading frames. This unusual feature was

suggested to be important for coordinate regulation of these gene products in response to stress signals (Gil & Peters, 2006; Popov & Gil, 2010). The tumor suppressor functions of INK4A and ARF are not redundant since animals with individually inactivated products of the locus showed less spontaneous tumors than the double knockouts (Sharpless et al., 2004). It is well established that the function of INK4A is to inhibit cyclin D-dependent protein kinases and thus suppress proliferation, loss of function contributing to signal-independent cell cycle entry (Serrano et al., 1993). Recently, INK4A was also implicated in regulation of thymocyte apoptosis in response to oxidative stress or gamma irradiation (Bianchi et al., 2006). An excellent review summarizing the biological functions of the INK4 family of proteins has been published (Canepa et al., 2007). Our focus will be on the ARF protein since, as shown in Figure 1, the best characterized activity of ARF is activation of p53. The mechanisms include direct inhibition of the enzymatic activity of the MDM2 E3 ubiquitin ligase and sequestering of the protein in nucleoli (Llanos et al., 2001; Honda & Yasuda, 1999): MDM2 binding is one of the major mechanisms keeping the apoptotic and growth-arresting function of p53 in check. MDM2 blocks p53 transactivation function and enforces its nuclear export and proteasomal degradation (Honda et al., 1997; Zhang & Xiong, 2001; Weber et al., 1999) (See Figure 1 for more details). In addition ARF was suggested to have p53-independent functions; for example, modulation of activity of transcription factors such as MYC, E2F and NF κ B family members leads to their enhanced apoptotic activity or stress-induced inhibition of protein synthesis (Sherr, 2006; Qi et al., 2004).

ARF is regulated at multiple levels. Its protein stability and subcellular localization is controlled by the nucleolar phosphoprotein NPM1. ARF-NPM1 complexes are predominantly localized in nucleoli where ARF is more stable (Bertwistle et al., 2004; Brady et al., 2004). Its half life significantly decreases in the nucleoplasm via ubiquitination and proteasome-mediated degradation (Rodway et al., 2004; Kuo et al., 2004). Perhaps the most important aspect of ARF regulation is at the level of transcription. ARF is not expressed in most normal tissues; however, it can be activated in response to stress or aberrant hyperproliferative signals (e.g. RAS mutations, MYC overexpression, BCR-ABL translocation).

In adult hematopoietic stem cells and in immature thymocytes, the INK4A-ARF locus is silenced by the Polycomb-group gene BMI1 (Jacobs et al., 1999; Bracken et al., 2007). This epigenetic regulation is required for survival of normal T cell precursors (Miyazaki et al., 2008). However, ARF expression can be induced at this stage and activates apoptosis in cells with aberrant T cell receptor gene rearrangements or other genetic lesions leading to ectopic activation of protooncogenes. The most frequent T-ALL-associated oncogenic events, involving NOTCH1 (50%) and TAL1 (60%), as well as other less frequent genetic abnormalities, such as activation of LMO2 (9%), were directly demonstrated to cooperate with the loss of INK4A-ARF function (Shank-Calvo et al., 2006; Treanor et al., 2011). The response of the locus may be developmentally specific. For example, the responsiveness of the INK4A-ARF locus to the constitutively-active truncated form of NOTCH1 depends on the stage of development, with the locus being silent in hematopoietic stem cells but inducible in thymocytes (Volanakis et al., 2009). From that perspective, it is noteworthy that in immature T-ALL cases, the INK4A-ARF locus is predominantly found intact but kept inactive by epigenetic mechanisms (Ferrando et al., 2002). Thus these silencing mechanisms may serve as potential therapeutic targets in immature T-ALLs.




Fig. 1. Regulation of p53 protein stability via ARF and MDM2/MDMX. From left to right: nucleolar phosphoprotein NPM1 controls ARF protein stability and subcellular localization. ARF-NPM1 complexes are predominantly localized in nucleoli where ARF is more stable. Liberated ARF may be degraded by proteasomes or may associate with MDM2 and prevent degradation of p53 via direct binding to MDM2 and inhibition of its ubiquitin ligase activity. p53 is stabilized in MDMX complexes, but for full activation of p53-dependent transcription MDMX is replaced by transcriptional cofactors. MDM2 and MDMX are subjected to stress-induced proteasomal degradation.

Because the immature T-ALL subset represents the highest risk among T-ALL, we will describe in more detail BMI1 function and regulation to highlight potential therapeutic strategies aiming to reactivate the INK4A-ARF locus. In thymocytes, BMI1 binds directly to the locus and specifically maintains trimethylation of histone H3 at lysine 27 (3mH3K27 modification) (Miyazaki et al., 2008). For maintenance of the repressed state through multiple rounds of cell divisions, DNA methylation markers are necessary. Polycomb proteins interpret DNA methylation marks and translate them into histone modifications to initiate/maintain repression (Spivakov & Fisher, 2007). Thus it is not surprising, but may be very important for potential clinical translation, that inhibition of DNA methyltransferase (DNMT) was shown to derepress the INK4A-ARF locus by affecting this mechanism in human cord blood-derived multipotent stem cells. The authors used 5-azacytidine, an inhibitor of DNMT analogous to cytidine; they showed that loss of DNA methylation marks caused diminished recruitment of EZH2, a key histone methyltransferase, and decreased 3mH3K27 modification of the INK4A-ARF locus (So et al., 2011). Another attractive candidate for therapeutic intervention is Hedgehog signaling, required for survival and proliferation of early thymocyte precursors (El Andaloussi et al., 2006). Moreover, it was recently shown that Sonic hedgehog activates BMI1 expression during cerebellar development (Leung et al., 2004). Thus testing Hedgehog inhibitors such as cyclopamine or vismodegib on immature T-ALL samples might be a promising approach. The strategy of reactivating the INK4A-ARF locus is complicated, however, by the fact that the locus responds to the same oncogenic signals that support the survival and proliferation of

malignant cells. For example, a powerful pro-survival kinase AKT1 targets EZH2 and suppresses methylation of histone H3 at lysine 27 (Cha et al., 2005) while MAPKAP kinase 3, a convergence point downstream of activated ERK and p38, inhibits BMI1 association with chromatin (Voncken et al., 2005). For these reasons and because the INK4A-ARF locus is deleted in the majority of T-ALL cases, current therapeutic strategies are based on activation of p53 in an ARF-independent manner.

3. NOTCH-governed network affects ARF and p53

Mutations activating the developmental regulator NOTCH1 occur in more than 50% of T-ALL cases (Weng et al., 2004). NOTCH1 was initially implicated in T-cell leukemogenesis by the finding of rare chromosomal translocations that place a constitutively-active truncated form of NOTCH1 (NIC) under the T cell receptor (TCR) beta chain promoter (Ellisen et al., 1991). NOTCH1 is a transmembrane receptor (Kopan & Ilagan, 2009). The Delta-like and Jagged ligands activate proteolytic processing of NOTCH1 that releases its cytoplasmic portion allowing it to translocate to the nucleus. In the nucleus, NOTCH1 activates transcription via a DNA-bound protein CSL (Aster et al., 2008). Subsequently, a high frequency of NOTCH1-activating mutations were characterized that enhance its proteolytic processing and/or stabilize its intracellular portion (Weng et al., 2004). Thus in T-ALL, NOTCH signaling is represented by a broad spectrum of levels of activation that may still be ligand dependent and also inhibited by drugs targeting the NOTCH1 processing machinery (Lewis et al., 2007; Sulis et al., 2011).

NOTCH signaling is required for several consecutive stages of normal thymocyte development, from the earliest stage of T-cell fate commitment until the late cortical thymocyte stage with fully rearranged TCR genes (Tanigaki & Honjo, 2007). NOTCH1 provides survival and stimulates growth of normal thymocytes and leukemic T cells (Grabher et al., 2006; Ciofani & Zuniga-Pflucker, 2005). That said, the functional role of NOTCH1 in T-ALL cells undergoing chemotherapeutic treatment is less clear. As NOTCH1 promotes their survival, NOTCH1 mutations would be expected to confer enhanced drug resistance. Interestingly, however, mutations activating NOTCH1 were found to associate with good initial response to treatment (Kox et al., 2010; Asnafi et al., 2009). In this context, and consistent with these observations, it is worth mentioning that we and others observed that NOTCH1 inhibition decreases sensitivity of T-ALL cell lines to selected chemotherapeutic agents while NOTCH1 activation enhances the response (De Keersmaecker et al., 2008; Liu et al., 2009; Riz et al., 2011).

The function of NOTCH1 is mediated by several signaling hubs that in turn impact ARF and p53 function: among them, mTOR kinase, a key growth regulator constitutively activated in many cancers; eIF4E, a selective regulator of translation initiation; SKP2, an E3 ubiquitin ligase; and transcription factors such as MYC and NF κ B (Chan et al., 2007; Mungamuri et al., 2006; Hsieh & Ruggero, 2010; Kao et al., 2009; Dohda et al., 2007; Murphy et al., 2008).

The accumulated evidence indicates that ectopic activation of NOTCH1 in premalignant thymocytes may lead to ARF induction. First, in T-ALL, NOTCH1 mutations frequently coincide with INK4A-ARF inactivation (Ferrando et al., 2002; Treanor et al., 2011). Second, in mice, progression to full malignancy in NIC-expressing thymocytes is associated with decreased ARF expression (Li et al., 2008). And finally, oncogenic cooperation of these

two genetic aberrations was directly demonstrated in a murine T-ALL model (Volanakis et al., 2009). Moreover, the authors showed indirect activation of the ARF promoter by expression of NIC. Thus these data demonstrated that NOTCH1 mutations may activate ectopic signaling that triggers the oncogene-induced stress response exemplified by ARF induction. However, there is also evidence suggesting that NOTCH1 may downregulate the tumor suppressor function of p53. In an elegant system allowing regulatable NIC expression, it was shown that downregulation of NIC levels in mouse lymphomas *in vivo* is associated with activation of p53 (Beverly et al., 2005). Other work also demonstrated that ectopic expression of NIC partially downregulates the p53-mediated apoptotic response to DNA-damaging drugs in human leukemic cell lines (Mungamuri et al., 2006). The effect was mediated by the mTOR kinase. The authors showed that mTOR inhibition by rapamycin prevented (and eIF4E overexpression restored) the NOTCH1 pro-survival effect. Subsequently, in experiments with loss or gain of NOTCH1 function, MYC was placed upstream of the mTOR-eIF4E pathway (Chan et al., 2007). In agreement with these findings, eIF4E-driven CAP-mediated translation was shown to be required for MYC transforming function (Ruggero et al., 2004; Lin et al., 2008; Barna et al., 2008). Other work indicated that the oncogenic function of the mTOR-eIF4E pathway is mediated at least in part by inhibition of p53 via enhanced translation of MDM2 (Kao et al., 2009). The authors reported that rapamycin increases the p53/MDM2 protein ratio by inhibition of MDM2 translation without affecting its mRNA expression or protein stability. Ectopic expression of eIF4E abrogated the effect. With some gaps yet to be filled, the accumulated data indicate the existence of a signaling axis in T-ALL cells connecting the following components NOTCH1-MYC-mTOR-eIF4E-MDM2-p53 (see Figure 2 for a complete representation of the pathway).

NOTCH1 was shown to positively regulate the ubiquitin ligase SKP2 and, as a result, downregulate the p27 Kip1 and p21 Cip1 cell cycle inhibitors in T-ALL cell lines (Dohda et al., 2007; Sarmiento et al., 2005). Moreover, SKP2 was shown to inhibit p53 function by targeting the acetyltransferase p300 (Kitagawa et al., 2008). Notably, MYC protein is targeted for degradation by SKP2 (Kim et al., 2003; von der Lehr et al., 2003) and SKP2 is not required for the transforming function of MYC (Old et al., 2010). However, as mentioned earlier, MYC gene expression is directly activated by NOTCH1 (Weng et al., 2006; Sharma et al., 2007). It is possible therefore that NOTCH1-induced SKP2 counteracts the NOTCH1-mediated transcriptional activation of MYC to keep MYC protein levels within a pro-survival range. Recently published data suggest that this is indeed the case. It was reported (although not discussed by the authors) that downregulation of NIC in murine T-cell lymphomas expressing MYC under a NOTCH1-independent constitutive promoter showed very strong activation of MYC protein levels without affecting mRNA levels [see Figure 4A and C in (Demarest et al., 2011)]; of note, the process coincided with strong upregulation of p53 protein and tumor regression. Because overexpression of MYC was documented to activate oncogene-induced stress via p53 in ARF-dependent and -independent manners (Murphy et al., 2008), it is tempting to propose that another potential NOTCH1 survival function is mediated by SKP2 control of MYC protein levels. This mechanism may play a particularly prominent role in those cases when, as a result of chromosomal rearrangements, MYC transcription is constitutive and no longer dependent on NOTCH1.

Finally, it is important to note that NOTCH was shown to activate NF κ B transcriptional function on multiple levels that includes upregulation of expression of NF κ B subunits,




Fig. 2. NOTCH and p53 network. The function of NOTCH1 is mediated by several signaling hubs that in turn impact ARF and p53 function: NOTCH1 activates AKT, AKT directly activates MDM2. NOTCH activates SKP2 and Skp2 suppresses p300-mediated acetylation of p53 and the transactivation ability of p53. MYC gene expression is directly activated by NOTCH; MYC protein levels are controlled by SKP2 and AKT; ectopic MYC activation may cause activation of p53. NOTCH and/or MYC activate mTOR and eIF4E; eIF4E mediates protein synthesis of MYC and activators of AKT kinase; MYC and eIF4E promote CAP-mediated translation of MDM2. Solid arrows indicate direct interaction, dashed arrows indicate functional interaction via one or more intermediaries.

direct interaction with its upstream regulatory components such as IKK kinase, and inhibition of a negative loop of regulation (Osipo et al., 2008; Shin et al., 2006; Espinosa et al., 2010). NF κ B and p53 exhibit a well documented history of cross-talk as well as synergistic interactions. For example, p53 and the p52 NF κ B subunit coordinately regulate SKP2 gene expression (Barre & Perkins, 2010). NF κ B is not only a pro-survival factor, it was shown to activate apoptosis in response to chemotherapeutic drug treatments (Radhakrishnan & Kamalakaran, 2006). Moreover, NF κ B can induce p53 function, while p53 was shown to selectively inhibit the survival function of NF κ B but to cooperate with the NF κ B-mediated transcriptional activation of apoptotic genes (Meley et al., 2010; Ryan et al., 2000). Thus the functional outcome of NOTCH1-NF κ B-p53 pathway interaction is not easy to predict; and, depending on the conditions (e.g., chemotherapy-induced stress levels), the NOTCH1-NF κ B-p53 pathways may cooperate in promoting apoptosis. In addition, our data indicate that NOTCH1 may positively contribute to NF κ B apoptotic function in a p53-independent manner.

We believe therefore that additional studies should be carried out to address the conflicting laboratory and clinical findings about the role of NOTCH1 activation in the regulation of T-

ALL cell survival in response to therapy. Briefly, as mentioned above, a number of preclinical studies have demonstrated a pro-survival role of NOTCH signaling in T-ALL; based on these studies, T-ALL clinical trials investigating the therapeutic potential of γ -secretase inhibitors have been initiated (ClinicalTrials.gov, NCT01088763; NCT00878189); so far, the drugs tested have demonstrated low anti-leukemic efficacy. A novel approach targeting NOTCH1 processing via inhibition of ADAM10 (a disintegrin and metalloprotease 10) was also suggested (Sulis et al., 2011). Moreover, on the basis of results obtained with T-ALL cell lines, in which inhibition of NOTCH signaling was reported to enhance glucocorticoid sensitivity, combination therapies of γ -secretase inhibitors and glucocorticoids were proposed (Real & Ferrando, 2009; Real et al., 2009). Clearly, the inclusion of γ -secretase inhibitors in T-ALL protocols needs to be reevaluated in light of the new clinical data showing that activated NOTCH1 is associated with a better initial response regardless of the type of treatment and particularly to prednisone (Kox et al., 2010). On the other hand, perhaps if it is not combined with conventional therapy, NOTCH1 inhibition may prove to be a successful approach; for example, in combination with Sonic hedgehog inhibition (Okuhashi et al., 2011) or mTOR and PTEN-PI3K/AKT modulation (see below).

4. PTEN: AKT-dependent and -independent activation of p53

Recurring oncogenic events in T-ALL involve inactivation of the PTEN tumor suppressor gene (Zhang et al., 2006; Palomero et al., 2008). The frequency of PTEN mutations was estimated to be about 20%; however, its functional downregulation is more common (Jotta et al., 2010; Silva et al., 2008). PTEN is a lipid phosphatase hydrolyzing phosphate in position 3 from phosphoinositides. In primary T-ALL, PTEN was suggested to be a major factor contributing to elevated levels of phosphoinositides and thus indirectly contributing to MYC protein stability (Silva et al., 2010; Bonnet et al., 2011).

Phosphoinositides, such as phosphatidylinositol-(3,4,5)-trisphosphate (PIP3), are membrane second messengers connecting cytokine and growth factor signaling with intracellular components such as the serine threonine protein kinase AKT (Carracedo & Pandolfi, 2008). PTEN dephosphorylates PIP3 while phosphoinositide-3 kinase (PI3K) reverses the reaction such that the levels of PIP3 are controlled by the balance of these two enzymes (Figure 3). PI3K is activated by various genetic lesions, the most common of which activate NOTCH1 (Sade et al., 2004). In addition, about 20% of mutations are in genes encoding upstream activators of the PI3K pathway; among them, IGF signaling components are the most prominent (Remke et al., 2009) while mutations directly affecting PI3K subunits are more rare (Gutierrez et al., 2009). Thus, as a result of PTEN inhibition and/or PI3K activation, close to 90% of T-ALL show elevated PIP3 levels (Silva et al., 2008). The signal from PIP3 is transmitted via membrane recruitment and activation of a member of the pleckstrin homology domain protein family, one of the best characterized being AKT kinase. Indeed in T-ALL, AKT levels were shown to be activated very frequently (close to 90%) (Silva et al., 2008). AKT phosphorylates about 100 different proteins and mainly promotes survival (by inactivating BAD, MDM2 and forkhead transcription factors) and growth (by inhibition of p27, GSK3 kinase), and regulates glucose homeostasis (by enhancing the glucose transporter GLUT4). Importantly, AKT regulates protein translation and ribosome biogenesis via activation of the mTOR pathway (Carracedo & Pandolfi, 2008). Therefore, the contribution of the PTEN/PI3K-AKT pathway is of central importance to the pathobiology of T-ALL, and especially with regard to p53 regulation.




Fig. 3. Role of PTEN in p53 regulation. PTEN hydrolyses the phosphate group in the 3' position from phosphatidylinositol 3,4,5-triphosphate (PIP3) to form phosphatidylinositol 4,5-biphosphate that counteracts PI3K function. PIP3 activates AKT. AKT phosphorylates and activates MDM2 directly and via mTOR pathways contributes to its protein synthesis. PTEN directly binds p53, enhancing its stability by antagonizing the p53-MDM2 interaction and promoting p300/CBP-mediated acetylation of p53. On the other hand, ectopic AKT activation may induce ARF function, and ectopic MYC activation may induce p53 in ARF-dependent and -independent ways. Solid arrows indicate direct interaction, dashed arrows indicate functional interaction via one or more intermediaries.

Most of the characterized downstream effectors of PTEN are AKT dependent; in T-ALL, they include p53, mTOR and MYC. p53 is regulated by AKT phosphorylation of MDM2 that leads to its nuclear translocation (Mayo & Donner, 2001). In addition, PTEN may directly bind p53 protein, enhance its stability by antagonizing p53-MDM2 interaction, and promote p300/CBP-mediated acetylation of p53 (Zhou et al., 2003; Freeman et al., 2003). Gain or loss of function experiments in T-ALL demonstrated that the PTEN-mTOR axis is important for the growth of the leukemogenic cells (Yilmaz et al., 2006). As previously mentioned, mTOR was also shown to enhance translation of MDM2. A recent publication addressing the frequency of MYC deregulation in T-ALL demonstrated that downregulation of PTEN function is one of the predominant features associated with enhanced MYC protein levels (Bonnet et al., 2011). MYC is regulated via AKT-dependent inhibition of GSK3 kinase. If it is not inhibited by AKT, this constitutively active kinase directly targets MYC protein for degradation (Gregory et al., 2003).

PTEN is a haploinsufficient tumor suppressor since the presence of a single copy does not prevent cancers (Salmena et al., 2008). On the other hand, biallelic loss of PTEN in primary thymocytes causes a cellular stress response resulting in the AKT-dependent induction of cell cycle arrest (Xue et al., 2008) via elevated expression of ARF and p53 proteins (Lee et al., 2010). The data are consistent with earlier work observing that loss of PTEN induces p53 function in other cell types (Chen et al., 2005b; Kim et al., 2007a). Together, the data strongly demonstrate that complete loss of PTEN may activate an oncogene-induced stress response and explains why it usually occurs in advanced cancers with inactivated p53 genes and poor prognosis (Jotta et al., 2010; Gutierrez et al., 2009).

Thus cancer intervention via modulation of PTEN function requires a precise knowledge of PTEN and p53 statuses. For p53 positive cases, PTEN inhibition was proposed as a therapeutic strategy (Mak et al., 2010). A more widely applicable approach is therapeutic restoration/activation of PTEN function. The activity of PTEN is directly inhibited by reactive oxygen species (ROS) oxidation of its catalytic center. Since leukemic cells frequently show increased levels of ROS, antioxidants may contribute to the restoration of PTEN function (Silva et al., 2008). For example, ascorbic acid or resveratrol treatment of T-ALL cell lines was associated with activation of p53; however, PTEN function was not addressed (Harakeh et al., 2007; Cecchinato et al., 2007). The levels of PTEN protein expression are tightly regulated. In T-ALL cells, CK2 was shown to control its protein levels and thus CK2 inhibition was suggested as a therapeutic strategy (Silva et al., 2010). Inhibition of PI3K, AKT and mTOR kinases was also suggested (Chiarini et al., 2010; Chiarini et al., 2009; Evangelisti et al., 2011a; Evangelisti et al., 2011b). However, despite a significant overlap in downstream targets, PTEN loss cannot compensate for NOTCH1 oncogenic function (Medyouf et al., 2010). Thus inhibition of both pathways was shown to cooperate in primary leukemic T cells and in mouse tumor models (Guo et al., 2011; Cullion et al., 2009). Finally, dual inhibition of PI3K and mTOR has been suggested as a therapeutic option for T-ALL (Chiarini et al., 2009).

5. TP53 coexpressed genes — potential chemotherapeutic targets

As discussed above, genetic aberrations may cause activation of a stress response. For example, deregulation of the NOTCH, PTEN, TAL1 or LYL1 loci causes activation of ARF in normal thymocytes. TP53 gene expression can also be induced in response to stress (Vilborg et al., 2010). Despite the secondary adaptive mutations preventing oncogene-induced apoptosis or senescence, cancer cells still frequently show elevated stress levels. The elevated stress together with enhanced growth is exploited in cancer therapies, which aim to selectively kill tumor cells while sparing normal cells. Knowing the transcription programs mediating the stress phenotype of cancer cells is important for the rational design of new “targeted” treatment strategies (Luo et al., 2009).

While analyzing publicly available expression profiles of primary T-ALL cells, we noticed that expression levels of TP53 varied greatly between patient samples. Moreover, in the majority of cases, TP53 was expressed at higher levels than in normal thymocytes, indicating that the complexity of aberrations in T-ALL manifests in various levels of oncogene-induced stress. We asked what transcripts are coregulated with TP53 with the expectation of identifying genes that functionally interact with it. We analyzed expression

profiles of primary T-ALL samples and normal thymocytes (Soulier et al., 2005). Genes were selected for analysis that were coexpressed with TP53 (Affymetrix probe set 211300_s_at U133A) based on similarity of their expression patterns ($r > 0.65$) within about 100 T-ALL patient samples. As expected, the set of TP53-profile neighbors contained a number of genes encoding proteins that interact with TP53 at the mRNA or protein levels, regulating its stability and function (Figure 4). Interestingly, among them were genes that counteract p53 function, possibly reflecting neoplastic adaptation to high levels of TP53 gene expression. For example, inactivation of p53 at the level of protein stability is illustrated by PA2G4, PSME3 and HUWE1. Briefly, PA2G4 promotes p53 polyubiquitination and degradation; PSME3, encoding the 26S proteasome subunit, was shown to be required for p53 degradation (Zhang & Zhang, 2008); and HUWE1 (or ARF-BP1) E3 ubiquitin ligase directly binds to p53 and targets it for degradation in an MDM2-independent manner (Chen et al., 2005a). Other examples include G3BP1, which facilitates redistribution of p53 from the nucleus to the cytoplasm (Kim et al., 2007b); UBE2N (or UBC13), which inhibits formation of transcriptionally active p53 tetramers (Topisirovic et al., 2009); and CHD4, which deacetylates p53 and blocks p21 induction (Polo et al., 2010). Finally, inhibition of p53 transcriptional outcome is exemplified by YBX1, which is a component of the repressor complex blocking expression of p53 target genes (Shiota et al., 2008; Kim et al., 2008). Proteins cooperating with p53 function were also found among the set of p53-profile neighbors; for example, HNRNPF promotes p53 mRNA 3'-end formation (Decorsiere et al., 2011); DKC1 facilitates p53 translation; heat shock-induced stabilization of p53 occurs via direct binding to HSP90AA1; the purine biosynthesis enzyme GART is involved in p53-activating posttranslational modifications (Bronder & Moran, 2003); and SSRP1 is a component of the p53 transcriptional complex (Keller & Lu, 2002; Keller et al., 2001). There were also examples of genetic cooperation such as NOLC1 and SMARCC1. TP53 and NOLC1 cooperate in snoRNA-mediated ribosomal RNA editing, an important process for stress-induced stabilization of ribosomes (Krastev et al., 2011). Haploinsufficiency of both SMARCC1 and p53 cooperate to induce tumorigenesis in a mouse model (Ahn et al., 2011).

Because p53 is known to interact in some manner with a large portion of the genome, we asked if the enrichment of the p53-interacting genes in the set of TP53-profile neighbors is statistically significant. An empirical approach was used to determine the p-value. We generated 100 sets of 100 genes randomly selected from all annotated genes. For each set of 100 genes, TP53 was added to the set. The resulting simulated gene sets were subjected to Ingenuity Pathway Analysis and treated the same way as the list of T-ALL TP53-profile neighbors. The tabulated results of the number of p53 connections identified were used to estimate the p-value: this statistical approach indicated that the number of p53 connections within the set of T-ALL TP53-profile neighbors significantly exceeded the number of those within the simulated randomly-selected gene sets (p -value < 0.01). Importantly, our statistical analysis suggests that other less-studied genes among the T-ALL TP53-profile neighbors may also be important regulators of p53. This idea warrants further investigation because novel regulators of p53 might be found that may serve as potential therapeutic targets (Cheok et al., 2011). In support of this notion, certain of the known p53-interacting proteins, such as HUWE1, have already been suggested as possibilities for therapeutic intervention to restore p53 function in cancer cells (Chen et al., 2006).




Fig. 4. TP53 profile neighbors in T-ALL regulate its function. Shown are examples of the TP53 profile neighbors known to target sequential steps of p53 activation.

The enrichment of p53-interacting genes and common expression signature also indicated that the set of TP53-profile neighbors is deregulated as a result of leukemogenesis and may be targeted by common T-ALL oncogenes. To ask which oncogenes may be involved in regulation of the set of TP53-coexpressed genes in T-ALL, we used genome-wide chromatin immunoprecipitation data previously published by others (Margolin et al., 2009; De Keersmaecker et al., 2010) to identify direct MYC or TLX1 targets. We found that the set of T-ALL TP53-coexpressed genes is significantly enriched for direct MYC and/or TLX1 targets, accounting for 70% of the set. Specifically, out of 16,697 genes represented on the array, 8,404 of them were bound by either MYC or TLX1. On the other hand, out of the 99 genes found to be coexpressed with TP53 in T-ALL, 69 of them were bound by either MYC or TLX1. Using the hypergeometric distribution, we determined that the p-value for the frequency of MYC or TLX1 target genes within the set of T-ALL p53-profile neighbors is 7×10^{-5} . Notably, NOTCH1 targets were within the subsets of MYC and/or

TLX1 targets. Moreover, the remaining 30% of the genes showed similar expression profiles and functional classification, arguing that they are regulated by one of NOTCH1, MYC or TLX1, but indirectly via downstream transcription factors. Promoter analysis revealed that the incidence of glucocorticoid receptor (GR) binding sites was 10 times more frequent in this set versus the genes directly targeted by MYC and TLX1, suggesting that these genes are potential targets of GR, and that GR may be the transcription factor contributing to the oncogene-induced stress response. A number of observations support this hypothesis. First, TLX1-positive T-ALL cases are characterized by high GR mRNA expression levels (Ferrando et al., 2002). In this regard, we found that shRNA-mediated knockdown of TLX1 in the T-ALL-derived ALL-SIL cell line was associated with increased resistance to glucocorticoid-induced cell death (manuscript in preparation) indicating that TLX1 may contribute positively to GR function. Moreover, recent clinical studies have demonstrated that activation of the NOTCH1 oncogene is associated with a superior initial therapeutic response to glucocorticoids (Kox et al., 2010) indicating that NOTCH1 may also cooperate with GR-induced killing. Finally, we and others observed that ectopic activation of MYC may cause transcriptional induction of pro-apoptotic BIM (Riz et al., 2011), which is a known mediator of GR induced apoptosis in T-ALL cells. Thus we hypothesize based on our analysis that an interacting network of transcription factors – NOTCH1-MYC-TLX1 – may activate the TP53-anchored transcriptional program of an oncogene-induced stress response, predominantly via direct binding to promoters and in part via activation of GR function. We hope that our hypothesis will help to stimulate further studies seeking novel therapeutic targets to restore p53 function and to understand the intricate relationships between NOTCH1 and GR, two major targets of T-ALL therapeutic intervention.

6. Interaction of T-ALL mutations and p53 downstream targets

It is important to appreciate that not only p53 function but also execution of p53-governed transcriptional programs is often compromised by T-ALL mutations. For example, NOTCH1 via activation of SKP2 decreases the levels of p21 and thus counteracts one of the best characterized activities of p53 (Sarmiento et al., 2005). As discussed above, the pathways downstream of NOTCH and PTEN intersect at the level of PI3K-mTOR. A set of p53 target genes controls this pathway as well (Feng et al., 2005). PTEN itself was shown to be a p53 target gene about 10 years ago (Stambolic et al., 2001). Since then, it has become appreciated that PTEN is activated by p53 in response to high-dose chemotherapy as part of a p53-governed transcriptional program committing cells to apoptosis. NOTCH1 was shown to inhibit PTEN via upregulation of the transcription factor HES1, which directly represses the PTEN promoter (Palomero et al., 2007). Among upstream modulators of PI3K, p53 induces IGF-BP3 (Buckbinder et al., 1995). IGF-BP3 binds to IGF1 or IGF2 and prevents their interaction with the receptor. Mutations involving components of IGF signaling are frequent contributors to PI3K activation in T-ALL (Remke et al., 2009). Other p53 targets affecting the mTOR pathway include TSC2, AMPK beta1, sestrins 1 and 2 and REDD1, all of which contribute to negative regulation of mTOR by targeting the TORC1 complex (which counteracts AKT function) (Feng et al., 2007) (Figure 5).




Fig. 5. T-ALL-associated genetic lesions compromise the function of p53 downstream targets. Alterations of NOTCH1 and PTEN loci are complemented by less frequent mutations in the growth-promoting IGF-PI3K/AKT/mTOR network; mutant proteins are indicated by 'm'. By inhibiting this network, the p53 target genes shown adjust metabolic rates in response to stress conditions and stall cell cycle progression.

7. Conclusion

A plethora of inactivating mutations notwithstanding there is still a possibility for therapeutic restoration of p53 apoptotic function because of two major features typical to this cellular regulator: multiplicity of activating stimuli and redundancy of the activating modifications. p53 is activated in response to a variety of stresses such as lack of nutrients, energy deprivation, DNA damage, heat shock, hypoxia or enhanced oxidation, and ER protein overload. Importantly, the apoptotic function of p53 may be activated only in the presence of persistent irreparable stress. For example, while reparable DNA damage activates p53 only partially via Ser-15/20 phosphorylation (which is sufficient for cell cycle arrest), PTEN induction by p53 is triggered by persistent DNA damage and has an additional checkpoint that requires phosphorylation of Ser46 in p53 (Mayo et al., 2005; Zhang et al., 2011). Thus chemotherapeutic agents activating p53 beyond its growth-arresting function should be considered as an aid to p53 protein stabilizers such as blockers of MDM2 function (Hasegawa et al., 2009). Among these regulators, WIP1 phosphatase (PPM1D), a p53 target and negative loop of autoregulation of p53 was suggested as a possibility, however, not tested in T-ALL (Lu et al., 2008; Yoda et al., 2008). Even though the Ser46 phosphorylated form of p53 was shown to associate with apoptotic activity, point mutation substituting this amino acid to alanine did not prevent activation of p53 apoptotic function (Kurihara et al., 2007). This illustration, together with studies showing that the p53-

MDM2 interaction is not affected by single point mutations, supports the idea of p53 regulatory redundancy (Kruse & Gu, 2009).

At the onset of neoplastic development, p53 is often activated as part of an oncogene-induced stress response. It is noteworthy that TP53 is rarely mutated in T-ALL (De Keersmaecker et al., 2005). Despite adaptive genetic and epigenetic mechanisms disrupting its functional outcome [discussed herein and (Vilas-Zornoza et al., 2011)], the p53 pathway still stays partially activated in fully developed T-ALL. Thus selective reactivation of p53 tumor suppressor function in the malignant cells is possible in principle by overcoming the disrupted links of the pathway. The development of personalized medicine providing knowledge of the patient's cancer genome should facilitate efforts to devise the appropriate strategy to activate wild-type p53 function in T-ALL. We believe that combining conventional cytotoxic therapy with molecular targeted approaches restoring p53 activity to its full potential will improve current protocols and prevent relapse. In this regard, there are several small molecule inhibitors of the p53-MDM2 interaction that are currently being investigated (Cheek et al., 2011), at least one of which (RG7112) is undergoing clinical trials in T-ALL patients (ClinicalTrials.gov, NCT00623870).

In spite of over 30 years of p53 research and investigation into the molecular basis of T-ALL, surprisingly little is known about the role and function of p53 in T-ALL. At time of writing, 43,847 PubMed articles were found by searching for "p53 and cancer" whereas only 146 articles could be retrieved for "p53 and T-ALL". T-ALL represents 15% of pediatric hematological malignancies which are the most common cancers in children. So, we believe that the subject is significantly underrepresented. We hope that by summarizing the current state of the art, this chapter will bring more attention to this issue and pave the way for new therapeutic strategies for patients with this disease.

8. Acknowledgments

We thank Sana Ahmad for assistance determining the statistical significance of p53 connections within the set of TP53-profile neighbors and Ian Toma for help preparing the figures. This work was supported in part by an American Cancer Society Pilot Grant for Junior Faculty from The George Washington University Cancer Institute (IRG-08-091-01) and a Pilot Project Award from the Clinical and Translational Science Institute at Children's National Medical Center (NIH UL1RR031988) (I.R.); and by NIH Grants R01HL65519 and R01HL66305, an Elaine H. Snyder Cancer Research Award and a King Fahd Endowment from The George Washington University School of Medicine (R.G.H.). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the National Center for Research Resources or the National Institutes of Health. Correspondence: I.R. (anaixr@gwumc.edu) or R.G.H. (rghawley@gwu.edu)

9. References

- Ahn, J., Ko, M., Lee, C., Kim, J., Yoon, H., & Seong, R.H. (2011). Srg3, a mouse homolog of BAF155, is a novel p53 target and acts as a tumor suppressor by modulating p21(WAF1/CIP1) expression. *Oncogene* 30, 445-456.
- Asnafi, V., Buzyn, A., Le, N.S., Baleyrier, F., Simon, A., Beldjord, K., Reman, O., Witz, F., Fagot, T., Tavernier, E., et al. (2009). NOTCH1/FBXW7 mutation identifies a large subgroup with favorable outcome in adult T-cell acute lymphoblastic leukemia (T-

- ALL): a Group for Research on Adult Acute Lymphoblastic Leukemia (GRAALL) study. *Blood* 113, 3918-3924.
- Aster, J.C., Pear, W.S., & Blacklow, S.C. (2008). Notch signaling in leukemia. *Annu. Rev. Pathol.* 3, 587-613.
- Barna, M., Pusic, A., Zollo, O., Costa, M., Kondrashov, N., Rego, E., Rao, P.H., & Ruggero, D. (2008). Suppression of Myc oncogenic activity by ribosomal protein haploinsufficiency. *Nature* 456, 971-975.
- Barre, B. & Perkins, N.D. (2010). The Skp2 promoter integrates signaling through the NF-kappaB, p53, and Akt/GSK3beta pathways to regulate autophagy and apoptosis. *Mol. Cell* 38, 524-538.
- Bertwistle, D., Sugimoto, M., & Sherr, C.J. (2004). Physical and functional interactions of the Arf tumor suppressor protein with nucleophosmin/B23. *Mol. Cell Biol.* 24, 985-996.
- Beverly, L.J., Felsher, D.W., & Capobianco, A.J. (2005). Suppression of p53 by Notch in lymphomagenesis: implications for initiation and regression. *Cancer Res.* 65, 7159-7168.
- Bianchi, T., Rufer, N., MacDonald, H.R., & Migliaccio, M. (2006). The tumor suppressor p16Ink4a regulates T lymphocyte survival. *Oncogene* 25, 4110-4115.
- Bonnet, M., Loosveld, M., Montpellier, B., Navarro, J.M., Quilichini, B., Picard, C., Di Cristofaro, J., Bagnis, C., Fossat, C., Hernandez, L., et al. (2011). Posttranscriptional deregulation of MYC via PTEN constitutes a major alternative pathway of MYC activation in T-cell acute lymphoblastic leukemia. *Blood* 117, 6650-6659.
- Bracken, A.P., Kleine-Kohlbrecher, D., Dietrich, N., Pasini, D., Gargiulo, G., Beekman, C., Theilgaard-Monch, K., Minucci, S., Porse, B.T., Marine, J.C., et al. (2007). The Polycomb group proteins bind throughout the INK4A-ARF locus and are disassociated in senescent cells. *Genes Dev.* 21, 525-530.
- Brady, S.N., Yu, Y., Maggi, L.B., Jr., & Weber, J.D. (2004). ARF impedes NPM/B23 shuttling in an Mdm2-sensitive tumor suppressor pathway. *Mol. Cell Biol.* 24, 9327-9338.
- Bronder, J.L. & Moran, R.G. (2003). A defect in the p53 response pathway induced by de novo purine synthesis inhibition. *J. Biol. Chem.* 278, 48861-48871.
- Buckbinder, L., Talbott, R., Velasco-Miguel, S., Takenaka, I., Faha, B., Seizinger, B.R., & Kley, N. (1995). Induction of the growth inhibitor IGF-binding protein 3 by p53. *Nature* 377, 646-649.
- Canepa, E.T., Scassa, M.E., Ceruti, J.M., Marazita, M.C., Carcagno, A.L., Sirkin, P.F., & Ogara, M.F. (2007). INK4 proteins, a family of mammalian CDK inhibitors with novel biological functions. *IUBMB. Life* 59, 419-426.
- Carracedo, A. & Pandolfi, P.P. (2008). The PTEN-PI3K pathway: of feedbacks and cross-talks. *Oncogene* 27, 5527-5541.
- Cecchinato, V., Chiaramonte, R., Nizzardo, M., Cristofaro, B., Basile, A., Sherbet, G.V., & Comi, P. (2007). Resveratrol-induced apoptosis in human T-cell acute lymphoblastic leukaemia MOLT-4 cells. *Biochem. Pharmacol.* 74, 1568-1574.
- Cha, T.L., Zhou, B.P., Xia, W., Wu, Y., Yang, C.C., Chen, C.T., Ping, B., Otte, A.P., & Hung, M.C. (2005). Akt-mediated phosphorylation of EZH2 suppresses methylation of lysine 27 in histone H3. *Science* 310, 306-310.
- Chan, S.M., Weng, A.P., Tibshirani, R., Aster, J.C., & Utz, P.J. (2007). Notch signals positively regulate activity of the mTOR pathway in T-cell acute lymphoblastic leukemia. *Blood* 110, 278-286.

- Chen, D., Brooks, C.L., & Gu, W. (2006). ARF-BP1 as a potential therapeutic target. *Br. J. Cancer* 94, 1555-1558.
- Chen, D., Kon, N., Li, M., Zhang, W., Qin, J., & Gu, W. (2005a). ARF-BP1/Mule is a critical mediator of the ARF tumor suppressor. *Cell* 121, 1071-1083.
- Chen, Z., Trotman, L.C., Shaffer, D., Lin, H.K., Dotan, Z.A., Niki, M., Koutcher, J.A., Scher, H.I., Ludwig, T., Gerald, W., et al. (2005b). Crucial role of p53-dependent cellular senescence in suppression of Pten-deficient tumorigenesis. *Nature* 436, 725-730.
- Cheok, C.F., Verma, C.S., Baselga, J., & Lane, D.P. (2011). Translating p53 into the clinic. *Nat. Rev. Clin. Oncol.* 8, 25-37.
- Chiarini, F., Fala, F., Tazzari, P.L., Ricci, F., Astolfi, A., Pession, A., Pagliaro, P., McCubrey, J.A., & Martelli, A.M. (2009). Dual inhibition of class IA phosphatidylinositol 3-kinase and mammalian target of rapamycin as a new therapeutic option for T-cell acute lymphoblastic leukemia. *Cancer Res.* 69, 3520-3528.
- Chiarini, F., Grimaldi, C., Ricci, F., Tazzari, P.L., Evangelisti, C., Ognibene, A., Battistelli, M., Falciari, E., Melchionda, F., Pession, A., et al. (2010). Activity of the novel dual phosphatidylinositol 3-kinase/mammalian target of rapamycin inhibitor NVP-BEZ235 against T-cell acute lymphoblastic leukemia. *Cancer Res.* 70, 8097-8107.
- Ciofani, M. & Zuniga-Pflucker, J.C. (2005). Notch promotes survival of pre-T cells at the beta-selection checkpoint by regulating cellular metabolism. *Nat. Immunol.* 6, 881-888.
- Cullion, K., Draheim, K.M., Hermance, N., Tammam, J., Sharma, V.M., Ware, C., Nikov, G., Krishnamoorthy, V., Majumder, P.K., & Kelliher, M.A. (2009). Targeting the Notch1 and mTOR pathways in a mouse T-ALL model. *Blood* 113, 6172-6181.
- De Keersmaecker, K., Lahortiga, I., Mentens, N., Folens, C., Van Neste, L., Bekaert, S., Vandenberghe, P., Odero, M.D., Marynen, P., & Cools, J. (2008). *In vitro* validation of gamma-secretase inhibitors alone or in combination with other anti-cancer drugs for the treatment of T-cell acute lymphoblastic leukemia. *Haematologica* 93, 533-542.
- De Keersmaecker, K., Marynen, P., & Cools, J. (2005). Genetic insights in the pathogenesis of T-cell acute lymphoblastic leukemia. *Haematologica* 90, 1116-1127.
- De Keersmaecker, K., Real, P.J., Gatta, G.D., Palomero, T., Sulis, M.L., Tosello, V., Van Vlierberghe, P., Barnes, K., Castillo, M., Sole, X., et al. (2010). The TLX1 oncogene drives aneuploidy in T cell transformation. *Nat. Med.* 16, 1321-1327.
- Decorsiere, A., Cayrel, A., Vagner, S., & Millevoi, S. (2011). Essential role for the interaction between hnRNP H/F and a G quadruplex in maintaining p53 pre-mRNA 3'-end processing and function during DNA damage. *Genes Dev.* 25, 220-225.
- Demarest, R.M., Dahmane, N., & Capobianco, A.J. (2011). Notch is oncogenic dominant in T-cell acute lymphoblastic leukemia. *Blood* 117, 2901-2909.
- Dohda, T., Maljukova, A., Liu, L., Heyman, M., Grander, D., Brodin, D., Sangfelt, O., & Lendahl, U. (2007). Notch signaling induces SKP2 expression and promotes reduction of p27Kip1 in T-cell acute lymphoblastic leukemia cell lines. *Exp. Cell Res.* 313, 3141-3152.
- El Andaloussi, A., Graves, S., Meng, F., Mandal, M., Mashayekhi, M., & Aifantis, I. (2006). Hedgehog signaling controls thymocyte progenitor homeostasis and differentiation in the thymus. *Nat. Immunol.* 7, 418-426.

- Ellisen, L.W., Bird, J., West, D.C., Soreng, A.L., Reynolds, T.C., Smith, S.D., & Sklar, J. (1991). TAN-1, the human homolog of the Drosophila notch gene, is broken by chromosomal translocations in T lymphoblastic neoplasms. *Cell* 66, 649-661.
- Espinosa, L., Cathelin, S., D'Altri, T., Trimarchi, T., Statnikov, A., Guiu, J., Rodilla, V., Ingles-Esteve, J., Nomdedeu, J., Bellosillo, B., et al. (2010). The Notch/Hes1 pathway sustains NF-kappaB activation through CYLD repression in T cell leukemia. *Cancer Cell* 18, 268-281.
- Evangelisti, C., Ricci, F., Tazzari, P., Chiarini, F., Battistelli, M., Falcieri, E., Ognibene, A., Pagliaro, P., Cocco, L., McCubrey, J.A., et al. (2011a). Preclinical testing of the Akt inhibitor triciribine in T-cell acute lymphoblastic leukemia. *J. Cell Physiol* 226, 822-831.
- Evangelisti, C., Ricci, F., Tazzari, P., Tabellini, G., Battistelli, M., Falcieri, E., Chiarini, F., Bortul, R., Melchionda, F., Pagliaro, P., et al. (2011b). Targeted inhibition of mTORC1 and mTORC2 by active-site mTOR inhibitors has cytotoxic effects in T-cell acute lymphoblastic leukemia. *Leukemia* 25, 781-791.
- Feng, Z., Hu, W., de Stanchina, E., Teresky, A.K., Jin, S., Lowe, S., & Levine, A.J. (2007). The regulation of AMPK beta1, TSC2, and PTEN expression by p53: stress, cell and tissue specificity, and the role of these gene products in modulating the IGF-1-AKT-mTOR pathways. *Cancer Res.* 67, 3043-3053.
- Feng, Z., Zhang, H., Levine, A.J., & Jin, S. (2005). The coordinate regulation of the p53 and mTOR pathways in cells. *Proc. Natl. Acad. Sci. USA* 102, 8204-8209.
- Ferrando, A.A., Neuberg, D.S., Staunton, J., Loh, M.L., Huard, C., Raimondi, S.C., Behm, F.G., Pui, C.H., Downing, J.R., Gilliland, D.G., et al. (2002). Gene expression signatures define novel oncogenic pathways in T cell acute lymphoblastic leukemia. *Cancer Cell* 1, 75-87.
- Freeman, D.J., Li, A.G., Wei, G., Li, H.H., Kertesz, N., Lesche, R., Whale, A.D., Martinez-Diaz, H., Rozengurt, N., Cardiff, R.D., et al. (2003). PTEN tumor suppressor regulates p53 protein levels and activity through phosphatase-dependent and -independent mechanisms. *Cancer Cell* 3, 117-130.
- Gardie, B., Cayuela, J.M., Martini, S., & Sigaux, F. (1998). Genomic alterations of the p19ARF encoding exons in T-cell acute lymphoblastic leukemia. *Blood* 91, 1016-1020.
- Gil, J. & Peters, G. (2006). Regulation of the INK4b-ARF-INK4a tumour suppressor locus: all for one or one for all. *Nat. Rev. Mol. Cell Biol.* 7, 667-677.
- Grabher, C., von Boehmer, H., & Look, A.T. (2006). Notch 1 activation in the molecular pathogenesis of T-cell acute lymphoblastic leukaemia. *Nat. Rev. Cancer* 6, 347-359.
- Gregory, M.A., Qi, Y., & Hann, S.R. (2003). Phosphorylation by glycogen synthase kinase-3 controls c-myc proteolysis and subnuclear localization. *J. Biol. Chem.* 278, 51606-51612.
- Guo, D., Teng, Q., & Ji, C. (2011). NOTCH and phosphatidylinositide 3-kinase/phosphatase and tensin homolog deleted on chromosome ten/AKT/mammalian target of rapamycin (mTOR) signaling in T-cell development and T-cell acute lymphoblastic leukemia. *Leuk. Lymphoma*.
- Gutierrez, A., Sanda, T., Grebliunaite, R., Carracedo, A., Salmena, L., Ahn, Y., Dahlberg, S., Neuberg, D., Moreau, L.A., Winter, S.S., et al. (2009). High frequency of PTEN, PI3K, and AKT abnormalities in T-cell acute lymphoblastic leukemia. *Blood* 114, 647-650.

- Harakeh, S., Diab-Assaf, M., Khalife, J.C., Abu-el-Ardatt, K.A., Baydoun, E., Niedzwiecki, A., El-Sabban, M.E., & Rath, M. (2007). Ascorbic acid induces apoptosis in adult T-cell leukemia. *Anticancer Res.* 27, 289-298.
- Hasegawa, H., Yamada, Y., Iha, H., Tsukasaki, K., Nagai, K., Atogami, S., Sugahara, K., Tsuruda, K., Ishizaki, A., & Kamihira, S. (2009). Activation of p53 by Nutlin-3a, an antagonist of MDM2, induces apoptosis and cellular senescence in adult T-cell leukemia cells. *Leukemia* 23, 2090-2101.
- Hebert, J., Cayuela, J.M., Berkeley, J., & Sigaux, F. (1994). Candidate tumor-suppressor genes MTS1 (p16INK4A) and MTS2 (p15INK4B) display frequent homozygous deletions in primary cells from T- but not from B-cell lineage acute lymphoblastic leukemias. *Blood* 84, 4038-4044.
- Honda, R., Tanaka, H., & Yasuda, H. (1997). Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53. *FEBS Lett.* 420, 25-27.
- Honda, R. & Yasuda, H. (1999). Association of p19(ARF) with Mdm2 inhibits ubiquitin ligase activity of Mdm2 for tumor suppressor p53. *EMBO J.* 18, 22-27.
- Hsieh, A.C. & Ruggero, D. (2010). Targeting eukaryotic translation initiation factor 4E (eIF4E) in cancer. *Clin. Cancer Res.* 16, 4914-4920.
- Jacobs, J.J., Kieboom, K., Marino, S., DePinho, R.A., & van Lohuizen, M. (1999). The oncogene and Polycomb-group gene bmi-1 regulates cell proliferation and senescence through the ink4a locus. *Nature* 397, 164-168.
- Jotta, P.Y., Ganazza, M.A., Silva, A., Viana, M.B., da Silva, M.J., Zambaldi, L.J., Barata, J.T., Brandalise, S.R., & Yunes, J.A. (2010). Negative prognostic impact of PTEN mutation in pediatric T-cell acute lymphoblastic leukemia. *Leukemia* 24, 239-242.
- Kao, C.L., Hsu, H.S., Chen, H.W., & Cheng, T.H. (2009). Rapamycin increases the p53/MDM2 protein ratio and p53-dependent apoptosis by translational inhibition of mdm2 in cancer cells. *Cancer Lett.* 286, 250-259.
- Keller, D.M. & Lu, H. (2002). p53 serine 392 phosphorylation increases after UV through induction of the assembly of the CK2.hSPT16.SSRP1 complex. *J. Biol. Chem.* 277, 50206-50213.
- Keller, D.M., Zeng, X., Wang, Y., Zhang, Q.H., Kapoor, M., Shu, H., Goodman, R., Lozano, G., Zhao, Y., & Lu, H. (2001). A DNA damage-induced p53 serine 392 kinase complex contains CK2, hSpt16, and SSRP1. *Mol. Cell* 7, 283-292.
- Kim, J.S., Lee, C., Bonifant, C.L., Resson, H., & Waldman, T. (2007a). Activation of p53-dependent growth suppression in human cells by mutations in PTEN or PIK3CA. *Mol. Cell Biol.* 27, 662-677.
- Kim, K., Choi, J., Heo, K., Kim, H., Levens, D., Kohno, K., Johnson, E.M., Brock, H.W., & An, W. (2008). Isolation and characterization of a novel H1.2 complex that acts as a repressor of p53-mediated transcription. *J. Biol. Chem.* 283, 9113-9126.
- Kim, M.M., Wiederschain, D., Kennedy, D., Hansen, E., & Yuan, Z.M. (2007b). Modulation of p53 and MDM2 activity by novel interaction with Ras-GAP binding proteins (G3BP). *Oncogene* 26, 4209-4215.
- Kim, S.Y., Herbst, A., Tworkowski, K.A., Salghetti, S.E., & Tansey, W.P. (2003). Skp2 regulates Myc protein stability and activity. *Mol. Cell* 11, 1177-1188.
- Kitagawa, M., Lee, S.H., & McCormick, F. (2008). Skp2 suppresses p53-dependent apoptosis by inhibiting p300. *Mol. Cell* 29, 217-231.

- Kopan, R. & Ilagan, M.X. (2009). The canonical Notch signaling pathway: unfolding the activation mechanism. *Cell* 137, 216-233.
- Kox, C., Zimmermann, M., Stanulla, M., Leible, S., Schrappe, M., Ludwig, W.D., Koehler, R., Tolle, G., Bandapalli, O.R., Breit, S., et al. (2010). The favorable effect of activating NOTCH1 receptor mutations on long-term outcome in T-ALL patients treated on the ALL-BFM 2000 protocol can be separated from FBXW7 loss of function. *Leukemia* 24, 2005-2013.
- Krastev, D.B., Slabicki, M., Paszkowski-Rogacz, M., Hubner, N.C., Junqueira, M., Shevchenko, A., Mann, M., Neugebauer, K.M., & Buchholz, F. (2011). A systematic RNAi synthetic interaction screen reveals a link between p53 and snoRNP assembly. *Nat. Cell Biol.* 13, 809-818.
- Kruse, J.P. & Gu, W. (2009). Modes of p53 regulation. *Cell* 137, 609-622.
- Kuo, M.L., den Besten, W., Bertwistle, D., Roussel, M.F., & Sherr, C.J. (2004). N-terminal polyubiquitination and degradation of the Arf tumor suppressor. *Genes Dev.* 18, 1862-1874.
- Kurihara, A., Nagoshi, H., Yabuki, M., Okuyama, R., Obinata, M., & Ikawa, S. (2007). Ser46 phosphorylation of p53 is not always sufficient to induce apoptosis: multiple mechanisms of regulation of p53-dependent apoptosis. *Genes Cells* 12, 853-861.
- Lee, J.Y., Nakada, D., Yilmaz, O.H., Tothova, Z., Joseph, N.M., Lim, M.S., Gilliland, D.G., & Morrison, S.J. (2010). mTOR activation induces tumor suppressors that inhibit leukemogenesis and deplete hematopoietic stem cells after Pten deletion. *Cell Stem Cell* 7, 593-605.
- Leung, C., Lingbeek, M., Shakhova, O., Liu, J., Tanger, E., Saremaslani, P., van Lohuizen, M., & Marino, S. (2004). Bmi1 is essential for cerebellar development and is overexpressed in human medulloblastomas. *Nature* 428, 337-341.
- Lewis, H.D., Leveridge, M., Strack, P.R., Haldon, C.D., O'Neil, J., Kim, H., Madin, A., Hannam, J.C., Look, A.T., Kohl, N., et al. (2007). Apoptosis in T cell acute lymphoblastic leukemia cells after cell cycle arrest induced by pharmacological inhibition of notch signaling. *Chem. Biol.* 14, 209-219.
- Li, X., Gounari, F., Protopopov, A., Khazaie, K., & von Boehmer, H. (2008). Oncogenesis of T-ALL and nonmalignant consequences of overexpressing intracellular NOTCH1. *J. Exp. Med.* 205, 2851-2861.
- Lin, C.J., Cencic, R., Mills, J.R., Robert, F., & Pelletier, J. (2008). c-Myc and eIF4F are components of a feedforward loop that links transcription and translation. *Cancer Res.* 68, 5326-5334.
- Liu, S., Breit, S., Danckwardt, S., Muckenthaler, M.U., & Kulozik, A.E. (2009). Downregulation of Notch signaling by gamma-secretase inhibition can abrogate chemotherapy-induced apoptosis in T-ALL cell lines. *Ann. Hematol.* 88, 613-621.
- Llanos, S., Clark, P.A., Rowe, J., & Peters, G. (2001). Stabilization of p53 by p14ARF without relocation of MDM2 to the nucleolus. *Nat. Cell Biol.* 3, 445-452.
- Lu, X., Nguyen, T.A., Moon, S.H., Darlington, Y., Sommer, M., & Donehower, L.A. (2008). The type 2C phosphatase Wip1: an oncogenic regulator of tumor suppressor and DNA damage response pathways. *Cancer Metastasis Rev.* 27, 123-135.
- Luo, J., Solimini, N.L., & Elledge, S.J. (2009). Principles of cancer therapy: oncogene and non-oncogene addiction. *Cell* 136, 823-837.


- Mak, L.H., Vilar, R., & Woscholski, R. (2010). Characterisation of the PTEN inhibitor VO-OHpic. *J.Chem.Biol.* 3, 157-163.
- Margolin, A.A., Palomero, T., Sumazin, P., Califano, A., Ferrando, A.A., & Stolovitzky, G. (2009). CHIP-on-chip significance analysis reveals large-scale binding and regulation by human transcription factor oncogenes. *Proc. Natl. Acad. Sci. USA* 106, 244-249.
- Mayo, L.D. & Donner, D.B. (2001). A phosphatidylinositol 3-kinase/Akt pathway promotes translocation of Mdm2 from the cytoplasm to the nucleus. *Proc. Natl. Acad. Sci. USA* 98, 11598-11603.
- Mayo, L.D., Seo, Y.R., Jackson, M.W., Smith, M.L., Guzman, J.R., Korgaonkar, C.K., & Donner, D.B. (2005). Phosphorylation of human p53 at serine 46 determines promoter selection and whether apoptosis is attenuated or amplified. *J. Biol. Chem.* 280, 25953-25959.
- Medyouf, H., Gao, X., Armstrong, F., Gusscott, S., Liu, Q., Gedman, A.L., Matherly, L.H., Schultz, K.R., Pflumio, F., You, M.J., et al. (2010). Acute T-cell leukemias remain dependent on Notch signaling despite PTEN and INK4A/ARF loss. *Blood* 115, 1175-1184.
- Meley, D., Spiller, D.G., White, M.R., McDowell, H., Pizer, B., & See, V. (2010). p53-mediated delayed NF-kappaB activity enhances etoposide-induced cell death in medulloblastoma. *Cell Death. Dis.* 1, e41.
- Miyazaki, M., Miyazaki, K., Itoi, M., Katoh, Y., Guo, Y., Kanno, R., Katoh-Fukui, Y., Honda, H., Amagai, T., van Lohuizen, M., et al. (2008). Thymocyte proliferation induced by pre-T cell receptor signaling is maintained through polycomb gene product Bmi-1-mediated Cdkn2a repression. *Immunity.* 28, 231-245.
- Mungamuri, S.K., Yang, X., Thor, A.D., & Somasundaram, K. (2006). Survival signaling by Notch1: mammalian target of rapamycin (mTOR)-dependent inhibition of p53. *Cancer Res.* 66, 4715-4724.
- Murphy, D.J., Junttila, M.R., Pouyet, L., Karnezis, A., Shchors, K., Bui, D.A., Brown-Swigart, L., Johnson, L., & Evan, G.I. (2008). Distinct thresholds govern Myc's biological output *in vivo*. *Cancer Cell* 14, 447-457.
- Okuhashi, Y., Itoh, M., Nara, N., & Tohda, S. (2011). Effects of combination of notch inhibitor plus hedgehog inhibitor or Wnt inhibitor on growth of leukemia cells. *Anticancer Res.* 31, 893-896.
- Old, J.B., Kratzat, S., Hoellein, A., Graf, S., Nilsson, J.A., Nilsson, L., Nakayama, K.I., Peschel, C., Cleveland, J.L., & Keller, U.B. (2010). Skp2 directs Myc-mediated suppression of p27Kip1 yet has modest effects on Myc-driven lymphomagenesis. *Mol. Cancer Res.* 8, 353-362.
- Omura-Minamisawa, M., Diccianni, M.B., Batova, A., Chang, R.C., Bridgeman, L.J., Yu, J., Pullen, J., Bowman, W.P., & Yu, A.L. (2000). Universal inactivation of both p16 and p15 but not downstream components is an essential event in the pathogenesis of T-cell acute lymphoblastic leukemia. *Clin. Cancer Res.* 6, 1219-1228.
- Osipo, C., Golde, T.E., Osborne, B.A., & Miele, L.A. (2008). Off the beaten pathway: the complex cross talk between Notch and NF-kappaB. *Lab Invest.* 88, 11-17.
- Palomero, T., Dominguez, M., & Ferrando, A.A. (2008). The role of the PTEN/AKT pathway in NOTCH1-induced leukemia. *Cell Cycle* 7, 965-970.

- Palomero, T., Sulis, M.L., Cortina, M., Real, P.J., Barnes, K., Ciofani, M., Caparros, E., Buteau, J., Brown, K., Perkins, S.L., et al. (2007). Mutational loss of PTEN induces resistance to NOTCH1 inhibition in T-cell leukemia. *Nat. Med.* 13, 1203-1210.
- Polo, S.E., Kaidi, A., Baskcomb, L., Galanty, Y., & Jackson, S.P. (2010). Regulation of DNA-damage responses and cell-cycle progression by the chromatin remodelling factor CHD4. *EMBO J.* 29, 3130-3139.
- Popov, N. & Gil, J. (2010). Epigenetic regulation of the INK4b-ARF-INK4a locus: in sickness and in health. *Epigenetics* 5, 685-690.
- Qi, Y., Gregory, M.A., Li, Z., Brousal, J.P., West, K., & Hann, S.R. (2004). p19ARF directly and differentially controls the functions of c-Myc independently of p53. *Nature* 431, 712-717.
- Quelle, D.E., Zindy, F., Ashmun, R.A., & Sherr, C.J. (1995). Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest. *Cell* 83, 993-1000.
- Radhakrishnan, S.K. & Kamalakaran, S. (2006). Pro-apoptotic role of NF-kappaB: implications for cancer therapy. *Biochim. Biophys. Acta* 1766, 53-62.
- Real, P.J. & Ferrando, A.A. (2009). NOTCH inhibition and glucocorticoid therapy in T-cell acute lymphoblastic leukemia. *Leukemia* 23, 1374-1377.
- Real, P.J., Tosello, V., Palomero, T., Castillo, M., Hernando, E., de Stanchina, E., Sulis, M.L., Barnes, K., Sawai, C., Homminga, I., et al. (2009). Gamma-secretase inhibitors reverse glucocorticoid resistance in T cell acute lymphoblastic leukemia. *Nat. Med.* 15, 50-58.
- Remke, M., Pfister, S., Kox, C., Toedt, G., Becker, N., Benner, A., Werft, W., Breit, S., Liu, S., Engel, F., et al. (2009). High-resolution genomic profiling of childhood T-ALL reveals frequent copy-number alterations affecting the TGF-beta and PI3K-AKT pathways and deletions at 6q15-16.1 as a genomic marker for unfavorable early treatment response. *Blood* 114, 1053-1062.
- Riz, I., Zweier-Renn, L.A., Toma, I., Hawley, T.S., & Hawley, R.G. (2011). Apoptotic role of IKK in T-ALL therapeutic response. *Mol. Cancer Res.* 9, 979-984.
- Rodway, H., Llanos, S., Rowe, J., & Peters, G. (2004). Stability of nucleolar versus non-nucleolar forms of human p14(ARF). *Oncogene* 23, 6186-6192.
- Ruggero, D., Montanaro, L., Ma, L., Xu, W., Londei, P., Cordon-Cardo, C., & Pandolfi, P.P. (2004). The translation factor eIF-4E promotes tumor formation and cooperates with c-Myc in lymphomagenesis. *Nat. Med.* 10, 484-486.
- Ryan, K.M., Ernst, M.K., Rice, N.R., & Vousden, K.H. (2000). Role of NF-kappaB in p53-mediated programmed cell death. *Nature* 404, 892-897.
- Sade, H., Krishna, S., & Sarin, A. (2004). The anti-apoptotic effect of Notch-1 requires p56lck-dependent, Akt/PKB-mediated signaling in T cells. *J. Biol. Chem.* 279, 2937-2944.
- Salmena, L., Carracedo, A., & Pandolfi, P.P. (2008). Tenets of PTEN tumor suppression. *Cell* 133, 403-414.
- Sarmiento, L.M., Huang, H., Limon, A., Gordon, W., Fernandes, J., Tavares, M.J., Miele, L., Cardoso, A.A., Classon, M., & Carlesso, N. (2005). Notch1 modulates timing of G1-S progression by inducing SKP2 transcription and p27 Kip1 degradation. *J. Exp. Med.* 202, 157-168.
- Serrano, M., Hannon, G.J., & Beach, D. (1993). A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/CDK4. *Nature* 366, 704-707.

- Shank-Calvo, J.A., Draheim, K., Bhasin, M., & Kelliher, M.A. (2006). p16Ink4a or p19Arf loss contributes to Tal1-induced leukemogenesis in mice. *Oncogene* 25, 3023-3031.
- Sharma, V.M., Draheim, K.M., & Kelliher, M.A. (2007). The Notch1/c-Myc pathway in T cell leukemia. *Cell Cycle* 6, 927-930.
- Sharpless, N.E., Ramsey, M.R., Balasubramanian, P., Castrillon, D.H., & DePinho, R.A. (2004). The differential impact of p16(INK4a) or p19(ARF) deficiency on cell growth and tumorigenesis. *Oncogene* 23, 379-385.
- Sherr, C.J. (2006). Divorcing ARF and p53: an unsettled case. *Nat. Rev. Cancer* 6, 663-673.
- Shin, H.M., Minter, L.M., Cho, O.H., Gottipati, S., Fauq, A.H., Golde, T.E., Sonenshein, G.E., & Osborne, B.A. (2006). Notch1 augments NF-kappaB activity by facilitating its nuclear retention. *EMBO J.* 25, 129-138.
- Shiota, M., Izumi, H., Onitsuka, T., Miyamoto, N., Kashiwagi, E., Kidani, A., Hirano, G., Takahashi, M., Naito, S., & Kohno, K. (2008). Twist and p53 reciprocally regulate target genes via direct interaction. *Oncogene* 27, 5543-5553.
- Silva, A., Jotta, P.Y., Silveira, A.B., Ribeiro, D., Brandalise, S.R., Yunes, J.A., & Barata, J.T. (2010). Regulation of PTEN by CK2 and Notch1 in primary T-cell acute lymphoblastic leukemia: rationale for combined use of CK2- and gamma-secretase inhibitors. *Haematologica* 95, 674-678.
- Silva, A., Yunes, J.A., Cardoso, B.A., Martins, L.R., Jotta, P.Y., Abecasis, M., Nowill, A.E., Leslie, N.R., Cardoso, A.A., & Barata, J.T. (2008). PTEN posttranslational inactivation and hyperactivation of the PI3K/Akt pathway sustain primary T cell leukemia viability. *J. Clin. Invest* 118, 3762-3774.
- So, A.Y., Jung, J.W., Lee, S., Kim, H.S., & Kang, K.S. (2011). DNA methyltransferase controls stem cell aging by regulating BMI1 and EZH2 through microRNAs. *PLoS One* 6, e19503.
- Soulier, J., Clappier, E., Cayuela, J.M., Regnault, A., Garcia-Peydro, M., Dombret, H., Baruchel, A., Toribio, M.L., & Sigaux, F. (2005). HOXA genes are included in genetic and biologic networks defining human acute T-cell leukemia (T-ALL). *Blood* 106, 274-286.
- Spivakov, M. & Fisher, A.G. (2007). Epigenetic signatures of stem-cell identity. *Nat. Rev. Genet.* 8, 263-271.
- Stambolic, V., MacPherson, D., Sas, D., Lin, Y., Snow, B., Jang, Y., Benchimol, S., & Mak, T.W. (2001). Regulation of PTEN transcription by p53. *Mol. Cell* 8, 317-325.
- Sulis, M.L., Saftig, P., & Ferrando, A.A. (2011). Redundancy and specificity of the metalloprotease system mediating oncogenic NOTCH1 activation in T-ALL. *Leukemia* doi:10.1038/leu.2011.130 [Epub ahead of print].
- Sulong, S., Moorman, A.V., Irving, J.A., Strefford, J.C., Konn, Z.J., Case, M.C., Minto, L., Barber, K.E., Parker, H., Wright, S.L., et al. (2009). A comprehensive analysis of the CDKN2A gene in childhood acute lymphoblastic leukemia reveals genomic deletion, copy number neutral loss of heterozygosity, and association with specific cytogenetic subgroups. *Blood* 113, 100-107.
- Tanigaki, K. & Honjo, T. (2007). Regulation of lymphocyte development by Notch signaling. *Nat. Immunol.* 8, 451-456.
- Topisirovic, I., Gutierrez, G.J., Chen, M., Appella, E., Borden, K.L., & Ronai, Z.A. (2009). Control of p53 multimerization by Ubc13 is JNK-regulated. *Proc. Natl. Acad. Sci. USA* 106, 12676-12681.

- Treanor, L.M., Volanakis, E.J., Zhou, S., Lu, T., Sherr, C.J., & Sorrentino, B.P. (2011). Functional interactions between Lmo2, the Arf tumor suppressor, and Notch1 in murine T-cell malignancies. *Blood* 117, 5453-5462.
- Van Vlierberghe, P., Homminga, I., Zuurbier, L., Gladdines-Buijs, J., van Wering, E.R., Horstmann, M., Beverloo, H.B., Pieters, R., & Meijerink, J.P. (2008). Cooperative genetic defects in TLX3 rearranged pediatric T-ALL. *Leukemia* 22, 762-770.
- Vilas-Zornoza, A., Agirre, X., Martin-Palanco, V., Martin-Subero, J.I., San Jose-Eneriz, E., Garate, L., Alvarez, S., Miranda, E., Rodriguez-Otero, P., Rifon, J., et al. (2011). Frequent and simultaneous epigenetic inactivation of TP53 pathway genes in acute lymphoblastic leukemia. *PLoS One* 6, e17012.
- Vilborg, A., Wilhelm, M.T., & Wiman, K.G. (2010). Regulation of tumor suppressor p53 at the RNA level. *J. Mol. Med. (Berl)* 88, 645-652.
- Volanakis, E.J., Williams, R.T., & Sherr, C.J. (2009). Stage-specific Arf tumor suppression in Notch1-induced T-cell acute lymphoblastic leukemia. *Blood* 114, 4451-4459.
- von der Lehr, N., Johansson, S., Wu, S., Bahram, F., Castell, A., Cetinkaya, C., Hydbring, P., Weidung, I., Nakayama, K., Nakayama, K.I., et al. (2003). The F-box protein Skp2 participates in c-Myc proteasomal degradation and acts as a cofactor for c-Myc-regulated transcription. *Mol. Cell* 11, 1189-1200.
- Voncken, J.W., Niessen, H., Neufeld, B., Rennefahrt, U., Dahlmans, V., Kubben, N., Holzer, B., Ludwig, S., & Rapp, U.R. (2005). MAPKAP kinase 3pK phosphorylates and regulates chromatin association of the polycomb group protein Bmi1. *J. Biol. Chem.* 280, 5178-5187.
- Weber, J.D., Taylor, L.J., Roussel, M.F., Sherr, C.J., & Bar-Sagi, D. (1999). Nucleolar Arf sequesters Mdm2 and activates p53. *Nat. Cell Biol.* 1, 20-26.
- Weng, A.P., Ferrando, A.A., Lee, W., Morris, J.P., Silverman, L.B., Sanchez-Irizarry, C., Blacklow, S.C., Look, A.T., & Aster, J.C. (2004). Activating mutations of NOTCH1 in human T cell acute lymphoblastic leukemia. *Science* 306, 269-271.
- Weng, A.P., Millholland, J.M., Yashiro-Ohtani, Y., Arcangeli, M.L., Lau, A., Wai, C., Del, B.C., Rodriguez, C.G., Sai, H., Tobias, J., et al. (2006). c-Myc is an important direct target of Notch1 in T-cell acute lymphoblastic leukemia/lymphoma. *Genes Dev.* 20, 2096-2109.
- Xue, L., Nolla, H., Suzuki, A., Mak, T.W., & Winoto, A. (2008). Normal development is an integral part of tumorigenesis in T cell-specific PTEN-deficient mice. *Proc. Natl. Acad. Sci. USA* 105, 2022-2027.
- Yilmaz, O.H., Valdez, R., Theisen, B.K., Guo, W., Ferguson, D.O., Wu, H., & Morrison, S.J. (2006). Pten dependence distinguishes haematopoietic stem cells from leukaemia-initiating cells. *Nature* 441, 475-482.
- Yoda, A., Toyoshima, K., Watanabe, Y., Onishi, N., Hazaka, Y., Tsukuda, Y., Tsukada, J., Kondo, T., Tanaka, Y., & Minami, Y. (2008). Arsenic trioxide augments Chk2/p53-mediated apoptosis by inhibiting oncogenic Wip1 phosphatase. *J. Biol. Chem.* 283, 18969-18979.
- Zhang, J., Grindley, J.C., Yin, T., Jayasinghe, S., He, X.C., Ross, J.T., Haug, J.S., Rupp, D., Porter-Westpfahl, K.S., Wiedemann, L.M., et al. (2006). PTEN maintains haematopoietic stem cells and acts in lineage choice and leukaemia prevention. *Nature* 441, 518-522.

- Zhang, X.P., Liu, F., & Wang, W. (2011). Two-phase dynamics of p53 in the DNA damage response. *Proc. Natl. Acad. Sci. USA* 108, 8990-8995.
- Zhang, Y. & Xiong, Y. (2001). Control of p53 ubiquitination and nuclear export by MDM2 and ARF. *Cell Growth Differ.* 12, 175-186.
- Zhang, Z. & Zhang, R. (2008). Proteasome activator PA28 gamma regulates p53 by enhancing its MDM2-mediated degradation. *EMBO J.* 27, 852-864.
- Zhou, M., Gu, L., Findley, H.W., Jiang, R., & Woods, W.G. (2003). PTEN reverses MDM2-mediated chemotherapy resistance by interacting with p53 in acute lymphoblastic leukemia cells. *Cancer Res.* 63, 6357-6362.



Novel Aspects in Acute Lymphoblastic Leukemia

Edited by Dr. Stefan Faderl

ISBN 978-953-307-753-6

Hard cover, 258 pages

Publisher InTech

Published online 16, November, 2011

Published in print edition November, 2011

Acute lymphoblastic leukemia (ALL) has turned from a universally fatal to a highly curable disease in little more than four decades. Even though differences in outcome continue to exist between children and adults, intense efforts are under way to overcome this discrepancy and improve the prognosis of adult patients as well. This exemplary progress in ALL therapy has been possible by the combination of an increasingly better understanding of the biology of the disease, availability of a range of effective drugs, and astute designs and relentless executions of many clinical trials. ALL is a complex disease requiring complex therapy. Whereas this book cannot provide a comprehensive review of every one of its many facets, the chapters from many investigators from around the world nevertheless cover a number of relevant topics: aspects of the epidemiology of ALL in Hispanics, ophthalmologic manifestations of ALL, overviews of current therapy and drug-resistance mechanisms, novel biological pathways and targets, new drugs in development, and long-term consequences of CNS prophylaxis and therapy. The publishers and editor therefore hope that the prospective readers will find enough insight and information for their own endeavors.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Irene Riz, Wenjing Yang, Weiqun Peng and Robert G. Hawley (2011). p53 as a Therapeutic Target in T-ALL, Novel Aspects in Acute Lymphoblastic Leukemia, Dr. Stefan Faderl (Ed.), ISBN: 978-953-307-753-6, InTech, Available from: <http://www.intechopen.com/books/novel-aspects-in-acute-lymphoblastic-leukemia/p53-as-a-therapeutic-target-in-t-all>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen