

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Marker Assisted Characterization in *Tigridia pavonia* (L.f) DC

José Luis Piña-Escutia, Luis Miguel Vázquez-García
and Amaury Martín Arzate-Fernández

*Centro de Investigación y Estudios Avanzados en Fitomejoramiento, Facultad de Ciencias
Agrícolas, Universidad Autónoma del Estado de México,
México*

1. Introduction

Conventional plant breeding programs largely depend on phenotypic selection, breeders' experience, and knowledge of plant genetics for traits of agronomic importance. A large amount of biological data is available from genetic studies of a crop of interest related to important target traits in crop breeding, which may in turn directly assist genotypic selection (Xu & Crouch, 2008). However, conventional breeding is time consuming and often testing procedures may be complex, unreliable or expensive due to the nature of the target traits (e.g. abiotic stresses) or the target environment (Choudhary et al., 2008).

Marker assisted characterization offers such a possibility by adopting a wide range of novel approaches to improving the selection strategies in horticultural plant breeding (Ibitoye & Akin-Idowu, 2010). The genetic markers involved in marker assisted characterization include morphological, biochemical and DNA fingerprints which can be employed in selection and identification of closely related genotypes.

Many variants of an enzyme, referred to as isozymes, can be resolved by electrophoresis and are very useful genetic markers. The alleles of most isozyme markers segregate in a codominant manner and rarely show epistatic interactions, which allows accumulation of many polymorphic isozyme loci in a single F₂ population and increases the efficiency of gene mapping. Once the map location of isozyme genes are known, they can be used efficiently as biochemical markers to tag other genes for morphological, physiological and phytopathological traits (Tanksley & Rick, 1980). However, only a relatively small number of protein variants may exist between the two parents and this limits the total number of protein loci that can actually be scored in a given mapping population (Young, 1999).

DNA markers have been used in the identification of varieties and characterization of genotypes because they offer a fast screening and a more precise discriminatory power (Vicente & Fulton, 2003).

With the discovery of restriction enzymes and the polymerase chain reaction (PCR), different markers systems such as Restriction Fragment Length Polymorphisms (RFLP), Random Amplified Polymorphic DNAs (RAPD), Amplified Fragment Length Polymorphisms (AFLP),

Simple Sequence Repeats or microsatellites (SSR), Single Nucleotide Polymorphism (SNP), Inter-Simple Sequence Repeats (ISSR) and others have been developed to visualize the composition of organisms at the DNA level and obtain their genetic fingerprints. Among these markers, RFLPs, AFLPs, SSRs and SNPs have been reported as highly polymorphic and reproducible. However RFLPs require large amount of DNA, are laborious intensive, time consuming and mostly require radioactively labeled probes, whereas AFLPs and SSRs are quite costly also require high resolution electrophoresis or automated sequences. On the other hand, although SNPs are less time consuming against rest of the markers and highly amenable to automation, initial cost involved is quite high (Jehan and Lakhanpaul, 2006). So, although marker informativeness is an important element when comparing different assay systems, other factors such as cost per assay, level of skills required, and reliability of assays should also be considered. PCR molecular markers like ISSR and RAPD would be an option because of the lower level of skill required, low costs per assays and the ready availability of primers can scan the entire genome and make more efficient the genotype characterization. Thus, due to their characteristics and efficiency for detecting polymorphism, the ISSR and RAPD markers have been successfully used to calculate the intra or inter-specific genetic diversity in different domesticated and wild species (Arzate-Fernández et al., 2005b; Escandón et al., 2005a; Tapia et al., 2005; Kumar et al., 2006; Luna-Paez et al., 2007; Marotti et al., 2007; Muthusamy et al., 2008; Ye et al., 2008).

The genus *Tigridia* comprises plants with highly colored flowers that exhibit great morphological variation, making many species potentially valuable as cultivated plants. *Tigridia pavonia* (L.f.) DC, also known as tiger flower or oceloxóchitl, is a species native to México which was used by the Aztecs for ornamental, nutritional and medicinal purposes (Hernández, 1959). Due to the great variability of the coloring of its flower and its ornamental potential, it is one of the principal phylogenetic resources of this country, considered the center of greatest genetic diversity of this species. In spite of fact that in Mexico, the potential for sustainable *T. pavonia* utilization is enormous, its remains largely unexploited due to the little information on the resource abundance and distribution, whereas in different countries of Europe, Asia and Australia this species is widely distributed and commercialized as a garden plant (Vázquez-García et al., 2001). Thus, genetic diversity studies are needed to provide information for its efficient conservation as well as for possible future use in breeding programs.

The widely distributed species generally present morphological and physiological variation, as well as in the genetic structure of their populations (Wen & Hsiao, 2001). Likewise, the dissemination in different regions could cause that the same variety or cultivar may be known under different names in different countries (and even within a given country) and different varieties sometimes may appear under the same name. Therefore, the study of phenotypic and genetic diversity to identify groups with similar genotypes is important for conserving, evaluating and exploiting genetic resources. Also, these studies are useful for determining the uniqueness and distinctness of the phenotypic and genetic constitution of genotypes with the purpose of protecting the breeder's intellectual property rights (Franco et al., 2001).

The focus of this chapter is in a more applied direction, encompassing the potential of the morphological, isozymes and DNA markers for characterization of nine *T. pavonia* varieties to their management into selection and breeding programme.

2. Materials

Nine botanical varieties of *Tigridia pavonia* (L.f.) DC, collected in three localities of the State of Mexico and registered in the National Seed Inspection and Certification Service (SNICS) were used in this study (Figure 1). The Sandra variety (TGD-008-030408) was collected in the municipality of Tenancingo, Carolina (TGD-002-030408), Trinidad (TGD-009-030408), Penelope (TGD-006-030408), Angeles (TGD-001-030408), Dulce (TGD-003-030408) and Mariana (TGD-005-030408), in the municipality of Temascaltepec and Samaria (TGD-007-030408) and Gloria (TGD-004-030408) in the municipality of Temoaya. All of the varieties were cultivated in a rustic greenhouse, at the Faculty of Agricultural Sciences (FCAgr) of the Autonomous University of State of Mexico (UAEM).



Fig.1. Nine botanical varieties of *Tigridia pavonia* (L. f.) DC used in the morphological and molecular characterization assay

3. Methodology

3.1 Morphological markers

To make the matrix of morphological data, 20 quantitative and one qualitative characters (Table 1) were evaluated among the nine *T. pavonia* varieties. Nine characters were previously published by Vázquez-García et al. (2001). Each character was evaluated in 14 individuals of each variety.

Character	Mean	Max	Min	Standar Deviation	CV (%)	P
1. Number of flowers per shoot	9.32	17.00	3.00	2.22	23.23	0.0753
2. Length of external tepal	8.65	11.70	7.00	0.88	7.87	<0.0001
3. Length of internal tepal	4.42	5.20	3.70	0.31	6.60	<0.0004
4. Type of internal tepal (oval formed, sharp pointed, lanceolated, rounded)	1.92	4.00	1.00	1.23	22.72	<0.0001
5. Length of staminal column	4.49	5.40	3.80	0.29	4.48	<0.0001
6. Distance from the base of the anther to the stigma	1.05	2.50	0.30	0.54	20.46	<0.0001
7. Length of anther	1.59	2.00	1.00	0.21	7.40	<0.0001
8. Length of the reproductive part	6.31	7.30	5.10	0.49	4.42	<0.0001
9. Number of fruits per shoot	4.84	13.00	1.00	2.32	34.44	<0.0001
10. Number of shoot per bulb	4.00	6.00	3.00	0.98	18.64	<0.0001
11. Length of shoot	60.03	96.50	38.60	18.68	7.03	<0.0001
12. Nodes per shoot	3.84	5.00	2.00	0.44	11.14	0.0240
13. Length of the internode	8.18	22.00	0	6.15	39.24	<0.0001
14. Number of branches per shoot	1.98	3.00	0.90	0.72	25.03	<0.0001
15. Length of the leaf	34.13	46.00	16.00	6.78	8.23	<0.0001
16. Width of the leaf	3.53	5.60	2.50	0.88	7.46	<0.0001
17. Length of the pseudopetiole	7.55	11.00	6.00	1.32	6.16	<0.0001
18. Floral scapus length	11.21	28.00	3.00	7.29	10.89	<0.0001
19. Floral scapus thickness	0.55	1.00	0.30	0.22	6.33	<0.0001
20. Length of the bract	11.29	24.50	8.00	3.88	16.60	<0.0001
21. Bract width	2.85	3.50	2.20	0.38	4.03	<0.0001

Table 1. Morphological characters and basic statistical data of nine varieties of *T. pavonia* (L. f.) DC.

3.2 Isozyme markers

For isozyme markers, two individuals of each *T. pavonia* variety were used. Leaf sections were taken from each individual eight to 12 weeks after budding initiated. All of the samples were processed in the laboratory of Plant Molecular Biology at the FCAgr of the UAEM.

Approximately 50 mg of fresh leaf tissue from each sample was homogenized using a plastic chopstick in 50 μ L extraction buffer Tris-HCl 0.1M, 2% EDTA-2Na, 50.4% glycerol, 3.2% Tween-80, and 0.8% DTT, pH 7.5 (Arzate-Fernández et al., 2005a).

The samples were preserved at $-20\text{ }^{\circ}\text{C}$ until analysis. Two buffer systems of histidine-citric acid, pH 5.7 and pH 6.5 (H-AC) and Tris-citrate/Tris-histidine, pH 8.5 (T-C/T-H) were used following the procedures of Glaszmann et al. (1988) and Stuber et al. (1988).

The 18 isozyme systems evaluated were aspartate amino transferase (AAT; EC 2.6.1.1), acid phosphatase (ACP; EC 13.1.3.2), dehydrogenase alcohol (ADH; EC 1.1.1.1), aminopeptidase (AMP; EC 3.4.11.1), catalase (CAT; EC 1.11.1.6), endopeptidase (ENP; EC 3.4.23.6), esterase (EST; EC 3.1.1.1), formate dehydrogenase (FDH; EC 1.2.1.2), glutamate dehydrogenase (GDH; EC 1.4.1.2), glucose-6-phosphate dehydrogenase (G-6-PDH; EC 1.1.1.49), malic enzyme (MAL; EC 1.1.1.40), malate dehydrogenase (MDH; EC 1.1.1.37), phosphoglucose dehydrogenase (PGD; EC 1.1.1.44), phosphoglucose isomerase (PGI; EC 5.3.1.9), phosphoglucomutase (PGM; EC 2.7.5.1), phosphohexose isomerase (PHI; EC 5.3.1.8), peroxidase (POX; EC 1.11.1.7), and shikimate dehydrogenase (SKD; EC 1.1.1.25). The samples were loaded into a 12% hydrolyzed potato starch gel and electrophoresis was conducted at $4\text{ }^{\circ}\text{C}$ for a mean duration of 3 h, 40-50 mA and 100-150 v. Each sample was run at least three times to verify reproducibility. Enzyme staining was performed according to the procedures described by Torres et al. (1978), Vallejos (1983), Glaszmann et al. (1988), Stuber et al. (1988) Wendel & Weeden (1990) and Ishikawa (1994).

3.3 DNA markers

Young leaves were collected of two individuals of each variety (in the same development stage), and stored at $-20\text{ }^{\circ}\text{C}$ prior to DNA extraction. The genomic DNA was extracted of approximately 100 mg of leaf tissue of each variety of *T. pavonia*. The extraction procedure was the CTAB as reported by Zhou et al. (1999). The DNA samples were stored to $-20\text{ }^{\circ}\text{C}$ prior to analysis. The polymerase chain reaction (PCR) was made in a final volume of $10\text{ }\mu\text{L}$ with: $1\text{ }\mu\text{L}$ of DNA (10 ng), $1\text{ }\mu\text{L}$ of 10X PCR buffer with ammonium (15 mM), $0.5\text{ }\mu\text{L}$ of MgCl_2 (15 mM), $1\text{ }\mu\text{L}$ of dNTPs (10 mM), $1\text{ }\mu\text{L}$ of the primer (20 mM) and 0.1 units of the enzyme Taq DNA polymerase.

3.3.1 ISSR markers

For ISSR markers, five primers of the anchored microsatellites type were used (3'-ASSR) (Yamagishi et al., 2002). In each primer, the anchor consisted of a triplicate of distinct sequence (Table 4). The amplification conditions for the primers 3'-ASSR02 and 3'-ASSR15 were those described by Arzate-Fernández et al. (2005b), and consisted of an initial cycle of 9 min at $94\text{ }^{\circ}\text{C}$, 1 min at $46\text{ }^{\circ}\text{C}$ and 1 min at $72\text{ }^{\circ}\text{C}$ and a final cycle of 9 min at $94\text{ }^{\circ}\text{C}$, 1 min at $46\text{ }^{\circ}\text{C}$ and 10 min at $72\text{ }^{\circ}\text{C}$. For primers 3'-ASSR20, 3'-ASSR29 and 3'-ASSR35, the amplification cycles were those used by Yamagishi et al. (2002) and consisted of an initial cycle of 9 min at $94\text{ }^{\circ}\text{C}$, 1 min at $46\text{ }^{\circ}\text{C}$ and 1 min at $72\text{ }^{\circ}\text{C}$ and a final cycle of 10 min at $72\text{ }^{\circ}\text{C}$.

3.3.2 RAPD markers

For RAPD markers, five 10 base primers (Yamagishi, 1995), five 15 base primers (Yamagishi et al., 2002), and five 20 base primers (Debener & Mattiesch, 1998) were used (Table 5). The amplification conditions for RAPD primers of 10 base were the reported by Yamagishi (1995). The program for 15 base primers used in this study was followed according to Yamagishi et al. (2002) with minor modifications (40 cycles of $94\text{ }^{\circ}\text{C}$ for 1 min, $53\text{ }^{\circ}\text{C}$ for 3

min and 72 °C for 2 min). The PCR cycle conditions for 20 base primers were performed according to Debener & Mattiesch (1998).

The amplification of DNA fragments for ISSR and RAPD markers was made in a thermocycler (Mastercycler gradient, Eppendorf, Germany). The separation of the fragments was made in horizontal electrophoresis. A molecular marker of 100 to 3000 pb molecular weight was used. The running conditions for each sample were 100 V and 120 mA for 80 min, and the observation of the fragments was made in a transilluminator UVP.

3.4 Statistical analysis

For morphological markers, the 21 morphological characters were evaluated through an analysis of variance and a completely randomized design with 14 replicates, with the statistical program SAS version 8.0. With the data generated of the morphological characters, a binary matrix was made following the criteria described by Vicente & Fulton (2003); that is, the value of 1 was assigned if the character was present and 0 if it was absent.

For isozyme markers, a record of banding patterns was constructed and each band was assigned to a binary matrix as absent, 0, or present, 1.

For ISSR and RAPD markers, each band generated by each primer was considered as an independent locus calculated manually; that is, the value of 1 was assigned for the presence of a band and 0 for its absence. The total number fragments (FT), polymorphic fragments (FP), percentage of polymorphism (%P) and the Nei's genetic distance (G_D) were calculated using the program POPGENE version 1.32 (Yeh & Boyle, 1999). Also, capacity of each primer to differentiate the nine varieties under study was evaluated through the resolution power (R_p), according to Prevost & Wilkinson (1999).

In RAPD markers, a simple correlation analysis was performed to investigate the correlation between the length primer and the polymorphism generated by each primer group.

In all cases (morphological, isozyme and DNA markers), to determine genetic relationships and variety grouping, a dendrogram was constructed with the data using POPGENE with the UPGMA method (modified from the NEIGHBOR procedure of PHYLIP, version 3.5) (Felsenstein, 1990) based on the Nei (1972) matrix genetic distances. UPGMA is one of the simplest and most commonly used hierarchical clustering algorithms. It receives as input a set of elements and a dissimilarity matrix which contains pairwise distances between all elements, and returns a hierarchy of clusters on this set.

4. Results and discussion

4.1 Morphological markers

The values of the mean, maximum, minimum, standard deviation (SD) and coefficient of variation (CV) were calculated for each one of the 21 morphological characters evaluated in the nine varieties (Table 1). According to the values of the CV, the characters number of flowers per shoot, type of internal tepal, distance from the base of the anther to the stigma, number of fruits per shoot, length of the internode and number of branches per shoot showed a slightly high variation level. In contrast, the characters that presented low variation levels among the varieties were the length of the staminal column, length of the reproductive part and bract width.

The statistical analysis showed that the number of flowers per shoot and nodes per shoot were not significant. However, for the other 19 characters there were highly significant differences ($p \leq 0.0001$) (Table 1). These results show the high phenotypic diversity among the varieties analyzed, and can be used for the selection of parental lines within a plant breeding program.

The varieties formed two groups based on their morphological characteristics (Figure 2). Group I with average of $G_D = 0.42$ consisted of the varieties Carolina, Trinidad, Mariana, Angeles, Sandra, Penelope and Dulce, while group II included the varieties Gloria and Samaria, with an average of $G_D = 0.32$.

The morphological characters (18 of 21 equal) showed the close genetic relationship between the varieties Trinidad and Mariana ($G_D = 0.28$). In contrast, the least related varieties were Carolina and Samaria, as well as Penélope and Samaria ($G_D = 0.80$), with 4 and 5 of 21 similar morphological characters, respectively.

According to Vázquez-García et al. (2001), the varieties Gloria and Samaria can be easily distinguished by shoot length. In the present study, the results were similar and both varieties showed the highest values for length of shoot, length of internode, floral scapus length and length of the bract, as well as bract width.




Fig. 2. Dendrogram of nine varieties of *T. pavonia* based on the genetic distance of Nei (1972), with the method UPGMA. The bar indicates the genetic distance among the varieties and the groups, calculated from morphological data.

4.2 Isozyme markers

The nine *T. pavonia* varieties evaluated were wild and free-pollinating. Therefore, they were considered to be heterozygotic; this agrees with the diverse isozyme banding patterns (IBPs) observed. Of the 18 isozymes tested, 12 could be stained, but only nine had sufficient resolution and clarity to be recorded and studied. Some bands were omitted because of their poor resolution. The isozymes ACP, MDH, PGI, PHI and PGM had better resolution when examined with the H-AC system, pH 5.7, and the isozymes AAT and PGD in the same system, but at pH 6.5. In contrast, the isozymes CAT and POX were better in the T-C/T-H system. Of the selected isozymes, nine zymograms in which 32 bands arranged in 20 IBP were identified: two for AAT, two for ACP, two for CAT, four for MDH, one for PGD, one for POX, three for PGI, three for PHI, and two for PGM (Figure 3).

In this study most of the varieties could be identified by the IBP obtained in each isozyme, as we describe in the following sections.

4.2.1 Aspartate amino transferase

AAT showed five active bands (Figure 3A) and varieties with four bands arranged in two banding patterns were detected. Pattern 1 was observed in the Angeles, Carolina, Dulce, Mariana, Penelope, and Sandra varieties, while the Gloria, Samaria, and Trinidad varieties exhibited pattern 2 (Table 2).

4.2.2 Acid phosphatase

ACP revealed four active bands (Figure 3B) and varieties with one or four bands arranged in two banding patterns were detected. The Angeles, Carolina, Dulce, Gloria, Mariana, Penelope, Sandra, and Trinidad varieties exhibited pattern 1 with four bands. Pattern 2 had a single band and was seen only in the Samaria variety (Table 2).

4.2.3 Catalase

CAT produced two active bands (Figure 3C) and varieties with one or two bands arranged in two banding patterns were detected. Pattern 1 with two bands was observed in Carolina, Dulce, Mariana, Penelope, and Samaria. Pattern 2 was observed in Angeles, Gloria, Sandra, and Trinidad (Table 2).

4.2.4 Malate dehydrogenase

MDH showed six active bands (Figure 3D) and varieties with three or four bands in four different arrangements were detected to produce four banding patterns. Patterns 1, 2, and 3 were composed of four bands. Pattern 1 was observed in Angeles, Carolina, Dulce, Mariana, Sandra, and Trinidad, while Penelope and Gloria exhibited patterns 2 and 3. Pattern 4 had three active bands and was observed in Samaria (Table 2).

4.2.5 Phosphoglucose dehydrogenase

PGD produced two active bands (Figure 3E) arranged in a single pattern, so that it was not possible to differentiate any of the varieties (Table 2).

4.2.6 Peroxidase

POX produced two active bands (Figure 3F) arranged in a single pattern, and thus it was not possible to differentiate varieties (Table 2).

4.2.7 Phosphoglucose isomerase

PGI produced four active bands (Figure 3G) and varieties with two, three or four bands arranged to produce three banding patterns were detected. Pattern 1 had four bands, and was observed in Angeles, Carolina, Dulce, Penelope, Samaria, Sandra, and Trinidad. Pattern 2 had three bands and was observed in Mariana. Pattern 3 had two bands and was observed in Gloria (Table 2).

4.2.8 Phosphohexose isomerase

PHI produced three active bands (Figure 3H) and varieties with two or three active bands in three arrangement producing three banding patterns were detected. Pattern 1 had three bands and was observed in ANG, CAR, DUL, PEN, SAN, and TRI. Patterns 2 and 3 had two bands each. Varieties GLO and SAM exhibited pattern 2 and variety MAR pattern 3 (Table 2).

4.2.9 Phosphoglucomutase

PGM produced four active bands (Figure 3I) and varieties with three or four bands arranged in two banding patterns were observed. Pattern 1 had four bands and was observed only in Gloria. Pattern 2 had three bands and was observed in Angeles, Carolina, Dulce, Mariana, Penelope, Samaria, Sandra, and Trinidad (Table 2).

As can be seen in Table 2, the nine isozymes produced sufficient polymorphic bands to distinguish most of the *T. pavonia* varieties evaluated. Of all the isozymes tested, MDH produced the highest number (4) of IBP and, therefore, the highest percentage of polymorphism (18.75%). Based on the observed IBP, this isozyme allowed to differentiate a larger number of varieties. In contrast, the isozymes PGD and POX produced only one IBP, and therefore neither isozyme was useful for differentiating the varieties used.

Twenty-nine bands derived from seven isozymes were only considered: five for AAT, four for ACP, two for CAT, six for MDH, four for PGI, three for PHI, and four for PGM (Figure 3). A dendrogram (Figure 4) was generated from the genetic relationship among these bands, clustering the varieties in two main groups. Group 1 comprises seven varieties: Angeles, Sandra, Carolina, Dulce, Mariana, Penelope, and Trinidad, with an average G_D of 0.074. Group II only includes the Gloria and Samaria varieties, with a G_D of 0.374.

Genetic analysis yielded an average genetic distance G_D of 0.194 among the nine *T. pavonia* varieties (Table 3). As observed in Table 3, the G_D between Angeles and Sandra varieties and between Carolina and Dulce was 0.0, revealing genetic similarity and, in spite of the number of isozymes tested, it was not possible to distinguish one variety from the other. In contrast, Vázquez-García et al. (2001) characterized these four varieties morphologically, differentiating them by flower color: white (Angeles), red (Sandra), pink (Carolina), and yellow (Dulce). The varieties with the highest G_D value (0.421) were Gloria-Mariana, Gloria-Penelope, and Mariana-Samaria.




Fig. 3. Schematic representation of isozyme banding patterns (IBP) detected in nine *T. pavonia* (L.f.) DC. varieties, produced by A) AAT; B) ACP; C) CAT; D) MDH; E) PGD; F) POX; G) PGI; H) PHI; I) PGM. (m=monomer, d=dimer, t=trimer. "o" represents origin of electrophoresis).

Variety	Isozyme and pattern								
	AAT	ACP	CAT	MDH	PGD	POX	PGI	PHI	PGM
Ángeles ¹	1	1	2	1	1	1	1	1	2
Carolina ²	1	1	1	1	1	1	1	1	2
Dulce ²	1	1	1	1	1	1	1	1	2
Gloria	2	1	2	3	1	1	3	2	1
Mariana	1	1	1	1	1	1	2	3	2
Penélope	1	1	1	2	1	1	1	1	2
Samaria	2	2	1	4	1	§	1	2	2
Sandra	1	1	2	1	1	1	1	1	2
Trinidad	2	1	2	1	1	1	1	1	2

Table 2. Isozyme characterization of nine *T. pavonia* (L.f) DC. varieties using nine isozyme systems. The number indicates the isozyme banding pattern (IBP) of each variety. AAT, aspartate amino transferase; ACP, acid phosphatase; CAT, catalase; MDH, malate dehydrogenase; PGD, phosphoglucose dehydrogenase; PGI, phosphoglucose isomerase; PGM, phosphoglucomutase; PHI, phosohexose isomerase; POX, peroxidase. ^{1,2} varieties with the same number can not be distinguished with the evaluated isozymes. § It was not possible determinate the IBP.




Fig. 4. Dendrogram of nine varieties of *Tigridia pavonia* based on Nei (1972) genetic distance, UPGMA method. The bar indicates the genetic distance among the varieties and the groups obtained with isozyme markers.

Variety	ANG	CAR	DUL	GLO	MAR	PEN	SAM	SAN
ANG								
CAR	0.031							
DUL	0.031	0.000						
GLO	0.287	0.330	0.330					
MAR	0.098	0.064	0.064	0.421				
PEN	0.098	0.064	0.064	0.421	0.133			
SAM	0.374	0.330	0.330	0.374	0.421	0.246		
SAN	0.000	0.031	0.031	0.287	0.098	0.098	0.374	
TRI	0.064	0.098	0.098	0.207	0.169	0.169	0.287	0.064

Table 3. Estimation of genetic distance (G_D), according to the formula of Nei (1972), for the nine varieties of *T. pavonia*.

4.3 DNA markers

4.3.1 ISSR markers

The total number of fragments reproducible with the five ASSR primers was 40, with an interval of 350 to 1900 pb in the size of the amplified fragments. Of the 40 fragments, 35 were polymorphic, with an average of seven per primer. The percentage of polymorphic fragments varied from 66.6 to 100%, with an average of 87.5% (Table 4). These results are similar with those reported by Yamagishi et al. (2002) and Arzate-Fernández et al. (2005b), who used these ASSR primers and proved their efficiency for detecting high polymorphism percentage in different species of the genus *Lilium*.

One aspect that favors the effectiveness of the ASSR primers as well as the type of the motive replicate in its sequence, is the sequence of its anchor. The CT motive sequences produce higher polymorphism with respect to the AT replicates (Pradeep et al., 2002; Hu et al., 2003), which despite being the most abundant in the plant genomes, have the disadvantage that the amplification of DNA fragments is low; this may be due to the semi-complementarity of the primer in the alignment stage of the PCR (Fang & Roose, 1997). In this study, the level of distinction among the variables depended on the sequence of the anchor of the primer. Thus, it was possible to obtain 100% polymorphism among the varieties of *T. pavonia* with primers ASSR02 and ASSR35 (Table 4).

Primer	Sequence (5'-3')	AF	PF	%P	R_p	FS (pb)
3'-ASSR02	CTCTCTCTCTCT ATC	9	9	100	3.12	500-1300
3'-ASSR15	CTCTCTCTCTCT ATG	8	6	85.7	2.2	350-1200
3'-ASSR20	CTCTCTCTCTCT GCA	7	6	85.7	2.2	600-1800
3'-ASSR29	CTCTCTCTCTCT GTA	6	4	66.6	1.8	550-1400
3'-ASSR35	CTCTCTCTCTCT TGA	10	10	100	5.54	350-1900
Total or mean		40	35	87.5	1.97	450

Table 4. Characteristics of the ISSR fragments amplified in nine varieties of *T. pavonia* (L.f.) DC.: total amplified fragments (AF), polymorphic fragments (PF), percentage of polymorphism (%P), resolving power (R_p) and size of the amplified fragments (FS) for each ASSR primer.

Prevost & Wilkinson (1999) described the resolution power (R_p) as a useful tool for evaluating the capacity of a primer in the distinction of various genotypes. In our results, the highest values of R_p were for the primers ASSR02 and ASSR35 (Table 4), with which the nine varieties were distinguished from each other. Escandón et al. (2005a) obtained similar results with just two ASSR primers and generated specific profiles for 18 of 21 collections of *Jacaranda mimosifolia*. The efficiency of the ASSR primers for discriminating genotypes in the inter-varietal level has also been reported in *Solanum tuberosum* (Prevost & Wilkinson, 1999), *Jacaranda mimosifolia* (Pérez de la Torre et al., 2003), *Nierembergia linaeriefolia* (Escandón et al., 2005b) and *Ficus carica* (Guasmi et al., 2006), among others.

The dendrogram generated with the ISSR data (Figure 5) formed two groups of varieties: group I included Carolina, Ángeles, Trinidad, Sandra, Dulce, Mariana and Penélope with an average $G_D = 0.19$ whereas the group II included Gloria and Samaria with a $G_D = 0.39$. The highest genetic association ($G_D = 0.07$) was found between the varieties Carolina and Ángeles, while the least related were Gloria and Dulce ($G_D = 0.91$).




Fig. 5. Dendrogram of nine varieties of *T. pavonia* based on the genetic distance of Nei (1972), with the method UPGMA. The bar indicates the genetic distance among the varieties and the groups, calculated from ISSR data.

4.3.2 RAPD markers

In the present study, PCR amplification with 10, 15 and 20 length base RAPD primers led to reproducible fragment patterns for all varieties of *T. pavonia* evaluated. The majority of those RAPD fragments ranged from 250 to 2800 bp (Figure 6). For 10 base primers, the average of

total and the polymorphic fragments generated per primer were 8.8 and 8.2, respectively, whereas for 15 base primers, those values were 11 and 10.4, respectively, and for 20 base primers, 9.4 and 9, respectively (Table 5).

Generally, in RAPD analysis 10 base primers are preferred, nevertheless, because of the high annealing temperature applied in PCR reaction than their original, some 10 base primers could not hybridize with template DNA, generating only a few minor bands or none (Yamagishi et al., 2002), while the advantage of long primers is the smaller number of fragments containing repetitive DNA, thereby increasing the specificity and reproducibility of DNA fragments (Debener & Mattiesch, 1998). In the present study, the positive correlation ($r = 0.99$; $p \leq 0.05$) between primer length and the percentage of polymorphism generated per each primer was observed, indicating that the efficiency of primers to generate polymorphic fragments it increased by primer length. Thus, 15 or 20 base primers generated more DNA fragments (55), and a greater number of polymorphic fragments (52), in comparison with those obtained by the 10 base primers (44, and 41, respectively) (Table 5). The high efficiency of long primers to generate a large number of RAPD markers has been also reported in other species as *Lilium* spp. (Yamagishi et al., 2002), *Vitis vinifera* (Solouki et al., 2007) and *Matricaria chamomilla* (Solouki et al., 2008).

Primer	%GC	AF	PF	%P	R _p
Y24*	70	8	6	75	1.6
Y29*	70	10	10	100	5.1
Y37*	70	3	2	66.6	0.9
Y38*	70	15	15	100	7.5
Y41*	80	8	8	100	3.3
Average	72	8.8	8.2	93.1	3.6
P619**	66	13	13	100	5.8
P625**	66	8	6	75	2.6
P628**	60	4	3	75	1.3
P635**	73	17	17	100	8.9
P647**	60	13	13	100	6.4
Average	65	11	10.4	94.5	5
P495***	60	11	10	90.9	3.9
P496***	60	6	5	83.3	2.6
P497***	60	8	8	100	3.8
P498***	60	12	12	100	4.9
P500***	60	10	10	100	3.8
Average	60	9.4	9	95.7	3.8

Table 5. Percentage of GC (%GC), total amplified fragments (AF), polymorphic fragments (PF), percentage of polymorphism (%P) and resolving power (R_p) for each RAPD primer used. *10 base (Yamagishi, 1995). **15 base (Yamagishi et al., 2002). ***20 base (Debener & Mattiesch 1998).




Fig. 6. RAPD profiles of nine varieties of *T. pavonia* generated by the 10 base primer Y38 (a), 15 base primer P647 (b) and 20 base primer P497 (c).

Although it is not clear exactly why the long primers produced more polymorphic bands, it has been reported that the GC content may be a factor that determines the efficiency of a primer (Solouki et al., 2007), since GC content is associated with annealing temperature and related to generation of more DNA fragments. Thus, it has been observed that with long primers with a lower GC content, the efficiency for amplifying polymorphic bands is higher, in comparison with short primers with a higher GC content (Ye et al., 1996; Solouki et al., 2007). According to this, it is possible that the GC content has favored the major efficiency of the 15 (65 % GC) and 20 (60 % GC) base primers used in the present study, because more DNA fragments were amplified and also a greater percentage of polymorphism, in comparison with those amplified with 10 base primers with 72 % GC content (Table 5). Our results are closely similar with those reported by Solouki et al., (2008) where it was obtained more DNA fragments and 100 % of polymorphism, using a long primer with low GC content.

The high efficiency of long primers in the genetic differentiation of *T. pavonia* varieties was also confirmed with the measurement of resolving power. The highest values of R_p belonged to the 15 and 20 base primers (5 and 3.8, respectively) (Table 5), indicating a better distinction of the varieties. So, these results also confirm the utility of the R_p as a measure of capacity of a primer to discriminate among closely related individuals as was pointed out by Prevost & Wilkinson (1999) and Escandón et al. (2007).

The dendrograms based on UPGMA analysis of the 10, 15, 20 and the pooled (10, 15 and 20) RAPD data showed the genetic differentiation of the nine varieties of *T. pavonia* (Figure 7). In the dendrogram generated with decamers, the G_D among the varieties ranged from 0.20 to 0.69, with an average of 0.24. For the 15 base dendrogram, the G_D ranged from 0.13 to 0.78, with an average of 0.32, while the dendrogram of the 20 base showed a range of G_D of 0.11 to 0.67, with an average of 0.28, and finally the dendrogram with pooled data showed a range of G_D of 0.20 to 0.68, with an average of 0.24.

All four dendrograms clearly grouped the varieties in two major clusters. In the dendrograms of 10 and 15 bases, and in the pooled, the cluster I consisted of the varieties Carolina, Dulce, Trinidad, Penelope, Angeles, Mariana, and Sandra, whereas the cluster II grouped the varieties Gloria and Samaria. The close relationship observed in these dendrograms, among the varieties collected in Tenancingo and Temascaltepec municipalities, might be the result of having some common morphological characters such as: number of shoots by bulb, number of nodes by shoot, length of leaf, and number flowers by shoot. The separation of Gloria and Samaria collected in Temoaya, could be due to the highest values for length of shoot, length of internode, and length of floral escape, besides that the flowering of these varieties is delayed in comparison with the rest of them, as it was reported in our previous study (Piña-Escutia et al., 2010).

On the other hand, the dendrogram developed with 20 base primers also grouped the cluster I similar to the dendrograms generated with other primers sets, except for the variety Sandra, which was grouped with the varieties Gloria and Samaria in the cluster II. Thus, this dendrogram showed a best differentiation of the varieties evaluated, confirming the utility of long primers in generation high polymorphism and genetic discrimination of plant species, which was also reported by Solouki et al. (2007), and Solouki et al. (2008).




Fig. 7. Dendrogram generated using UPGMA analysis, showing relationships between nine varieties of *T. pavonia*, based on RAPD data of 10 (a), 15 (b), 20 (c) and the pooled (10, 15 and 20) base of length primer (d). The bar indicates the genetic distance among the varieties and the groups.

5. General conclusions and perspectives

The assessment of genetic variability of Mexican native species is fundamental for the conservation, and plant variety protection. In this review we have conducting the morphological analysis in combination with the molecular assays to distinguish *T. pavonia* varieties. The analyses of the morphological and DNA markers (ISSR and RAPD) seem to be more efficient than the isozymes, because whereas the former showed polymorphism among the nine varieties evaluated and permit the distinction among them, the biochemical characterization revealed that the varieties Angeles and Sandra as well as Carolina and Dulce were the same, indicating the low discrimination power of isozymes which can be due to inadequate genome coverage. So, our results indicate that the marker assisted characterization can be applied to complement existing conventional breeding programmes and, despite its application in genetic improvement of horticultural crops is limited, their use as balance between molecular and non-molecular methods can offers opportunities of improved phenotypic selection in the future.

6. Acknowledgment

We are grateful to RED TIGRIDIA (SINAREFI, México) for covering the publishing payment.

7. References

- Arzate-Fernández, A.M.; Mejía G, C.O.; Nakazaki, T.; Okumoto, Y. & Tanisaka, T. (2005a). Isozyme electrophoretic characterization of 29 related cultivars of lily (*Lilium* spp.). *Plant Breeding* Vol.124, No. 1, (February 2005), pp. 71-78. ISSN 1437-0523
- Arzate-Fernández, A.M., Miwa, M.; Shimada, T.; Tonekura, T. & Ogawa, K. (2005b). Genetic diversity of miyamasukashi-yuri (*Lilium maculatum* Thumb. Var. Bukosanense), an endemic populations endangered species at Mount Buko, Saitama, Japan. *Plant Species Biology* Vol.20, No.1, (March 2005), pp. 57-65. ISSN 1442-1984
- Choudhary, K.; Choudhary O.P. & Shekhawat, N.S. (2008). Marker assisted selection: a novel approach for crop improvement. *American-Eurasian Journal of Agronomy* Vol.1, No.2, (August 2008) pp. 26-30. ISSN 1995-896X
- Debener, T. & Mattiesch, I. (1998). Effective pairwise combination of long primers for RAPD analysis in roses. *Plant Breeding* Vol.117, No.2, (May 1998), pp. 147-151. ISSN 1437-0523
- Escandón, A.; Pérez de la Torre, M.; Acevedo, A.; Marucci-Poltri, S. & Mijayima, I. (2005a). Anchored ISSR as molecular marker to characterize accesions of *Jacaranda mimosifolia* L. Don. *Acta Horticulturae* Vol.683, No.1, pp. 121-127. ISSN 0567-7572
- Escandón, A.; Pérez de la Torre, M.; Soto, M.S. & Zelener, N. (2005b). Identificación de clones selectos de *Nierembergia linariaefolia* mediante microsatélites anclados. *Revista de Investigaciones Agropecuarias* Vol.34, No.1, (Abril 2005) pp. 5-17. ISSN 1669-2614, Available from http://www.inta.gov.ar/ediciones/ria/34_1/01.pdf

- Fang, D.Q. & Roose, M.L. (1997). Identification of closely related citrus cultivars with inter-simple sequence repeat markers. *Theoretical and Applied Genetics* Vol.95, No.3, (August 1997), pp. 408-417. ISSN 1432-2242
- Felsenstein, J. (1990). *Phylyp Manual*, version 3.3. University Herbarium, University of California, Berkley, CA, USA.
- Franco, J.; Crossa, J.; Ribaut, J.M.; Betran, J; Warburton, M.L. & Khairallah, M. (2001). A method for combining molecular markers and phenotypic attributes for classifying plant genotypes. *Theoretical and Applied Genetics* Vol.103, No.6-7, (November 2001), pp. 944-952. ISSN 1432-2242
- Glaszmann, J.C.; De los Reyes B.G. & Khush, G.S. (1988). Electrophoretic variation or isozymes in plumules of rice (*Oriza sativa* L.) a key to the identification of 76 alleles at 24 loci. *IRRI Research Paper Series* Vol.134, No.11, (November 1988), pp. 1-14. ISSN 0115-3862
- Guasmi, F.; Ferchichi, A.; Farés, K. & Touil, L. (2006). Identification and differentiation of *Ficus carica* L. cultivars using Inter simple sequence repeat markers. *African Journal of Biotechnology* Vol.5, No.15, (August 2006), pp. 1370-1374. ISSN 1684-5315 Available from <http://www.academicjournals.org/AJB/PDF/pdf2006/3Aug/Guasmi%20et%20al.pdf>
- Hernández, F. (1959). *Historia natural de la Nueva España*. Universidad Nacional Autónoma de México. Tomo II. Vol. I. 476.
- Hu, J.; Nakatani, M.; García, L.A.; Kuranouchi, T. & Fujimura, T. (2003). Genetic analysis of sweetpotato and wild relatives using inter-simple sequence repeats (ISSRs). *Breeding Science* Vol.53, No.4, (December 2003), pp. 297-304. ISSN 1437-3735 Available from http://www.jstage.jst.go.jp/article/jsbbs/53/4/297/_pdf
- Ibitoye, D.O. & Akin-Idowu P. E. (2010). Marker-assisted-selection (MAS): A fast track to increase genetic gain in horticultural crop breeding. *African Journal of Biotechnology* Vol.9, No.52, (December 2010), pp. 8889-8895. ISSN 1684-5315 Available from <http://www.academicjournals.org/AJB/PDF/pdf2010/21DecConf/Ibitoye%20and%20Akin-Idowu.pdf>
- Ishikawa, R. (1994). Genetical studies on isozyme genes in rice (in japanese with english summary). *Bulletin Faculty of Agriculture Hirosaki University* Vol.57, pp. 105-180. ISSN 0073-229X
- Jehan, T. & Lakhanpaul, S. (2006). Single nucleotide polymorphism (SNP)-Methods and applications in plant genetics: A review. *Indian Journal of Biotechnology* Vol.5, No.4, (October 2006), pp. 435-439. ISSN 0972-5849 Available from <http://nopr.niscair.res.in/bitstream/123456789/5608/1/IJBT%205%284%29%20435-459.pdf>
- Kumar, S.; Prasad, K.V. & Choudhary, M.L. (2006). Detection of genetic variability among *Chrysanthemum* radiomutants using RAPD markers. *Current Science* Vol.90, No.8 (April 2006), pp. 1108-1113. ISSN 001-3891 Available from <http://www.ias.ac.in/currsci/apr252006/1108.pdf>
- Luna-Paez, A.; Valadez-Moctezuma, E.; Barrientos-Priego, A.F. & Gallegos-Vázquez, C. (2007). Caracterización de *Opuntia* spp. mediante semilla con marcadores RAPD e

- ISSR y su posible uso para diferenciación. *Journal of the Professional Association for Cactus Development* Vol.9, (June 2007), pp. 43-59. ISSN 1938-6648 Available from <http://www.jpacd.org/?modulo=JS&ID=10>
- Marotti, I.; Bonetti, A.; Minelli, M.; Catizone, P. & Dinelli, G. (2007) Characterization of some Italian common bean (*Phaseolus vulgaris* L.) landraces by RAPD, semi-random and ISSR molecular markers. *Genetic Resources and Crop Evolution* Vol.54, No.1, (February 2007), pp. 175-188. ISSN 1573-5109
- Muthusamy, S.; Kanagarajan, S. & Ponnusamy, S. (2008). Efficiency of RAPD and ISSR markers system in accessing genetic variation of rice bean (*Vigna umbellata*) landraces. *Electronic Journal of Biotechnology* Vol.11, No.3, (July 2008), pp. 1-10. ISSN 0717-3458 Available from <http://www.ejbiotechnology.cl/content/vol11/issue3/full/8/index.html>
- Nei M. 1972 Genetic distance between populations. *American Naturalist* 106: 282-283.
- Pérez de la Torre, M.; Acevedo, A.; Serpa, J.C.; Mijayima, I. & Escandón, A.S. (2003). Puesta a punto de la técnica de microsatélites anclados para la caracterización de individuos selectos de jacaranda. En: Mascarini, L., F. Vilella., and E. WRIGHT, (eds). *Floricultura en la Argentina*. Investigación y Tecnología de Producción. 3-12.
- Piña-Escutia, J.L.; Vences-Contreras, C.; Gutiérrez-Martínez, M.G.; Vázquez-García, L.M. & Arzate-Fernández, A.M. (2010). Morphological and molecular characterization of nine botanical varieties of *Tigridia pavonia* (L.f.) DC. *Agrociencia* Vol.44, No.2, (Febrero-Marzo 2010), pp. 147-158. ISSN 1405-3195 Available from <http://www.colpos.mx/agrocien/Bimestral/2010/feb-mar/art-3.pdf>
- Pradeep, R.M.; Sarla, N. & Siddiq, E.A. (2002). Inter simple sequence repeat (ISSR) polymorphism and its application in plant breeding. *Euphytica* Vol.128, No.1, (November 2002), pp. 9-17. ISSN 1573-5060
- Prevost, A. & Wilkinson, M.J. (1999). A new system of comparing PCR primers applied to ISSR fingerprint of potato cultivars. *Theoretical and Applied Genetics* Vol.98, No.1 (January 1999), pp. 107-112. ISSN 1432-2242
- Solouki, M.; Rigi Nazhad, N.; Vignani, R.; Ali Siahisar, B.; Kamaladini, H. & Emamjomeh, A. (2007). Polymorphism of some native sistán grapes assessed by long and short primers for RAPD markers. *Pakistan Journal of Biological Sciences* Vol. 10, No.12, (August 2007), pp. 1996-2001. ISSN 1028-8880 Available from <http://scialert.net/pdfs/pjbs/2007/1996-2001.pdf>.
- Solouki, M.; Mehdikhani, H.; Zeinali H. & Emamjomeh, A. A. (2008). Study of genetic diversity in chamomile (*Matricaria chamomilla*) based on morphological traits and molecular markers. *Scientia Horticulturae* Vol.117, No. 3, (July 2008), pp. 281-287. ISSN 0304-4238
- Stuber, C.W.; Wendel, J.; Goodman, M. & Smith, J. (1988). Techniques and scoring procedures for starch gel electrophoresis of enzymes from maize (*Zea mays* L.) Tech. Bull. 286. North Carolina State University. Raleigh, North Carolina. 217.
- Tanksley, S.D. & Rick, C.M. (1980). Isozyme gene linkage map of the tomato: applications in genetics and breeding. *Theoretical and Applied Genetics* Vol.58, no.2 (March 1980), pp. 161-170. ISSN 1432-2242

- Tapia, C.E.; A. Guillén, H. & Gutiérrez, E.M.A. (2005). Caracterización genética de materiales de piña (*Ananas* spp.) mediante RAPD e ISSR. *Revista Fitotecnia Mexicana* Vol.28, No.3, (Julio-Septiembre 2005), pp. 187-194. ISSN 0187-7380 Available from <http://www.revistafitotecniamexicana.org/documentos/28-3/2a.pdf>
- Torres, A.M., Soost, R.K., & Diedenhofen, U. (1978). Leaf isozymes as genetic markers in citrus. *American Journal of Botany* Vol.65, No.8, (September 1978), pp. 869-881. ISSN 1537-2197
- Vallejos, E. (1983). Enzyme activity staining. In: Tanksley, S. D., and T. J. Orton (eds). *Isozymes in Plant Genetics and Breeding*, Part A. Elsevier, Amsterdam, Netherlands. 459-416.
- Vázquez-García, L.M.; Przybyla, A.A.; De la Cruz, T.E.; Torres, N.H. & Rodríguez, G. (2001). Morphological description of nine botanical varieties of *Tigridia pavonia* (L. F.) Ker. Gawl. *Journal of Applied Botany* Vol.75, No.1-2, pp. 14-19. ISSN 1439-040X
- Vicente, M.C. & Fulton, T. (2003). Tecnologías de Marcadores Moleculares para Estudios de Diversidad Genética de Plantas: Módulo de Aprendizaje. Illus. Nelly Giraldo. Instituto Internacional de Recursos Fitogenéticos (IPGRI), Roma, Italia.1: 1-52.
- Wen, C.S. & Hsiao, J.Y. (2001). Altitudinal genetic differentiation and diversity of Taiwan Lily (*Lilium longiflorum* var. *formosanum*; liliaceae) using RAPD markers and morphological characters. *International Journal of Plant Science* Vol.162, No.2, (March 2001), pp. 287-295. ISSN 1058-5893
- Wendel, J.F. & Weeden, N.F. (1990). Visualization and interpretation of plant isozymes. In: Soltis D. E., and P. S. Soltis. (eds). *Isozymes in plant biology*. Dioscorides Press. Portland, Oregon. pp: 5-45.
- Xu, Y. & Crouch J.H. (2008). Marker-assisted selection in plant breeding: from publications to practice. *Crop Science* Vol.48, No.2, (March-April 2008), pp. 391-407. ISSN 1435-0653 Available from doi: 10.2135/cropsci2007.04.0191
- Yamagishi, M.; Abe, H.; Nakano, M. & Nakatsuka, A. (2002). PCR-based molecular markers in Asiatic hybrid lily. *Scientia Horticulturae* Vol.96, No.1-4, (December 2002), pp. 225-234. ISSN 0304-4238
- Yamagishi, M. (1995). Detection of section-specific random amplified polymorphic DNA (RAPD) markers in *Lilium*. *Theoretical and Applied Genetics* Vol.91, No.6-7, (November 1995), pp. 830-835. ISSN 1432-2242
- Ye, G.N.; Hemmat, M.; Lodhi, M.A.; Weeden, N.F. & Reisch, B.I. (1996). Long primers for RAPD mapping and fingerprinting of grape and pear. *Biotechniques* Vol.20, No.3, (March 1996), pp. 368-371. ISSN 0736-6205 Available from http://www.biotechniques.com/multimedia/archive/00053/19962003368_53006a.pdf
- Ye, Y.M.; Zhang, J.W.; Ning, G.G. & Bao, M.Z. (2008). A comparative analysis of the genetic diversity between inbred lines of *Zinnia elegans* using morphological traits and RAPD and ISSR markers. *Scientia Horticulturae* Vol.118, No.1, (September 2008), pp. 1-7. ISSN 0304-4238
- Yeh, F.C. & Boyle, T.J.B. (1999). Population genetics analysis of codominant and dominant markers and quantitative traits. *Belgium Journal of Botany* Vol.129, pp. 157.

- Young, N.D. (1999). A cautiously optimistic vision for marker assisted breeding. *Molecular Breeding* Vol.5, No.6, (December 1999), pp. 505-510. ISSN 1572-9788
- Zhou, Z.; Miwa, M. & Hogetsu, T. (1999). Analysis of genetic structure of a *Suillus grevillei* population in a *Larix kaemferi* stand by polymorphism of inter-simple sequence repeat (ISSR). *New Phytologist* Vol.144, No.1, (October 1999), pp. 55-63. ISSN 1469-8137 Available from <http://onlinelibrary.wiley.com/doi/10.1046/j.1469-8137.1999.00504.x/pdf>

IntechOpen

IntechOpen



Plant Breeding

Edited by Dr. Ibromkhim Abdurakhmonov

ISBN 978-953-307-932-5

Hard cover, 352 pages

Publisher InTech

Published online 11, January, 2012

Published in print edition January, 2012

Modern plant breeding is considered a discipline originating from the science of genetics. It is a complex subject, involving the use of many interdisciplinary modern sciences and technologies that became art, science and business. Revolutionary developments in plant genetics and genomics and coupling plant "omics" achievements with advances on computer science and informatics, as well as laboratory robotics further resulted in unprecedented developments in modern plant breeding, enriching the traditional breeding practices with precise, fast, efficient and cost-effective breeding tools and approaches. The objective of this Plant Breeding book is to present some of the recent advances of 21st century plant breeding, exemplifying novel views, approaches, research efforts, achievements, challenges and perspectives in breeding of some crop species. The book chapters have presented the latest advances and comprehensive information on selected topics that will enhance the reader's knowledge of contemporary plant breeding.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

José Luis Piña-Escutia, Luis Miguel Vázquez-García and Amaury Marfín Arzate-Fernández (2012). Marker Assisted Characterization in *Tigridia pavonia* (L.f) DC, Plant Breeding, Dr. Ibromkhim Abdurakhmonov (Ed.), ISBN: 978-953-307-932-5, InTech, Available from: <http://www.intechopen.com/books/plant-breeding/marker-assisted-characterization-in-tigridia-pavonia-l-f-dc>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen