

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

**4,800**

Open access books available

**122,000**

International authors and editors

**135M**

Downloads

Our authors are among the

**154**

Countries delivered to

**TOP 1%**

most cited scientists

**12.2%**

Contributors from top 500 universities



**WEB OF SCIENCE™**

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



# The Phytochemical and *In Vitro* Pharmacological Testing of Maltese Medicinal Plants

Everaldo Attard<sup>1</sup> and Pierpaolo Pacioni<sup>2</sup>

<sup>1</sup>University of Malta, Institute of Earth Systems,  
Division of Rural Sciences and Food Systems,

<sup>2</sup>Universita' degli Studi di Perugia, Facoltà di Agraria,

<sup>1</sup>Malta

<sup>2</sup>Italy

## 1. Introduction

### 1.1 General background

The Maltese archipelago is composed of a small number of islands with a total surface area of approximately 457 km<sup>2</sup>. Albeit this small size the Maltese islands host a vast number of plant and animal species. Plant biodiversity, with its 1264 vascular species, is mainly attributed to the strategic position of Malta within the Mediterranean, in which throughout the years several conquerors and civilisations sought to possess Malta particularly for military purposes. In part, the plant diversity of Malta is attributed to introductions brought about by various military forces, as an aid during injury and sickness. Naturally, the phytodiversity has an inclination towards the Mediterranean type of flora with an approximately 66% of the Maltese flora pertaining to this region (E. Attard, 2004). Typical Mediterranean medicinal plants include conifers (*Pinus halepensis* and *Cupressus sempervirens*), broad-leaved trees (*Laurus nobilis*, *Morus nigra* and *Tamarix gallica*), fruit trees (*Ceratonia siliqua*, *Citrus* trees, *Nerium oleander*, *Olea europaea* and *Punica granatum*), and others (*Allium sativum*, *Aloe ferox*, *Capparis spinosa*, *Opuntia ficus-indica*, *Origanum vulgare*, *Papaver somniferum*, *Phytolacca decandra* and *Pistacia lentiscus*). The other portion (34%) is attributed to plants originating from the warm North African (*Cynomorium coccineum*, *Ficus carica* and *Myrtus communis*) and the colder South European regions (*Crataegus monogyna*, *Populus alba* and *Salix species*).

There are approximately 458 medicinal taxa, used in the past to treat one or more ailments (Lanfranco 1993; Lanfranco 1975). Most popular treatments were for the gastrointestinal system, nervous system, cardiovascular system and dermatological conditions. The most predominating plant family within this group is the Asteraceae family, followed by the Lamiaceae and Fabaceae families (Attard, 2004). In spite of their use, these medicinal plants were administered on a trial and error basis. Today, with the advent of modern scientific techniques, the ethnobotanical attributes of a medicinal plant can be challenged by phytochemical and pharmacological testing.

## 1.2 Scientific evaluation of local medicinal and aromatic plants in relation to pharmacology

Locally, only 8 % out of the 458 taxa have been studied scientifically. However, the studies conducted were rather fragmented and covering one or two extracts from a specific plant. Plants include *Ecballium elaterium* (E. Attard et al., 2005; E. Attard & H. Attard, 2008), *Crataegus monogyna* (E. Attard & H. Attard, 2006), *Olea europaea* (Mangion Randon & E. Attard, 2007), *Ephedra fragilis* (E. Attard & Vella, 2009) *Urtica dubia* (Rossi & E. Attard, 2011), *Tetralinis articulata* (Buhagiar et al., 1999) and *Ricinus communis* (Darmanin, 2003) amongst others.

## 1.3 Technical approaches

The evaluation of plant species using different solvent systems has been widely exploited in previous studies (Rodriguez-Lopez et al., 2003; Kumarasamy et al., 2002; Calderon et al., 2003; Konning et al., 2004). A wide spectrum of solvents may be employed when a small number of plants (1-15) are investigated, but when investigating larger numbers or a new group of plants for the first time, the solvents used in ethno-medicine are preferentially selected (Punjani and Kumar, 2003; Guarrera, 2003).

Phytochemical analysis for major classes of metabolites is an important first step in pharmacological evaluation of plant extracts. Some journals require that pharmacological studies be accompanied by a comprehensive phytochemical analysis. Details of such analysis are found in several text books (Harborne, 1984; Evans, 2009). The main secondary metabolite classes include flavonoids, terpenoids and alkaloids, which have been widely tested by the acidified vanillin test, the Salkowski test and the Dragendorff's test, respectively.

Bench top bioassays have been devised to facilitate screening of a large number of samples (Meyer et al., 1982; Carballo et al., 2002). They are based on the principle that pharmacology is simply toxicology at low doses, while toxicology is pharmacology at high doses. Several researchers have used these bioassays for primary pharmacological screening of medicinal plants (Franssen et al., 1997; Kanegusuku et al., 2001; Javidnia et al., 2003). The brine shrimp lethality test (BST), which involves the exposure of brine shrimps to different extract concentrations, is considered as a useful tool for preliminary assessment of cytotoxicity (Jaki et al., 1999). It is a rapid (24 hours), inexpensive and simple technique. A positive correlation has been found between the brine shrimp test and cytotoxicity of the 9KB human nasopharyngeal carcinoma, and other cell lines (Meyer et al., 1982; Kim et al., 2000).

The DNA methyl green bioassay is a simple and comprehensive technique with a high throughput. Methyl green, binds quantitatively to DNA forming a DNA-methyl green complex, hence identifying agents with a high affinity for the DNA. This affinity determines the displacement of methyl green, hence leading to a colourless carbinol (N. Kurnick, 1950; B. Kurnick and Foster, 1950; Krey and Hahn, 1975).

## 1.4 Aims of study

We believe that Maltese medicinal and aromatic plants have a great pharmacological potential. This is based on the concept that, in the past, these plants had important medicinal uses. Therefore, we aimed our study at ethnobotanical research by:

1. Identifying plants cited in ethnobotanical research as active medicinal plants
2. Preparing five extracts using different solvents from each medicinal plant, and the subsequent determination of the classes of metabolites present in the different extracts.
3. Determining whether or not, the extracts obtained eventually possess pharmacological activity employing a primary screening programme.
4. Identifying plant extracts that possess DNA binding.

## 2. Materials and methods

### 2.1 Plant materials

Fifty-five authenticated plant specimens were collected locally during different seasons of the year. The plants were selected on their relative abundance, and collected during their flowering period. The plants were further identified at the Rural Sciences and Food Systems Division, Institute of Earth Systems. Voucher specimens are stored within the Institute. The botanical and ethnobotanical details of the medicinal plants and their voucher specimen code numbers are listed in Table 1.

### 2.2 Preparation of plant extracts

Fresh plants were cut and oven-dried for 48 hours at 35-40°C in a hot air convection oven. Five 300g samples of the dried plants were ground in a heavy duty blender for 20 minutes. 500 ml of solvent (distilled water, distilled water and ethanol (1:1) mixture, ethanol or chloroform or petroleum ether) were added to the respective sample, shaken for 48 hours at 210 rpm, and filtered through a Buchner funnel. Each filtered extract was concentrated at 38 °C under reduced pressure, and finally dried in an oven at 38 °C.

### 2.3 Phytochemical analysis: quantitative colorimetric assays

Although most phytochemical analysis carried out may have a qualitative importance, the methods were modified according to other authors to read absorbance values at a wavelength of 405 nm rather than visual examination. The MTP reader gave more concrete results, in the form of absorbance values. Therefore semi-quantification is possible through this process.

Four colorimetric tests were quantitatively used to determine the presence or absence of metabolites:

1. The Salkowski test for terpenoids. After the addition of chloroform and concentrated sulphuric acid, a reddish brown colouration at the interface forms, hence showing a positive result for the presence of terpenoids (Edeoga et al., 2005);
2. The Dragendorff's test for alkaloids (Steinberg et al., 1997) gives a brown coloration;
3. The Acidified Vanillin test for flavonoids. Under acidic conditions, vanillaldehyde condenses to flavan-3,4-diols, flavan-3-ol monomers and proanthocyanidins to give a cherry-red product (Deshpande et al., 1986);
4. The ninhydrin test was used for proteins (Delhaye & Landry, 1992). The  $\alpha$ -amino acids typically give a blue-purple product.

Voucher specimen number	Botanical Name, family	Maltese, (English) Names	Part/s used, preparation and Maltese Traditional uses
IOA-AMP-002	<i>Acanthus mollis</i> L., Acanthaceae	Ħannewija (Common bear's breeches)	Herb/Emollient as skin softener (Borg, 1927)
IOA-AMP-015	<i>Aloe vera</i> L., Liliaceae	Sabbara (Yellow aloe)	Leaf Juice in child weaning, laxative, increases menstruation (Penza, 1969; Cassar Pullicino, 1947; Cassar, 1964)
IOA-AMP-026	<i>Anagallis arvensis</i> L., Primulaceae	Ħarira ħamra or Ħarira kaħla (Scarlet pimpernel or Blue pimpernel)	Herb and seeds as sudorific and in rabies (Penza, 1969; Gulia, 1855)
IOA-AMP-037	<i>Antirrhinum siculum</i> , Mil., Scrophulariaceae	Papoċċi bojod (Sicilian snapdragon)	Leaves as astringent, diuretic and in chest problems (Penza, 1969; Borg, 1927)
IOA-AMP-036	<i>Antirrhinum tortuosum</i> L., Scrophulariaceae	Papoċċi ħomor (Red snapdragon)	Leaves as astringent, diuretic and in chest problems (Penza, 1969; Borg, 1927)
IOA-AMP-049	<i>Asparagus aphyllus</i> L., Liliaceae	Spraġ selvaġġ (Wild asparagus)	Herb as diuretic (Penza, 1969)
IOA-AMP-453	<i>Aster squamatus</i> (Sprengel) Hieron, Asteraceae	Settembrina selvaġġa (Narrow leaved aster)	A very abundant plant, said to be introduced to the Maltese Islands sometime around the 1930s
IOA-AMP-068	<i>Calendula arvensis</i> L., Asteraceae	Suffejra Selvaġġa (Wild or woody marigold)	Herb in coughs and colds, chiblain, sudorific, warts and calluses, jaundice (Lanfranco, 1993; Penza, 1969)
IOA-AMP-071	<i>Calendula suffruticosa</i> L., Asteraceae	Suffejra Selvaġġa (Wild or woody marigold)	Herb in jaundice (Penza, 1969)
IOA-AMP-081	<i>Carlina gummifera</i> (L.) Les., Asteraceae	Xewk tal-miskta (Stemless atractylis)	Herb is poisonous (Lanfranco, 1993)
IOA-AMP-091	<i>Ceratonia siliqua</i> L., Mimosaceae	Ħarruba (Carob)	Decoction of unripe pods as astringent for the gums and in cough (Penza, 1969; Lanfranco, 1980)
IOA-AMP-145	<i>Cynoglossum creticum</i> Miller, Boraginaceae	Ilsien il-kelb (Southern hound's tongue)	Root decoction and leaf poultice for joint pain and burn relief (Penza, 1969)
IOA-AMP-153	<i>Diplotaxis eruroides</i> (L.) DC., Brassicaceae	Ġarġir (White rocket)	Herb as a stimulant (Penza, 1969)
IOA-AMP-463	<i>Diplotaxis tenuifolia</i> , Brassicaceae	Ġarġir (perennial wall rocket)	Herb as a stimulant (Penza, 1969)
IOA-AMP-223	<i>Dittrichia viscosa</i> (L.) Greut., Asteraceae	Tulliera Komuni (Sticky Fleabane)	Leaf decoction, liquid preparation and oil as haemeostatic, wound healing, itching, improve eye sight; pain, depurative and venereal diseases (Penza, 1969; Lanfranco, 1980; Gulia, 1855; Cassar Pullicino, 1947)
IOA-AMP-460	<i>Eucalyptus globulus</i> , Myrtaceae	Ewkaliptus (Tasmanian Blue)	Oil as astringent and expectorant (Lanfranco, 1993)



Voucher specimen number	Botanical Name, family	Maltese, (English) Names	Part/s used, preparation and Maltese Traditional uses
		Gum)	
IOA-AMP-459	<i>Ferula communis</i> , Apiaceae	Ferla (Giant fennel)	Herb (Penza, 1969)
IOA-AMP-185	<i>Foeniculum vulgare</i> Miller, Apiaceae	Busbies (fennel)	Seeds and herb as flavouring agent in liquid preparations and treatment of colic pain (Penza, 1969)
IOA-AMP-191	<i>Fumaria capreolata</i> , Fumariaceae	Dahnet l-art (Fumitory)	Herb infusion as tonic, taenifuge, stomachic, kidney stones, in bath for sick children (Borg, 1927; Penza, 1969; Gulia, 1855)
IOA-AMP-190	<i>Fumaria officinalis</i> L., Fumariaceae	Dahnet l-art (Fumitory)	Herb infusion as tonic, taenifuge, stomachic, kidney stones, in bath for sick children (Borg, 1927; Penza, 1969; Gulia, 1855)
IOA-AMP-454	<i>Galactites tomentosa</i> Moench, Asteraceae	Xewka bajda (Boar thistle)	Herb consumed as a monofloral boar thistle honey
IOA-AMP-197	<i>Gladiolus italicus</i> Gaud., Iridaceae	Gladjoli salvaġġ (Common cornflag)	Leaves and bulb as galactagogue, aphrodisiac and emmenagogue (Penza, 1969; Borg, 1927)
IOA-AMP-101	<i>Glebionis coronaria</i> Tzvelev, Asteraceae	Lellux or Żigland (Crown daisy)	Herb (Lanfranco, 1993)
IOA-AMP-202	<i>Hedera helix</i> L., Araliaceae	Liedna (Ivy)	Gum and leaves in wound healing and as astringent (Penza, 1969)
IOA-AMP-461	<i>Holoschoenus vulgaris</i> , Cyperaceae	Simar tal-boċċi (roundhead bulrush)	A common plant in halophytic environments
IOA-AMP-213	<i>Hyoscyamus albus</i> L., Solanaceae	Mammażejża (White henbane)	Leaf poultice and ointment as sedative, in haemorrhoids and wound healing (Penza, 1969)
IOA-AMP-217	<i>Hypericum aegyptiacum</i> L., Guttiferae	Fexfiex il-baħar (Egyptian St. John's wort)	Herb Juice in wound healing, urinary tract infections and increases menstrual flow (Penza, 1969)
IOA-AMP-450	<i>Inula crithmoides</i> L., Asteraceae	Xorbett (Golden samphire)	Herb (Gulia, 1855)
IOA-AMP-462	<i>Lactuca sativa</i> , Asteraceae	ħassa salvaġġa (Wild lettuce)	Leaf poultice as sedative (Penza, 1969)
IOA-AMP-236	<i>Lactuca virosa</i> , Asteraceae	ħassa salvaġġa (Wild lettuce)	White latex as sedative (Penza, 1969)
IOA-AMP-234	<i>Laurus nobilis</i> L., Lauracea	Rand (Laurel)	Seed oil and leaf decoction in rheumatic pain and neuralgia; stomachic; diaphoretic, depurative (Penza, 1969; Cassar Pullicino, 1947; Lanfranco, 1980; Cremona, 1971)
IOA-AMP-238	<i>Leontodon tuberosus</i> , Asteraceae	Żigland (Tuberous hawkbit)	Herb as diuretic and tonic (Lanfranco, 1993)
IOA-AMP-254	<i>Malva sylvestris</i> L., Malvaceae	ħubbejża (Common mallow)	Leaf/flower poultices and root decoction in vaginitis, intestinal problems, depurative, skin and throat inflammation (Penza, 1969; Lanfranco, 1980)
IOA-AMP-268	<i>Mercurialis annua</i> L., Euphorbiaceae	Burikba (Annual mercury)	Juice as tonic and galactofuge (Penza, 1969; Lanfranco, 1975)

Voucher specimen number	Botanical Name, family	Maltese, (English) Names	Part/s used, preparation and Maltese Traditional uses
IOA-AMP-285	<i>Nerium oleander</i> L., Apocynaceae	Oljandru (Oleander)	Herb for skin itching (Cassar Pullicino, 1947)
IOA-AMP-290	<i>Olea europaea</i> L., Oleaceae	Żebbuġa (Olive)	Olive oil and leaves as laxative, wound healing, sunburn, antihypertensive, aching muscles (Penza 1969; Lanfranco, 1980)
IOA-AMP-286	<i>Opuntia ficus-indica</i> (L.) Mill., Cactaceae	Bġħajtar tax-xewk (Prickly pear)	Cladode/flower poultice in stomach pain, burnt skin, joint pain/headaches; astringent and antidiarrhoeal (Cassar Pullicino, 1947; Lanfranco, 1980; Lanfranco, 1975)
IOA-AMP-291	<i>Oxalis pes-caprae</i> L., Oxaliaceae	Ħaxixa ingliza, Cape sorrel	Herb juice as emetic and for acne (Lanfranco, 1975)
IOA-AMP-090	<i>Palaeocyanus crassifolius</i> (Bert.) Dost., Asteraceae	Widnet il-baħar (Maltese rock centauray)	National Plant of Malta
IOA-AMP-294	<i>Papaver somniferum</i> L. Papaveraceae	Xaħxieh (Opium poppy)	Poppy heads and latex as sedative (Penza, 1969)
IOA-AMP-296	<i>Parietaria judaica</i> , Urticaceae	Xeħt ir-riħ (Pellitory of the wall)	Herb, decoction; herb boiled with garlic and chamomile in bronchitis, pharyngitis, pulmonitis and cough; catarrh; kidney stones; haemorrhoids (Borg, 1927; Penza, 1969; Cassar Pullicino, 1947)
IOA-AMP-304	<i>Phlomis fruticosa</i> L., Lamiaceae	Salvja tal-Madonna (Jerusalem sage)	Boiled leaves as cough remedy (Penza, 1969)
IOA-AMP-317	<i>Pinus halepensis</i> Miller, Pinaceae	Żnuber (Aleppo pine)	Inhalation and ointment for catarrh and as diuretic (Lanfranco, 1975)
IOA-AMP-319	<i>Pistacia lentiscus</i> L., Anacardiaceae	Deru (Mastic tree)	Mastic resin for filling of teeth (Gulia, 1855)
IOA-AMP-318	<i>Plantago lagopus</i> L., Plantaginaceae	Beżbula komuni (Hare's foot plantain)	Boiled leaves for wound healing, eye diseases and increases urination (Penza, 1969; Cassar Pullicino, 1947)
IOA-AMP-331	<i>Prasium majus</i> L., Lamiaceae	Te Sqalli (Mediterranean Prasium)	Infused leaves as diuretic (Penza, 1969; Gulia, 1855; Cremona, 1971)
IOA-AMP-345	<i>Psoralea bituminosa</i> L., Mimosaceae	Silla tal-blat (Bitumen pea)	Herb in rheumatic pain (Penza, 1969)
IOA-AMP-308	<i>Reicardia picroides</i> , Asteraceae	Kanċlita (Common reichardia)	Herb as diuretic and tonic (Lanfranco, 1993)
IOA-AMP-348	<i>Reseda alba</i> L., Resedaceae	Denb il-haruf (White mignonette)	Roots for painful gums (Borg, 1927; Penza, 1969)
IOA-AMP-360	<i>Ricinus communis</i> L., Euphorbiaceae	Riġnu (Castor oil tree)	Decoction of seeds, roots or leaves as laxative, rheumatism, neuralgic affections, ophthalmia; galactorrhoea (Penza, 1969)
IOA-AMP-374	<i>Schinus terebinthifolius</i> , Anacardiaceae	Bżar Falz (Drooping false pepper)	Ground fruit (Borg, 1927)
IOA-AMP-392	<i>Silybum marianum</i> (L.) Gaertn., Asteraceae	Xewk Bagħli (Milk thistle)	Herb as tonic, urinary tract, fever (Penza, 1969)

Voucher specimen number	Botanical Name, family	Maltese, (English) Names	Part/s used, preparation and Maltese Traditional uses
IOA-AMP-388	<i>Smyrniolum olusatrum</i> L., Apiaceae	Karfus il-ħmir (Alexanders)	Herb as stimulant (Penza, 1969)
IOA-AMP-393	<i>Sonchus oleraceus</i> L., Asteraceae	Tfief (Sow thistle)	Herb as diuretic and purgative (Penza, 1969)
IOA-AMP-443	<i>Verbena officinalis</i> L., Verbenaceae	Buqexrem (Vervain)	Poultice/decoction for wound healing, astringent, diarrhoea, dysentery, diabetes (Penza, 1969)

Table 1. Botanical, ethnobotanical and voucher specimen code numbers for the fifty-five plants studied.

## 2.4 Brine Shrimp Test (BST)

In a set of 12-well plates, each well contained 10 nauplii, 1 ml sea water and 1 ml of extract diluted to final concentrations of 1%, 0.1%, 0.01%, 0.001% and 0.0001% respectively. The tests were set out in triplicate so that a total of fifteen wells per extract were used. Numbers of living nauplii were counted after 24 hours. The  $LC_{50}$  values and 95 % confidence intervals were determined in  $\mu\text{g/ml}$ , using the Finney probit analysis computer program. A median lethal concentration ( $LC_{50}$ ) smaller than 1000  $\mu\text{g/ml}$  (Alkofahi et al., 1997) indicates pharmacological activity.

## 2.5 DNA-methyl green (intercalation) tests

DNA intercalation assay for DNA activity. Samples were incubated with 200  $\mu\text{l}$  of DNA-methyl green in the dark at 25 °C for 24 h. The decrease in absorbance at 650 nm was calculated as a percentage of the untreated DNA-methyl green absorbance value. The median inhibitory concentration ( $IC_{50}$ ) was calculated (Desmarchelier et al., 1996) through regression analysis. Cucurbitacin E and Dexamethasone were used as potent and moderate positive controls, respectively. Data was analyzed using Student's t-test.

## 3. Results and discussion

In this study, 55 plant species from 31 plant families were studied. The plant families ranked in the following order: Asteraceae (15 species), Apiaceae (3 species), Liliaceae, Scrophulariaceae, Mimosaceae, Brassicaceae, Fumariaceae, Euphorbiaceae, Lamiaceae and Anacardiaceae (2 species), Acanthaceae, Primulaceae, Boraginaceae, Iridaceae, Araliaceae, Solanaceae, Guttiferae, Lauracea, Malvaceae, Apocynaceae, Cactaceae, Oleaceae, Oxaliaceae, Papaveraceae, Urticaceae, Pinaceae, Plantaginaceae, Resedaceae, Verbenaceae, Myrtaceae and Cyperaceae (1 species). The distribution of plants within families was as broad as possible. However, the most abundant plant family of the Maltese flora (Attard, 2004) was given more importance than the other families.

### 3.1 Phytochemical analysis

The results for the four phytochemical classes are illustrated in table 2 and a generalised picture of the number of extracts, containing phytochemicals for each solvent system used, is illustrated in figure 1.



PLANT NAME	P. N°	Aqueous	Aqueous-ethanol	Ethanol	Chloroform	Petroleum ether
<i>Acanthus mollis</i>	002	TP	TP	TFP	TP	T
<i>Aloe vera</i>	015	-	-	TF	AF	-
<i>Anagallis arvensis</i>	028	P	P	P	FP	P
<i>Antirrhinum siculum</i>	037	P	TP	TP	TP	T
<i>Antirrhinum tortuosum</i>	036	TFP	TFP	TFP	T	TP
<i>Arum italicum</i>	046	TFP	TFP	TFP	T	T
<i>Asparagus aphyllus</i>	049	TFP	TFP	TFP	TFP	TAFP
<i>Aster squamatus</i>	453	FP	AFP	P	-	F
<i>Calendula arvensis</i>	068	TAP	TFP	AFP	-	F
<i>Calendula suffruticosa</i>	073	F	TFP	TFP	-	P
<i>Carlina gummifera</i>	081	TAFP	FP	A	-	TF
<i>Ceratonia siliqua</i>	091	AF	F	FP	TAF	AF
<i>Cynoglossum creticum</i>	145	FP	TFP	TFP	-	TP
<i>Diplotaxis eruroides</i>	153	P	P	T	-	-
<i>Diplotaxis tenuifolia</i>	463	TF	FP	FP	TF	T
<i>Dittrichia viscosa</i>	223	TFP	FP	TF	FP	AFP
<i>Eucalyptus globulus</i>	460	TFP	TFP	TFP	TFP	-
<i>Ferula communis</i>	459	TP	TP	TP	-	FP
<i>Foeniculum vulgare</i>	185	T	TP	TP	-	F
<i>Fumaria capreolata</i>	191	TP	TFP	TFP	TF	TP
<i>Fumaria officinalis</i>	190	TP	TP	TP	F	P
<i>Galactites tomentosa</i>	454	TFP	TFP	AP	-	TFP
<i>Gladiolus italicus</i>	197	-	-	TAF	-	TF
<i>Glebionis coronaria</i>	101	TFP	T	TAFP	-	TF
<i>Hedera helix</i>	202	-	TP	T	-	TF
<i>Holoschoenus vulgaris</i>	461	TFP	FP	F	F	T
<i>Hyoscyamus albus</i>	213	P	A	TP	-	TP
<i>Hypericum aegyptiacum</i>	217	FP	F	-	F	TP
<i>Inula crithmoides</i>	450	TFP	TFP	TFP	F	TF
<i>Lactuca sativa</i>	462	T	T	P	P	F
<i>Lactuca virosa</i>	236	-	P	-	TP	F
<i>Laurus nobilis</i>	234	F	T	T	-	TAF
<i>Leontodon tuberosus</i>	238	TFP	TFP	TF	F	F
<i>Malva sylvestris</i>	254	T	TP	T	-	-
<i>Mercurialis annua</i>	268	TF	TF	F	TFP	TF
<i>Nerium oleander</i>	285	TF	TFP	TFP	P	TF
<i>Olea europaea</i>	290	TFP	TFP	TF	TF	TFP
<i>Opuntia ficus-indica</i>	286	T	TP	TFP	P	TP
<i>Oxalis pes-caprae</i>	291	TF	TFP	TF	TF	A
<i>Palaeocyanus crassifolius</i>	90	TFP	TP	TP	TP	TFP
<i>Papaver somniferum</i>	294	FP	TP	TP	FP	TF

PLANT NAME	P. N°	Aqueous	Aqueous-ethanol	Ethanol	Chloroform	Petroleum ether
<i>Parietaria judaica</i>	296	F	FP	TFP	FP	AP
<i>Phlomis fruticosa</i>	304	TP	TFP	TFP	T	TFP
<i>Pinus halepensis</i>	317	P	-	TF	F	AF
<i>Pistacia lentiscus</i>	319	TFP	TF	P	-	-
<i>Plantago lagopus</i>	318	-	FP	AFP	TP	AP
<i>Prasium majus</i>	331	TFP	TFP	TFP	TP	P
<i>Psoralea bituminosa</i>	345	TP	TFP	T	T	P
<i>Reichardia picroides</i>	308	TFP	TFP	TP	F	TFP
<i>Ricinus communis</i>	360	TFP	TP	TFP	TF	-
<i>Schinus terebinthifolius</i>	374	TFP	TP	TP	P	TF
<i>Smyrniium olusatrum</i>	388	TFP	P	TFP	F	P
<i>Sonchus oleraceus</i>	393	TF	TF	TF	-	TF
<i>Verbena officinalis</i>	443	P	TP	TP	-	P

Table 2. The phytochemical analysis of the extracts under investigation for the main phytochemical classes: Flavonoids (F), Terpenoids (T), Alkaloids (A) and Proteins (P)




Fig. 1. A generalised profile of the number of extracts containing terpenoids, alkaloids, flavonoids and proteins for each solvent system used (n=280).

The predominating compound classes were terpenoids (56.07 %), followed by proteins (53.57 %) and flavonoids (48.93 %). Alkaloids were limited to a smaller number of extracts (7.50 %). The majority of the polar solvents, aqueous, aqueous-ethanol and ethanol contained terpenoids and proteins ( $p < 0.05$ ,  $n = 4$ ). The chloroform extract contained mainly flavonoids ( $p < 0.05$ ,  $n = 4$ ), while the petroleum ether extracts contained predominantly flavonoids and terpenoids.

The highest terpenoid contents were found in the ethanol and aqueous-ethanol extracts. In fact, it was observed that 70.70 % of the positive extracts were polar extracts, i.e. using

water, water-ethanol and ethanol as extracting solvents. This is due to the fact that most terpenoids are present in the glycosidic form rather than the non-polar or low polarity terpene aglycone form. Some plants exhibited the presence of terpenes and related compounds in all solvent systems. Typical examples included *Acanthus mollis*, which mainly contains  $\beta$ -sitosterol as the triterpene-like compound (Loukis & Philianos, 1980), *Antirrhinum tortuosum*, with mono and sesquiterpene volatile derivatives (Nagegowda et al., 2008), *Arum italicum*, with the tetraterpene carotenoids (Bonora et al., 2000), *Asparagus aphyllus* with saponins and sapogenins (Shao et al., 1996), *Olea europaea* containing mainly triterpenoids (Caputo et al., 1974; Elamrani, 2011), *Palaeocyanus crassifolius* containing sesquiterpene lactones (Koukoulitsa et al., 2002) and *Phlomis fruticosa*, mainly containing mono- and sesquiterpenes (Amor et al., 2009). In the case of *Fumaria capreolata* the main constituents mentioned in previous studies were the alkaloids (Soušek et al., 1999; Maiza-Benabdesselam et al., 2007). In this present study, there was the strong presence of terpenoids.

The distribution of alkaloids in polar and non-polar solvents was almost equal (52.38 % and 47.62 %, respectively). Alkaloids may be present either as the non-polar organic form or as the polar ionised alkaloid salt. The highest content was recorded in the ethanol extract of *Gladiolus italicus* and in the chloroform extract of *Aloe vera*. For *Gladiolus*, this result goes in accordance with that obtained by Ameh and coworkers (2011) and for *Aloe*, a similar result was obtained by Waller and coworkers (1978). Other plants with an alkaloidal content include *Asparagus aphyllus* (Negi et al., 2010), *Calendula arvensis* (Shamsa et al., 2008), *Glactites tomentosa*, *Glebionis coronaria*, *Oxalis pes-caprae*, *Parietaria judaica*, *Carlina gummifera*, *Hyoscyamus albus* (Doerk-Schmitz et al., 1993), *Laurus nobilis* (Nayak et al., 2006), *Pinus halepensis* (Tawara et al., 1993) and *Plantago lagopus* (Hultin & Torssell, 1965). *Fumaria* species are known to contain alkaloids (Soušek et al., 1999; Maiza-Benabdesselam et al., 2007). However, no alkaloids were detected for *Fumaria officinalis* and *Fumaria capreolata* in this present study. Although *Ceratonia siliqua* is claimed to contain no alkaloids (El Hajaji et al., 2011), in this present study, alkaloids were detected in the aqueous, chloroform and petroleum ether extracts. It was also observed that for *Papaver somniferum* no alkaloids were detected in the leaves. This depends on several factors. Primarily, the wild variety might have a low potential for the production of morphinan alkaloids, and other plant parts such as the stem, roots and capsules, tend to accumulate more alkaloids than the leaves (Williams & Ellis, 1989).

For the flavonoid group, out of the positive responses, 65.69 % were polar extracts while the rest (34.41 %) were extracts derived from non-polar solvents. Typically, flavonoids are polyphenolic compounds that are highly soluble in aqueous and aqueous-alcohol solvents. However, flavonoids have been reported to be also extracted by chloroform and petroleum ether (Gudej & Czapski, 2009; Rajendran & Krishnakumar, 2010). Plants containing flavonoids in all extracts, consistent with other studies, include *Asparagus aphyllus* (Sun et al., 2007), *Ceratonia siliqua* (Papagiannopoulos et al., 2004; Vaya & Mahmood, 2006), *Dittrichia viscosa* (M.J. Martin et al., 1988), *Leontodon tuberosus* (Zidorn & Stuppner, 2001), *Mercurialis annua* (Aquino et al., 1987) and *Olea europaea* (Benavente-García et al., 2000). Almost all plant species exhibited the presence of flavonoids in one or more extracts, except for four plants, namely, *Antirrhinum siculum*, *Diplotaxis eruroides*, *Malva sylvestris* and *Verbena officinalis*. Other studies report the presence of flavonoids in *Diplotaxis eruroides* (Bennett et al., 2006), *Malva sylvestris* (Billeter et al., 1991) and *Verbena officinalis* (Rehecho et al., 2011).

The absence of flavonoids in these species for the current study may be due to several factors that include a different chemotype, different environmental conditions and the presence of these compounds below the detection limit, amongst others. *Antirrhinum siculum* is palely pigmented and this may contribute to the insignificant content of flavonoids (C. Martin et al., 2010).

Proteins prevail in many plants. Within the positive response group, 74.67 % were polar extracts while 25.33 % were non-polar extracts. This indicated that three-fourths of the detected proteins were functional proteins including enzymes. *Anagallis arvensis*, *Asparagus aphyllus*, *Palaeocyanus crassifolius* and *Prasium majus* exhibited the presence of proteins in all their extracts. This goes in accordance with previous studies carried out on these plants (Alignier et al., 2008; King et al., 1990). Plants that were devoid of proteins in all their extracts include *Aloe vera*, *Gladiolus italicus*, *Laurus nobilis* and *Sonchus oleraceus*. In previous studies, *Aloe vera* revealed the presence of glutathione peroxidase (Sabeih et al., 1993), *Gladiolus italicus* contained arabinogalactan-protein (Gleeson & Clarke, 1979) and *Laurus nobilis* contained lipase (Isbilir et al., 2008). Although most plant material was collected at flowering time, the inclusion of seed protein in the extract would have been possible in cases where fruit were harvested alongside the flowers.

### 3.2 The Brine Shrimp Test

The results for the tested extracts are given in Table 3. Primary screening involves the use of bench-top bioassays. Extracts exhibiting  $LC_{50}$  values above 1000  $\mu\text{g}/\text{ml}$  are generally regarded as ineffective extracts. In this study, 42.26 % of the extracts were therefore inactive (Table 4). The most inactive were the petroleum ether extracts, while the most active were the ethanolic extracts. Correlating the BST lethal concentrations to phytochemical classes, it was observed that inactive extracts contained several phytochemicals. The reason may be due to the low concentration or possible antagonistic activity between the phytochemicals from the different classes. 55.68 % of the extracts exhibited  $LC_{50}$  values below 1000  $\mu\text{g}/\text{ml}$ . The most active were the ethanolic extracts (72.97 %), while the least active were the petroleum ether extracts (35.14 %). Four plants exhibited activity for all their five extracts. These were *Nerium oleander*, *Olea europaea*, *Opuntia ficus-indica* and *Pinus halepensis*, all exhibiting  $LC_{50}$  values below 0.01  $\mu\text{g}/\text{ml}$ . These four plant species are amongst the most popular Maltese traditional medicinal plants. It was also observed that some extracts with non-detectable phytochemicals exhibited significant  $LC_{50}$  values. Typical examples include the aqueous extract of *Lactuca virosa*, the aqueous-ethanol extract of *Pinus halepensis*, the ethanolic extracts of *Hypericum aegypticum* and *Lactuca virosa*, and the chloroform extracts of *Ferula communis*, *Foeniculum vulgare* and *Pistacia lentiscus*. On the other hand, there were extracts that exhibited significant  $LC_{50}$  values as opposed to other studies. For example, for *Fumaria officinalis* aqueous-ethanol and ethanol extracts, in the present study, exhibited significant effects on brine shrimps as opposed to the ethanol extract reported in the study by Erdoğan (2009).

### 3.3 The DNA-methyl green assay

Table 5 shows the  $IC_{50}$  values obtained for the DNA-methyl green assay. Although low  $IC_{50}$  values have been reported for pure compounds (Burres et al., 1992), such as rubiflavin and

distamycin A (17 and 18  $\mu\text{g}/\text{ml}$ , respectively), it is reasonable that in the case of extracts higher  $\text{IC}_{50}$  values are acceptable as for pyrido[2,3-d]pyrimidin-4(1H)-one and pyrido[2,3-d]triazolo[3,4-b]pyrimidine analogs (40 – 53  $\mu\text{g}/\text{ml}$ ) (Goda & Badria, 2005). Since plant extracts are complex matrices with several phytochemicals,  $\text{IC}_{50}$  values are expected to be higher than for pure compounds. Therefore, extracts with  $\text{IC}_{50}$  values below 70  $\mu\text{g}/\text{ml}$  were considered as active (Figure 2). Only 15 % of the extracts displaced methyl green from the methyl green DNA complex. It is likely that these compounds act as intercalating agents at the DNA level. 86.67 % were active polar extracts with proteins predominating in these extracts. The other extracts either exhibited an  $\text{IC}_{50}$  value higher than 70  $\mu\text{g}/\text{ml}$  or else a 50 % activity was never achieved. From the remaining 85 %, only one-third of the extracts exhibited values above 70  $\mu\text{g}/\text{ml}$ . Alkaloids only featured in one active aqueous extract of *Ceratonia siliqua*. Terpenoids, flavonoids and proteins predominated mainly in aqueous and aqueous-ethanol extracts. For a few extracts, there was no correlation between the phytochemical class and DNA-methyl green activity. These include the aqueous-ethanol extract of *Gladiolus italicus*, the aqueous extract of *Hedera helix* and the ethanolic extract of *Hypericum aegyptiacum*. For example *H. aegyptiacum* contains hypericin that can inhibit DNA topoisomerase II (Peebles et al., 2001), but the naphthodianthrone was not detected by the phytochemical tests.




Fig. 2. The number of extracts classified as (a) below 70  $\mu\text{g}/\text{ml}$ , (b) above 70  $\mu\text{g}/\text{ml}$  range and (c) non-active extracts with the different solvent types for the DNA methyl green assay.



PLANT NAME	P. N°	Aqueous	Aqueous-ethanol	Ethanol	Chloroform	Petroleum ether
<i>Acanthus mollis</i>	002	<0.01	<0.01	<0.01	<0.01	>1000
<i>Anagallis arvensis</i>	028	460	<0.01	<0.01	>1000	>1000
<i>Antirrhinum tortuosum</i>	036	<0.01	>1000	<0.01	>1000	<0.01
<i>Antirrhinum siculum</i>	037	10	<0.01	<0.01	>1000	<0.01
<i>Asparagus aphyllus</i>	049	>1000	<0.01	<0.01	<0.01	>1000
<i>Aster squamatus</i>	453	>1000	>1000	>1000	>1000	>1000
<i>Calendula arvensis</i>	068	>1000	63	>1000	>1000	796
<i>Carlina gummifera</i>	081	>1000	>1000	>1000	>1000	>1000
<i>Dittrichia viscosa</i>	223	>1000	>1000	>1000	<0.01	>1000
<i>Ferula communis</i>	459	<0.01	<0.01	<0.01	<0.01	>1000
<i>Foeniculum vulgare</i>	185	>1000	<0.01	<0.01	<0.01	>1000
<i>Fumaria officinalis</i>	190	>1000	<0.01	<0.01	>1000	<0.01
<i>Fumaria capreolata</i>	191	>1000	>1000	<0.01	<0.01	>1000
<i>Galactites tomentosa</i>	454	>1000	>1000	>1000	>1000	>1000
<i>Glebionis coronaria</i>	101	>1000	>1000	93	131	>1000
<i>Hyoscyamus albus</i>	213	>1000	ND	<0.01	>1000	>1000
<i>Hypericum aegyptiacum spreng</i>	217	<0.01	<0.01	<0.01	>1000	>1000
<i>Inula crithmoides</i>	450	>1000	344	>1000	>1000	562
<i>Lactuca sativa</i>	462	<0.01	<0.01	<0.01	>1000	<0.01
<i>Lactuca virosa</i>	236	<0.01	<0.01	<0.01	>1000	>1000
<i>Leontodon tuberosus</i>	238	>1000	>1000	>1000	>1000	>1000
<i>Nerium oleander</i>	285	<0.01	<0.01	<0.01	<0.01	<0.01
<i>Opuntia ficus- indica</i>	286	<0.01	<0.01	<0.01	<0.01	<0.01
<i>Olea europaea</i>	290	<0.01	<0.01	<0.01	<0.01	<0.01
<i>Oxalis pes-caprae</i>	291	<0.01	<0.01	<0.01	<0.01	>1000
<i>Palaeocyanus crassifolius</i>	090	>1000	<0.01	<0.01	>1000	10
<i>Papaver somniferum</i>	294	<0.01	<0.01	<0.01	<0.01	>1000
<i>Pinus halepensis</i>	317	<0.01	<0.01	<0.01	<0.01	<0.01
<i>Plantago lagopus</i>	318	>1000	10	<0.01	>1000	0.07
<i>Pistacia lentiscus</i>	319	<0.01	<0.01	<0.01	<0.01	ND
<i>Prasium majus</i>	331	<0.01	<0.01	<0.01	<0.01	>1000
<i>Psoralea bituminosa</i>	345	<0.01	<0.01	<0.01	10	<0.01
<i>Reichardia picroides</i>	308	>1000	>1000	>1000	>1000	>1000
<i>Reseda alba</i>	348	>1000	<0.01	ND	10	>1000
<i>Ricinus communis</i>	360	>1000	<0.01	<0.01	<0.01	>1000
<i>Schinus terebinthifolius</i>	374	<0.01	<0.01	<0.01	<0.01	>1000
<i>Sonchus oleraceus</i>	393	>1000	>1000	>1000	>1000	>1000

Table 3. The result for the effect of extracts on the Brine Shrimp Test

BST result	Percentage per extract type					Percentage of Total Extracts
	Aqueous	Aqueous-ethanol	Ethanol	Chloroform	Petroleum ether	
>1000	51.35	27.03	24.32	48.65	59.46	42.16 %
0.01-1000	5.41	8.11	2.70	8.11	10.81	7.03 %
0.01	43.24	62.16	70.27	43.24	24.32	48.65 %
ND	0.00	2.70	2.70	0.00	5.41	2.16 %

Table 4. The percentage of results classified as (a) above 1000 µg/ml, (b) 0.01 – 1000 µg/ml range, (c) less than 0.01 µg/ml and (d) not determined (ND) with the different solvent types for the brine shrimp test.

PLANT NAME	P. N°	Aqueous	Aqueous-ethanol	Ethanol	Chloroform	Petroleum ether
<i>Acanthus mollis</i>	2	30.399	34.102	NA	131.005	34.354
<i>Aloe vera</i>	15	NA	278.589	270.983	NA	NA
<i>Anagallis arvensis</i>	28	28.658	43.534	NA	NA	NA
<i>Antirrhinum tortuosum</i>	36	63.354	156.171	314.838	38.065	354.278
<i>Antirrhinum siculum</i>	37	NA	NA	NA	NA	NA
<i>Asparagus aphyllus</i>	49	NA	52.985	NA	305.865	NA
<i>Calendula suffruticosa</i>	73	70.296	144.921	350.003	261.826	364.324
<i>Ceratonia siliqua</i>	91	23.230	NA	NA	134.563	NA
<i>Cynoglossum creticum</i>	145	45.941	50.063	NA	NA	NA
<i>Eucalyptus globulus</i>	460	NA	NA	NA	NA	NA
<i>Ferula communis</i>	459	NA	NA	NA	71.158	92.264
<i>Foeniculum vulgare</i>	185	NA	NA	NA	NA	NA
<i>Fumaria capreolata</i>	191	133.655	77.458	NA	596.272	NA
<i>Fumaria officinalis</i>	190	NA	346.108	NA	NA	79.496
<i>Gladiolus italicus</i>	197	NA	54.931	NA	143.109	251.228
<i>Hedera helix</i>	202	60.086	231.921	NA	NA	NA
<i>Holoschoenus vulgaris</i>	461	NA	91.229	NA	NA	NA
<i>Hyoscyamus albus</i>	213	NA	NA	195.094	NA	NA
<i>Hypericum aegyptiacum</i>	217	NA	NA	66.803	NA	443.799
<i>Diplotaxis tenuifolia</i>	463	NA	30.688	NA	NA	NA
<i>Dittrichia viscosa</i>	223	26.945	NA	108.418	171.989	NA
<i>Laurus nobilis</i>	234	50.835	NA	324.334	189.539	NA
<i>Lactuca sativa</i>	462	NA	166.722	326.359	NA	NA
<i>Lactuca virosa</i>	236	NA	NA	NA	NA	57.791
<i>Malva sylvestris</i>	254	NA	31.530	NA	NA	54.922
<i>Mercurialis annua</i>	268	NA	165.974	NA	NA	NA
<i>Nerium oleander</i>	285	NA	62.957	NA	105.401	163.761
<i>Opuntia ficus-indica</i>	286	41.899	85.339	NA	115.870	NA
<i>Olea europaea</i>	290	53.908	95.687	80.988	227.594	NA
<i>Oxalis pes-caprae</i>	291	NA	NA	64.089	223.200	171.288
<i>Palaeocyanus crassifolius</i>	90	122.931	NA	NA	NA	NA
<i>Papaver somniferum</i>	294	147.538	NA	NA	NA	131.727

<i>Parietaria judaica</i>	296	NA	NA	89.115	NA	NA
<i>Phlomis fruticosa</i>	304	137.404	NA	159.786	130.591	171.195
<i>Pinus halepensis</i>	317	NA	NA	NA	NA	NA
<i>Plantago lagopus</i>	318	NA	49.252	338.260	NA	NA
<i>Pistacia lentiscus</i>	319	NA	NA	NA	NA	NA
<i>Prasium majus</i>	331	48.142	103.372	273.826	92.489	279.194
<i>Psoralea bituminosa</i>	345	NA	59.332	NA	NA	177.895
<i>Reseda alba</i>	348	193.473	NA	NA	NA	NA
<i>Ricinus communis</i>	360	48.912	NA	NA	103.224	NA
<i>Schinus terebinthifolius</i>	374	NA	NA	NA	NA	181.849
<i>Smyrniium olusatrum</i>	388	104.605	NA	NA	NA	NA
<i>Silybum marianum</i>	392	NA	46.528	NA	NA	NA
<i>Verbena officinalis</i>	443	67.656	104.453	145.529	158.008	NA
Cucurbitacin E	-	19.12				
Dexamethasone	-	32.74				

Table 5. The median inhibitory concentration (IC<sub>50</sub> in µg/ml) values obtained for the DNA-methyl green assay (NA no activity – 50% effect was never reached).

#### 4. Conclusions

This study has confirmed the presence of useful phytochemicals and biological activities of several extracts from selected Mediterranean plants. It is expected that these results will serve as a stimulus for further investigations into the active phytochemicals.

#### 5. Acknowledgment

This work was supported and funded by the Malta Council for Science and Technology, Malta (Project code: RTDI-2004-074).

#### 6. References

- Alignier, A., Meiss, H., Petit, S. & Reboud, X. (2008). Variation of post-dispersal weed seed predation according to weed species, space and time. *Journal of Plant Diseases and Protection*, Vol. 21, pp. 221-228, ISSN 1861-3829
- Ameh, S.J., Obodozie, O.O., Olorunfemi, P.O., Okoliko, I.E. & Ochekepe, N.A. (2011). Potentials of *Gladiolus* corms as an antimicrobial agent in food processing and traditional medicine. *Journal of Microbiology and Antimicrobials*, Vol. 3, No. 1, pp. 8-12, ISSN 2141-2308
- Amor, I.L.B., Boubakera, J., Sgaiera, M.B., Skandrania, I., Bhouria, W., Neffati, A., Kilani, S., Bouhlel, I., Ghedira, K. & Chekir-Ghedira, L. (2009). Phytochemistry and biological activities of *Phlomis* species. *Journal of Ethnopharmacology*, Vol. 125, No. 2, pp. 183-202, ISSN 0378-8741
- Aquino, R., Behar, I., D'agostino, M., De Simone, F., Schettino, O. & Pizza, C. (1987). Phytochemical investigation on *Mercurialis annua*. *Biochemical Systematics and Ecology*, Vol. 15, No. 6, pp. 667-669, ISSN 0305-1978

- Attard, E. & Attard, H. (2006). The Potential Angiotensin-Converting Enzyme Inhibitory Activity of Oleanolic Acid in the Hydroethanolic Extract of *Crataegus monogyna* Jacq. *Natural Product Communications*, Vol. 1, No. 5, pp. 381-386, ISSN 1934-578X
- Attard, E. & Attard, H. (2008) Antitrypsin activity of extracts from *Ecballium elaterium* seeds. *Fitoterapia*, Vol. 79, No. 3, pp. 226-228, ISSN 0367-326X
- Attard, E. & Vella, K. (2009). The Effects of Ephedrine and *Ephedra fragilis* Crude Extracts on Human Peripheral Lymphocytes. *Pharmacognosy Research*, Vol. 1, No. 2, pp. 38-42, ISSN 0974-8490
- Attard, E., Brincat, M.P. & Cuschieri, A. (2005). Immunomodulatory activity of cucurbitacin E isolated from *Ecballium elaterium*. *Fitoterapia*, Vol. 76, pp. 439-441, ISSN 0367-326X
- Attard, E. (2004). Status of Medicinal and Aromatic Plants in Malta, In: Baricevic, D., Bernath, J., Maggioni, L. and Lipman E. *Report of a Working Group on Medicinal and Aromatic Plants*, First Meeting, Gozd Martuljek, Slovenia, 12-14 September 2002, International Plant Genetic Resources Institute, Rome, Italy, pp. 85 - 87, ISBN 978-92-9043-812-0
- Benavente-García, O., Castillo, J., Lorente, J., Ortuño, A. & Del Rio, J.A. (2000). Antioxidant activity of phenolics extracted from *Olea europaea* L. leaves. *Food Chemistry*, Vol. 68, No. 4, pp. 457-462. ISSN 0308-8146
- Bennett, R.N., Rosa, E.A.S, Mellon, F.A. & Kroon, P.A. (2006). Ontogenic profiling of glucosinolates, flavonoids, and other secondary metabolites in *Eruca sativa* (salad rocket), *Diplotaxis eruroides* (wall rocket), *Diplotaxis tenuifolia* (wild rocket), and *Bunias orientalis* (Turkish rocket). *Journal of Agricultural and Food Chemistry*, Vol. 54, No. 11, pp. 4005-4015, ISSN 0021-8561
- Billeter, M, Meier, B. & Sticher, O. (1991). 8-hydroxyflavonoid glucuronides from *Malva sylvestris*. *Phytochemistry*, Vol. 30, No. 3, pp. 987-990, ISSN 0031-9422
- Bonora, A., Pancaldi, S., Gualandri, R. & Palmira Fasulo, M. (2000). Carotenoid and ultrastructure variations in plastids of *Arum italicum* Miller fruit during maturation and ripening. *Journal of Experimental Botany*, Vol. 51, No. 346, pp. 873-884, ISSN 0022-0957
- Borg, J. (1927). *Descriptive Flora of the Maltese Islands*, Government Printing Office, Malta
- Buhagiar, J.A., Podesta, M.T., Wilson, A.P., Micallef, M.J. & Ali, S. (1999). The induction of apoptosis in human melanoma, breast and ovarian cancer cell lines using an essential oil extract from the conifer *Tetraclinis articulata*. *Anticancer Research*, Vol. 19, No. 6B, pp. 5435-5443, ISSN 0250-7005
- Burres, N.S., Frigo, A., Rasmussen, R.R. & McAlpine, J.B (1992). A colorimetric microassay for the detection of agents that interact with DNA. *Journal of Natural Products*, Vol. 55, No. 11, pp. 1582-1587, ISSN 0163-3864
- Carballo, J.L., Hernandez-Inda, Z.L., Perez, P. & Garcia-Gravalos, M.D. (2002). A comparison between two brine shrimp assays to detect in vitro cytotoxicity in marine natural products. *BMC Biotechnology*, Vol. 2, pp. 17-25, ISSN 1472-6750
- Cassar, P. (1964). *Medical History of Malta*, Wellcome Historical Medical Library, London
- Cassar Pullicino, J. (1947). *An Introduction to the Maltese Folklore*, Malta
- Calderon, A.I., Terreaux, C., Gupta, M.P. & Hostettmann, K. (2003). *In vitro* cytotoxicity of 11 Panamanian plants. *Fitoterapia*, Vol. 74, pp. 378 - 383, ISSN 0367-326X
- Caputo, R., Mangoni, L., Monaco, P. & Previtera L. (1974). New triterpenes from the leaves of *Olea europaea*. *Phytochemistry*, Vol. 13, No. 12, pp. 2825-2827, ISSN 0031-9422


- Cremona, A. (1971). *Hxejjex ta' Medicina Popolari fil-Gzejjer ta' Malta u Ghawdex*, Lehen il-Malti, No. 16
- Darmanin, S. (2003). Investigations of the Anti-Neoplastic Activity of a Bioactive Extract derived from *Ricinus communis*. University of Malta. Faculty of Medicine and Surgery. Department of Anatomy. (Unpublished)
- Delhaye, S. & Landry, J. (1992). Determination of tryptophan in pure proteins and plant material by three methods. *Analyst*, Vol. 117, pp. 1875 - 1877, ISSN 0003-2654
- Deshpande, S.S., Cheryan, M. & Salunkhe, D.K. (1986). Tannin analysis of food products. *CRC Critical Reviews in Food Science & Nutrition*, Vol. 24, pp. 401-449, ISSN 0099-0248
- Doerk-Schmitz, K., Witte, L. & Alfermann, A.W. (1993). Tropane alkaloid patterns in plants and hairy roots of *Hyoscyamus albus*. *Phytochemistry*, Vol. 35, No. 1, pp. 107-110, ISSN 0031-9422
- Edeoga, H.O., Okwu, D.E. & Mbaebie, B.O. (2005). Phytochemical constituents of some Nigerian medicinal Plants. *African Journal of Biotechnology*, Vol. 4, pp. 685-688, ISSN 1684-5315
- El Hajaji, H., Lachkar, N., Alaoui Cherrah, Y., Farah, A., Ennabili, A., El Bali, B. & Lachkar, M. (2011). Antioxidant activity, phytochemical screening, and total phenolic content of extracts from three genders of carob tree barks growing in Morocco. *Arabian Journal of Chemistry*, Vol. 4, No. 3, pp. 321-324, ISSN 1878-5352
- Elamrani, A. (2011). The Antitumoral Activity and the Cytotoxicity on Renal Cells of Ethanolic Extracts from the Leaves of Four Varieties of *Olea europaea* L. Grown in Morocco. *Analytical Chemistry Letters*, Vol. 1, No. 1, pp. 63 - 69, ISSN 2229-7928
- Erdoğan, T.F. (2009). Brine Shrimp Lethality Bioassay of *Fumaria densiflora* Dc. and *Fumaria officinalis* L. Extracts. *Hacettepe University Journal of the Faculty of Pharmacy*, Vol. 28, No. 2, pp. 125-132, ISSN 1300-0608
- Evans, W.C. (2009). *Trease and Evans' Pharmacognosy* (16<sup>th</sup> Edition), Saunders, ISBN 978-0-7020-2933-2, London.
- Franssen, F.F.J., Sweijsters, L.L.J., Berger, I. & Medinilla Aldana, B.E. (1997). *In vivo* and *In vitro* Antiplasmodial Activities of some plants traditionally used in Guatemala against Malaria. *Antimicrobial Agents and Chemotherapy*, Vol. 41, No. 7, pp. 1500-1503, ISSN 0066-4804
- Gleeson, P.A. & Clarke, A.E. (1979). Structural studies on the major component of *Gladiolus* style mucilage, an arabinogalactan-protein. *Biochemical Journal*, Vol. 181, No. 3, pp. 607-621, ISSN 0264-6021
- Goda, F.E. & Badria, F.A. (2005). Synthesis and Biological Evaluation of Certain New Substituted Pyrido[2,3-D]Pyrimidin-4(1h)-One and Pyrido [2,3-D]Triazolo[3,4-B]Pyrimidine Analogs. *Saudi Pharmaceutical Journal*, Vol. 13, No. 2-3, pp. 65-73, ISSN 1319-0164
- Gudej, J. and Czapski, P. (2009). Components of the petroleum ether and chloroform extracts of *Chrysosplenium alternifolium*. *Chemistry of Natural Compounds*. Vol. 45, No. 5, pp. 717-719. ISSN 0009-3130
- Guarrera, P.M. (2003). Food medicine and minor nourishment in the folk traditions of Central Italy (Marche, Abruzzo and Latium), *Fitoterapia*, Vol. 74, pp. 515-544. ISSN 0367-326X
- Gulia, G. (1855). *Repertorio Botanico Maltese*, Ex Libris Publii, Malta.



- Harborne, J.B. (1984). *Phytochemical Methods* (Second Edition), Chapman and Hall, ISBN 0-412-25550-2, London and New York.
- Hultin, E. & Torssell, K. (1965). Alkaloid-screening of Swedish plants. *Phytochemistry*, Vol. 4, No. 3, pp. 425-433, ISSN 0031-9422
- Isbilir, S.S., Ozcan, H.M. & Yagar, H. (2008). Some Biochemical Properties of Lipase from Bay Laurel (*Laurus nobilis* L.) Seeds. *Journal of the American Oil Chemists' Society*, pp. 227-233, ISSN 0003-021X
- Jaki, B., Orjala, J., Burgi, H.R. & Sticher, O. (1999). Biological screening of cyanobacteria for antimicrobial and molluscicidal activity, brine shrimp lethality and cytotoxicity. *Pharmaceutical Biology*, Vol. 37, pp. 138-143, ISSN 1388-0209
- Javidnia, K, Miri, R, Najifi, RB & Jagromi, NK (2003). A preliminary study on the Biological Activiy of *Daphne mucronata* royle. *DARU Journal of the School of Pharmacy*, Vol 11, pp. 28-31, ISSN 1560-8115
- Kanegusuku, M., Benassi, J.C., Pedrosa, R.C., Yunes, R.A., Filho, V.C., Maia, A.A., de Sousa, M.M., Delle Monache, F. & Niero, R. (2001). Cytotoxic, Hypoglycemic Activity and Phytochemical Analysis of *Rubus imperialis* (Rosaceae), *Zeitschrift für Naturforschung*, Vol. 57c, pp. 272-276, ISSN 0932-0784
- Kim, E.J., Tian, F. & Woo, M.H. (2000). Asitrocin, (2,4)-cis- and trans-asitrocinones: novel bioactive mono-tetrahydrofuran acetogenins from *Asimina triloba* seeds, *Journal of Natural Products*, Vol. 63, No. 11, pp. 1503-1506, ISSN 0163-3864
- King, G.A., Woollard, D.C., Irving, D.E. & Borst, W.M. (1990). Physiological changes in asparagus spear tips after harvest, *Physiologia Plantarum*, Vol. 80, No. 3, pp. 393-400, ISSN 0031-9317
- Konning, G.H., Agyare, C. & Ennison, B. (2003). Antimicrobial activity of some medicinal plants from Ghana, *Fitoterapia*, Vol. 75, pp. 65-67, ISSN 0367-326X
- Koukoulitsa, E., Skaltsa, H., Karioti, A., Demetzos, C. & Dimas, K. (2002). Bioactive Sesquiterpene Lactones from *Centaurea* Species and their Cytotoxic/Cytostatic Activity Against Human Cell Lines *in vitro*, *Planta Medica*, Vol. 68, No. 7, pp. 649-652, ISSN 0032-0943
- Krey, A.K. & Hahn, F.E. (1975). Studies on the methyl green-DNA complex and its dissociation by drugs. *Biochemistry*, Vol. 14, No. 23, pp. 5061 - 5067, ISSN 0006-2960
- Kumarasamy Y., Fergusson M.E., Nahar L. & Sarker S.D. (2002) Bioactivity of Moschamindole from *Centaurea moschata*. *Pharmaceutical Biology*, Vol. 40, No. 4, pp. 307 - 310, ISSN 1388-0209
- Kurnick B. & Foster M. (1950). Methyl green. III. Reaction with desoxyribonucleic acid: Stoichiometry and behavior of the reaction product, *Journal of General Physiology*, Vol. 34, pp. 147-159, ISSN 0022-1295
- Kurnick, N. (1950) The determination of desoxyribonuclease activity by methyl green: Application to Serum, *Archives of Biochemistry*, Vol. 29, pp. 41-53, ISSN 0570-6963
- Lanfranco, G. (1975). *Duwa u Semm il-Ħxejjex Maltin*. Edizzjoni Klabb Kotba Maltin, Malta.
- Lanfranco, G. (1980). Some Recent Communications on the Folk Medicine of Malta, *L-Imnara*, Vol. 3
- Lanfranco, G. (1993). *Ħxejjex mediċinali u oħrajn fil-Għejjer Maltin*. Media Centre Publications, Malta
- Loukis, A. & Philianos, S. (1980). Phytochemical investigation of *Acanthus mollis* L. *Fitoterapia*, Vol. 51, No. 4, pp. 183-185, ISSN 0367-326X

- Maiza-Benabdesselam, F., Chibane, M., Madani, K., Max, H. & Adach, S. (2007). Determination of isoquinoline alkaloids contents in two Algerian species of *Fumaria* (*Fumaria capreolata* and *Fumaria bastardi*), *African Journal of Biotechnology*, Vol. 6, No. 21, pp. 2487-2492, ISSN 1684-5315
- Mangion Randon, A. & Attard, E (2007). The *in vitro* Immunomodulatory Activity of Oleuropein, a Secoiridoid Glycoside from *Olea europaea* L. *Natural Product Communications*, Vol. 2, No. 5, pp. 515-519, ISSN 1934-578X
- Martin, C., Ellis, N. & Rook F. (2010). Do Transcription Factors Play Special Roles in Adaptive Variation? *Plant Physiology*, Vol. 154, No. 2, pp. 506-511, ISSN 0032-0889
- Martin, M.J., Alarcón de la Lastra, C., Marhuenda, E., Delgado, F. & Torreblanca J. (1988). Anti-ulcerogenicity of the flavonoid fraction from *Dittrichia viscosa* (L.) W. Greuter in rats. *Phytotherapy Research*. Vol. 2, No. 4, pp. 183-186, ISSN 0951-418X
- Meyer, B.N., Ferrigni, N.R., Putnam, J.E., Jacobsen, L.B., Nichols, D.E. & McLaughlin, J.L. (1982). Brine Shrimp: A convenient general bioassay for active plant constituents. *Planta Medica*, Vol. 45, pp. 31-34, ISSN 0032-0943
- Nagegowda, D.A., Gutensohn, M., Wilkerson, C.G. & Dudareva N. (2008). Two nearly identical terpene synthases catalyze the formation of nerolidol and linalool in snapdragon flowers. *The Plant Journal*, Vol. 55, No. 2, pp. 224-239, ISSN 0960-7412
- Nayak, S., Nalabothu, P., Sandiford, S., Bhogadi, V. & Adogwa, A. (2006). Evaluation of wound healing activity of *Allamanda cathartica*. L. and *Laurus nobilis*. L. extracts on rats. *BMC Complementary and Alternative Medicine*, Vol. 6, p. 12, ISSN 1472-6882
- Negi, J.S., Singh, P., Joshi, G.P., Rawat, M.S. & Bisht, V.K. (2010). Chemical constituents of *Asparagus*. *Pharmacognosy Review*, Vol. 4, No. 8, pp. 215-220, ISSN 0973-7847
- Papagiannopoulos, M., Wollseifen, H.R., Mellenthin, A., Haber, B. & Galensa, R. (2004). Identification and Quantification of Polyphenols in Carob Fruits (*Ceratonia siliqua* L.) and Derived Products by HPLC-UV-ESI/MS. *Journal of Agricultural and Food Chemistry*, Vol. 52, No. 12, pp. 3784-3791, ISSN 0021-8561
- Peebles, K.A., Baker, R.K., Kurz, E.U., Schneider, B.J. & Kroll, D.J. (2001). Catalytic inhibition of human DNA topoisomerase II $\alpha$  by hypericin, a naphthodianthrone from St. John's wort (*Hypericum perforatum*). *Biochemical Pharmacology*. Vol. 62, No. 8, pp. 1059-1070. ISSN 0006-2952
- Penza, C. (1969). *Flora Maltija Medicinali*, Progress Press Co. Ltd., Malta
- Punjani, B.L. & Kumar, V. (2003) Ethnomedicinal plants specially used for liver disorders in the Aravalli ranges of Gujarat, India. *Journal of Natural Remedies*, Vol. 3, No. 2, pp. 195-198, ISSN 0972-5547
- Rajendran, R. & Krishnakumar, E. (2010). Hypolipidemic Activity of Chloroform Extract of *Mimosa pudica* Leaves. *Avicenna Journal of Medical Biotechnology*. Vol. 2, No. 4, pp. 215-221, ISSN 2008-4625
- Rehecho, S., Hidalgo, O., García-Iñiguez de Cirano, M., Navarro, I., Astiasarán, I., Ansorena, D., Cavero, R.Y. & Calvo, M.I. (2011). Chemical composition, mineral content and antioxidant activity of *Verbena officinalis* L. *LWT- Food Science and Technology*, Vol. 44, No. 4, pp. 875-882, ISSN 0023-6438
- Rodriguez-Lopez, V, Salazar, L & Estrada, S. (2003). Spasmolytic activity of several extracts obtained from some Mexican medicinal plants. *Fitoterapia*, Vol. 74, pp. 725-728, ISSN 0367-326X

- Rossi, B. & Attard, E. (2011). The Haemagglutination potential of extracts from *Urtica dubia* plant parts. *Journal of Natural Remedies*, Vol. 11, No. 1, pp. 76-78, ISSN 0972-5547
- Sabeh, F., Wright, T. & Norton, S.J. (1993). Purification and characterization of a glutathione peroxidase from the *Aloe vera* plant. *Enzyme & Protein*, Vol. 47, No. 2, pp. 92-98, ISSN 1019-6773
- Shamsa, F., Monsef, H., Ghamooshi, R. & Verdian-rizi, M. (2008). Spectrophotometric determination of total alkaloids in some Iranian medicinal plants. *Thai J. Pharm. Sci.* Vol. 32, pp. 17-20, ISSN 0125-4685
- Shao, Y., Chin, C.K., Ho, C.T., Ma, W., Garrison, S.A. & Huang, M.T. (1996). Anti-tumor activity of the crude saponins obtained from asparagus. *Cancer Letters*, Vol. 104, No. 1, pp. 31-36, ISSN 0304-3835
- Soušek, J., Guédon, D., Adam, T., Bochořáková H., Táborská E., Válka I. & Šimánek V. (1999). Alkaloids and organic acids content of eight *Fumaria* species. *Phytochemical Analysis*. Vol. 10, No. 1, pp. 6-11, ISSN 0958-0344
- Steinberg, D.M., Sokoll, L.J., Bowles, K.C., Nichols, J.H., Roberts, R., Schultheis, S.K. & O'Donnell, C.M. (1997). Clinical evaluation of Toxi-Prep™: A semiautomated solid-phase extraction system for screening of drugs in urine. *Clinical Chemistry*, Vol. 43, pp. 2099-2105, ISSN 0009-9147
- Sun, T., Powers, J.R. & Tang, J. (2007). Evaluation of the antioxidant activity of asparagus, broccoli and their juices. *Food Chemistry*. Vol. 105, No. 1, pp. 101-106, ISSN 0308-8146
- Tawara, J.N., Blokhin, A., Foderaro, T. A., Stermitz, F. R. & Hope, H. (1993). Toxic piperidine alkaloids from pine (*Pinus*) and spruce (*Picea*) trees. New structures and a biosynthetic hypothesis. *Journal of Organic Chemistry*, Vol. 58, No. 18, pp. 4813-4818, ISSN 0022-3263
- Vaya, J. & Mahmood, S. (2006). Flavonoid content in leaf extracts of the fig (*Ficus carica* L.), carob (*Ceratonia siliqua* L.) and pistachio (*Pistacia lentiscus* L.). *BioFactors*, Vol. 28, No. 3-4, pp. 169-175, ISSN 0951-6433
- Waller, G.R., Mangiafico, S. & Ritchey, C.R. (1978). A Chemical Investigation of *Aloe barbadensis* Miller. *Proceedings of the Oklahoma Academy of Science*, Vol. 58, pp. 69-76, ISSN: 0078-4303.
- Williams, R.D. & Ellis, B.E. (1989). Age and tissue distribution of alkaloids in *Papaver somniferum*. *Phytochemistry*, Vol. 28, No. 8, pp. 2085-2088, ISSN 0031-9422
- Zidorn, C. & Stuppner, H. (2001). Evaluation of chemosystematic characters in the genus *Leontodon* (Asteraceae). *Taxon*, Vol. 50, pp. 115-133, ISSN 0040-0262



## **Bioactive Compounds in Phytomedicine**

Edited by Prof. Iraj Rasooli

ISBN 978-953-307-805-2

Hard cover, 218 pages

**Publisher** InTech

**Published online** 18, January, 2012

**Published in print edition** January, 2012

There are significant concerns regarding the potential side effects from the chronic use of conventional drugs such as corticosteroids, especially in children. Herbal therapy is less expensive, more readily available, and increasingly becoming common practice all over the world. Such practices have both their benefits and risks. However, herbal self-therapy might have serious health consequences due to incorrect self-diagnosis, inappropriate choice of herbal remedy or adulterated herbal product. In addition, absence of clinical trials and other traditional safety mechanisms before medicines are introduced to the wider market results in questionable safe dosage ranges which may produce adverse and unexpected outcomes. Therefore, the use of herbal remedies requires sufficient knowledge about the efficacy, safety and proper use of such products. Hence, it is necessary to have baseline data regarding the use of herbal remedies and to educate future health professionals about various aspects of herbal remedies.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Everaldo Attard and Pierpaolo Pacioni (2012). The Phytochemical and In Vitro Pharmacological Testing of Maltese Medicinal Plants, *Bioactive Compounds in Phytomedicine*, Prof. Iraj Rasooli (Ed.), ISBN: 978-953-307-805-2, InTech, Available from: <http://www.intechopen.com/books/bioactive-compounds-in-phytomedicine/the-phytochemical-and-in-vitro-pharmacological-testing-of-maltese-medicinal-plants>

**INTECH**  
open science | open minds

#### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

#### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen