

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Demographic and Epidemiologic Features of Acute Appendicitis

Barlas Sulu

*Kafkas University Faculty of Medicine,
Department of General Surgery, Kars,
Turkey*

1. Introduction

Acute Appendicitis (AA) is the most commonly encountered disease in emergency clinics, with about 250,000 cases of appendicitis reported in The United States and 40,000 in England each year (Deng et al., 2010; Simpson et al., 2008). The key form of treatment for the disease is surgery and the mortality, morbidity, and economic cost ratio rise the longer it remains untreated.

Despite the high prevalence of the disease, the uncertainty of just how many appendicitis patients will present to emergency clinics each day still remains an unpredictable situation for us surgeons, with some days bringing a large number of cases, and others relatively few. Ever since Amyand Claudius performed the first appendectomy in 1735 at St George's Hospital in London, this decrease and increase in cases has led researchers to conduct studies into both the etiology of AA and the disease's epidemiological and demographic characteristics.

The etiology of AA has, as of yet, not been clarified. Clarification of the etiology has been the main point of epidemiological studies and advancements in this area will influence the incidence of the disorder as well. There are various theories on the frequency and incidence of appendicitis at present and these are still debated, despite some having strong scientific backing.

2. Demographic features of appendicitis

Studies have demonstrated that AA is seen most commonly in western societies, particularly in youths and males (Addiss et al., 1990; Al-Omran et al., 2003; Noudeh et al., 2007; Sulu et al., 2010). Research carried out in our own region has also shown that AA is seen more commonly in young people aged 10-19 and in males (**Figure 1**).

The appendix tissue possesses the features of a lymphoid organ and there is a larger amount of lymphoid tissue in young subjects. Lymphoid hyperplasia can be caused by any obstruction occurring in the lumen of the appendix and this can develop into appendicitis if the condition continues. Appendicitis is therefore seen more frequently in young people. Lymphoid tissue is vague. Lumen of the appendix enlarges after lymphoid tissue atrophies and probability of obstruction by decreases fall of in over aged people. For that reason, incidence of acute appendicitis decreases with age. Consequently only 5% to 10% of

acute appendicitis cases are seen in the elderly (Jones et al., 1985). The more common involvement of young people and men is easier to discern in Western societies than other regions. The lower incidence of AA and the less pronounced gender gap in regions such as Africa and Asia is worth mentioning (Ajao, 1979; Arnbjörnsson, 1983; Oguntola et al., 2010; Walker & Segal, 1995). This may be because people living in these regions are less influenced by the western-type diet (fast-food diet) with the majority of foods consumed being high in carbohydrates and low in fiber. The height of males as a factor in the development of the disease is not clear (Addiss et al., 1990). However, the effect of sex hormones in females alongside the predisposition of males to consume 'fast-food' are considered to be important (Walker & Segal, 1979).




Fig. 1. The distribution of acute appendicitis according to age groups and sex (2004-2007).

3. Epidemiological theories for the development of appendicitis

The most widely accepted theories today are the diet and hygiene hypotheses (Walker & Segal, 1995). At the same time, these hypotheses have also formed the groundwork of AA's epidemiological features.

The dietary theory was first developed by Rendle Short in 1920. He stated that the AA incidence was higher with a lower ratio of cellulose in the diet and that this was the reason why Britain had been seeing an increase in the incidence of AA since the turn of the 20th century, as well as why the rate differed by country (Barker, 1985; Walker & Segal, 1995). This theory was further developed to reveal a positive correlation between AA and a diet poor in fiber but rich in such foods as meat, potatoes, and sugar, and a negative correlation between AA and a diet rich in fiber containing green vegetables, fruits, and tomatoes (Morris et al., 1987). In other words, a diet devoid of sufficient fiber triggers the formation of AA. A diet featuring low consumption of fiber reduces the colonic transit time by reducing

the lumen of the appendix, and the resulting lymphoid hyperplasia causes obstruction, ultimately leading to infection and appendicitis.

The hygiene theory was promoted by Barker in articles in the 1980s (Barker, 1985; Barker et al., 1988). Barker felt that the diet did not explain the decrease in incidence of appendicitis in England in the 1930's. He found that the appendicitis incidence had decreased, together with that of many diseases, such as tuberculosis, in relation to the introduction of better housing and water supply following World War II. He felt that these improvements had decreased exposure of young children to enteric microorganisms and that the number of deaths due to diarrhea in childhood had been somewhat reduced. As a result, children and adults now avoided the effect of bacteria and viruses causing enteric and respiratory infections which pave the way for appendicitis and instead developed immunity against them. An insufficient number of bathrooms, toilets, hot water, and sewage systems in communal areas can promote the enteric bacterial, viral, and parasitic infectious agents that are responsible for the formation of appendicitis. These improvements will decrease the incidence of deaths due to enteric organisms in children and adults and also decrease the risk of appendicitis. This outcome is strongly related with the incidence of infectious disease and socioeconomic status of the patients with appendicitis. For example, Schistosomiasis is a waterborne parasitic disease that can be prevented by following principles of good hygiene and preventing, by the management of sewage networks, the spread of parasite eggs in water that is used for consumption. As a result, the incidence of Schistosomal appendicitis is 0.2% in the USA, where sewage networks are well maintained, while it is 20 times this rate in Nigeria (Terada, 2009). Interestingly, the patients who suffered from this illness in the USA were African Americans.

4. Factors influencing appendicitis incidence and current tendencies

Epidemiological and demographic studies report the appendicitis incidence to vary according to age, gender, race, socioeconomic status, food culture, and seasonal changes (Adamidis et al., 2000; Ergul, 2007; Noudeh et al., 2007; Oguntola et al., 2010; Sulu et al., 2010). Therefore, the frequency of AA is different in each country. For example, the rate of appendicitis in Europe during the 1980s was 116 per 100,000 while this rate was 96-120 in the USA, 75 in Ontario, 200 in Hong Kong, and 32-37 in Thailand for the same years (Al-Omran et al., 2003; Chatbanchai et al., 1989; Luckmann & Davis, 1991; Zoguéréh et al., 2001).

Epidemiological studies report that the incidence of AA within a single country tends to increase or decrease at different times of the year. In fact, differences have even been reported between people living in the same country when the society is formed from individuals from various geographical locations, cultures, and races. Walker et al. reported the prevalence of appendicitis as 0.5% in rural blacks, 1.2% in urban blacks, and 14% in urban whites (Walker & Segal, 1979). Walker et al. also evaluated the relationship between ethnicity and appendicitis in a study of 56 high school age (16-18 years) young people in South Africa and found that the rate of appendectomies was 0.6% in rural Blacks, 0.7% in urban Blacks, 2.9% in Indians, 1.7% in Coloureds (Eur-African-Malay), and 10.5% in Caucasians (Walker et al., 1982). This situation is similar within different ethnic communities in western societies, where the gap between gender and ethnic origins has shown similar distributions. In California, the incidence of appendicitis was 137.5 per 100,000 for Caucasian males while this incidence was 162.7 for Hispanics, 98.0 for Asian/others, and 70.7 for blacks. The same was true in female patients (Andersson, 2008)

with rates per 100,000 of 98.8, 97.5, 64.6, and 49.6 for the above groups respectively. The authors reported that the difference observed between whites and blacks was associated with their consumption of different amounts of fiber.

However, the fact that studies performed in the last decade have reported a decrease in the incidence of AA in western countries (Andersson, 2008; Andersen et al., 2009; Andreu-Ballester et al., 2009; Cirocchi et al., 2008) but an increase in some African and Asian countries is interesting (Chavda et al., 2005; Lee et al., 2010). The rate of patients undergoing an appendectomy has decreased in the UK, the USA, and Europe (Williams et al., 1998). Even though there has been no change in the amount of fiber consumed in food in England and Wales during the 20st century, the mortality rate from appendicitis, which was 40 to 70 per million in the first decade, dropped to 5 per million in the last decade (Barker, 1985). A study on Danish children has found a decreased appendicitis incidence in children from every age group (Andersen et al., 2009). The effect of better socioeconomic conditions created as a result of improving water supplies and hygienic conditions, has been found to be the reason for this decrease in Western societies. A recent study from Spain found a decrease in appendicitis in the last 10 years (Andreu-Ballester et al., 2009). A study from Greece evaluating the last 30 years found the age-standardized appendicitis rate to fall 75% from 652/100.000 to 164/100.000 (Papadopoulos et al., 2008). Both countries eat a Mediterranean diet rich in fiber but the decrease in Spain may be due to improved hygienic conditions with socioeconomic development. It is believed that the spread of the western-type diet in African and Asian societies today is responsible for the increase in the number of AA cases observed. However, we found in a previous study that short-term dietary changes had no effect on an increase in AA incidence (Papadopoulos et al., 2008). Studies conducted on the effects of fasting seem to support this conclusion. More than one billion Muslims consider the month of 'Ramadan' to be sacred and abstain from food or drink from sunrise to sunset for one month. During this month, devotees avoid performing any daily habitual action, such as the consumption of any food, drinks, drug therapy, smoking, or having sexual intercourse for a total of 12-19 hours each day. Since the body is used to receiving two meals a day, these changes in diet and the number of meals consumed each day change the metabolism (Sulu et al., 2010). This behavior has provided an ideal model for many researchers to investigate the effect that long-term hunger has on the human body. We have also used this opportunity to investigate the effect of changes in hunger and dietary habits on the development of AA in a study conducted throughout the month of Ramadan. We compared the frequency of AA before and after Ramadan in a 4-year study that included 4288 AA cases in two different cities. Of the 992 patients investigated, 37.1% developed AA in the period before Ramadan, 32.1% during the month of Ramadan, and 30.8% were diagnosed after the month of Ramadan. There was no significant difference between the values of these three periods. Based on these results, it is possible to infer that there is no short-term increase in the frequency of AA in societies starting to follow a western-type diet nor is there expected to be an increase in cases where migrants have moved to a new geographical location.

One further subject of discussion is whether environmental conditions have an influence on the frequency of the development of AA. It has been reported that seasonal changes, in particular, and moisture in the air (humidity) do have an effect (Al-Omran et al., 2003; Gallerani et al., 2006; Lee et al., 2010; Noudeh et al., 2007; Oguntola et al., 2010; Trepanowski & Bloomer, 2010). An increase in the frequency of cases in the AA summer months has been reported in many studies. However, studies have shown that this increase is observed in regions situated in low altitudes and close to the seaside (**Table 1**).

Altitude (meters)		Appendicitis	Perforated
Ferrara	9	Summer	
Jersey City	25	Summer	
Ontario	86	Summer	
Beer Sheva	260	Autumn-Summer	
Hail	988	Spring-Summer	
Shahr-e-Rey	1050	Summer	
Kirman	1749	Winter	
İstanbul	100	Spring-Summer	Summer-Autumn
Kars	1750	Winter	

Table 1. The seasonal tendency of total and perforated appendicitis according to the altitude of different regions

In a study conducted in two Turkish cities with different climatic characteristics and altitudes, we found that the number of patients with acute appendicitis increases at low altitude in Istanbul during spring and summer ($p < 0.05$). At high altitude in Kars, this increase is seen during winter ($p > 0.05$). (Table 2).

Hospital	Season				Age Groups					
	1*	2*	3*	4*	0-9	10-19	20-29	30-39	40-49	≥50
KSH	503	476	458	434	300	814	349	204	113	91
HNTEH	712	846	762	686	22	664	1133	606	289	292

KSH : Kars State Hospital, HNTEH: Haydarpasa Numune Teaching and Research Hospital 1*: Winter, 2*: Spring, 3*: Summer, 4*: Autumn

Table 2. A comparison of the seasonal and age distribution of patients with appendicitis in Kars and Istanbul (2004-2007)

A further study reported that AA was seen more frequently in the winter months in Kerman, an Iranian city with an altitude similar to that of Kars (Nabipour & Mohammad, 2005; Sulu et al., 2010). In other words, an increase in altitude resulted in more appendicitis cases being seen during the winter months. The reason for this trend is unclear, but it has been reported that several factors may play a part: 1) the varying effects of bacterial or viral pathogens that cause infections at different temperatures, 2) the effect of allergens occurring in summer and warmer months, 3) changes in the form of nutrition, and 4) the effect of migration for touristic purposes during the summer. Another controversial environmental

factor is daytime humidity. In a study by Brummer, a significant relationship was observed between humidity and AA, and it was reported, in their study on the physiology of hunger, that a decrease in body fluid loss, fecal stasis, and fecal dehydration prepared the ground for inflammation (Brumer, 1970). In contrast, van Nieuwenhoven et al. reported that changes in intestinal system functions such as intestinal permeability and orocaecal transit time were not the reason for dehydration occurring (van Nieuwenhoven et al., 2000). We also determined, during a 4-year (2004-2007) study using meteorological data in our region, that such data as moisture and amount of rain did not have an effect on the frequency of AA (Table 3).

	Appendicitis	Perforated	z value	p value
Humidity (%)				
Winter	76.7±7.95 (54.0 – 94.0)	76.7±8.01 (60.7 – 94.0)	-0.054	0.957
Spring	68.2±10.24 (30.0 – 93.3)	67.7±10.77 (40.0 – 89.3)	-0.047	0.963
Summer	63.9±8.75 (35.0 – 87.8)	64.7±10.33 (36.0 – 87.8)	-1.191	0.234
Autumn	71.9±9.60 (47.3 – 94.0)	71.7±9.91 (48.7 – 93.3)	-0.148	0.882
General	70.4±10.32 (30.0 – 94.0)	69.6±10.75 (36.0 – 94.0)	-0.548	0.584
Temperature (°C)				
Winter	-5.8±6.41 (-21.1 – 8.5)	-6.5±7.19 (-19.7 – 7.2)	-0.751	0.453
Spring	10.7±5.14 (-4.6 – 19.7)	9.6±5.63 (-2.8 – 18.5)	-1.209	0.227
Summer	17.2±3.34 (8.7 – 28.6)	16.6±3.38 (6.2 – 23.3)	-1.031	0.302
Autumn	0.2±7.90 (-21.2 – 14.4)	0.8±9.48 (-21.2 – 14.4)	-0.912	0.362
General	5.3±10.81 (-21.2 – 28.6)	6.2±10.77 (-21.2 – 23.3)	-1.193	0.233

Table 3. Seasons, Humidity, and Temperature Distribution Levels by Group

Other factors said to influence AA development include vascular disorders, non-specific viral infections, depression and emotional problems due to a stressful lifestyle, being the child of a mother who smoked while pregnant, air pollution, and anemic diseases these have not, however, been widely accepted (Ahmed et al., 2005; Butland & Strachan, 1999; Ewald et al., 2001; Kaplan et al., 2009; Walker & Segal, 1995). As with the development of many diseases, the effect of genetics on the development of AA is unknown. In a survey of 282 patients, it was discovered that 21% of patients undergoing appendectomies had first-degree relatives (siblings, parents, and children), 12% had second-degree relatives (grandparents, grandchildren, uncles, aunts, nieces, and nephews), and 7% had third-degree relatives with a history of appendicitis (Basta et al., 1990). However, more research to reveal the transitional property of genetics in many bowel diseases is clearly needed.

5. Conclusion

As I have discussed briefly above, the development of AA is multifactorial. Many of these reasons are not clear and require further discussion (Figure 2).

The most accepted are diet and hygiene. Appendicitis is considered a preventable disease due to the effect of factors such as diet and hygiene on its development. The morbidities that can result from this disease, as well as mortality rate, may therefore be reduced by improving the socioeconomic status of poorer communities, as well as by the members of

these communities modifying their dietary habits. The cost per patient for the surgical treatment of appendicitis in the United States ranges from \$ 11,577 to \$ 13,965 (Long et al., 2001). Therefore, a reduction in appendicitis will benefit not only public health but will also make a substantial contribution to the economy.




Fig. 2. Factors affecting the formation of appendicitis

6. References


- Adamidis, D., Roma-Giannikou, E., Karamolegou, K., Tselalidou, E., & Constantopoulos, A. (2000). Fiber intake and childhood appendicitis. *Int J Food Sci Nutr*, Vol.51, No.3, pp.153-157, ISSN: 0963-7486
- Addiss, DG., Shaffer, N., Fowler, BS., & Tauxe, RV. (1990). The epidemiology of appendicitis and appendectomy in the United States. *Am J Epidemiol*, Vol.132, No.5, pp.910-925, ISSN: 0002-9262
- Ahmed, S., Shahid, RK., & Russo, LA. (2005). Unusual causes of abdominal pain: sickle cell anemia. *Best Pract Res Clin Gastroenterol*, Vol.19, No.2, pp.297-310, ISSN: 1521-691
- Ajao, OG. (1979). Appendicitis in a tropical African population. *J Natl Med Assoc*, Vol.71, No.10, pp.997-999, ISSN: 0027-9684.
- Al-Omran, M., Mamdani, M., & McLeod, RS. (2003). Epidemiologic features of acute appendicitis in Ontario, Canada. *Can J Surg*, Vol.46, No.4, pp.263-268, ISSN: 0008-428X
- Andersson, RE. (2008). Changing trends in surgery for acute appendicitis (*Br J Surg* 2008; 95: 363-368). *Br J Surg*. Vol.95, No.9, pp.1187-1188, ISSN: 0007-1323

- Andersen, SB., Paerregaard, A., & Larsen, K. (2009). Changes in the epidemiology of acute appendicitis and appendectomy in Danish children 1996-2004. *Eur J Pediatr Surg*, Vol.19, No.5, pp.286-289, ISSN: 0939-7248
- Andreu-Ballester, JC., González-Sánchez, A., Ballester, F., Almela-Quilis, A., Cano-Cano, MJ., Millan-Scheiding, M., & del Castillo, JR. (2009). Epidemiology of appendectomy and appendicitis in the Valencian community (Spain), 1998-2007. *Dig Surg*, Vol.26, No.5, pp.406-412, ISSN: 0253-4886.
- Arnbjörnsson, E. (1983). Acute appendicitis risk in various phases of the menstrual cycle. *Acta Chir Scand*, Vol.149, No.6, pp.603-605, ISSN: 0001-5482
- Barker, DJ. (1985). Acute appendicitis and dietary fibre: an alternative hypothesis. *Br Med J (Clin Res Ed)*, Vol. 290, No.6475, pp.1125-1127, ISSN: 0267-0623
- Barker, D.J, Morris, JA., Simmonds, SJ., & Oliver, RH. (1988). Appendicitis epidemic following introduction of piped water to Anglesey. *J Epidemiol Community Health*, Vol.42, No.2, pp.144-148, ISSN: 0143-005X
- Basta, M., Morton, NE., Mulvihill, JJ., Radovanović, Z., Radojčić, C., & Marinković, D. (1990). Inheritance of acute appendicitis: familial aggregation and evidence of polygenic transmission. *Am J Hum Genet*, Vol.46, No.2, pp.377-382, ISSN: 0002-9297
- Brumer, M. (1970). Appendicitis. Seasonal incidence and postoperative wound infection. *Br J Surg*, Vol.57, No.2, pp.93-99, ISSN: 0007-1323
- Butland, BK., & Strachan, DP. (1999). Smoking and acute appendicitis. *Lancet*, Vol.353, No.9165, pp.1712, ISSN: 0140-6736
- Chatbanchai, W., Hedley, AJ., Ebrahim, SB., Areemit, S., Hoskyns, EW., & de Dombal, FT. (1989). Acute abdominal pain and appendicitis in north east Thailand. *Paediatr Perinat Epidemiol*, Vol.3, No.4, pp.448-459, ISSN: 0269-5022
- Chavda, SK., Hassan, S., & Magoha, GA. (2005). Appendicitis at Kenyatta National Hospital, Nairobi. *East Afr Med J*, Vol.82, No.10, pp.526-530, ISSN: 0012-835X
- Cirocchi, R., Morelli, U., La Mura, F., Cattorini, L., Napolitano, V., Galanov, I., Covarelli, P., Giustozzi, G., & Sciannameo, F. (2008). Acute appendicitis: a descending trend? *Minerva Chir*, Vol.63, No.2, pp.109-113. ISSN: 0026-4733
- Deng, Y., Chang, DC., Zhang, Y., Webb, J., Gabre-Kidan, A., & Abdullah, F. (2010). Seasonal and day of the week variations of perforated appendicitis in US children. *Pediatr Surg Int*, Vol.26, No.7, pp.691-696, ISSN: 0179-0358
- Ergul, E. (2007). Heredity and familial tendency of acute appendicitis. *Scand J Surg*, Vol.96, No.4, pp.290-292, ISSN: 1457-4969
- Ewald, H., Mortensen, PB., & Mors, O. (2001). Decreased risk of acute appendicitis in patients with schizophrenia or manic-depressive psychosis. *Schizophr Res*, Vol.49, No.3, pp.287-293, ISSN: 0920-9964
- Freud, E., Pilpel, D., & Mares, AJ. (1988). Acute appendicitis in childhood in the Negev region: some epidemiological observations over an 11-year period (1973-1983). *J Pediatr Gastroenterol Nutr*, Vol.7, No.5, pp.680-684. ISSN: 0277-2116
- Gallerani, M., Boari, B., Anania, G., Cavallesco, G., & Manfredini, R. (2006). Seasonal variation in onset of acute appendicitis. *Clin Ter*. Vol.157, No.2, pp.123-127. ISSN: 0009-9074
- Jones, BA., Demetriades, D., Segal, I., & Burkitt, DP. (1985). The prevalence of appendiceal fecaliths in patients with and without appendicitis. A comparative study from Canada and South Africa. *Ann Surg*. Vol.202, No.1, pp.80-82. ISSN: 0003-4932

- Kaplan, GG., Dixon, E., Panaccione, R., Fong, A., Chen, L., Szyszkowicz, M., Wheeler, A., MacLean, A., Buie, WD., Leung, T., Heitman, SJ., & Villeneuve, PJ. (2009). Effect of ambient air pollution on the incidence of appendicitis. *CMAJ*, Vol.181, No.9, pp.591-597, ISSN: 0008-4409
- Lee, JH., Park, YS., & Choi, JS. (2010). The epidemiology of appendicitis and appendectomy in South Korea: national registry data. *J Epidemiol*, Vol.20, No.2, pp.97-105, ISSN: 0917-5040
- Long, KH., Bannon, MP., Zietlow, SP., Helgeson, ER., Harmsen, WS., Smith, CD., Ilstrup, DM., Baerga-Varela, Y., & Sarr, MG. (2001). Laparoscopic Appendectomy Interest Group. A prospective randomized comparison of laparoscopic appendectomy with open appendectomy: Clinical and economic analyses. *Surgery*, Vol.129, No.4, pp.390-400. ISSN: 0263-9319
- Luckmann, R., & Davis, P. (1991). The epidemiology of acute appendicitis in California: racial, gender, and seasonal variation. *Epidemiology*, Vol.2, No.5, pp.323-330, ISSN: 1044-3983
- Morris, J., Barker, D.J, & Nelson, M. (1987). Diet, infection, and acute appendicitis in Britain and Ireland. *J Epidemiol Community Health*, Vol.41, No.1, pp.44-49, ISSN: 0143-005X
- Nabipour, F., & Mohammad, BD. (2005). Histopathological features of acute appendicitis in Kerman-Iran from 1997 to 2003. *American J. Environmental Sci*, Vol.1, No.2, pp.130-132, ISSN: 1553-345X
- Noudeh, YJ., Sadigh, N., & Ahmadnia, AY. (2007). Epidemiologic features, seasonal variations and false positive rate of acute appendicitis in Shahr-e-Rey, Tehran. *Int J Surg*, Vol.5, No.2, pp.95-98, ISSN: 1743-9191
- Oguntola, AS., Adeoti, ML., & Oyemolade, TA. (2010). Appendicitis: Trends in incidence, age, sex, and seasonal variations in South-Western Nigeria. *Ann Afr Med*, Vol. 9, No.4, pp.213-217, ISSN: 1596-3519
- Papadopoulos, AA., Polymeros, D., Kateri, M., Tzathas, C., Koutras, M., & Ladas, SD. (2008). Dramatic decline of acute appendicitis in Greece over 30 years: index of improvement of socioeconomic conditions or diagnostic aids? *Dig Dis*, Vol.26, No.1, pp.80-84. Epub 2008 Feb 15. ISSN: 0257-2753
- Simpson, J., Samaraweera, AP., Sara, RK., & Lobo, DN. (2008). Acute appendicitis--a benign disease? *Ann R Coll Surg Engl*, Vol.90, No.4, pp.313-316, ISSN: 0035-8843
- Sulu, B., Gunerhan, Y., Ozturk, B., & Arslan, H. (2010). Is long-term hunger (Ramadan model) a risk factor for acute appendicitis? *Saudi Med J*, Vol.31, No.1, pp.59-63, ISSN: 0379-5284
- Sulu, B., Günerhan, Y., Palanci, Y., İşler, B., & Çağlayan, K. (2010). Epidemiological and demographic features of appendicitis and influences of several environmental factors. *Ulus Travma Acil Cerrahi Derg*, Vol.16, No.1, pp.38-42, ISSN: 1306-696X
- Terada, T. (2009). Schistosomal appendicitis: incidence in Japan and a case report. *World J Gastroenterol*, Vol.15, No.13, pp.1648-1649. ISSN: 1007-9327
- Trepanowski, JF., & Bloomer, RJ. (2010). The impact of religious fasting on human health. *Nutr J*, Vol.9, No.57, ISSN: 1475-2891
- van Nieuwenhoven, MA., Vriens, BE., Brummer, RJ., & Brouns, F. (2000). Effect of dehydration on gastrointestinal function at rest and during exercise in humans. *Eur J Appl Physiol*, Vol.83, No.6, pp.578-584, ISSN: 1439-6319

- Walker, AR., & Segal, I. (1979). Is appendicitis increasing in South African blacks? *S Afr Med J*, Vol.56, No.13, pp.503-504, ISSN: 0256-9574
- Walker, AR., Walker, BF., Duvenhage, A., Jones, J., Ncongwane, J., & Segal, I. (1982). Appendicectomy prevalences in South African adolescents. *Digestion*, Vol.23, No.4, pp.274-278, ISSN: 0012-2823
- Walker, AR., & Segal, I. (1995). Appendicitis: an African perspective. *J R Soc Med*, Vol. 88, No.11, pp.616-619, ISSN: 0141-0768
- Williams, NM., Jackson, D., Everson, NW., & Johnstone, JM. (1998). Is the incidence of acute appendicitis really falling? *Ann R Coll Surg Engl*, Vol.80, No.2, pp.122-124, ISSN: 0035-8843
- Zoguéréh, DD., Lemaître, X., Ikoli, JF., Delmont, J., Chamlian, A., Mandaba, JL., & Nali, NM. (2001). [Acute appendicitis at the National University Hospital in Bangui, Central African Republic: epidemiologic, clinical, paraclinical and therapeutic aspects]. *Sante*, Vol.11, No.2, pp.117-125, ISSN: 1157-5999

IntechOpen



Appendicitis - A Collection of Essays from Around the World

Edited by Dr. Anthony Lander

ISBN 978-953-307-814-4

Hard cover, 226 pages

Publisher InTech

Published online 11, January, 2012

Published in print edition January, 2012

This book is a collection of essays and papers from around the world, written by surgeons who look after patients of all ages with abdominal pain, many of whom have appendicitis. All general surgeons maintain a fascination with this important condition because it is so common and yet so easy to miss. All surgeons have a view on the literature and any gathering of surgeons embraces a spectrum of opinion on management options. Many aspects of the disease and its presentation and management remain controversial. This book does not answer those controversies, but should prove food for thought. The reflections of these surgeons are presented in many cases with novel data. The chapters encourage us to consider new epidemiological views and explore clinical scoring systems and the literature on imaging. Appendicitis is discussed in patients of all ages and in all manner of presentations.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Barlas Sulu (2012). Demographic and Epidemiologic Features of Acute Appendicitis, *Appendicitis - A Collection of Essays from Around the World*, Dr. Anthony Lander (Ed.), ISBN: 978-953-307-814-4, InTech, Available from: <http://www.intechopen.com/books/appendicitis-a-collection-of-essays-from-around-the-world/demographic-and-epidemiologic-features-of-acute-appendicitis>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen