

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Microbial Weed Control and Microbial Herbicides

Tami L. Stubbs and Ann C. Kennedy

*Washington State University and United States Department of Agriculture –
Agricultural Research Service
United States of America*

1. Introduction

Microbial weed control represents an innovative means to manage troublesome weeds and utilize the naturally occurring biological herbicides produced by soil microorganisms. These compounds kill or hinder the growth of weeds so that beneficial plant species can gain a competitive advantage. The vast diversity of microorganisms in our environment is largely untapped, and the potential discovery and characterization of these microbial compounds represents an opportunity to complement chemical herbicides, or reduce the potential for erosion or soil degradation due to tillage for weed control. Invasive weeds continue to threaten the productivity of agricultural lands and natural areas; however, for many weeds adequate, cost-effective control measures presently are not available (Jones & Sforza, 2007). Discovery of biological controls for invasive plants represents an alternative way to slow the spread of these weeds using natural enemies (Jones & Sforza, 2007). Further advances in microbial genetics will continue to improve our understanding of the wealth of genetic diversity and potential in the soil and to better use plant-microbe interactions. The development of biocontrol agents would lessen the need for chemical herbicides and provide greater options for weed management. Microbes have a place in integrated, ecologically based weed management and their potential is only just being realized.

The concept of utilizing microbial herbicides has been explored for more than a quarter century, but there remain many challenges to overcome before they can be widely used in agricultural, range and forest lands, or waterways. Those challenges include improving the efficacy of the microbial activity, survival of microorganisms, persistence of the suppressive compound, delivery systems, determining host range, and avoiding injury to non-target organisms. Other considerations are interactions with chemical herbicides, regulations, commercialization and mass production, and economic feasibility.

Biological controls for weeds can generally be divided into one of three general types: classical, augmentative or inundative, and cultural. The classical approach involves the introduction of a control agent into an area where it did not previously exist, and where the agent eventually becomes able to sustain itself. An example of this is the release of *Puccinia chondrillina* to control *Chondrilla juncea* (rush skeletonweed; Barton, 2004). Augmentative or inundative biological control refers to repeated application of a foreign agent with the intent to reduce weed densities to a level where beneficial plant species can compete. An example

of an augmentative control is *Colletotrichum gloeosporioides* for control of sicklepod (*Senna obtusifolia* L.; Boyette, 2006). Cultural weed control might include crop rotation, fallow periods, sanitation to prevent the introduction and spread of weed seeds, and maintaining soil fertility to produce healthy crop plants. The successful history of insects in weed control and the ever increasing list of successful insect biological controls placed in the hands of land managers offers hope for microbial weed control. Use of biological herbicides requires a shift in thinking from the use of chemical herbicides, as biological controls will most likely not eliminate a weed problem as quickly or as thoroughly as some herbicides on an annual basis; the goal is to inhibit the weed pest below an economically damaging threshold over a long time period in order for beneficial species to gain a competitive advantage (Ghosheh, 2005). Among the criteria that Charudattan (2005) lists for determining which invasive plants make suitable candidates for microbial herbicides are those that have a number of available pathogens that might be suitable for biological control, and where the cost of using that control will be competitive with other control measures. There are a number of reasons for developing microbial herbicides, and those include the potential for herbicide resistant weeds; chemical herbicides may persist in soil for longer than one growing cycle, thus limiting options for crop rotations; there may be limitations on herbicide registrations for certain crops; there is the potential for fewer undesirable effects to the environment than from chemical herbicides; and finally there is the potential for injury to non-target organisms (Guske et al., 2004). Special considerations are needed for management of parasitic weeds (Sauerborn et al., 2007). Herbicide use to control parasitic weeds is difficult, because the greatest damage to the host plant has often occurred before the parasitic weed emerges above the soil surface. Another challenge is that parasitic weeds are closely related to their hosts, and selective control with herbicides is difficult (Sauerborn et al., 2007).

The various regions of the world may be plagued by specific invasive weed problems, while other weeds are problematic worldwide. No matter whether they are worldwide problems or regional challenges, many are elusive to management efforts. These weeds vary by region and ecosystem, and several microorganisms have been studied or are under development in greenhouse or field studies as potential sources for microbial herbicides (Table 1A and B).

Weed Pest	Ecosystem	Region	Biocontrol agent	Reference
<i>Abutilon theophrasti</i> (Velvetleaf)	Croplands	U.S., Canada	<i>Fusarium oxysporum</i> ; <i>Colletotrichum</i> <i>coccodes</i> ; <i>Pseudomonas</i> <i>putida</i> & <i>Acidovorax</i> <i>delafieldii</i>	Kremer & Schulte, 1989; Wymore & Watson, 1989; Owen & Zdor, 2001
<i>Ailanthus altissima</i> (Tree- of-heaven)	Native forests, urban areas	North America, Europe	<i>Aecidium ailanthi</i> J.Y. Zhuang sp. nov.; <i>Coleosporium</i> sp.; <i>Fusarium oxysporum</i> f. sp. <i>perniciosum</i> ; <i>Verticillium albo-atrum</i>	Review by Ding et al., 2006; Schall & Davis, 2009
<i>Alternanthera</i> <i>philoxeroides</i> (Alligatorweed)	Aquatic, upland sites	Worldwide	<i>Nimbya alternantherae</i>	Pomella et al., 2007
<i>Amaranthus</i> spp. (Amaranthus)	Croplands	Europe, North America	<i>Phomopsis</i> <i>amaranthicola</i>	Roskopf et al., 2006
<i>Amaranthus retroflexus</i> (Redroot pigweed)	Croplands	Europe, North America	<i>Alternaria alternata</i>	Lawrie et al., 2002a

Weed Pest	Ecosystem	Region	Biocontrol agent	Reference
<i>Amaranthus hybridus</i> (Pigweed), <i>Senna obtusifolia</i> (Sicklepod), <i>Crotalaria spectabilis</i> (Showy crotalaria)	Croplands	North America	<i>Phomopsis amaranthicola</i> , <i>Alternaria cassia</i> , <i>Colletotrichum dematium</i> , <i>Fusarium udum</i>	Chandramohan & Charudattan, 2003
<i>Arceuthobium tsugense</i> (Hemlock dwarf mistletoe)	Coniferous forests	Vancouver Island, Canada	<i>Neonectria neomacrospora</i>	Rietman et al., 2005
<i>Brunnichia ovata</i> (Redvine), <i>Campis radicans</i> (Trumpet creeper), <i>Pueraria lobata</i> (Kudzu)	Croplands, wastelands, natural areas	Southern U.S.	<i>Myrothecium verrucaria</i>	Boyette et al., 2006, 2008a
<i>Cannabis sativa</i> (Ditchweed)	Croplands, grasslands	Kazakhstan	<i>Fusarium oxysporum</i> f. sp. <i>cannabis</i>	Tiourebaev et al., 2001
<i>Carduus pycnocephalus</i> (Italian thistle)	Grasslands, woodlands	Tunisia	<i>Puccinia carduorum</i>	Mejri et al., 2010a
<i>Centaurea diffusa</i> (Diffuse knapweed), <i>Centaurea maculosa</i> (Spotted knapweed)	Rangelands	Western U.S., Canada	<i>Fusarium</i> spp.	Caesar et al., 2002
<i>Chenopodium album</i> (Common lambsquarters)	Croplands	Worldwide	<i>Ascochyta caulina</i>	Ghorbani et al., 2002; Vurro et al., 2001
<i>Chrysanthemoides monilifera</i> ssp. <i>monilifera</i> (Boneseed)	Natural areas	Southeastern Australia	<i>Endophyllum osteospermi</i>	Wood & Crous, 2005
<i>Cirsium arvense</i> (Canada thistle)	Croplands, rangelands, pastures, roadsides	Temperate regions of northern hemisphere	<i>Phyllosticta cirsii</i> ; <i>Stagonospora cirsii</i> ; <i>Alternaria cirsinoxia</i> ; <i>Pseudomonas syringae</i> pv. <i>tagetis</i> ; Mix of <i>Phoma destructiva</i> , <i>Phoma hedericola</i> , <i>Mycelia sterila</i> , <i>Phoma nebulosa</i> , <i>Phomopsis cirsii</i>	Evidente et al., 2008; Yuzikhin et al., 2007; Bailey, 2004; Gronwald et al., 2002; Tichich & Doll, 2006; Guske et al., 2004; Leth et al., 2008
<i>Cirsium arvense</i> (Canada thistle), <i>Ranunculus acris</i> (Tall buttercup)	Pasture	New Zealand	<i>Sclerotinia sclerotiorum</i>	Bourdot et al., 2006a
<i>Damasonium minus</i> (Starfruit)	Croplands (rice), aquatic	Australia	<i>Plectosporium alismatis</i>	Jahromi, 2007
<i>Eichhornia crassipes</i> (Water hyacinth)	Aquatic	Tropical, sub-tropical regions	<i>Alternaria eichhorniae</i> isolate 5; <i>Cercospora piaropi</i>	Shabana & Mohamed, 2005; Shabana, 2005; Tessman et al., 2008
<i>Euphorbia esula/virgata</i> (Leafy spurge)	Rangelands, natural areas	North America	<i>Uromyces scutellatus</i>	Caesar, 2006

Weed Pest	Ecosystem	Region	Biocontrol agent	Reference
<i>Euphorbia heterophylla</i> (Wild poinsettia)	Croplands (soybean)	Brazil	<i>Sphaceloma poinsettiae</i>	Nechet et al., 2004
<i>Galium spurium</i> (False cleavers)	Croplands	Canada	<i>Plectosporium tabacinum</i>	Zhang et al., 2002
<i>Gaultheria shallon</i> (Salal)	Native forests	Canadian and American Pacific Coast	<i>Phoma exigua</i> ; <i>Valdensinia heterodoxa</i>	Zhao & Shamoun, 2006; Vogelgsang & Shamoun, 2004; Wilkin et al., 2005
<i>Hydrilla verticillata</i> (Hydrilla)	Aquatic	U.S. and worldwide	<i>Fusarium culmorum</i> ; <i>Mycocleptodiscus terrestris</i>	Shabana et al., 2003; Shearer & Jackson, 2006
<i>Isatis tinctoria</i> (Dyer's woad)	Natural areas	Western North America	<i>Puccinia thlaspeos</i>	Kropp et al., 2002; Kropp & Darrow, 2006
<i>Lantana camara</i> (Lantana)	Natural and cultivated lands	South Africa	<i>Mycovellosiella lantanae</i> var. <i>lantanae</i> ; <i>Corynespora cassiicola</i> f. sp. <i>lantanae</i>	Den Breeyen & Morris, 2003; Pereira et al., 2003
<i>Matricaria perforata</i> (Scentless chamomile)	Croplands	Canada	<i>Colletotrichum truncatum</i>	Graham et al., 2006, 2007; Peng et al., 2005a
<i>Miconia calvescens</i> (Velvet tree; miconia)	Natural areas	Hawaii and French Polynesia	<i>Ditylenchus drepanocercus</i>	Seixas et al., 2004
<i>Nasella neesiana</i> (Chilean needle grass)	Pastures, grasslands	Australia, New Zealand	<i>Uromyces pencanus</i>	Anderson et al., 2010
<i>Orobancha cumana</i> (Sunflower broomrape)	Croplands	Mediterranean region, southeast Europe	<i>Fusarium oxysporum</i> f. sp. <i>orthoceras</i>	Müller-Stöver & Sauerborn, 2007
<i>Orobancha ramosa</i> (Broomrape; branched broomrape)	Croplands	Central & western Europe	<i>Fusarium oxysporum</i> (FOG); <i>Fusarium</i> spp.	Müller-Stöver et al., 2009; Kohlschmid et al., 2009; Boari & Vurro, 2004
<i>Orobancha aegyptiaca</i> (Egyptian broomrape)	Croplands	India, Israel	<i>Fusarium solani</i>	Sharma et al., 2011; Dor & Hershenhorn, 2009
<i>Orobancha crenata</i> , <i>Orobancha foetida</i> (Broomrape)	Croplands	Northern Tunisia	<i>Pseudomonas fluorescens</i> Bf7-9	Zermane et al., 2007
<i>Papaver somniferum</i> (Opium poppy)	Illicit plants		<i>Pleospora papaveracea</i>	Bailey et al., 2004
<i>Portulaca oleracea</i> (Common purslane), <i>Trianthema portulacastrum</i> (Horse purslane), <i>Euphorbia maculata</i> (Spotted spurge), <i>Euphorbia supina</i> (Prostrate spurge)	Tomato fields	Southeastern U.S.	<i>Myrothecium verrucaria</i>	Boyette et al., 2007a
<i>Raphanus raphanistrum</i> (Wild radish)	Croplands, vineyards	Australia	<i>Hyaloperonospora parasitica</i> ; <i>Pseudomonas fluorescens</i>	Maxwell & Scott, 2008; Flores-Vargas & O'Hara, 2006

Weed Pest	Ecosystem	Region	Biocontrol agent	Reference
<i>Salsola kali</i> (Russian thistle)	Croplands, pastures, rangelands	Western U.S., Eurasia	<i>Uromyces salsolae</i>	Hasan et al., 2001
<i>Schinus terebinthifolius</i> (Brazilian peppertree)	Forests	Florida, U.S.	<i>Neofusicoccum batangarum</i>	Shetty et al., 2011
<i>Senecio vulgaris</i> (Common groundsel)	Croplands (carrots)	Switzerland	<i>Puccinia lagenophorae</i>	Frantzen & Müller-Scharer, 2006
<i>Senna obtusifolia</i> (Sicklepod)	Croplands	Southeastern U.S.	<i>Colletotrichum gloeosporioides</i>	Boyette et al., 2007b
<i>Sesbania exaltata</i> (Hemp sesbania)	Croplands	Southern U.S.	<i>Colletotrichum truncatum</i>	Boyette et al., 2007c, 2008b
<i>Sonchus arvensis</i> (Perennial sowthistle)	Croplands	North America, Europe	<i>Alternaria sonchi</i>	Evidente et al., 2009a
<i>Striga hermonthica</i> (Striga, Witchweed)	Croplands	Africa	<i>Fusarium oxysporum</i> (Foxy 2; PSM 197); <i>F. oxysporum</i> (4-3-B); <i>F. nygamai</i> ; <i>Pseudomonas fluorescens</i> ; <i>P. putida</i>	Venne et al., 2009; Yonli et al., 2004; Ahonsi et al., 2002
<i>Taraxacum officinale</i> (Dandelion)	Turfgrass	Worldwide	<i>Phoma herbarum</i> ; <i>Phoma macrostoma</i> ; <i>Sclerotinia minor</i>	Stewart-Wade & Boland, 2004; Zhou et al., 2004; Abu-Dieyeh & Watson, 2006, 2007
<i>Ulex europaeus</i> (Gorse)	Native areas, pastures, forests	New Zealand	<i>Chondrostereum purpureum</i> ; <i>Fusarium tumidum</i>	Bourdot et al., 2006b; Yamoah et al., 2008
<i>Xanthium strumarium</i> (Common cocklebur)	Croplands	Southern U.S.	<i>Alternaria helianthi</i>	Abbas et al., 2004
<i>Xanthium occidentale</i> (Noogoora burr)	Croplands, rangelands	Australia	<i>Puccinia xanthii</i>	VanKlinken & Julien, 2003

Table 1A. List of current biological control research projects on dicotyledonous weed species that have shown promise in greenhouse and/or field trials, including the invasive weed species, ecosystem, region of importance, biological control agent and reference.

Biological herbicides represent a means to reduce dependence on synthetic herbicides; focus on ecologically grounded methods of management; reduce weed seed bank populations through environmentally friendly practices; and potentially reduce costs of weed control in crop production, rangeland restoration, forestry and aquatic systems (Bailey et al., 2010; Kennedy & Stubbs, 2007).

2. Challenges for weed control with microbial herbicides

The list of challenges in developing a successful microbial herbicide is long. Once a potential biological control agent is identified, the first challenge lies in reproducing lab and/or greenhouse results successfully in the field. The potential biological control agent and the toxin responsible for weed inhibition must be able to survive the harsh, and often unpredictable, environmental conditions that exist in the field. Li & Kremer (2006) showed inhibition of several weed species using rhizobacteria isolated from various weed hosts in

Weed Pest	Ecosystem	Region	Biocontrol agent	Reference
<i>Aegilops cylindrica</i> (Jointed goatgrass)	Croplands	Western U.S.	<i>Pseudomonas</i> spp.; <i>Xanthomonas</i> spp.	Kennedy & Stubbs, 2007
<i>Avena fatua</i> (Wild oat)	Cereal crops, croplands, native areas	North America, Europe, Australia	<i>Fusarium avenaceum</i> ; <i>F. culmorum</i> ; <i>Drechslera avenacea</i> (conidial state of <i>Pyrenophora chaetomioides</i>); <i>Puccinia coronata</i> f. sp. <i>avenae</i>	deLuna et al., 2011; Ghajar et al., 2006; Carsten et al., 2001
<i>Bromus diandrus</i> (Great brome)	Croplands	Tunisia	<i>Pseudomonas trivialis</i> strain X33D	Mejri et al., 2010b
<i>Bromus tectorum</i> (Downy brome; cheatgrass)	Croplands, rangelands, natural areas	Western North America	<i>Pseudomonas fluorescens</i> strain D7; <i>Pyrenophora semeniperda</i> ; <i>Ustilago bullata</i>	Kennedy et al., 1991, 2001; Meyer et al., 2001, 2007
<i>Digitaria sanguinalis</i> (Giant crabgrass; large crabgrass)	Croplands	China	<i>Curvularia eragrostidis</i> QZ-2000; <i>C. intermedia</i>	Zhu & Qiang, 2004; Jiang et al., 2008; Tilley & Walker, 2002
<i>Elytrigia repens</i> (Quackgrass)	Croplands	Temperate regions of northern & southern hemisphere	<i>Ascochyta agropyrina</i> var. <i>nana</i>	Evidente et al., 2009b
(Grasses)	Croplands	Australia	<i>Pyrenophora semeniperda</i>	Medd & Campbell, 2005
<i>Microstegium vimineum</i> (Japanese stiltgrass)	Forests	Eastern U.S.	<i>Bipolaris</i> sp.	Kleczewski & Flory, 2010
<i>Poa annua</i> (Annual bluegrass)	Turfgrass	Japan	<i>Xanthomonas campestris</i> pv. <i>poae</i> (JT-P482)	Imaizumi et al., 1999
<i>Setaria viridis</i> (Green foxtail)	Croplands	Worldwide	<i>Drechslera gigantea</i> ; <i>Exserohilum rostratum</i> ; <i>E. longirostratum</i> ; <i>Pyricularia setariae</i>	Casella et al., 2010; Green et al., 2004

Table 1B. List of current biological control research projects on monocotyledonous weed species that have shown promise in greenhouse and/or field trials, including the invasive weed species, ecosystem, region of importance, biological control agent and reference.

greenhouse studies. They noted the importance of greenhouse studies as a step toward identifying isolates that may be suitable for testing under the more variable conditions of the field, where there is greater competition from indigenous organisms and unpredictable environmental factors. In any case, the conditions under which a microorganism best survives depend on the microorganism itself and it cannot be assumed to be the same for all biocontrol agents. Cool, moist conditions are required for survival of the deleterious

rhizobacteria (DRB) used to control jointed goatgrass (*Aegilops cylindrica*) and downy brome (*Bromus tectorum*) in the field (Kennedy & Stubbs, 2007). For greatest biocontrol success, field application of organisms should be timed when rains are expected. On the other hand, Boguena et al. (2007) found that extremely cold temperatures reduced the ability of *Ustilago bullata* to infect downy brome and may limit its use as a biological control. One challenge to the development of fungal bioherbicides is the inability to infect the weed pest without a period of free water retention (Chittick & Auld, 2001). Favorable moisture and temperature conditions are critical to the efficacy of many mycoherbicides. Tichich et al. (2006) found that for optimum population growth of the pathogen *Pseudomonas syringae* pv. *tagetis* targeting Canada thistle (*Cirsium arvense*), periods of wet weather were required. This moisture requirement often limits candidates for biocontrol. After investigating the dew period requirements of three fungal pathogens that infected green foxtail (*Setaria viridis*), Peng & Boyetchko (2006) found that *Drechslera gigantea* was the best choice for control of green foxtail because this pathogen was the most virulent and it did not require a specific dew temperature for efficacy.

For successful biocontrol, the microorganism's growth stage needs to be matched with the period of greatest vulnerability of the weed. Studies are needed to ensure that the most virulent stage of biocontrol agent growth also coincides with the susceptible host habit and an active growth period of the host. The efficacy of *Plectosporium alismatis* to reduce starfruit (*Damasonium minus* (R. Br.) Buch) in rice was increased by using conidia and chlamydospores (Cliquet & Zeeshan, 2008) and applying the biocontrol agent on juvenile rather than older starfruit (Jahromi, 2007). Qiang et al. (2006) found that the mycelia of *Alternaria alternata* strain 501 were able to infect the host plant, *Eupatorium adenophorum*, in a much shorter time than with conidia. The *Pseudomonas fluorescens* strain D7 has the greatest efficacy on downy brome when the bacterium is applied in the fall. In addition, populations of *P. fluorescens* D7 in soil are greatest in the fall and spring, which coincides with active root growth of downy brome (Kennedy et al., 1991).

Another challenge to successful microbial herbicides is developing strains that are effective against weed populations with high genetic diversity. Diversity within a species can lead to inconsistent results with biocontrol measures (Bailey, 2004; Ward et al., 2008). Biological control isolates may be specific to the region where they were first isolated, for example Ash et al. (2008) studied fungal pathogens from Korea and Australia, and found that Korean isolates showed less pathogenicity on Australian weeds than the Australian isolates. The efficacy of *Sclerotinia minor* on dandelion (*Taraxacum officinale*) was dependent upon weed accession, age and plant competition (Abu-Dieyeh & Watson, 2007). Both physiological and ecological considerations need to be examined for successful biocontrol interactions.

Prior to commercialization of a biological herbicide, extensive host-range studies must be completed to determine effects of the agent on non-target organisms in order to minimize any harmful effects. Potential microbial herbicides must be virulent on the target species, while non-target plants remain disease-free (Bailey, 2004). Wapshere's (1974) concept of concentric spheres of related plant species is a starting point for investigations of non-target plant species. In host-range studies of *Alternaria alternata*, a potential biological control of water hyacinth (*Eichhornia crassipes*), the biocontrol agent also inhibited the weed water lettuce (*Pistia stratiotes* L.; Mohan Babu et al., 2002). They evaluated 29 economically and environmentally significant species that encompassed more than 18 families. After finding that only two plant species were

inhibited, they concluded that *A. alternata* would not be harmful to any economically important plants. Kennedy et al. (2001) found that *Pseudomonas fluorescens* strain D7 inhibited the target weed, downy brome, a few other *Bromus* species, jointed goatgrass and medusahead (*Taeniatherum caput-medusae*) in bioassays, greenhouse and field studies. They investigated representatives from a wide array of families that included native, crop and weed species, while concentrating on families, tribes, subtribes and accessions closely related to *Bromus* species. No dicots and only a few monocots were negatively affected by the bacterium. While a few other grasses were suppressed slightly by *P. fluorescens* D7 in bioassays, they were not suppressed in greenhouse studies using nonsterile soil. These two examples illustrate the importance of extensive testing of other plant species. The lack of host-range testing often leads to the early demise of a potential biocontrol agent.

Other challenges are large-scale production and storage, survival of the organism, requirement of the agent for formulation, shelf-life of the organism, and delivery system of the biocontrol agent to the host plant. The ability to produce large quantities of microbial products, and maintain survival of the organisms, are the major obstacles to field-scale application of biological control agents (Amsellem et al., 1999). Teshler et al. (2007) examined shelf life of the *Sclerotinia minor* bioherbicide and found that storage temperatures less than 11°C increased *S. minor* shelf life, but CO₂ did not affect shelf life. Cooler storage temperatures helped to prolong the shelf life of *Fusarium oxysporum* (Foxy2 and PSM197) for control of *Striga*; however, vacuum packing was not helpful. The fungicide Apron XL enhanced shelf life, but another fungicide, Ridomil Gold (both fungicides Metalaxyl-M, Syngenta, Basle, Switzerland & Germany), did not (Elzein et al., 2009). The authors attributed this difference to the higher recommended application rate for Ridomil Gold, or possibly that Apron XL is ineffective against ascomycetes such as *Fusarium oxysporum*. While most of the microbial herbicides developed to-date have been mycoherbicides, bacterial herbicides are becoming available and have different challenges and advantages. These include simpler fermentation processes, ease of upscaling and mass-production, production of secondary metabolites and lack of spore formation that may require specific growth conditions (Li et al., 2003).

Application technology and cultural practices also affect the efficacy of microbial herbicides applied in field situations. Aerosol applications in the field may need different inoculum levels than what are found to be successful in greenhouse studies. In addition, droplet size, and spray direction may need to be readjusted to reach the weed or soil for greatest weed reduction (Lawrie et al., 2002b). Byer et al. (2006) found that finer droplet size led to greater spray retention of *Colletotrichum truncatum* on scentless chamomile and *Colletotrichum gloeosporioides* f. sp. *malvae* on round-leaved mallow. Peng et al. (2005b) looked at spray retention for efficacy of the biological control agent *Pyricularia setariae* on green foxtail. They looked at sprayer type and nozzle size for varying application rates and droplet sizes. In general, finer droplet size was more advantageous, and the authors suggested further study with other factors such as use of adjuvants and formulations. Boyette et al. (2007b) found that *Colletotrichum gloeosporioides* was an effective biocontrol agent of sicklepod in soybean; however, the wider row spacings required repeat applications of the agent for adequate weed suppression. The challenges to successful biocontrol are many, and continued research and development are needed to offer the alternative of microbial weed management.

3. Successful microbial herbicides

The majority of biological herbicides developed to-date are mycoherbicides; however, several bacterial herbicides are under development as well. A screening procedure for a bacterial biological control agent has been developed by the authors (Figure 1). Our screening procedure takes advantage of the soil microbial community and the diversity within. This screening includes sampling the soil and plant material at the peak of population growth and when the suppression naturally occurs. An initial assay can separate the weed-suppressive microorganisms from the bulk of the native population. Further multiple screenings on various plants are suggested to get rid of those isolates that may also inhibit beneficial plants. While inhibition of the target weed is critical, non-host range must be determined early in the process to ensure the product progresses through the registration process and eventually becomes marketable. Both steps need to be thorough to ensure that no lesser known, but economically important, host is detected later in the process. Souissi and Kremer (1998) utilized a multiple-well plate procedure with leafy spurge (*Euphorbia esula* L.) callus to rapidly determine phytotoxicity of rhizobacterial isolates, and Vidal et al. (2004) have developed a successful method to produce yellow starthistle (*Centaurea solstitialis* L.) calli for bioassay screening of biocontrol pathogens such as *Phoma exigua*. A quick substitute bioassay will reduce the length of time needed for the initial screenings. In many cases, a quick screen is not possible (Kennedy et al., 1991), but automating data recording can often make data collection easier (Doty et al., 1994).




Fig. 1. Flow of assays to obtain a biological control organism from soil that successfully suppresses the growth of target organisms but has a limited host range and does not inhibit beneficial plant species.

Since the early 1980's, there have been several successful biological herbicides released to the market. Biomal® (ATCC 20767) *Colletotrichum gloeosporioides* f. sp. *malvae*, was developed as a weed biocontrol agent against round-leaved mallow (*Malva neglecta*) and registered in 1992. DeVine was marketed as a control for stranglervine (*Morrenia odorata*) in citrus, and Collego was used to control northern jointvetch (*Aeschynomene virginica*) in rice. Velgo is a potential mycoherbicide for control of velvetleaf in the U.S. and Canada (Mortensen, 1998; Owen & Zdor, 2001). DeVine and Collego were registered for use in the United States, and each controls a single weed species. Four bioherbicides released in Canada included two strains of *Chondrostereum perpureum* (HQ1 (Myco-Tech Paste) and PFC 2139 (Chontrol Paste)) for control of trees and shrubs (Becker et al., 2005), *Colletotrichum gloeosporioides* f. sp. *malvae* to control round-leaved mallow (Cross and Polonenko, 1996; no longer available due to small market size), and *Sclerotinia minor* (IMI 3144141) for control of dandelions (Abu-Dieyeh & Watson, 2007; Bailey et al., 2010). CAMPERICO (*Xanthomonas campestris* pv. *poae* isolate JT-P482) was registered in Japan for control of annual bluegrass (*Poa annua* L.) in 1997 (Imaizumi et al., 1999). The fungal pathogen *Alternaria destruens* strain 059 (Smolder G and Smolder WP) was registered in the U.S. in 2005 for control of dodder (*Cuscuta* sp.) in field crops and ornamental plants (USEPA, 2005).

The mode of action of each biocontrol agent is as varied as the microorganisms themselves (de Luna et al., 2011). They range from simple but effective compounds like cyanide (Kremer & Souissi, 2001; Owen & Zdor, 2001) and organic acids to complex molecules with tertiary structure (Bouizgarne et al., 2006; Gurusiddaiah et al., 1994), and from secondary metabolites (Kroschel & Elzein, 2004) to plant growth regulators, such as auxins and ethylene (de Luna et al., 2005). Pedras et al. (2003) isolated toxins from *Pseudomonas fluorescens* strain BRG100. One of the toxins, pseudophomin A, showed greater inhibition of green foxtail than did the other, pseudophomin B, which showed greater activity against several plant pathogens. There are sufficient successful products out in the market or in development that indicate continued efforts will provide more microbial herbicides and better weed management options.

4. Formulations to improve success of microbial herbicides

In order to overcome the obstacles associated with development of a microbial herbicide, and dramatically improve the chances for success of microbial herbicides, numerous researchers have investigated combining the biological control organism with a formulation designed to improve application, survivability, and efficacy. Several research projects have studied the use of a formulation or carrier in combination with a biological control organism (Table 2).

As mentioned earlier, appropriate temperatures and length of dew period are critical to the success of fungal bioherbicides. Formulations can extend the period of time before dew is required, and adjuvants such as unrefined corn oil and Silwet L-77 can improve chances for success of mycoherbicides (Abbas et al., 2004; Boyette, 2006; Boyette et al., 2007a). Elzein et al. (2004) found that *Fusarium oxysporum* 'Foxy 2' could be encapsulated in a pesta formulation to improve shelf life. Amsellem et al. (1999) utilized the 'Stabileze' formulation (containing starch, sucrose, corn oil and silica) to enhance preservation of mycelia from two *Fusarium* spp. and found that they remained viable for over one year. The 'Stabileze' method has also been utilized to enhance survival of bacteria. Zidack & Quimby (2002) used this

Biological Control Agent	Weed Pest	Formulation / Carrier	Reference
<i>Alternaria eichhorniae</i> (Ae5)	Water hyacinth (<i>Eichhornia crassipes</i>)	Nine oil emulsions	Shabana, 2005
<i>Alternaria helianthi</i>	Common cocklebur (<i>Xanthium strumarium</i>)	Unrefined corn oil; Silwet L-77	Abbas et al., 2004
<i>Bipolaris</i> sp.	Japanese stiltgrass (<i>Microstegium vimineum</i>)	Tween 20	Kleczewski & Flory, 2010
<i>Colletotrichum gloeosporioides</i>	Sicklepod (<i>Senna obtusifolia</i>)	Adjuvants (unrefined corn oil; invert emulsion-MSG 8.25; Silwet L-77)	Boyette, 2006
<i>Colletotrichum truncatum</i>	Hemp sesbania (<i>Sesbania exaltata</i>)	Unrefined corn oil; Silwet L-77	Boyette et al., 2007c
<i>Curvularia eragrostidis</i> (QZ-2000)	Large crabgrass (<i>Digitaria sanguinalis</i>)	Tween 80; rapeseed oil	Zhu & Qiang, 2004
<i>Curvularia intermedia</i>	Large crabgrass (<i>Digitaria sanguinalis</i>)	Silwet L-77	Tilley & Walker, 2002
<i>Fusarium oxysporum</i> f. sp. <i>cannabis</i>	Ditchweed (<i>Cannabis sativa</i>)	Birch sawdust; wheat seeds; oat seeds	Tiourebaev et al., 2001
<i>Fusarium oxysporum</i> f. sp. <i>orthoceras</i>	Sunflower broomrape (<i>Orobanche cumana</i>)	Pesta granules; commercial iron fertilizers	Müller-Stöver & Sauerborn, 2007
<i>Fusarium oxysporum</i> (FOG)	Branched broomrape (<i>Orobanche ramosa</i>)	Pesta granules; alginate pellets	Kohlschmid et al., 2009
<i>Fusarium oxysporum</i> ; <i>F. arthrosporioides</i>	Broomrapes (<i>Orobanche</i> spp.)	Alginate beads; 'Stabileze' (starch, sucrose, corn oil, silica)	Amsellem et al., 1999
<i>Fusarium oxysporum</i> f. sp. <i>strigae</i> (Foxy 2)	<i>Striga</i> (<i>Striga hermonthica</i>)	Film-coat on sorghum seeds (gum arabic, 40%); Pesta granules; seed treatment (gum arabic, SUET binder)	Elzein et al., 2004, 2006, 2009
<i>Fusarium tumidum</i>	Gorse (<i>Ulex europaeus</i>)	Tween 80; 5% Triton X-100	Yamoah et al., 2008
<i>Helminthosporium gramineum</i> subsp. <i>echinochloae</i> ; <i>Curvularia lunata</i>	Barnyardgrass (<i>Echinochloa crus-galli</i>)	Tween 20	Zhang et al., 2007
<i>Mycleptodiscus terrestris</i>	Hydrilla (<i>Hydrilla verticillata</i>)	Diatomaceous earth	Shearer & Jackson, 2006
<i>Myrothecium verrucaria</i>	<i>Portulaca</i> spp., <i>Euphorbia</i> spp.	Silwet L-77	Boyette et al., 2007a
<i>Myrothecium verrucaria</i>	Kudzu (<i>Pueraria lobata</i>)	Silwet L-77; unrefined corn oil	Hoagland et al., 2007
<i>Neonectria neomacrospora</i>	Hemlock dwarf mistletoe (<i>Arceuthobium tsugense</i>)	'Stabileze'	Rietman et al., 2005
<i>Phomopsis amaranthicola</i>	<i>Amaranthus</i> spp.	16 adjuvants	Wyss et al., 2004
<i>Pseudomonas fluorescens</i> strain G2-11	Green foxtail (<i>Setaria viridis</i>); velvetleaf (<i>Abutilon theophrasti</i>)	Corn gluten meal; semolina flour	Zdor et al., 2005

Biological Control Agent	Weed Pest	Formulation / Carrier	Reference
<i>Pseudomonas syringae</i> pv. <i>tabaci</i> ; <i>P. syringae</i> pv. <i>tagetis</i>	Multiple weeds	'Stabileze' method (bacteria, oil, sucrose; silica)	Zidack & Quimby, 2002
<i>Pseudomonas syringae</i> pv. <i>tagetis</i>	Canada thistle (<i>Cirsium arvense</i>)	Silwet L-77; Canada thistle sap	Gronwald et al., 2002; Tichich et al., 2006; Tichich & Doll, 2006
<i>Puccinia thlaspeos</i>	Dyer's woad (<i>Isatis tinctoria</i>)	Surfactants Sylgard, IFA-S90, Regulaid	Kropp & Darrow, 2006
<i>Sclerotinia minor</i>	Dandelion (<i>Taraxacum officinale</i>)	Barley (<i>Hordeum vulgare</i>) grits	Teshler et al., 2007

Table 2. Research projects examining formulations or carriers to improve survival and efficacy of microbial herbicides.

method with two *Pseudomonas* spp., and showed that populations were still high after one year, and that components of the formulation could be varied depending on bacterial species response.

Elzein et al. (2006, 2010) studied seed coatings containing *Fusarium oxysporum* isolates to control *Striga*, and found a 40% gum arabic seed coating combined with dried chlamydo spores to be the most effective combination of seed coating and inoculum type for causing disease in *Striga* (Elzein et al., 2006). Zhao & Shamoun (2005) tested combinations of gelatin and potato dextrose broth concentrations for optimum efficacy of *Phoma exigua* to control salal (*Gaultheria shallon*), a perennial evergreen shrub. *Fusarium oxysporum* f. sp. *orthoceras* (FOO) is known to suppress the root parasitic weed broomrape (*Orobancha cumana*) in sunflower. In addition, Benzo(1,2,3)thiadiazole-7-carbothioic acid S-methyl ester (BTH) induces sunflower resistance to *Orobancha cumana*. Using FOO and BTH, Müller-Stöver et al. (2005) improved the efficacy of *Fusarium oxysporum* f. sp. *orthoceras* (FOO) to reduce broomrape infection in greenhouse studies. They also found FOO incorporated into a wheat-kaolin and iron mix further improved efficacy by increasing FOO survival (Müller-Stöver & Sauerborn, 2007).

Chittick & Auld (2001) examined the use of hydrophilic polymers as a formulation for a mycoherbicide to improve efficacy of *Colletotrichum orbiculare* on *Xanthium spinosum* (Bathurst burr) in Australia. Hoagland et al. (2007) studied formulation, application method and growth media for control of kudzu (*Pueraria lobata*) using *Myrothecium verrucaria* fungi. Shabana (2005) found that the efficacy of *Alternaria eichhorniae* isolate #5 could be improved, and the requirement for a dew-period avoided, by applying the fungi using an oil emulsion. With that formulation, complete control of water hyacinth under field conditions in Egypt was achieved. Hurrell et al. (2001) found a granular mycelium-wheat formulation of *Sclerotinia sclerotiorum* controlled *Cirsium arvense* best in pasture lands of New Zealand when the formulation was a water-miscible powder applied as a slurry rather than a dry product. They also found that spring and early summer applications, when some moisture was present, were more effective at reducing the weed than late summer or early autumn. While some moisture was needed for efficacy, too much rain was thought to wash the agent off the leaf. Kohlschmid et al. (2009) found that a combination of alginate pellets and pesta granules formulated with the *Fusarium oxysporum* isolate FOG was more efficacious and reliable for

controlling the parasitic weed branched broomrape (*Phelipanche ramosa*) than the untreated control under field conditions. Boyette (2006) examined several adjuvants in combination with the mycoherbicide *Colletotrichum gloeosporioides* for control of sicklepod. These compounds reduced length of the dew period requirement, and improved the performance of this organism in controlling sicklepod. Boyette et al. (2007c) showed that *Colletotrichum truncatum* formulated with unrefined corn oil and the surfactant Silwet L-77 was able to effectively control hemp sesbania in the greenhouse and field by reducing the dew period requirement. Zhang et al. (2007) utilized protoplast fusion as a means to improve the biocontrol efficacy of *Helminthosporium gramineum* subsp. *echinochloae* strain HM1 against barnyardgrass (*Echinochloa crus-galli*) in rice.

Most of the research conducted to develop formulations that improve survival and efficacy of microbial herbicides has been directed at fungal herbicides; however, there have been a few studies aimed at improving delivery of bacterial herbicides. Zdor et al. (2005) studied effects of DRB (*Pseudomonas fluorescens* strain G2-11) in combination with corn gluten meal and semolina flour in soil assays using weed and crop species. Zhang et al. (2010) studied the stability of pyoluteorin, a polyketide metabolite produced by fluorescent pseudomonads that has shown potential to control weeds, among other pests. Tichich and Doll (2006) examined a novel application approach for the pathogen *Pseudomonas syringae* pv. *tagetis* where the sap of infected Canada thistle is extracted, combined with water and Silwet-77, and sprayed. While there was disease expression in the treated plants, it was not enough to control the Canada thistle and further work is needed on this approach.

Various technologies have been used and will continue to be used to enhance biological weed control (Cohen et al., 2002). The protoplast fusion technique was used to create new strains using *Helminthosporium gramineum* subsp. *echinochloae* strain HM1 (high pathogenicity, low spore formation) and *Curvularia lunata* (low pathogenicity, high spore formation) to create strains that effectively control barnyardgrass and other weeds in rice production (Zhang et al., 2007). Hypervirulence selection or manipulation may improve efficacy of biological control agents. Cohen et al. (2002) transformed genes of the indole-3-acetamide (IAM) pathway to cause an auxin imbalance that increased the virulence of *Fusarium oxysporum* and *F. arthrosporioides*, pathogenic on broomrape (*Orobancha aegyptiaca*). Sands and Pilgeram (2009) outline the steps to enhance virulence of the biocontrol agent using amino acid overproduction. They discuss control of the parasitic weeds *Orobancha* and *Striga*, which are especially challenging to control due to the close relationship they develop with their hosts. Economic formulations and genetic manipulations to alter phenotype will assist in the understanding and development of microbial herbicides.

5. Integrating microbial herbicides with other control measures

In many cases, microbial herbicides alone will not be enough to remedy invasive weed problems. Researchers worldwide have shown that an integrated approach utilizing microbial weed management in a synergistic or additive manner with chemical herbicides or in combination with cultural practices or biological controls with insects is more successful than any of the control measures alone. Innovative approaches for microbial management of several different weed species in various regions of the world have been employed to sustainably manage some of the world's worst weed problems in diverse systems.

Denoth et al. (2002) reviewed biological control projects on weeds and insects, and determined that for weeds, those projects where a number of agents were released showed greater success. Wandeler et al. (2008) used weevils as an insect vector to apply *Puccinia punctiformis* to creeping thistles (*Cirsium arvense*) in the field, causing systemic rust infection. Another integrated approach to biological control with microorganisms utilizes mixtures of organisms, rather than a single pathogen to control weed growth. Chandramohan & Charudattan (2003) propose the use of a mixture of four fungal plant pathogens (*Phomopsis amaranthicola*, *Alternaria cassia*, *Colletotrichum dematium* f.sp. *crotalariae*, *Fusarium udum* f.sp. *crotalariae*) to control pigweed (*Amaranthus hybridus* L.), sicklepod and showy crotalaria (*Crotalaria spectabilis* Roth.). In their greenhouse study, they showed that it was possible to control these three weeds together using several fungal strains without losing efficacy or host-specificity. Some potential biological control agents work best as individual applications, and do not exhibit synergy when applied together. Dooley & Beckstead (2010) found no improvement in downy brome inhibition using *Pseudomonas fluorescens* strain D7 in combination with the fungal pathogen *Pyrenophora semeniperda*. These two microorganisms suppress downy brome at different growth stages. Another example of this is *Chondrostereum purpureum*, which is applied in spring and *Fusarium tumidum*, which is applied in early winter to reduce gorse (*Ulex europaeus*) regrowth in New Zealand forests (Bourdote et al., 2006b). These mycoherbicides inhibit gorse at different growth stages, but together they did not further reduce the regrowth of stumps. Further studies are needed to understand the weed reduction and ecological implications of consortia in biocontrol efforts.

In greenhouse studies, Caesar (2003) found that the combined effects of *Fusarium oxysporum* and *Rhizoctonia solani* with flea beetle adults and larvae resulted in greater inhibition of *Euphorbia esula/virgata* than any of the biological control agents alone. In their survey of *Lepidium draba* throughout Europe, Caesar et al. (2010) showed that plants sustaining both insect damage and disease were being colonized by the root pathogen *Rhizoctonia solani*. They concluded that when determining potential biological control agents, the synergistic relationships between plant pathogens and insects should be considered. Likewise, Kremer et al. (2006) found that the most effective biocontrol of leafy spurge occurred with the synergistic effect of “plant-associated microorganisms and root-damaging insects”. There was a higher incidence of *Fusarium* and *Rhizoctonia* isolates in *Euphorbia* plants that had injury caused by insect feeding. Rayamajhi et al. (2010) noted the additive effects of using combinations of a weevil, psyllid, and rust fungus (*Puccinia psidii*) to reduce regrowth of tree stumps of *Melaleuca quinquenervia* in southern Florida. The use of multiple enemies against invasive weeds is another combination that could be successful in biocontrol programs.

Babalola et al. (2007) examined the use of trap crops such as cowpea (*Vigna unguiculata*) combined with application of bacteria (*Enterobacter sakazakii* and *Pseudomonas* spp.) to stimulate germination and cause the subsequent death of the parasitic weed *Striga hermonthica*. Similarly, Ahonsi et al. (2003) used ethylene-producing *Pseudomonas syringae* pv. *glycinea* in combination with nitrogen-fixing *Bradyrhizobia japonicum* strains to induce germination and death in *Striga hermonthica* seeds in the presence of cowpea or soybean. No matter how successful a biocontrol agent may be, the control is never considered to be 100% and additional practices and management efforts need to be integrated with these microorganisms or insects to attain weed management or control.

Two methods have been employed to determine whether a synergistic relationship exists between a chemical herbicide and a microbial herbicide (Gressel, 2010): 1) random testing of herbicides, and 2) screening chemical herbicides with the intent to alter the target plant's defenses, leaving it more vulnerable to attack by pathogens. Better control of *Chenopodium album* was achieved by using low rates of herbicides in combination with toxins from *Ascochyta caulina* (Vurro et al., 2001). Jahromi et al. (2006) studied interactions between herbicides and *Plectosporium alismatis* to control starfruit in rice. Kropp & Darrow (2006) found that herbicides and surfactants did not negatively affect teliospore viability of *Puccinia thlaspeos* when it was sprayed on to *Isatis tinctoria* (Dyer's woad) plants in the field. Abu-Dieyeh & Watson (2006) examined the relationship between the fungal pathogen *Sclerotinia minor* Jagger applied with different turfgrass mowing heights compared to herbicide alone to control dandelion. In the greenhouse, *S. minor* caused greater damage to dandelions than the herbicide at all mowing heights; however, under field conditions, close mowing had an unfavorable effect on *Sclerotinia minor* for dandelion control. Magani et al. (2009) demonstrated that the fungal mycoherbicide *Fusarium oxysporum*, when applied in granular form followed by a post-emergence herbicide treatment, was successful in controlling the parasitic plant *Striga* in Nigeria.

Peng and Byer (2005) tested seven herbicides at reduced rates to determine whether there might be synergistic effects with *Pyricularia setariae* for control of green foxtail. Responses were variable and depended on weed growth stage, herbicide and application rates. However, each of the herbicides at a one-quarter rate applied with a sub-lethal dose of the biocontrol organism succeeded in controlling green foxtail in the greenhouse (Peng & Byer, 2005). Jahromi et al. (2006) studied interactions between the fungal pathogen *Plectosporium alismatis* and herbicides for control of starfruit in Australian rice production. In glasshouse experiments, there was no synergistic effect with the pathogen and 2-methyl-4-chlorophenoxyacetic acid (MCPA); however, there was a synergistic effect when the pathogen was applied after the sublethal dose of the herbicide Londax® (bensulfuron methyl, DuPont, Wilmington, DE).

Gressel (2010) notes that glyphosate is most commonly utilized in synergistic relationships with plant pathogens, and hypothesizes that this is due to the capability of glyphosate to affect multiple weed defense mechanisms. Boyette et al. (2008a) in a field study controlled redivine (*Brunnichia ovata*) and trumpetcreeper (*Campis radicans*) using the synergistic relationship between glyphosate and the fungus *Myrothecium verrucaria*. The combination of the chemical herbicide and the microbial herbicide was able to control the weeds better than either treatment alone. Boyette et al. (2008b) concluded that hemp sesbania (*Sesbania exaltata*) control might be augmented by utilizing the biological control agent *Colletotrichum truncatum* in combination with reduced rates of the herbicide glyphosate when the fungus is applied after the herbicide. In a previous experiment under controlled conditions, Boyette et al. (2006) tested *Myrothecium verrucaria* in combination with glyphosate on kudzu (*Pueraria lobata*), redivine and trumpetcreeper at various temperatures. Greatest disease development was achieved at higher temperatures, and weed inhibition was greatest when the fungus was applied after the herbicide rather than prior to or with the glyphosate. Cook et al. (2009) found that dodder (*Cuscuta pentagona*) was more effectively controlled in greenhouse studies using a mixture of the pathogen *Alternaria destruens*, glyphosate, oil and an ammonium sulfate surfactant than when using any of the treatments alone, while not harming the host citrus (*Citrus* spp.) plant.

Care must be taken when integrating biological controls with herbicides so that the herbicide does not affect microbial survival and efficacy. Ray et al. (2008) found that dose and herbicide type may be detrimental to the water hyacinth pathogen *Alternaria alternata*. The effects of four herbicides at recommended and reduced rates were tested on the rust fungus *Puccinia lagenophorae* to be used against *Senecio vulgaris* (common groundsel). Wyss & Müller-Scharer (2001) found that the herbicides they tested were either too toxic to the fungus, or did not increase plant susceptibility to the fungus, and so the combination of the rust fungus and herbicide was not an option for control of *Senecio vulgaris*. Shabana & Mohamed (2005) combined *Alternaria eichhorniae* isolate 5 with 3,6-dichloro-2-pyridinecarboxylic acid (MDCA) to weaken water hyacinth defenses, which increased disease severity and has the potential for greater biocontrol.

The soil possesses a wealth of genetic potential waiting to be discovered and used in weed management systems. Soil quality or the chemical, physical and biological properties of soil can influence plant-microbe interactions and the success of any biocontrol agent (Kennedy & Papendick, 1995). In addition, soil quality investigations often include weed populations as an indicator of soil quality, in part because a healthy soil and healthy desirable plant is the best weed control (Magdoff, 2001; Ryan et al., 2011). Soil investigations are needed in determining the impact of the soil environment on weed populations. The structure and function of soil microbial communities develop depending on the soil, location, climate, slope, aspect and vegetation. Management practices can influence the soil microbial community and plant-microbe interactions (Wander et al., 1995; Lupwayi et al., 1998; Kennedy & Schillinger, 2006). Cultural practices and application of amendments may also play a part in weed-suppressive soils, although the modes of action may be different for each practice. Cropping systems also influence weed-suppressive bacteria (Kremer & Li, 2003; Ryan et al., 2011). Kremer & Li (2003) found that high enzyme activity and greater volume of water stable aggregates correlated with more weed-suppressive bacteria. They found that uncultivated prairie and no-tillage systems contained the highest populations of deleterious rhizobacteria compared to other land use and soil quality indicators, and may be useful in selecting for weed-suppressive practices. In a similar study in Washington state, Kennedy & Stubbs (2007) could not find relationships among management systems and the prevalence of weed-suppressive bacteria.

The concept of weed-suppressive soils can be defined as soils, amendments, or management practices that have the capacity to reduce or limit specific weeds. Management for weed-suppressive soils is the ultimate goal for biological control efforts. The dynamics of weed-suppressive soils may be similar to what is seen with disease-suppressive soils (Mazzola, 2004). The biology, chemistry and physical properties of soil that comprise a weed-suppressive soil need to be characterized in order to manage for suppressive soils. As with all soil quality determinations, no one indicator will explain the complexity of weed suppression.

6. Regulatory issues

Rigorous testing is required prior to the release of a biological herbicide to ensure the safety of humans, animals and the environment. Host-range studies are needed to reduce potential risk and ensure that beneficial, non-target plant species are unaffected by the biocontrol agent. However, the length of time needed to complete assessments of new biological

herbicides adds to the costs and the length of time required before an agent can be released (Ghosheh, 2005). Non-host testing is important and the ranges of plant species tested depend on the areas of release, ecosystem variability and potential for dissemination of the biocontrol agent by wind or water. Testing should cover all economically important plant species of the area, and those plants known to be involved in ecosystem maintenance. In agronomic ecosystems, the major crop species are of interest. The U.S. Environmental Protection Agency (EPA) published a list of the top 25 major agricultural crops. Plants were placed on this list because of their economic importance, ecosystem activity or total production values (EPA, 2011). In aquatic systems, several aquatic plants are suggested that include algae, aquatic bacteria, marine and freshwater diatoms. In rangeland ecosystems the non-target species would include native or near native plant species. It is recommended to test six species covering at least four families in the Dicotyledonae, and at least four species of at least two families in the Monocotyledonae. Testing must be performed on all plants of economic importance in agriculture, horticulture or rangeland systems or known to be beneficial to maintenance of the ecosystem that have any reasonable likelihood of serving as hosts. This selection of additional plant species should be based upon a survey of plants closely related (same genus or, if not available, same family) to the target plant and a survey of known hosts of pathogens closely related to the microbial herbicide (EPA, 2011; Wapshere, 1974).

With thorough host-range testing, very few, if any, detrimental effects occur from the release of fungal herbicides to control weeds (Barton, 2004). In a review of pre- and post-release records from 26 projects, Barton (2004) found that there were no reports of a fungal biological control agent striking an unintended plant species. Additional animal, avian, fish and daphnia testing are also required in many countries before bioherbicides can be registered. In addition, as with all research and new products where there are safety concerns, buffer zones are often required to protect animal pastures and other non-target areas (Bourdot et al., 2006a). The risk of applying a microbial herbicide to the environment needs to be considered at the beginning and throughout the development of biocontrol agents.

7. Future prospects for microbial herbicides in sustainable ecosystems

The future of biocontrol is bright and full of possibilities with the many novel, successful biocontrol agents being studied. The advancements in microbial genetics, microbial community analyses and understanding plant-microbe interactions continue to accumulate and will be instrumental in helping microbial biocontrol of weeds move forward.

The area of biological control using soil microorganisms needs further investigations to discover additional isolate-host pairs that are a biocontrol match consisting of a biocontrol agent of highest virulence in contact with the host at its greatest susceptibility. Formulations are needed to increase shelf life of the living organisms to improve survival and efficacy. Research and development of each biocontrol agent are needed so that stakeholders and industry buy in to the marketing, economics and time investments of this approach to weed management. An understanding of microbial community, weed, and soil quality characteristics, and management practices is needed for the development of weed suppressive soils.

Investigations of the structure and function of soil microbial communities are needed to advance the area of biological control. Traditional techniques of microbial analyses to describe the composition and diversity of microbial populations in soils has commonly relied on phenotypic characteristics alone, and molecular investigations add to the information on structure and function of the soils (Mazzola, 2004). Profiling or fingerprinting of soil and soil microbial community structure using substrate utilization and fatty acid methyl ester analyses may be the first step in targeting weed-suppressive potential. There are several nucleic acid-based methods that can be used to probe soil and identify those microbes that produce similar compounds to those already known. Probes will assist field studies of known agents to follow survival in soils and explore soil for additional weed-suppressive factors. Nucleic acid technologies provide greater information on genetics, and possibly function of a given organism. Array, pyrosequencing and metagenomic investigations can provide information on the microbial community and the biological agent within that community. Selection for hypervirulence; construction of molecular probes; understanding the genetic material of the agent, weed-suppressive compounds, and host-microbe interactions can be investigated more thoroughly with these methods. The continual development of novel molecular methods to investigate genetics of a system will provide key information to better understanding of the plant-microbe phenomena. These methods are forever changing and improving to allow us to have increased knowledge of the microbial portion of the ecosystem and the various interactions that can occur.

Soil microbial ecology and the soil microbial community will affect weed ecosystem dynamics, diversity, function, and populations. As with soil quality, the compilation of indicators has been attempted often to hone in on a few indicators of importance. No one approach or method can be used to characterize and follow biocontrol agents, or to isolate and research additional novel plant-microbe interactions. The future is bright for continued development of microbial herbicides to reduce herbicide reliance and provide multiple options in weed management.

8. Conclusions

The wealth of genetic potential of microorganisms on this earth is boundless. There have been many investigations of potential products for weed management. Some have been successful at suppressing weeds in the field and a select few are marketed products that now reduce weed infestations. Further studies are needed to continue to search for additional tools to combat weeds. Increasing our understanding of plant-microbe interactions will assist in this effort. Biocontrol agents need to be specific, competitive and well-matched with the weed of interest. The search for biocontrol agents from the environment entails not only finding microorganisms that inhibit a weed, but that are specific for the weed or related plant species and have an economically viable market. Host-range testing and non-target species testing are needed early in the process. In addition, the development of formulations and delivery systems is necessary to prolong the shelf-life and efficacy of the biocontrol agents in a variety of environments. Biocontrol should not be considered a stand alone option, but may be best if integrated with other methods of control, especially with those that are ecologically sound. Biocontrol agents to reduce or complement chemical herbicides expand options in weed management and tend toward the

use of ecologically based systems. They add additional tools in the arsenal of weed management efforts. There is a wealth of genetic potential in the soil and the environment to be explored, screened and tested for weed suppression.

9. References

- Abbas, H., Johnson, B., Pantone, D., & Hines, R. (2004). Biological control and use of adjuvants against multiple seeded cocklebur (*Xanthium strumarium*) in comparison with several other cocklebur types. *Biocontrol science and technology*, Vol.14, No.8, (December 2004), pp. 855-860, ISSN 1360-0478
- Abu-Dieyeh, M. & Watson, A. (2006). Effect of turfgrass mowing height on biocontrol of dandelion with *Sclerotinia minor*. *Biocontrol science and technology*, Vol.16, No.5-6, (May 2006), pp. 509-524, ISSN 1360-0478
- Abu-Dieyeh, M. & Watson, A. (2007). Efficacy of *Sclerotinia minor* for dandelion control: effect of dandelion accession, age and grass competition. *Weed research*, Vol.47, No.1, (February 2007), pp. 63-72, ISSN 0043-1737
- Ahonsi, M., Berner, D., Emechebe, A., Lagoke, S., & Sanginga, N. (2003). Potential of ethylene-producing pseudomonads in combination with effective N₂-fixing bradyrhizobial strains as supplements to legume rotation for *Striga hermonthica* control. *Biological control*, Vol.28, No.1, (September 2003), pp.1-10, ISSN 1049-9644
- Ahonsi, M., Berner, D., Emechebe, A., & Lagoke, S. (2002). Selection of rhizobacterial strains for suppression of germination of *Striga hermonthica* (Del.) Benth. seeds. *Biological control*, Vol.24, No.2, (June 2002), pp. 143-152, ISSN 1049-9644
- Amsellem, Z., Zidack, N., Quimby, P. Jr., & Gressel, J. (1999). Long-term dry preservation of viable mycelia of two mycoherbicide organisms. *Crop protection*, Vol.18, No.10, (December 1999), pp. 643-649, ISSN 0261-2194
- Anderson, F., McLaren, D., & Barton, J. (2010). Studies to assess the suitability of *Uromyces penganus* as a biological control agent for *Nassella neesiana* (Poaceae) in Australia and New Zealand. *Australasian plant pathology*, Vol.39, No.1, (January 2010), pp. 69-78, ISSN 1448-6032
- Ash, G., Cother, E., McKenzie, C., & Chung, Y. (2008). A phylogenetic and pathogenic comparison of potential biocontrol agents for weeds in the family Alismataceae from Australia and Korea. *Australasian plant pathology*, Vol.37, No.4, (May 2008), pp. 402-405, ISSN 1448-6032
- Babalola, O., Sanni A., Odhiambo, G., & Torto, B. (2007). Plant growth-promoting rhizobacteria do not pose any deleterious effect on cowpea and detectable amounts of ethylene are produced. *World journal of microbiology & biotechnology*, Vol.23, No.6, (June 2007), pp. 747-752, ISSN 1573-0972
- Bailey, K. (2004). Microbial weed control: an off-beat application of plant pathology. *Canadian journal of plant pathology*, Vol.26, No.3, (July-Sept 2004), pp. 239-244, ISSN 0706-0661
- Bailey, B., Hebbbar, K., Lumsden, R., O'Neill, N., & Lewis, J. (2004). Production of *Pleospora papaveracea* biomass in liquid culture and its infectivity on opium poppy (*Papaver somniferum*). *Weed science*, Vol.52, No.1, (Jan-Feb 2004), pp. 91-97, ISSN 0043-1745
- Bailey, K., Boyetchko, S., & Langle, T. (2010). Social and economic drivers shaping the future of biological control: A Canadian perspective on the factors affecting the

- development and use of microbial biopesticides. *Biological control: theory and application in pest management*, Vol.52, No.3, (March 2010), pp.221-229, ISSN 1049-9644
- Barton, J. (2004). How good are we at predicting the field host-range of fungal pathogens used for classical biological control of weeds? *Biological control: theory and applications in pest management*, Vol.31, No.1, (September 2004), pp. 99-122, ISSN 1049-9644
- Becker, E., Shamoun, S., & Hintz, W. (2005). Efficacy and environmental fate of *Chondrostereum purpureum* used as a biological control for red alder (*Alnus rubra*). *Biological control: theory and application in pest management*. Vol.33, No.3, (June 2005) pp. 269-277, ISSN 1049-9644
- Boari, A. & Vurro, M. (2004). Evaluation of *Fusarium* spp. and other fungi as biological control agents of broomrape (*Orobanche ramosa*). *Biological control: theory and application in pest management*, Vol.30, No.2, (June 2004), pp. 212-219, ISSN 1049-9644
- Boguena, T., Meyer, S., & Nelson, D. (2007). Low temperature during infection limits *Ustilago bullata* (Ustilaginaceae, Ustilaginales) disease incidence on *Bromus tectorum* (Poaceae, Cyperales). *Biocontrol science and technology*, Vol.17, No.1-2, (January 2007), pp. 33-52, ISSN 1360-0478
- Bouizgarne, B., El-Maarouf-Bouteau, H., Madiona, K., Biligui, B., Monestiez, M., Pennarun, A., Amiar, Z., Rona, J., Ouhdouch, Y., El Hadrami, I., & Bouteau, F. (2006). A putative role for fusaric acid in biocontrol of the parasitic angiosperm *Orobanche ramosa*. *Molecular plant-microbe interactions*, Vol.19, No.5, (May 2006), pp. 550-556, ISSN 0894-0282
- Bourdot, G., Baird, D., Hurrell, G., & de Jong, M. (2006a). Safety zones for a *Sclerotinia sclerotiorum*-based mycoherbicide: Accounting for regional and yearly variation in climate. *Biocontrol science and technology*, Vol.16, No.3-4, (February 2006), pp. 345-358, ISSN 1360-0478
- Bourdot, G., Barton, J., Hurrell, G., Gianotti, A., & Saville, D. (2006b). *Chondrostereum purpureum* and *Fusarium tumidum* independently reduce regrowth in gorse (*Ulex europaeus*). *Biocontrol science and technology*, Vol.16, No.3-4, (January 2006), pp. 307-327, ISSN 1360-0478
- Boyette, C. (2006). Adjuvants enhance the biological control potential of an isolate of *Colletotrichum gloeosporioides* for biological control of sicklepod (*Senna obtusifolia*). *Biocontrol science and technology*, Vol.16, No.9-10, (October 2006), pp. 1057-1066, ISSN 1360-0478
- Boyette, C., Reddy, K., & Hoagland, R. (2006). Glyphosate and bioherbicide interaction for controlling kudzu (*Pueraria lobata*), redvine (*Brunnichia ovata*), and trumpet creeper (*Campsis radicans*). *Biocontrol science and technology*, Vol.16, No.9-10, (October 2006), pp. 1067-1077, ISSN 1360-0478
- Boyette, C., Hoagland, R., & Abbas, H. (2007a). Evaluation of the bioherbicide *Myrothecium verrucaria* for weed control in tomato (*Lycopersicon esculentum*). *Biocontrol science and technology*, Vol.17, No.1-2, (February 2007), pp. 171-178, ISSN 1360-0478
- Boyette, C., Hoagland, R., & Weaver, M. (2007b). Effect of row spacing on biological control of sicklepod (*Senna obtusifolia*) with *Colletotrichum gloeosporioides*. *Biocontrol science and technology*, Vol.17, No.9-10, (November 2007), pp. 957-967, ISSN 1360-0478

- Boyette, C., Hoagland, R., & Weaver, M. (2007c). Biocontrol efficacy of *Colletotrichum truncatum* for hemp sesbania (*Sesbania exaltata*) is enhanced with unrefined corn oil and surfactant. *Weed biology and management*, Vol.7, No.1, (March 2007), pp. 70-76, ISSN 1445-6664
- Boyette, C., Hoagland, R., Weaver, M., & Reddy, K. (2008a). Redvine (*Brunnichia ovata*) and trumpet creeper (*Campsis radicans*) controlled under field conditions by a synergistic interaction of the bioherbicide, *Myrothecium verrucaria*, with glyphosate. *Weed biology and management*, Vol.8, No.1, (March 2008), pp. 39-45, ISSN 1444-6162
- Boyette, C., Hoagland, R., & Weaver, M. (2008b) Interaction of a bioherbicide and glyphosate for controlling hemp sesbania in glyphosate-resistant soybean. *Weed biology and management*, Vol.8, No.1, (2008) pp. 18-24, ISSN 1444-6162
- Byer, K., Peng, G., Wolf, T., & Caldwell, B. (2006). Spray retention and its effect on weed control by mycoherbicides. *Biological control: theory and application in pest management*, Vol.37, No.3, (June 2006), pp. 307-313, ISSN 1049-9644
- Caesar, A. (2003). Synergistic interaction of soilborne plant pathogens and root-attacking insects in classical biological control of an exotic rangeland weed. *Biological control: theory and application in pest management*, Vol.28, No.1, (September 2003), pp. 144-153, ISSN 1049-9644
- Caesar, A. (2006). *Uromyces scutellatus* as a keystone species affecting *Euphorbia* spp. in Europe as shown by effects on density in the field. *Biocontrol science and technology*, Vol.16, No.9-10, (October 2006), pp. 1079-1086, ISSN 1360-0478
- Caesar, A., Caesar, T., & Maathuis, M. (2010). Pathogenicity, characterization and comparative virulence of *Rhizoctonia* spp. from insect-galled roots of *Lepidium draba* in Europe. *Biological control: theory and application in pest management*, Vol.52, No.2, (February 2010), pp. 140-144, ISSN 1049-9644
- Caesar, A., Campobasso, G., & Terragitti, G. (2002). Identification, pathogenicity and comparative virulence of *Fusarium* spp. associated with insect-damaged, diseased *Centaurea* spp. in Europe. *Biocontrol science and technology*, Vol.47, No.2, (April 2002), pp. 217-229, ISSN 1386-6141
- Carsten, L., Maxwell, B., Johnston, M., & Sands, D. (2001). Impact of crown rust (*Puccinia coronata* f. sp. *avenae*) on competitive interactions between wild oats (*Avena fatua*) and stipa (*Nassella pulchra*). *Biological control: theory and applications in pest management*, Vol.22, No.3, (November 2001), pp. 207-218, ISSN 1049-9644
- Casella, F., Charudattan, R., & Vurro, M. (2010). Effectiveness and technological feasibility of bioherbicide candidates for biocontrol of green foxtail (*Setaria viridis*). *Biocontrol science and technology*, Vol.20, No.9-10, (October 2010), pp. 1027-1045, ISSN 1360-0478
- Chandramohan, S. & Charudattan, R. (2003). A multiple-pathogen system for bioherbicidal control of several weeds. *Biocontrol science and technology*, Vol.13, No.2, (March 2003), pp. 199-205, ISSN 1360-0478
- Charudattan, R. (2005). Ecological, practical, and political inputs into selection of weed targets: what makes a good biological control target? *Biological control: theory and application in pest management*, Vol.35, No.3, (December 2005), pp. 183-196, ISSN 1049-9644

- Chittick, A. & Auld, B. (2001). Polymers in bioherbicide formulation: *Xanthium spinosum* and *Colletotrichum orbiculare* as a model system. *Biocontrol science and technology*, Vol.11, No.6, (December 2001), pp. 691-702, ISSN 1360-0478
- Cliquet, S. & Zeeshan, K. (2008). Impact of nutritional conditions on yields, germination rate and shelf-life of *Plectosporium alismatis* conidia and chlamydospores as potential candidates for the development of a mycoherbicide of weeds in rice crops. *Biocontrol science and technology*, Vol.18, No.7-8, (September 2008), pp. 685-695, ISSN 1360-0478
- Cohen, B., Amsellem, Z., Maor, R., Sharon, A., & Gressel, J. (2002). Transgenically enhanced expression of indole-3-acetic acid confers hypervirulence to plant pathogens. *Phytopathology*, Vol.92, No.6, (June 2002), pp.590-596, ISSN 0031-949X
- Cook, J., Charudattan, R., Zimmerman, T., Rosskopf, E., Stall, W., & MacDonald, G. (2009). Effects of *Alternaria destruens*, glyphosate, and ammonium sulfate individually and integrated for control of dodder (*Cuscuta pentagona*). *Weed technology*, Vol.23, No.4, (October-December 2009), pp. 550-555, ISSN 1550-2740
- Cross, J. & Polonenko, D. (1996). An industry perspective on registration and commercialization of biocontrol agents in Canada. *Canadian journal of plant pathology*, Vol.18, No.4, (December 1996), pp. 446-454, ISSN 0706-0661
- de Luna, L., Kennedy, A., Hansen, J., Paulitz, T., Gallagher, R. & Fuerst, E. (2011). Mycobiota on wild oat (*Avena fatua* L.) seed and their caryopsis decay potential. *Plant health progress*, Vol.10, No.1, (February 2011), pp.1-8, ISSN 1535-1025
- de Luna, L., Stubbs, T., Kennedy, A., & Kremer, R. (2005). Deleterious bacteria in the rhizosphere. In: *Roots and Soil Management: Interactions between Roots and the Soil*, R. Zobel and S. Wright, (Eds.), 233-261. Monograph no. 48. ISBN 978-0-89118-159-0 ASA. Madison WI.
- Denoth, M., Frid, L., & Myers, J. (2002). Multiple agents in biological control: improving the odds? *Biological control: theory and applications in pest management*, Vol.24, No.1, (May 2002), pp. 20-39, ISSN 1049-9644
- Den Breeyen, A. & Morris, M. (2003). Pathogenicity and host specificity of *Mycovellosiella lantanae* var. *lantanae*, a potential biocontrol agent for *Lantana camara* in South Africa. *Biocontrol science and technology*, Vol.13, No.3, (May 2003) pp. 313-322, ISSN 0958-3157
- Ding, J., Wu, Y., Zheng, H., Fu, W., Reardon, R., & Liu, M. (2006). Assessing potential biological control of the invasive plant, tree-of-heaven, *Ailanthus altissima*. *Biocontrol science and technology*, Vol.16, No.5-6, (May 2006), pp. 547-566, ISSN 1360-0478
- Dooley, S. & Beckstead, J. (2010). Characterizing the interaction between a fungal seed pathogen and a deleterious rhizobacterium for biological control of cheatgrass. *Biological control: theory and application in pest management*, Vol.53, No.1, (January 2010), pp. 197-203, ISSN 1049-9644
- Dor, E. & Hershenhorn, J. (2009). Evaluation of the pathogenicity of microorganisms isolated from Egyptian broomrape (*Orobancha aegyptiaca*) in Israel. *Weed biology and management*, Vol.9, No.3, (September 2009), pp. 200-208, ISSN 1445-6664
- Doty, J., Kennedy, A., & Pan, W. (1994). A rapid bioassay for inhibitory rhizobacteria using digital image analysis. *Soil science society of America journal*, Vol.58, No.6, (November 1994), pp.1699-1701, ISSN 1435-0661

- Elzein, A., Heller, A., Ndambi, B., De Mol, M., Kroschel, J., & Cadisch, G. (2010). Cytological investigations on colonization of sorghum roots by the mycoherbicide *Fusarium oxysporum* f. sp. *strigae* and its implications for Striga control using a seed treatment delivery system. *Biological control: theory and application in pest management*, Vol.53, No.3, (June 2010), pp. 249-257, ISSN 1049-9644
- Elzein, A., Kroschel, J., Marley, P., & Cadisch, G. (2009). Does vacuum-packaging or co-delivered amendments enhance shelf-life of Striga-mycoherbicide products containing *Fusarium oxysporum* f. sp. *strigae* during storage? *Biocontrol science and technology*, Vol.19, No.3-4, (March 2009), pp. 349-367, ISSN 1360-0478
- Elzein, A., Kroschel, J., & Leth, V. (2006). Seed treatment technology: An attractive delivery system for controlling root parasitic weed *Striga* with mycoherbicide. *Biocontrol science and technology*, Vol.16, No.1-2, (February 2006), pp. 3-26, ISSN 1360-0478
- Elzein, A., Kroschel, J., & Müller-Stöver, D. (2004). Optimization of storage conditions for adequate shelf-life of 'Pesta' formulation of *Fusarium oxysporum* 'Foxy 2', a potential mycoherbicide for Striga: effects of temperature, granule size and water activity. *Biocontrol science and technology*, Vol.14, No.6, (September 2004), pp. 545-559, ISSN 1360-0478
- EPA (2011). Microbial pesticide test guidelines, Series 885, Environmental Protection Agency Chemical Safety and Pollution Prevention, OCSPP, Harmonized test guideline series, http://www.epa.gov/ocspp/pubs/frs/publications/Test_Guidelines/series885.htm
- Evidente, A., Punzo, B., Andolfi, A., Berestetskiy, A., & Motta, A. (2009a). Alternethanoxins A and B, polycyclic ethanones produced by *Alternaria sonchi*, potential mycoherbicides for *Sonchus arvensis* biocontrol. *Journal of agricultural and food chemistry*, Vol.57, No.15, (August 2009), pp. 6656-6660, ISSN 0021-8561
- Evidente, A., Berestetskiy, A., Cimmino, A., Tuzi, A., Superchi, S., Melck, D., & Andolfi, A. (2009b). Papyracillic acid, a phytotoxic 1,6-dioxaspiro(4,4)nonene produced by *Ascochyta agropyrina* var. *nana*, a potential mycoherbicide for *Elytrigia repens*. *Biocontrol*, Vol.57, No.23, (December 2009), pp. 11168-11173, ISSN 0021-8561
- Evidente, A., Cimmino, A., Andolfi, A., Vurro, M., Zonno, M., & Motta, A. (2008). Phyllostoxin and phyllostin, bioactive metabolites produced by *Phyllosticta cirsii*, a potential mycoherbicide for *Cirsium arvense* biocontrol. *Journal of agricultural and food chemistry*, Vol.56, No.3, (February 2008), pp. 884-888, ISSN 0021-8561
- Flores-Vargas, R., & O'Hara, G. (2006). Isolation and characterization of rhizosphere bacteria with potential for biological control of weeds in vineyards. *Journal of applied microbiology*, Vol.100, No.5, (May 2006), pp. 946-954, ISSN 0973-7510
- Frantzen, J. & Müller-Scharer, H. (2006). Modeling the impact of a biocontrol agent, *Puccinia lagenophorae*, on interactions between a crop, *Daucus carota*, and a weed, *Senecio vulgaris*. *Biological control: theory and application in pest management*, Vol.37, No.3, (June 2006), pp. 301-306, ISSN 1049-9644
- Ghajar, F., Holford, P., Alhussaen, K., Beattie, A., & Cother, E. (2006). Optimising sporulation and virulence in *Drechslera avenacea*. *Biocontrol science and technology*, Vol.16, No.5-6, (May 2006), pp. 471-484, ISSN 1360-0478Z
- Ghorbani, R., Scheepens, P., Zweerde, W., Leifert, C., McDonald, A., & Seel, W. (2002). Effects of nitrogen availability and spore concentration on the biocontrol activity of

- Ascochyta caulina* in common lambsquarters (*Chenopodium album*). *Weed science*, Vol.50, No.5, (Sept-Oct 2002), pp. 628-633, ISSN 0043-1745
- Ghosheh, H. (2005). Constraints in implementing biological weed control: A review. *Weed biology and management*, Vol.5, No.3, (September 2005), pp.83-92, ISSN 1444-6162
- Graham, G., Peng, G., Bailey, K., & Holm, F. (2006). Effect of dew temperature, post-inoculation condition, and pathogen dose on suppression of scentless chamomile by *Colletotrichum truncatum*. *Biocontrol science and technology*, Vol.16, No.3-4, (February 2006), pp. 271-280, ISSN 1360-0478
- Graham, G., Peng, G., Bailey, K., & Holm, F. (2007). Effect of plant stage, *Colletotrichum truncatum* dose, and use of herbicide on control of *Matricaria perforata*. *BioControl*, Vol.52, No.4, (August 2007), pp. 573-589, ISSN 1386-6141
- Green, S., Peng, G., Connolly, T., & Boyetchko, S. (2004). Effect of moisture and temperature on disease of green foxtail caused by *Drechslera gigantea* and *Pyricularia setariae*. *Plant disease*, Vol.88, No.6, (June 2004), pp. 605-612, ISSN 0191-2917
- Gressel, J. (2010). Herbicides as synergists for mycoherbicides, and vice versa. *Weed science*, Vol.58, No.3, (July 2010), pp. 324-328, ISSN 0043-1745
- Gronwald, J., Plaisance, K., Ide, D., & Wyse, D. (2002). Assessment of *Pseudomonas syringae* pv. *tagetis* as a biocontrol agent for Canada thistle. *Weed science*, Vol.50 No.3, (May-June 2002), pp. 397-404, ISSN 0043-1745
- Gurusiddaiah, S., Gealy, D., Kennedy, A., & Ogg, A., Jr. (1994). Isolation and characterization of metabolites from *Pseudomonas fluorescens* strain D7 for control of downy brome (*Bromus tectorum* L.). *Weed science*, Vol.42, No.3, (July - September 1994), pp. 492-501, ISSN 0043-1745
- Guske, S., Schulz, B., & Boyle, C. (2004). Biocontrol options for *Cirsium arvense* with indigenous fungal pathogens. *Weed research*, Vol.44, No.2, (April 2004), pp. 107-116, ISSN 0043-1737
- Hasan, S., Sobhian, R., & Herard, F. (2001). Biology, impact and preliminary host-specificity testing of the rust fungus, *Uromyces salsolae*, a potential biological control agent for *Salsola kali* in the USA. *Biocontrol science and technology*, Vol.11, No.6, (December 2001), pp. 677-689, ISSN 1360-0478
- Hoagland, R., Boyette, C., & Abbas, H. (2007). *Myrothecium verrucaria* isolates and formulations as bioherbicide agents for kudzu. *Biocontrol science and technology*, Vol.17, No.7-8, (September 2007), pp. 721-731, ISSN 1360-0478
- Hurrell, G., Bourdot, G., & Saville, D. (2001). Effect of application time on the efficacy of *Sclerotinia sclerotiorum* as a mycoherbicide for *Cirsium arvense* control in pasture. *Biocontrol science and technology*, Vol.11, No.3, (June 2001), pp. 317-330, ISSN 1360-0478
- Imaizumi, S., Honda, M., & Fujimori, T. (1999) Effect of temperature on the control of annual bluegrass (*Poa annua* L.) with *Xanthomonas campestris* pv. *poae* (JT-P482). *Biological control*, Vol.16, No.1, (September 1999), pp.13-17, ISSN 1049-9644
- Jahromi, F. (2007). Effect of environmental factors on disease development caused by the fungal pathogen *Plectosporium alismatis* on the floating-leaf stage of starfruit (*Damasonium minus*), a weed of rice. *Biocontrol science and technology*, Vol.17, No.7-8, (September 2007), pp. 871-877, ISSN 1360-0478
- Jahromi, F., Van De Ven, R., Cother, E., & Ash, G. (2006). The interaction between *Plectosporium alismatis* and sublethal doses of bensulfuron-methyl reduces the

- growth of starfruit (*Damasonium minus*) in rice. *Biocontrol science and technology*, Vol.16, No.9-10, (October 2006), pp. 929-940, ISSN 1360-0478
- Jiang, S.-J., Qiang, S., Zhu, Y.-Z., & Dong, Y.-F. (2008). Isolation and phytotoxicity of a metabolite from *Curvularia eragrostidis* and characterisation of its modes of action. *Annals of applied biology*, Vol.152, No.1, (February 2008), pp. 103-111, ISSN 1744-7348
- Jones, W. & Sforza, R. (2007). The European Biological Control Laboratory: an existing infrastructure for biological control of weeds in Europe. *EPPO bulletin*, Vol.37, No.1, (April 2007), pp. 163-165, ISSN 0250-8052
- Kennedy, A. & Papendick, R. (1995). Microbial characteristics of soil quality. *Journal of Soil & Water Conservation*, Vol.50, No.3, (May-June 1995), pp. 243-248, ISSN 002-4561
- Kennedy, A. & Schillinger, W. (2006). Soil quality and water intake in conventional-till vs. no-till paired farms in Washington's Palouse region. *Soil science society of America journal*, Vol.70, No.3, (May-June 2006), pp.940-949, ISSN 0361-5995
- Kennedy, A. & Stubbs, T. (2007). Management effects on the incidence of jointed goatgrass inhibitory rhizobacteria. *Biological control: theory and application in pest management*, Vol.40, No.2, (February 2007) pp.213-221, ISSN 1049-9644
- Kennedy, A., Elliott, L., Young, F., & Douglas, C. (1991). Rhizobacteria suppressive to the weed downy brome. *Soil science society of America journal*, Vol.55, No.3, (May-June 1991). pp.722-727, ISSN 1435-0661
- Kennedy, A., Johnson, B., & Stubbs, T. (2001). Host range of a deleterious rhizobacterium for biological control of downy brome. *Weed science*, Vol.49, No.6, (Nov-Dec 2001), pp. 792-797, ISSN 0043-1745
- Kleczewski, N. & Flory, S. (2010). Leaf blight disease on the invasive grass *Microstegium vimineum* caused by a *Bipolaris* sp. *Plant disease*, Vol.94, No.7, (July 2010), pp. 807-811, ISSN 0191-2917
- Kohlschmid, E., Sauerborn, J., & Müller-Stöver, D. (2009). Impact of *Fusarium oxysporum* on the holoparasitic weed *Phelipanche ramosa*: biocontrol efficacy under field-grown conditions. *Weed research*, Vol.49, No.1, (November 2009), pp. 56-65, ISSN 0043-1737
- Kremer, R. & Souissi, T. (2001). Cyanide production by rhizobacteria and potential for suppression of weed seedling growth. *Current microbiology*, Vol. 43, No.3, (September 2001), pp. 182-186, ISSN 0343-8651
- Kremer, R., & Li, J. (2003). Developing weed-suppressive soils through improved soil quality management. *Soil & tillage research*, Vol.72, No.2, (August 2003), pp. 193-202, ISSN 0167-1987
- Kremer, R. & Schulte, L. (1989). Influence of chemical treatment and *Fusarium oxysporum* on velvetleaf (*Abutilon theophrasti*). *Weed technology*. Vol.3, No.2, (April 1989), pp. 369-374, ISSN 0890-037X
- Kremer, R., Caesar, A., & Souissi, T. (2006). Soilborne microorganisms of Euphorbia are potential biological control agents of the invasive weed leafy spurge. *Applied soil ecology*, Vol.32, No.1, (May 2006), pp. 27-37, ISSN 1744-7348
- Kropp, B. & Darrow, H. (2006). The effect of surfactants and some herbicides on teliospore viability in *Puccinia thlaspeos* (Schub.). *Crop protection*, Vol.25, No.4, (April 2006), pp. 369-374, ISSN 0261-2194
- Kropp, B., Hansen, D., & Thomson, S. (2002). Establishment and dispersal of *Puccinia thlaspeos* in field populations of Dyer's woad. *Plant disease*, Vol.86, No.3, (March 2002), pp. 241-246, ISSN 0191-2917

- Kroschel, J. & Elzein, A. (2004). Bioherbicidal effect of fumonisin B1, a phytotoxic metabolite naturally produced by *Fusarium nygamai*, on parasitic weeds of the genus *Striga*. *Biocontrol science and technology*, Vol.14, No.2, (March 2004), pp. 117-128, ISSN 1360-0478
- Lawrie, J., Greaves, M., Down, V., Morales-Aza, B., & Lewis, J. (2002a). Outdoor studies of the efficacy of *Alternaria alternata* in controlling *Amaranthus retroflexus*. *Biocontrol science and technology*, Vol.12, No.1, (February 2002), pp. 83-94, ISSN 1360-0478
- Lawrie, J., Greaves, M., Down, V., & Western, N. (2002b). Studies of spray application of microbial herbicides in relation to conidial propagule content of spray droplets and retention on target. *Biocontrol science and technology*, Vol.12, No.1, (February 2002), pp. 107-119, ISSN 1360-0478
- Leth, V., Netland, J., & Andreasen, C. (2008). *Phomopsis cirsi*: a potential biocontrol agent of *Cirsium arvense*. *Weed research*, Vol.48, No.6, (December 2008), pp. 533-541, ISSN 0043-1737
- Li, J. & Kremer, R. (2006). Growth response of weed and crop seedlings to deleterious rhizobacteria. *Biological control: theory and application in pest management*, Vol.39, No.1, (October 2006), pp. 58-65, ISSN 1049-9644
- Li, Y., Sun, Z., Zhuang, X., Xu, L., Chen, S., & Li, M. (2003). Research progress on microbial herbicides. *Crop protection*, Vol.22, No.2 (March 2003), pp. 247-252, ISSN 0261-2194
- Lupwayi, N., Rice, W., & Clayton, G. (1998). Soil microbial diversity and community structure under wheat as influenced by tillage and crop rotation. *Soil biology & biochemistry*, Vol.30, No.13 (November 1998). pp. 1733-1741, ISSN 0038-0717
- Magani, I., Ibrahim, A., & Avav, T. (2009). *Fusarium oxysporum* and post-emergence herbicide for the control of the parasitic plant *Striga hermonthica* in maize. *Biocontrol science and technology*, Vol.19, No.9-10, (October 2009), pp. 1023-1032, ISSN 1360-0478
- Magdoff, F. (2001). Concept, components, and strategies of soil health in agroecosystems. *Journal of nematology*. Vol. 33, No.4, (December 2001), pp. 169-172, ISSN 0022-300X
- Maxwell, A. & Scott, J. (2008). Pathogens on wild radish, *Raphanus raphanistrum* (Brassicaceae), in south-western Australia - implications for biological control. *Australasian Plant Pathology*, Vol.37, No.5, (2008), pp. 523-533, ISSN 1448-6032
- Mazzola, M. (2004). Assessment and management of soil microbial community structure for disease suppression. *Annual Review of Phytopathology*, 2004. Vol.42, No.1 (March 2004), pp. 35-59, ISSN 0066-4286
- Medd, R. & Campbell, M. (2005). Grass seed infection following inundation with *Pyrenophora semeniperda*. *Biocontrol science and technology*, Vol.15, No.1, (February 2005), pp. 21-36, ISSN 1360-0478
- Mejri, D., Souissi, T., & Berner, D. (2010a). Evaluation of *Puccinia carduorum* for biological control of *Carduus pycnocephalus* in Tunisia. *Biocontrol science and technology*, Vol.20, No.7-8, (August 2010), pp. 787-790, ISSN 1360-0478
- Mejri, D., Gamalero, E., Tombolini, R., Musso, C., Massa, N., Berta, G., & Souissi, T. (2010b). Biological control of great brome (*Bromus diandrus*) in durum wheat (*Triticum durum*): specificity, physiological traits and impact on plant growth and root architecture of the fluorescent pseudomonad strain X33d. *Biocontrol*, Vol.55, No.4, (August 2010), pp. 561-572, ISSN 1386-6141

- Meyer, S., Nelson, D., & Clement, S. (2001). Evidence for resistance polymorphism in the *Bromus tectorum-Ustilago bullata* pathosystem: implications for biocontrol. *Canadian journal of plant pathology*, Vol.23, No.1, (March 2001), pp. 19-27, ISSN 0706-0661
- Meyer, S., Quinney, D., Nelson, D., & Weaver, J. (2007). Impact of the pathogen *Pyrenophora semeniperda* on *Bromus tectorum* seedbank dynamics in North American cold deserts. *Weed research*, Vol.47, No.1, (February 2007), pp. 54-62, ISSN 0043-1737
- Mohan Babu, R., Sajeena, A., Seetharaman, K., Vidhyasekaran, P., Rangasamy, P., Som Prakash, M., Senthil Raja, A., & Biji, K. (2002). Host range of *Alternaria alternata*--a potential fungal biocontrol agent for water hyacinth in India. *Crop protection*, Vol.21, No.10, (December 2002), pp. 1083-1085, ISSN 0261-2194
- Mortensen, K. (1998). Biological control of weeds using microorganisms. In: *Plant-microbe interactions and biological control*, G. Boland and L. Kuykendall, (Eds.), 223-248, New York, Marcel Dekker, ISBN 0824700430 New York, NY
- Müller-Stöver, D., Kohlschmid, E., & Sauerborn, J. (2009). A novel strain of *Fusarium oxysporum* from Germany and its potential for biocontrol of *Orobanche ramosa*. *Weed research*, Vol.49, No.2, (April 2009), pp. 175-182, ISSN 0043-1737
- Müller-Stöver, D. & Sauerborn, J. (2007). A commercial iron fertilizer increases the survival of *Fusarium oxysporum* f. sp. *orthoceras* propagules in a wheat flour-kaolin formulation. *Biocontrol science and technology*, Vol.17, No.5-6, (May 2007), pp. 597-604, ISSN 1360-0478
- Müller-Stöver, D., Buschmann, H., & Sauerborn, J. (2005). Increasing control reliability of *Orobanche cumana* through integration of a biocontrol agent with a resistance-inducing chemical. *European journal of plant pathology*, Vol.111, No.3, (March 2005), pp. 193-202, ISSN 0929-1873
- Nechet, K. de L., Barreto, R., & Mizubuti, E. (2004). *Sphaceloma poinsettiae* as a potential biological control agent for wild poinsettia (*Euphorbia heterophylla*). *Biological control: theory and applications in pest management*, Vol.30, No.3, (July 2004), pp. 556-565, ISSN 1049-9644
- Owen, A. & Zdor, R. (2001). Effect of cyanogenic rhizobacteria on the growth of velvetleaf (*Abutilon theophrasti*) and corn (*Zea mays*) in autoclaved soil and the influence of supplemental glycine. *Soil biology & biochemistry*, Vol.33, No.6, (May 2001), pp. 801-809, ISSN 0038-0717
- Pedras, M., Ismail, N., Quail, J., & Boyetchko, S. (2003). Structure, chemistry, and biological activity of pseudophomins A and B, new cyclic lipodepsipeptides isolated from the biocontrol bacterium *Pseudomonas fluorescens*. *Phytochemistry*, Vol.62, No.7, (April 2003), pp. 1105-1114, ISSN 0031-9422
- Peng, G. & Boyetchko, S. (2006). Effect of variable dew temperatures on infection of green foxtail by *Pyricularia setariae*, *Drechslera gigantea*, and *Exserohilum rostratum*. *Biological control: theory and application in pest management*, Vol.39, No.3, (December 2006), pp. 539-546, ISSN 1049-9644
- Peng, G. & Byer, K. (2005). Interactions of *Pyricularia setariae* with herbicides for control of green foxtail (*Setaria viridis*). *Weed technology*, Vol.19, No.3, (July-September 2005), pp. 589-598, ISSN 550-2740
- Peng, G., Bailey, K., Hinz, H. & Byer, K. (2005a). *Colletotrichum* sp: a potential candidate for biocontrol of scentless chamomile (*Matricaria perforata*) in western Canada.

- Biocontrol science and technology*, Vol.15, No.5, (August 2005), pp. 497-511, ISSN 1360-0478
- Peng, G., Wolf, T., Byer, K., & Caldwell, B. (2005b). Spray retention on green foxtail (*Setaria viridis*) and its effect on weed control efficacy by *Pyricularia setariae*. *Weed technology*, Vol.19, No.1, (January 2005), pp. 86-93, ISSN 550-2740
- Pereira, J., Barreto, R., Ellison, C., & Maffia, L. (2003). *Corynespora cassiicola* f. sp. *lantanae*: a potential biocontrol agent from Brazil for *Lantana camara*. *Biological control: theory and applications in pest management*, Vol.26, No.1, (January 2003), pp. 21-31, ISSN 1049-9644
- Pomella, A., Barreto, R., & Charudattan, R. (2007). *Nimbya alternantherae* a potential biocontrol agent for alligatorweed, *Alternanthera philoxeroides*. *Biocontrol journal of the international organization for biological control*, Vol.52, No.2, (April 2007), pp. 271-288, ISSN 1386-6141
- Qiang, S., Zhu, Y., Summerell, B., & Li, Y. (2006). Mycelium of *Alternaria alternata* as a potential biological control agent for *Eupatorium adenophorum*. *Biocontrol science and technology*, Vol.16, No.7-8, (August 2006), pp. 653-668, ISSN 1360-0478
- Ray, P., Kumar, S., & Pandey, A. (2008). Deleterious effect of herbicides on water hyacinth biocontrol agents *Neochetina bruchi* and *Alternaria alternata*. *Biocontrol science and technology*, Vol.18, No.5-6, (May 2008), pp. 517-526, ISSN 1360-0478
- Rayamajhi, M., Pratt, P., Center T., & Van, T. (2010). Insects and a pathogen suppress *Melaleuca quinquenervia* cut-stump regrowth in Florida. *Biological control: theory and application in pest management*, Vol.53, No.1, (April 2010), pp.1-8, ISSN 1049-9644
- Rietman, L., Shamoun, S., & van der Kamp, B. (2005). Assessment of *Neonectria neomacrospora* (anamorph *Cylindrocarpon cylindroides*) as an inundative biocontrol agent against hemlock dwarf mistletoe. *Canadian journal of plant pathology*, Vol.27, No.4, (Oct-Dec 2005), pp. 603-609, ISSN 0706-0661
- Roskopf, E., Yandoc, C., & Charudattan, R. (2006). Genus-specific host range of *Phomopsis amaranthicola* (Sphaeropsidales), a bioherbicide agent for *Amaranthus* spp. *Biocontrol science and technology*, Vol.16, No.1-2, (February 2006), pp. 27-35, ISSN 1360-0478
- Ryan, M., Mirsky, S., Mortensen D., Teasdale, J., & Curran, W. (2011). Potential synergistic effects of cereal rye biomass and soybean planting density on weed suppression. *Weed science*, Vol.59, No.2, pp. 238-246, ISSN 0043-1745
- Sands, D. & Pilgeram, A. (2009). Methods for selecting hypervirulent biocontrol agents of weeds: why and how. *Pest management science*, Vol.65, No.5, (May 2009), pp. 581-587, ISSN 1049-9644
- Sauerborn, J., Müller-Stöver, D., & Hershenhorn, J. (2007). The role of biological control in managing parasitic weeds. *Crop protection*, Vol.26, No.3, (March 2007), pp. 246-254, ISSN 0261-2194
- Schall, M. & Davis, D. (2009). *Ailanthus altissima* wilt and mortality: etiology. *Plant disease*, Vol.93, No.7, (July 2009), pp. 747-751, ISSN 0191-2917
- Seixas, C., Barreto, R., Freitas, L., Maffia, L., & Monteiro, F. (2004). *Ditylenchus drepanocercus* (Nematoda), a potential biological control agent for *Miconia calvoescens* (Melastomataceae): host-specificity and epidemiology. *Biological control: theory and applications in pest management*, Vol.31, No.1, (September 2004), pp. 29-37, ISSN 1049-9644


- Shabana, Y. & Mohamed, Z. (2005). Integrated control of water hyacinth with a mycoherbicide and phenylpropanoid pathway inhibitor. *Biocontrol science and technology*, Vol.15, No.7, (November 2005), pp. 659-669, ISSN 1360-0478
- Shabana, Y. (2005). The use of oil emulsions for improving the efficacy of *Alternaria eichhorniae* as a mycoherbicide for water hyacinth (*Eichhornia crassipes*). *Biological control: theory and application in pest management*, Vol.32, No.1, (January 2005), pp. 78-89, ISSN 1049-9644
- Shabana, Y., Cuda, J., & Charudattan, R. (2003). Evaluation of pathogens as potential biocontrol agents of hydrilla. *Journal of phytopathology*, Vol.151, No.11-12, (Nov.-Dec 2003), pp. 607-613, ISSN 0973-7510
- Sharma, P., Rai, P., Siddiqui, S., & Chauhan, J. (2011). First report of *Fusarium* wilt in the broomrape parasite growing on *Brassica* spp. in India. *Plant disease: an international journal of applied plant pathology*, Vol.95, No.1, (January 2011), pp. 75-95, ISSN 0191-2917
- Shearer, J. & Jackson, M. (2006). Liquid culturing of microsclerotia of *Mycocleptodiscus terrestris*, a potential biological control agent for the management of hydrilla. *Biological control: theory and application in pest management*, Vol.38, No.3, (September 2006), pp. 298-306, ISSN 1049-9644
- Shetty, K., Minnis, A., Rossman, A., & Jayachandran, K. (2011). The Brazilian peppertree seed-borne pathogen, *Neofusicoccum batangarum*, a potential biocontrol agent. *Biological control: theory and application in pest management*. Vol.56, No.1, (January 2011), pp. 91-97, ISSN 1049-9644
- Souissi, T., & Kremer, R. (1998). A rapid microplate callus bioassay for assessment of rhizobacteria for biocontrol of leafy spurge (*Euphorbia esula* L.). *Biocontrol science and technology*. Vol.8, No.1, (January 1998), pp. 83-92, ISSN 0958-3157
- Stewart-Wade, S. & Boland, G. (2004). Selected cultural and environmental parameters influence disease severity of dandelion caused by the potential bioherbicidal fungi, *Phoma herbarum* and *Phoma exigua*. *Biocontrol science and technology*, Vol.14, No.6, (September 2004), pp. 561-569, ISSN 1360-0478
- Teshler, M., Ash, G., Zolotarov, Y., & Watson, A. (2007). Increased shelf life of a bioherbicide through combining modified atmosphere packaging and low temperatures. *Biocontrol science and technology*, Vol.17, No.3-4, (March 2007), pp. 387-400, ISSN 1360-0478
- Tessmann, D., Charudattan, R., & Preston, J. (2008). Variability in aggressiveness, cultural characteristics, cercosporin production and fatty acid profile of *Cercospora piaropi*, a biocontrol agent of water hyacinth. *Plant pathology*, Vol.57, No.5, (October 2008), pp. 957-966, ISSN 0032-08
- Tichich, R. & Doll, J. (2006). Field-based evaluation of a novel approach for infecting Canada thistle (*Cirsium arvense*) with *Pseudomonas syringae* pv. *tagetis*. *Weed science*, Vol.54, No.1, (January 2006), pp. 166-171, ISSN 0043-1745
- Tichich, R., Doll, J., & McManus, P. (2006). *Pseudomonas syringae* pv. *tagetis* population dynamics both on and in Canada thistle (*Cirsium arvense*) leaves as affected by rain events. *Weed science*, Vol.54, No.5, (September 2006), pp. 934-940, ISSN 0043-1745
- Tilley, A. & Walker, H. (2002). Evaluation of *Curvularia intermedia* (*Cochliobolus intermedius*) as a potential microbial herbicide for large crabgrass (*Digitaria sanguinalis*).

- Biological control: theory and applications in pest management*, Vol.25, No.1, (September 2002), pp. 12-21, ISSN 1360-0478
- Tiourebaev, K., Semenchenko, G., Dolgovskaya, M., McCarthy, M., Anderson, T., Carsten, L., Pilgeram, A., & Sands, D. (2001). Biological control of infestations of ditchweed (*Cannabis sativa*) with *Fusarium oxysporum* f.sp. *cannabis* in Kazakhstan. *Biocontrol science and technology*, Vol.11, No.4, (August 2001), pp. 535-540, ISSN 1360-0478
- United States Environmental Protection Agency (2005)
http://www.epa.gov/oppbppd1/biopesticides/ingredients/factsheets/factsheet_028301.htm
- Van Klinken, R. & Julien, M. (2003). Learning from past attempts: does classical biological control of Noogoora burr (Asteraceae: *Xanthium occidentale*) have a promising future? *Biocontrol science and technology*, Vol.13, No.2, (March, 2003), pp. 139-153, ISSN 1360-0478
- Venne, J., Beed, F., Avocanh, A., & Watson, A. (2009). Integrating *Fusarium oxysporum* f. sp. *strigae* into cereal cropping systems in Africa. *Pest management science*, Vol.65, No.5, (May 2009), pp. 572-580, ISSN 1526-498X
- Vidal, K., Guermache, F., & Widmer, T. (2004). In vitro culturing of yellow starthistle (*Centaurea solstitialis*) for screening biological control agents. *Biological control: theory and application in pest management*, Vol.30, No.2, (June 2004), pp. 330-335, ISSN 1049-9644
- Vogelgsang, S. & Shamoun, S. (2004). Evaluation of an inoculum production and delivery technique for *Valdensinia heterodoxa*, a potential biological control agent for salal. *Biocontrol science and technology*, Vol.14, No.8, (December 2004), pp. 747-756, ISSN 1360-0478
- Vurro, M., Zonno, M., Evidente, A., Andolfi, A., & Montemurro, P. (2001). Enhancement of efficacy of *Ascochyta caulina* to control *Chenopodium album* by use of phytotoxins and reduced rates of herbicides. *Biological control: theory and applications in pest management*, Vol.21, No.2, (June 2001), pp. 182-190, ISSN 1049-9644
- Wandeler, H., Nentwig, W., & Bacher, S. (2008). Establishing systemic rust infections in *Cirsium arvense* in the field. *Biocontrol science and technology*, Vol.18, No.1-2, (February 2008), pp. 209-214, ISSN 1360-0478
- Wander, M., Hedrick, D., Kaufman, D., Traina, S., Stinner, B., Kehrmeier, S., & White, D. (1995). The functional significance of the microbial biomass in organic and conventionally managed soils. In: *The significance and regulation of soil biodiversity*. H.P. Collins, G.P. Robertson, M.J. Klug (Eds). pp.87-97, Kluwer Academic Publishers, Netherlands, ISBN 079-233-1389
- Wapshere, A. (1974). A strategy for evaluating the safety of organisms for biological weed control. *Annals of applied biology*, Vol.77, No.2, (July 1974), pp. 201-211. ISSN 0003-4746
- Ward, S., Reid, S., Harrington, J., Sutton, J., & Beck, K. (2008). Genetic variation in invasive populations of yellow toadflax (*Linaria vulgaris*) in the Western United States. *Weed science*, Vol.56, No.3, (May 2008), pp. 394-399, ISSN 0043-1745
- Wilkin, J., Shamoun, S., Ritland, C., Ritland, K., & El-Kassaby, Y. (2005). Genetic diversity and population structure of *Valdensinia heterodoxa*, a potential biocontrol agent for salal in coastal British Columbia. *Canadian journal of plant pathology*, Vol.27, No.4, (Oct-Dec 2005), pp. 559-571, ISSN 0706-0661

- Wood, A. & Crous, P. (2005). Epidemic increase of *Endophyllum osteospermi* (Uredinales, Puccinaceae) on *Chrysanthemoides monilifera*. *Biocontrol science and technology*, Vol.15, No.2, (March 2005), pp. 117-125, ISSN 1360-0478
- Wymore, L. & Watson, A. (1989). Interaction between a velvetleaf isolate of *Colletotrichum coccodes* and thidiazuron for velvetleaf (*Abutilon theophrasti*) control in the field. *Weed science*, Vol. 37, No.3, (May 1989), pp.478-483.
- Wyss, G., Charudattan, R., Rosskopf, E., & Littell, R. (2004). Effects of selected pesticides and adjuvants on germination and vegetative growth of *Phomopsis amaranthicola*, a biocontrol agent for *Amaranthus* spp. *Weed research*, Vol.44, No.6, (December 2004), pp. 469-482, ISSN 0043-1737
- Wyss, G. & Müller-Scharer, H. (2001). Effects of selected herbicides on the germination and infection process of *Puccinia lagenophora*, a biocontrol pathogen of *Senecio vulgaris*. *Biological control: theory and applications in pest management*, Vol.20, No.2, (February 2001), pp. 160-166, ISSN 1049-9644
- Yamoah, E., Jones, E., Bourdot, G., Suckling, D., Weld, R., & Stewart, A. (2008). Factors influencing pathogenicity of *Fusarium tumidum* on gorse (*Ulex europaeus*). *Biocontrol science and technology*, Vol.18, No.7-8, (September 2008), pp. 779-792, ISSN 1360-0478
- Yonli, D., Traore, H., Hess, D., Abbasher, A., & Boussim, I. (2004). Effect of growth medium and method of application of *Fusarium oxysporum* on infestation of sorghum by *Striga hermonthica* in Burkina Faso. *Biocontrol science and technology*, Vol.14, No.4, (June 2004), pp. 417-421, ISSN 1360-0478
- Yuzikhin, O., Mitina, G., & Berestetskiy, A. (2007). Herbicidal potential of stagonolide, a new phytotoxic nonenolide from *Stagonospora cirsii*. *Journal of agriculture & food chemistry*, Vol.55, No.19, (September 2007), pp. 7707-7711, ISSN 1520-5118
- Zdor, R., Alexander, C. & Kremer, R. (2005). Weed suppression by deleterious rhizobacteria is affected by formulation and soil properties. *Communications in soil science and plant analysis*, Vol.36, No.9-10, (February 2007), pp. 1289-1299, ISSN 1532-2416
- Zermane, N., Souissi, T., Kroschel, J., & Sikora, R. (2007). Biocontrol of broomrape (*Orobancha crenata* Forsk. and *Orobancha foetida* Poir.) by *Pseudomonas fluorescens* isolate Bf7-9 from the faba bean rhizosphere. *Biocontrol science and technology*, Vol.17, No.5-6, (June 2007), pp. 483-497, ISSN 1360-0478
- Zhang, W., Sulz, M., Bailey, K., & Cole, D. (2002). Effect of epidemiological factors on the impact of the fungus *Plectosporium tabacinum* on false cleavers (*Galium spurium*). *Biocontrol science and technology*, Vol.12, No.2, (April 2002), pp. 183-194, ISSN 1360-0478
- Zhang, Z., Burgos, N., Zhang, J., & Yu, L. (2007). Biological control agent for rice weeds from protoplast fusion between *Curvularia lunata* and *Helminthosporium gramineum*. *Weed science*, Vol.55, No.6, (November 2007), pp. 599-605, ISSN 0043-1745
- Zhang, J., Wang, W., Lu, X., Xu, Y., & Zhang, X. (2010). The stability and degradation of a new biological pesticide, pyoluteorin. *Pest management science*, Vol.66, No.3, (March 2010), pp. 248-252, ISSN 1526-4998
- Zhao, S. & Shamoun, S. (2005). Effects of potato dextrose broth and gelatin on germination and efficacy of *Phoma exigua*, a potential biocontrol agent for salal (*Gaultheria shallon*). *Canadian journal of plant pathology*, Vol.27, No.2, (Apr-June 2005), pp. 234-244, ISSN 0706-0661

- Zhao, S. & Shamoun, S. (2006). Effects of culture media, temperature, pH, and light on growth, sporulation, germination, and bioherbicidal efficacy of *Phoma exigua*, a potential biological control agent for salal (*Gaultheria shallon*). *Biocontrol science and technology*, Vol.16, No.9-10, (October 2006), pp. 1043-1055, ISSN 1360-0478
- Zhou, L., Bailey, K., & Derby, J. (2004). Plant colonization and environmental fate of the biocontrol fungus *Phoma macrostoma*. *Biological control: theory and applications in pest management*, Vol.30, No.3, (July 2004), pp. 634-644, ISSN 1049-9644
- Zhu, Y. & Qiang, S. (2004). Isolation, pathogenicity and safety of *Curvularia eragrostidis* isolate QZ-2000 as a bioherbicide agent for large crabgrass (*Digitaria sanguinalis*). *Biocontrol science and technology*, Vol.14, No.8, (December 2004), pp. 769-782, ISSN 1360-0478
- Zidack, N. & Quimby, P. Jr. (2002). Formulation of bacteria for biological weed control using the Stabileze method. *Biocontrol science and technology*, Vol.12, No.1, (February 2002), pp. 67-74, ISSN 1360-0478

IntechOpen



Herbicides - Environmental Impact Studies and Management Approaches

Edited by Dr. Ruben Alvarez-Fernandez

ISBN 978-953-307-892-2

Hard cover, 248 pages

Publisher InTech

Published online 20, January, 2012

Published in print edition January, 2012

Weeds severely affect crop quality and yield. Therefore, successful farming relies on their control by coordinated management approaches. Among these, chemical herbicides are of key importance. Their development and commercialization began in the 1940's and they allowed for a qualitative increase in crop yield and quality when it was most needed. This book blends review chapters with scientific studies, creating an overview of some the current trends in the field of herbicides. Included are environmental studies on their toxicity and impact on natural populations, methods to reduce herbicide inputs and therefore overall non-target toxicity, and the use of bioherbicides as natural alternatives.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Tami L. Stubbs and Ann C. Kennedy (2012). Microbial Weed Control and Microbial Herbicides, *Herbicides - Environmental Impact Studies and Management Approaches*, Dr. Ruben Alvarez-Fernandez (Ed.), ISBN: 978-953-307-892-2, InTech, Available from: <http://www.intechopen.com/books/herbicides-environmental-impact-studies-and-management-approaches/microbial-weed-control-and-microbial-herbicides>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen