

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Echocardiography in Severe Aortic Stenosis

Gani Bajraktari

*Service of Cardiology, University Clinical Centre of Kosova, Prishtina  
Republic of Kosovo*

## 1. Introduction

Aortic stenosis (AS) is the most frequent valvular heart disease in west developed and developing countries, with prevalence between 0.02% in adults under 44 years and 3-9% in elderly over 80 years. Patients with this disease may remain asymptomatic for years, particularly in elderly with naturally limited exercise. If the patients remain untreated after they become symptomatic, the mortality at 10 years follow-up is 80-90%. Based on the etiology, mainly are three types of AS: 1) Calcific AS, which is most frequent type in adults of advanced age (2-7% of the population), 2) Congenital, which dominates in the younger patients, and 3) Rheumatic AS, which is becoming rare in developed countries.

Patient history and physical examination remain important in the diagnosis of AS. For the proper patient management, the evidence of the symptoms characteristic for AS: exertional shortness of breath, angina, dizziness, or syncope. Further diagnostic right direction is characteristic systolic murmur.

The disappearance of the second aortic sound is specific to severe AS.

Aortic valve replacement (AVR) is the only effective treatment for severe aortic AS. It is performed either isolated or concomitantly with coronary artery by-pass graft operation, which take place in almost 50% of patients with AS. The overall mortality of isolated AVR is 3-5% in patients below 70 years and 5-15% in elderly. After successful AVR, symptoms and quality of live improves significantly. The long term 10 years survival after successful AVR is very satisfied and it resulted till 75%. The most important factors that may affect the survival are old age, high NYHA functional class, associated aortic regurgitation, concomitant coronary aortic by-pass graft and atrial fibrillation.

## 2. Echocardiography in aortic stenosis patients

Echocardiography is the key diagnostic tool, not only to confirm the presence of AS, but also to assesses the degree of valve calcification, LV function and wall thickness. Today, echocardiography provides prognostic information in patients with AS.

The severity of AS is provided with a very high sensitivity and specificity by Doppler echocardiography. A valve area 1.0 cm<sup>2</sup> in a patient with AS is considered severe. The indexing of aortic valve area to body surface area is more powerful parameter, and a cut-off value of 0.6 cm<sup>2</sup>/m<sup>2</sup> is considered severe AS. However, valve area detected by Doppler

echocardiography cannot be the only parameter for clinical decision making for aortic valve replacement, and it should be considered in combination with flow rate, pressure gradient and ventricular function, as well as functional status of an individual patient.

In patients with AS and normal left ventricular (LV) ejection fraction (EF) the mean pressure gradient of 50 mmHg (Figure 1), was used as a cut-off for the decision making for aortic valve replacement. However, in patients with depressed global LV function, even in patients with severe AS, Doppler echocardiography may result with low pressure gradients (underestimated gradients). In these patients, stress echocardiography using low-dose dobutamine may be helpful to distinguish truly severe AS patients from the rare cases of pseudosevere AS. In patients with truly severe AS, only small changes in valve area, but significant increase in pressure gradients are shown, whereas in pseudosevere AS patients are registered significant increase of valve area surface, but only minor changes in pressure gradients, before and at peak dose of dobutamine. The dobutamine stress-echocardiography is useful also to detect the presence of contractile reserve, which has prognostic implications.




Fig. 1. Continues Doppler velocity of the aortic valve, in a patient with high pressure gradient and normal EF, before aortic valve replacement.

### 3. Outcome of patients with aortic stenosis

Aortic stenosis is a chronic progressive disease. Patients with AS may remain asymptomatic for a long period of time, and the duration of the asymptomatic phase varies widely among individuals. The most frequent cause of death in symptomatic patients is sudden cardiac death. However, sudden cardiac death in asymptomatic patients with AS is very rare.

Older age, presence of atherosclerotic risk factors, valve calcification, peak aortic jet velocity, low LV EF and increase of transvalvular pressure gradient with exercise, were shown as independent predictors of poor outcome in AS patients.

The development of symptoms on exercise testing, in physically active patients with AS, predicts a very high likelihood of symptom development within 12 months. The occurrence of symptoms, in these patients is a correlate of poor prognosis. The increased mortality in these patients has been reported within months of symptom onset, which is often not promptly reported by patients.

### 4. Echocardiographic predictors in patients with severe aortic stenosis and poor left ventricular systolic function

Left ventricular systolic function was shown as one of more important predictors of patients with AS. Patients with AS and LV systolic dysfunction have a poor prognosis if valve replacement is not performed. LV EF, as the most important conventional parameter for the LV global systolic function, was consistently reported as a postoperative prognostic factor in patients with severe AS. Patients with severe left ventricular dysfunction have increased intra-operative mortality, and there are yet contradictions about their improved outcomes after the AVR. Generally, the LV systolic dysfunction is not a contraindication to surgery. It was shown that patients who underwent AVR have a 5-year survival rate 60-70%, with a high operative mortality in the range of 10-15% for patients with LV systolic dysfunction. To predict the postoperative outcome of patients with severe AS and impaired LV function, the preoperative dobutamine stress echocardiography is useful technique. The presence of good contractile reserve in dobutamine stress echocardiography supports potential benefit from AVR and better outcome in these patients.

AVR decreases the LV afterload, through transvalvular pressure drop (Figure 2), resulting in regression of LV hypertrophy.

LV mass regression predominantly occurs within the first 6 months of surgery. Even there are few publications regarding the pre-operative echocardiographic predictors of LV functional recovery in AS patients with low EF, it justifies the statement to consider these patients for the operation, after individual assessment of the patient, considering comorbidities and general conditions.

Recovery of LV function was evident after aortic valve replacement in the majority of patients with aortic stenosis and pre-operative LV dysfunction.

Patients with increased LV end-systolic dimension and/or LV systolic volume index seem to have less chance for the LV functional recovery. It seems that these patients loaded

contractile reserve, and up to now there is no evidence that they may improve LV systolic function after operation, and therefore we should less encourage these patients for the AVR. However, there are studies that have shown that even in patients with poor LV systolic function, there is still ability for a LV function recovery after AVR, explaining it through the mechanism of the markedly reduction of outflow tract resistance.

Studies have shown that stented and stentless valves have similar effect on the LV mass reduction after AVR in all patients that underwent this procedure, despite significant differences in indexed effective orifice area and peak flow velocity in favor of the stentless valve. However, in patients with AS and markedly reduced ventricular function, there was shown more rapid LV mass and function normalization in stentless patients compared to similar patients receiving a stented valve. The lack of large randomized studies for these prostheses makes even more difficult decision. However, a numerous retrospective studies have shown improvement in symptoms and LV EF in about 70% of the survivors after AVR in patients with low LV EF.




Fig. 2. Continues Doppler velocity of the aortic valve, in the same patient, two weeks after aortic valve replacement.



They suggest that despite increased operative mortality, these patients should not be denied aortic valve replacement, given the substantial potential clinical benefit from AVR replacement.

In conclusion, in patients with severe aortic stenosis with impaired LV global systolic function, assessed by LV EF, AVR has significantly better outcome compared to those treated medically. These patients are likely to carry a high risk operation (up to 10%), than to have a very poor prognosis for 10 years survival in medical treatment.

### **5. Echocardiographic predictors in patients with severe aortic stenosis and preserved left ventricular systolic function**

Global LV function, assessed by conventional EF remains normal in most of AS patients. However, the long axis systolic function, assessed by M-mode echocardiography and/or tissue Doppler imaging (TDI) velocities, decreases even in patients with preserved EF. In AS patients with preserved EF, the longitudinal velocity, strain and strain rate are decreased and deteriorate further as AS become severe. These changes reflect that the LV myocardial dysfunction beginning at the subendocardium in early stages of AS and progress to mid-wall and to transmural contraction impairment in patients with severe AS. Recent studies have shown also that in patients with AS and preserved LV EF, the apical rotation and LV twist are increased and untwist is delayed compared to normals, as compensatory mechanisms for the increased intracavitary pressure overload and subendocardial ischaemia. Also, it was shown that these LV myocardial correlate with the severity of AS. However, these compensatory mechanisms are lost after the LV EF deterioration.

Strong evidence exists showing beneficial effect of AVR, not only in improving patients' symptoms but also in recovering, even partially, overall cardiac function. Improvement of LV ventricular function in these patients is interpreted on the basis of regression of myocardial hypertrophy, increased myocardial perfusion and hence overall cavity performance, at early and mid-term post-operative periods. While EF is the most popular measure of pre-operative LV systolic function in such patients, and surgical risk assessment it lacks representing subendocardial component of the LV function.

Severe aortic stenosis causes significant subendocardial dysfunction despite preserved ejection fraction. Aortic valve replacement surgery and removal of left ventricular afterload results in recovery of intrinsic subendocardial function within a week of surgery, well before mass regression and reverse remodeling. Such degree of pre-operative subendocardial disturbances may represent early changes that if ignored may substantiate and become irreversible. Thus, the presence of such abnormalities in symptomatic patients, even with normal ejection fraction, may suggest further evidence for a need for valve replacement in order to maintain overall integral ventricular function and to avoid potential clinical complications.

In patients with severe aortic stenosis and maintained LV EF, the left ventricular twist is increased as compared with normal subjects suggesting a wall motion compensation for the reduced long axis motion in the aim to preserve LVEF. These motions alter towards normal values within six months of aortic valve replacement (Figure 3). These findings are growing evidence that on LV dysfunction and their improvement after AVR, even in asymptomatic patients, and may assist in identifying patients needing surgery before LV damage becomes irreversible.




Fig. 3. Example of LV rotation and twist in control, pre AVR and post AVR. Purple line showing peak of apical rotation and green showing peak basal rotation. AVR, aortic valve replacement (Reproduced from Lindqvist P *et al.* Aortic valve replacement normalizes left ventricular twist function. *Interact CardioVasc Thorac Surg* 2011;12:701-706, doi:10.1510/icvts.2010.262303, with permission from the European Association for Cardio-Thoracic Surgery).

## 6. References

- [1] Iung B, Baron G, Butchart EG, Delahaye F, Gohlke-Barwolf C, Levang OW, Tornos P, Vanoverschelde JL, Vermeer F, Boersma E, Ravauud P, Vahanian A. A prospective survey of patients with valvular heart disease in Europe: the Euro Heart Survey on valvular heart disease. *Eur Heart J* 2003; 24: 1231-1243.

- [2] Stewart BF, Siscovick D, Lind BK, Gardin JM, Gottdiener JS, Smith VE, Kitzman DW, Otto CM. Clinical factors associated with calcific aortic valve disease. Cardiovascular Health Study. *J Am Coll Cardiol* 1997; 29: 630–634.
- [3] Vahanian A, Baumgartner H, Bax J, Butchart E, Dion R, Filippatos G, Flachskampf F, Hall R, Iung B, Kasprzak J, Nataf P, Tornos P, Torracca L, Wenink A; Task Force on the Management of Valvular Heart Disease of the European Society of Cardiology; ESC Committee for Practice Guidelines. Guidelines on the management of valvular heart disease: The Task Force on the Management of Valvular Heart Disease of the European Society of Cardiology. *Eur Heart J* 2007; 28: 230-68.
- [4] Quinones MA, Otto CM, Stoddard M, Waggoner A, Zoghbi WA, for the Doppler Quantification Task Force of the Nomenclature Standards Committee of the American Society of Echocardiography. Recommendations for quantification of Doppler echocardiography: a report from the Doppler Quantification Task Force of the Nomenclature and Standards Committee of the American Society of Echocardiography. *J Am Soc Echocardiogr* 2002; 15: 167-184.
- [5] deFilippi CR, Willett DL, Brickner ME, Appleton CP, Yancy CW, Eichhorn EJ, Grayburn PA. Usefulness of dobutamine echocardiography in distinguishing severe from nonsevere valvular aortic stenosis in patients with depressed left ventricular function and low transvalvular gradients. *Am J Cardiol* 1995; 75: 191-194.
- [6] Nishimura RA, Grantham JA, Connolly HM, Schaff HV, Higano ST, Holmes DR Jr. Low-output, low-gradient aortic stenosis in patients with depressed left ventricular systolic function: the clinical utility of the dobutamine challenge in the catheterization laboratory. *Circulation* 2002; 106: 809-813.
- [7] Monin JL, Quere JP, Monchi M, Petit H, Baleynaud S, Chauvel C, Pop C, Ohlmann P, Lelguen C, Dehant P, Tribouilloy C, Gueret P. Low-gradient aortic stenosis, operative risk stratification and predictors for long-term outcome: a multicenter study using dobutamine stress hemodynamics. *Circulation* 2003; 108:319-324.
- [8] Rosenhek R, Binder T, Porenta G, Lang I, Christ G, Schemper M, Maurer G, Baumgartner H. Predictors of outcome in severe, asymptomatic aortic stenosis. *N Engl J Med* 2000; 343: 611-617.
- [9] Pellikka PA, Sarano ME, Nishimura RA, Malouf JF, Bailey KR, Scott CG, Barnes ME, Tajik AJ. Outcome of 622 adults with asymptomatic, hemodynamically significant aortic stenosis during prolonged follow-up. *Circulation* 2005; 111: 3290-3295.
- [10] Amato MC, Moffa PJ, Werner KE, Ramirez JA. Treatment decision in asymptomatic aortic valve stenosis: role of exercise testing. *Heart* 2001; 86: 381-386.
- [11] Bruch C, Stypmann J, Grude M, Gradaus R, Breithardt G, Wichter T. Tissue Doppler imaging in patients with moderate to severe aortic valve stenosis: clinical usefulness and diagnostic accuracy. *Am Heart J* 2004; 148: 696-702.
- [12] Tarantini G, Buja P, Scognamiglio R, Razzolini R, Gerosa G, Isabella G, Ramondo A, Iliceto S. Aortic valve replacement in severe aortic stenosis with left ventricular dysfunction: determinants of cardiac mortality and ventricular function recovery. *Eur J Cardiothorac Surg* 2003; 24: 879-85.




- [13] Kupari M, Turto H, Lommi J. Left ventricular hypertrophy in aortic valve stenosis: preventive or promotive of systolic dysfunction and heart failure? *Eur Heart J* 2005; 26: 1790-6.
- [14] Das P, Rimington H, Chambers J. Exercise testing to stratify risk in aortic stenosis. *Eur Heart J* 2005; 26: 1309-1313.
- [15] Lancellotti P, Lebois F, Simon M, Tombeux C, Chauvel C, Pierard LA. Prognostic importance of quantitative exercise Doppler echocardiography in asymptomatic valvular aortic stenosis. *Circulation* 2005; 112(Suppl. I): I-377-I-382.
- [16] Lund O, Nielsen TT, Emmertsen K, Flo C, Rasmussen B, Jensen FT, Pilegaard HK, Kristensen LH, Hansen OK. Mortality and worsening of prognostic profile during waiting time for valve replacement in aortic stenosis. *Thorac Cardiovasc Surg* 1996; 44: 289-295.
- [17] Collinson J, Flather M, Coats AJ, Pepper JR, Henein M. Influence of valve prosthesis type on the recovery of ventricular dysfunction and subendocardial ischaemia following valve replacement for aortic stenosis. *Int J Cardiol* 2004; 97: 535-41.
- [18] Arshad W, Duncan AM, Francis DP, O'Sullivan CA, Gibson DG, Henein MY. Opposite effects of coronary artery disease and hypertrophic cardiomyopathy on left ventricular long axis function during dobutamine stress. *Int J Cardiol* 2005; 101: 123-8.
- [19] Pereira JJ, Lauer MS, Bashir M, Afridi I, Blackstone EH, Stewart WJ, McCarthy PM, Thomas JD, Asher CR. Survival after aortic valve replacement for severe aortic stenosis with low transvalvular gradients and severe left ventricular dysfunction. *J Am Coll Cardiol* 2002; 39: 1356-63.
- [20] Takeda S, Rimington H, Smeeton N, Chambers J. Long axis excursion in aortic stenosis. *Heart* 2001; 86: 52-6.
- [21] Blackstone EH, Cosgrove DM, Jamieson WR, Birkmeyer NJ, Lemmer JH Jr, Miller DC, Butchart EG, Rizzoli G, Yacoub M, Chai A. Prosthesis size and long-term survival after aortic valve replacement. *J Thorac Cardiovasc Surg* 2003; 126: 783-96.
- [22] Carabello BA. Evaluation and management of patients with aortic stenosis. *Circulation* 2002; 105:1746-50.
- [23] Vaquette B, Corbineau H, Laurent M, Lelong B, Langanay T, de Place C, Froger-Bompas C, Leclercq C, Daubert C, Leguerrier A. Valve replacement in patients with critical aortic stenosis and depressed left ventricular function: predictors of operative risk, left ventricular function recovery, and long term outcome. *Heart* 2005; 91: 1324-9.
- [24] Perez de Arenaza D, Lees B, Flather M, Nugara F, Husebye T, Jasinski M, Cisowski M, Khan M, Henein M, Gaer J, Guvendik L, Bochenek A, Wos S, Lie M, Van Nooten G, Pennell D, Pepper J; ASSERT (Aortic Stentless versus Stented valve assessed by Echocardiography Randomized Trial) Investigators. Randomized comparison of stentless versus stented valves for aortic stenosis: effects on left ventricular mass. *Circulation* 2005; 112: 2696-702.
- [25] Ding WH, Lam YY, Kaya MG, Li W, Chung R, Pepper JR, Henein MY. Echocardiographic predictors of left ventricular functional recovery following

- valve replacement surgery for severe aortic stenosis. *Int J Cardiol* 2008; 128: 178-84.
- [26] Lim E, Ali A, Theodorou P, Sousa I, Ashrafian H, Chamageorgakis T, Duncan A, Henein M, Diggle P, Pepper J. Longitudinal study of the profile and predictors of left ventricular mass regression after stentless aortic valve replacement. *Ann Thorac Surg* 2008; 85: 2026-9.
- [27] Cramariuc D, Gerdtts E, Davidsen ES, Segadal L, Matre K. Myocardial deformation in aortic valve stenosis: relation to left ventricular geometry. *Heart* 2010; 96: 106-12.
- [28] Dinh W, Nickl W, Smettan J, Kramer F, Krahn T, Scheffold T, Barroso MC, Brinkmann H, Koehler T, Lankisch M, FÜth R. Reduced global longitudinal strain in association to increased left ventricular mass in patients with aortic valve stenosis and normal ejection fraction: a hybrid study combining echocardiography and magnetic resonance imaging. *Cardiovasc Ultrasound* 2010; 8: 29.
- [29] Ding WH, Lam YY, Pepper JR, Kaya MG, Li W, Chung R, Henein MY. Early and long-term survival after aortic valve replacement in septuagenarians and octogenarians with severe aortic stenosis. *Int J Cardiol* 2010; 141: 24-31.
- [30] Ding WH, Lam YY, Duncan A, Li W, Lim E, Kaya MG, Chung R, Pepper JR, Henein MY. Predictors of survival after aortic valve replacement in patients with low-flow and high-gradient aortic stenosis. *Eur J Heart Fail* 2009; 11: 897-902.
- [31] Lindqvist P, Bajraktari G, Molle R, Palmerini E, Holmgren A, Mondillo S, Henein MY. Valve replacement for aortic stenosis normalizes subendocardial function in patients with normal ejection fraction. *Eur J Echocardiogr* 2010; 11: 608-13
- [32] Zhao Y, Lindqvist P, Nilsson J, Holmgren A, Näslund U, Henein MY. Trans-catheter aortic valve implantation--early recovery of left and preservation of right ventricular function. *Interact Cardiovasc Thorac Surg* 2011; 12: 35-9.
- [33] Lindqvist P, Zhao Y, Bajraktari G, Holmgren A, Henein MY. Aortic valve replacement normalizes left ventricular twist function. *Interact Cardiovasc Thorac Surg* 2011; 12: 701-6.
- [34] Owen A, Henein MY. Challenges in the management of severe asymptomatic aortic stenosis. *Eur J Cardiothorac Surg* 2011; 40: 848-50.
- [35] Lam YY, Bajraktari G, Lindqvist P, Holmgren A, Mole R, Li W, Duncan A, Ding WH, Mondillo S, Pepper JR, Henein MY. Prolonged total isovolumic time is related to reduced long-axis functional recovery following valve replacement surgery for severe aortic stenosis. *Int J Cardiol* 2011. [Epub ahead of print]
- [36] Zhao Y, Lindqvist P, Holmgren A, Henein MY. Accentuated left ventricular lateral wall function compensates for septal dyssynchrony after valve replacement for aortic stenosis. *Int J Cardiol*. 2011 Jul 30. [Epub ahead of print]
- [37] Strain analysis in patients with severe aortic stenosis and preserved left ventricular ejection fraction undergoing surgical valve replacement. Delgado V, Tops LF, van Bommel RJ, van der Kley F, Marsan NA, Klautz RJ, Versteegh MI, Holman ER, Schalij MJ, Bax JJ. *Eur Heart J* 2009; 30: 3037-47.
- [38] Miyazaki S, Daimon M, Miyazaki T, Onishi Y, Koiso Y, Nishizaki Y, Ichikawa R, Chiang SJ, Makinae H, Suzuki H, Daida H. Global longitudinal strain in relation to the severity of aortic stenosis: a two-dimensional speckle-tracking study. *Echocardiography* 2011; 28: 703-8.

- [39] Ng AC, Delgado V, Bertini M, Antoni ML, van Bommel RJ, van Rijnsoever EP, van der Kley F, Ewe SH, Witkowski T, Auger D, Nucifora G, Schuijf JD, Poldermans D, Leung DY, Schalij MJ, Bax JJ. Alterations in multidirectional myocardial functions in patients with aortic stenosis and preserved ejection fraction: a two-dimensional speckle tracking analysis. *Eur Heart J* 2011; 32: 1542-50.

IntechOpen

IntechOpen



## **Echocardiography - In Specific Diseases**

Edited by Prof. Gani Bajraktari

ISBN 978-953-307-977-6

Hard cover, 160 pages

**Publisher** InTech

**Published online** 18, January, 2012

**Published in print edition** January, 2012

The book "Echocardiography - In Specific Diseases" brings together contributions from well-known researchers from around the world, some of them specialized in imaging science in their clinical orientation, but also representatives from academic medical centers. Each chapter is structured and written to be accessible to those with a basic knowledge of echocardiography but also to be stimulating and informative to experts and researchers in the field of echocardiography. This book is primarily aimed at cardiology fellows during their basic echocardiography rotation, fellows of internal medicine, radiology and emergency medicine, but also experts in echocardiography. During the past few decades technological advancements in echocardiography have been developing rapidly, leading to improved echocardiographic imaging using new techniques. The authors of this book tried to explain the role of echocardiography in several special pathologies, which the readers may find in different chapters of the book.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Gani Bajraktari (2012). Echocardiography in Severe Aortic Stenosis, Echocardiography - In Specific Diseases, Prof. Gani Bajraktari (Ed.), ISBN: 978-953-307-977-6, InTech, Available from:

<http://www.intechopen.com/books/echocardiography-in-specific-diseases/echocardiography-in-severe-aortic-stenosis>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen