

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Non-Enzymatic Post-Translational Modifications in the Development of Age-Related Macular Degeneration

Yuichi Kaji¹, Tetsuro Oshika¹ and Noriko Fujii²

¹Department of Pathophysiology of Vision and Ophthalmology, University of Tsukuba Graduate School of Comprehensive and Human Sciences, Tsukuba, Ibaraki,

²Research Reactor Institute, Kyoto University, Kumatori, Sennan, Osaka, Japan

1. Introduction

Age-related macular degeneration (AMD) is a leading cause of blindness among Caucasians in various countries.¹⁻³ In addition, cases of AMD are increasing among non-Caucasians, and AMD has become one of the major causes of blindness worldwide. To address this problem, several etiological, pathological, and basic science studies are being conducted. Etiological studies have revealed that the risk factors of AMD include smoking,⁴ increasing age, and the presence of cardiovascular disorders.⁵ However, the molecular mechanisms linking these risk factors to AMD are still unclear.

Pathoclinical studies have revealed macular drusen, which are small lumps of abnormally accumulated proteins beneath the retinal pigment of epithelial cells, to be a sign of AMD (Figure 1).⁶ Proteomics analysis of drusen has revealed that it is composed of various proteins, including clusterin, albumin, TIMP3, vitronectin, complement components, and crystallin. However, the pathological role of drusen in the development of AMD is still unclear.




Fig. 1. Drusen as an early sign of age-related macular degeneration

Drusen are seen as yellow to white materials especially in the macular area (Left).

Histologically, drusen are recognized as extracellular deposits that form between the retinal pigmented epithelium (RPE) and Bruch's membrane (arrows in the right Fig.).

AMD and other aging-related changes in the body have certain common characteristics, including the aggregation of abnormal proteins, seen in the lens and brain of patients with cataract and Alzheimer's disease, respectively. Post-translational modifications of proteins are also aging-related changes commonly seen in the target organ. For example, advanced glycation of proteins and racemization of amino acids with resultant D-amino acid formation in proteins are well-documented changes related to aging. Post-translational modification of proteins occurs because of aging, and it contributes to the aging changes of organs.

In order to elucidate the molecular mechanism of AMD, we evaluated the role of post-translational modifications of proteins in the development of AMD, particularly the formation of advanced glycation end products (AGEs) and D-amino acids in the development of drusen.

2. Post-translational modifications in age-related disorders

Post-translational modification of proteins is a molecular characteristic of the aging process. This is of 2 types as follows: enzymatic post-translational modification, which includes phosphorylation and glycosylation and is essential for protein function; and non-enzymatic post-translational modification that includes advanced glycation, racemization (and the resultant D-amino acid formation), and truncation, which impairs protein function, contributing to the aging process at the molecular level in various organs (Figure 2).




Fig. 2. Aging of proteins at the molecular level

Non-enzymatic post-translational modifications of proteins, including formation of AGEs and D-amino acids, are recognized as an aging process of proteins at the molecular level.

2.1 Advanced glycation end products

Advanced glycation end products (AGEs) are the final reaction products of proteins and reducing sugars. The reaction of reducing sugars (glucose and fructose) with Lys and Arg residues in proteins leads to the formation of Schiff base and Amadori products, which slowly undergo oxidation, dehydration, and condensation to form AGEs (Figure 3). The final products vary depending on the proteins, sugars, and the reactions involved. However, common structures called AGE motifs are seen in the products, irrespective of the proteins and sugars involved. Recently, a number of AGE motifs, including N^{ϵ} -(carboxy) methyl-L-lysine, imidazoline, pyrrolidine, and pentosidine, have been revealed.




Fig. 3. **Multistep reactions of AGE formation**

Proteins and reducing sugars react with each other to form Amadori products. After multistep reactions, including oxidation, dehydration, and condensation, AGEs are generated.

Our body is like an incubator containing sugars and proteins, in which AGEs are naturally formed, particularly in the tissues with a low turnover rate such as bone, teeth, dura mater, and lens. AGEs tend to accumulate with age, and they are known to do so in the target organ of age-related disorders such as Alzheimer's disease and atherosclerosis. Thus, accumulation of AGEs in tissues is thought to be a biomarker of the aging process.

Formation of AGEs increases in diabetes partly because the concentration of reducing sugars increases in the blood. An accumulation of AGEs is seen in the thickened basement membrane and sclerotic lesions of the glomerulus in diabetic nephropathy. Furthermore, AGEs in the vitreous body and cornea are involved in the development of diabetic retinopathy and keratopathy, respectively.¹¹⁻¹⁴

AGEs, along with being the accessory products of the aging process and diabetes, also possibly contribute to the aging process and diabetic complications. AGEs are known to alter the structure of proteins by intra- and inter-molecular crosslinking, consequently disrupting their function. For example, naïve laminin has a figure of the cross, but AGE-modified laminin is deformed, and thus, loses its adhesion property to epithelial cells.

2.2 D-amino acids

Proteins of all living organism on earth are composed of 20 types of amino acids. Among them, 19 amino acids other than glycine have chiral carbon atoms in the molecule.

Depending on the configuration of the side chains around the chiral carbon atoms, amino acids are either L- or D-amino acids. If amino acids were synthesized chemically, the quantity of L- and D-amino acids would be equal. However, proteins in all living organisms are composed exclusively of L-amino acids.

L-amino acids are converted to D-amino acids with a half-life of several thousand years, thus creating an equal amount of L- and D-amino acids over a particular period. This amount of D-amino acids is used for the age determination of fossils. Thus, D-amino acids are regarded as the products of post-mortal change and are irrelevant to living organisms. However, biologically uncommon D-amino acids that are enantiomers of L-amino acids have been found in the lenses¹⁶⁻¹⁹, teeth²⁰, bones, brains²¹, skin²², aortas²³, erythrocytes²⁴, lungs²⁵, and ligaments of elderly donors.²⁶ The presence of D-amino acids in aged tissues of the living body is considered to be a result of racemization of L-amino acids in proteins in metabolically inert tissues during one's lifetime (Figure 4).




Fig. 4. Significance of D-amino acids in proteins

D-amino acids in proteins are seen in fossils as well as in the target organs of aging changes and age related disorders such as cataract, Alzheimer's disease, and atherosclerosis.

2.3 Co-localization of advanced glycation end products and D-amino acids in age-related macular degeneration

To reveal the role of AGEs and D-amino acids in the development of AMD, we analyzed the immunohistochemical localization of AGE and D-amino acid-containing proteins in human ocular samples of various ages.

Eye samples: Nine eyes from 9 donors of 18 to 88 years of age were obtained at the time of necropsy and used as samples for this study. Among them, 4 eyes from donors older than 68 years had drusen.

Antibodies: N^{ϵ} -(carboxy) methyl-L-lysine (CML) is the major component of AGEs in the body. A monoclonal antibody to CML was purchased (Transgenic Co. Ltd, Kumamoto, Japan).

The preparation and characterization of the primary antibody of D- β -Asp containing proteins was as described previously.²⁷ The polyclonal antibody to the synthetic peptide called peptide 3R(Gly-Leu-D- β -Asp-Ala-Thy-Gly-Leu-D- β -Asp-Ala-Thy-Gly-Leu-D- β -Asp-Ala-Thy) corresponding to 3 repeats of positions 149-153 of the human α -A-crystallin was prepared and purified.

Immunohistochemistry: Immunohistochemical localization of CML and D- β -Asp containing proteins was investigated using the antibodies mentioned above. After fixation with 10% formalin solution, 4- μ m-thick sections of the paraffin-embedded ocular samples were prepared. After deparaffinization, the sections were treated with 2 mg/mL of monoclonal antibody to CML or 1:500 diluted polyclonal antibody to D- β -Asp-containing proteins. After washing with phosphate-buffered saline, the sections were treated with secondary antibodies labeled with a polymer of horse radish peroxidase (Hitofine, Max-PO kit, Nichirei Co. Ltd, Tokyo, Japan). The final products were visualized using diaminobenzidine solution dissolved in phosphate-buffered saline.

No immunoreactivity to CML was seen in the retinas, choroids, or scleras of 5 eyes of donors younger than 18 years of age. In contrast, moderate immunoreactivity was seen in the retinal nerve fiber layers, and strong immunoreactivity was seen in the drusen and the thickened Bruch's membrane of the 5 eyes of donors older than 68 years (Figure 5).




Fig. 5. Immunohistochemical localization of AGE in human retina and choroid
No immunoreactivity to CML, a major component of AGEs, is noted in young donor eyes. In contrast, immunoreactivity to CML is seen in the retinal nerve fiber layer (*), drusen (arrows), and thickened Bruch's membrane (arrowheads).

Similarly, no immunoreactivity to the D- β -Asp containing proteins was seen in the retinas, choroids, or scleras of donors younger than 18 years. In contrast, strong immunoreactivity was seen in the drusen seen in donors older than 68 years. In addition, moderate immunoreactivity to D- β -Asp containing proteins was seen in the sclera, the internal limiting membrane of retinal vessels, and Bruch membranes (Figure 6).




Fig. 6. **Immunohistochemical localization of D-amino acid-containing proteins in human retina and choroid**

No immunoreactivity to D- β -Asp-containing proteins, one of the major components of D-amino acids, is noted in young donor eyes. In contrast, the immunoreactivity to D- β -Asp-containing proteins is seen in the vessel walls (*), drusen (arrows), and thickened Bruch's membrane (arrowheads).

3. Possible mechanism of age-related macular degeneration

Drusen, an early sign of AMD, are small lumps of aggregated proteins rich in AGEs and D-amino acid containing proteins, which, in addition to being accessory products of the aging process, also possibly accelerate the aging process, suggesting that AGEs and D-amino acids in drusen play a central role in the development of AMD via various mechanisms.

One possible mechanism involves the interaction of AGEs and AGE receptors, which increases inflammation and accelerates neovascularization. AGE-modified proteins are recognized by the receptor for AGE (RAGE),³⁰ galectin-3,³¹ macrophage scavenger receptors, and CD36.³² Particularly, the interaction of AGEs with RAGE induces inflammatory cytokines such as TNF- α and VEGF.³³ RAGEs are expressed on the surface of retinal pigment epithelial cells. Furthermore, the interaction of AGE and RAGE increases the expression of RAGE, which serves as a positive feedback for the reaction. Thus, the constant interaction of AGEs in drusen with RAGE on retinal pigment epithelial cells would increase the expression of VEGF and induce neovascularization, resulting in AMD (Figure 7).




Fig. 7. Possible mechanism of AMD in relation to AGEs in drusen

The constant interaction of RAGE with AGEs in retinal pigment epithelial cells and drusen leads to the expression of VEGF, reactive oxygen species, and the increased expression of RAGE. This process then leads to neovascularization of the retina, which is a typical clinicopathological finding of AMD.

Another possible mechanism involves autoantibodies to AGE-modified proteins that have been detected in the elderly or in patients with rheumatic arthritis, which may induce inflammatory changes. Thus, autoimmune reactions may occur in tissues containing AGEs. In fact, proteomic analysis of drusen in human and animal models of AMD has revealed the deposition of IgG and complement factors. In addition, Becerra *et al.* have reported that the pathogenesis of AMD involves inflammatory changes.³⁷ Based on these findings, intravitreal injection of corticosteroids to reduce the inflammation is clinically used to treat AMD.³⁸ At present, the pathological role of D-amino acid containing proteins in the development of AMD is unknown. However, AGEs and D-amino acids need to be targeted for the prevention and treatment of AMD. For example, pyridoxamine inhibits the formation of AGEs in the body, and it has been used in clinical trials for the treatment of diabetic nephropathy. In addition, D-aspartyl endopeptidase has been shown to digest some D-amino acid containing proteins,⁴¹ suggesting that an increased expression of the intrinsic D-aspartyl endopeptidase would decrease the amount of D-amino acids-containing proteins in the target organ of the aging process. We suggest that the accumulating data on AGEs and D-amino acids will pave the way for new therapies for AMD in the near future.

4. Acknowledgement

This work was supported by Grants for Scientific Research 21592216 (2009-2011) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.


5. References

- [1] Goldberg J, Flowerdew G, Smith E, Brody JA, Tso MO. Factors associated with age-related macular degeneration. An analysis of data from the first National Health and Nutrition Examination Survey. *Am J Epidemiol* 1988;128:700-710.
- [2] Klein R, Chou CF, Klein BE, Zhang X, Meuer SM, Saaddine JB. Prevalence of age-related macular degeneration in the US population. *Arch Ophthalmol* 2011;129:75-80.
- [3] Obisesan TO, Hirsch R, Kosoko O, Carlson L, Parrott M. Moderate wine consumption is associated with decreased odds of developing age-related macular degeneration in NHANES-1. *J Am Geriatr Soc* 1998;46:1-7.
- [4] Seddon JM, Willett WC, Speizer FE, Hankinson SE. A prospective study of cigarette smoking and age-related macular degeneration in women. *JAMA* 1996;276:1141-1146.
- [5] Vingerling JR, Dielemans I, Bots ML, Hofman A, Grobbee DE, de Jong PT. Age-related macular degeneration is associated with atherosclerosis. The Rotterdam Study. *Am J Epidemiol* 1995;142:404-409.
- [6] Sarks SH, Arnold JJ, Killingsworth MC, Sarks JP. Early drusen formation in the normal and aging eye and their relation to age related maculopathy: a clinicopathological study. *Br J Ophthalmol* 1999;83:358-368.
- [7] Crabb JW, Miyagi M, Gu X, et al. Drusen proteome analysis: an approach to the etiology of age-related macular degeneration. *Proc Natl Acad Sci U S A* 2002;99:14682-14687.
- [8] Umeda S, Suzuki MT, Okamoto H, et al. Molecular composition of drusen and possible involvement of anti-retinal autoimmunity in two different forms of macular degeneration in cynomolgus monkey (*Macaca fascicularis*). *FASEB J* 2005;19:1683-1685.
- [9] Grillari J, Grillari-Voglauer R, Jansen-Durr P. Post-translational modification of cellular proteins by ubiquitin and ubiquitin-like molecules: role in cellular senescence and aging. *Adv Exp Med Biol* 2010;694:172-196.
- [10] Harding JJ, Beswick HT, Ajiboye R, Huby R, Blakytyn R, Rixon KC. Non-enzymic post-translational modification of proteins in aging. A review. *Mech Ageing Dev* 1989;50:7-16.
- [11] Stitt AW. Advanced glycation: an important pathological event in diabetic and age related ocular disease. *Br J Ophthalmol* 2001;85:746-753.
- [12] Stitt AW. The maillard reaction in eye diseases. *Ann N Y Acad Sci* 2005;1043:582-597.
- [13] Kaji Y, Usui T, Ishida S, et al. Inhibition of diabetic leukostasis and blood-retinal barrier breakdown with a soluble form of a receptor for advanced glycation end products. *Invest Ophthalmol Vis Sci* 2007;48:858-865.
- [14] Kaji Y, Usui T, Oshika T, et al. Advanced glycation end products in diabetic corneas. *Invest Ophthalmol Vis Sci* 2000;41:362-368.
- [15] Charonis AS, Reger LA, Dege JE, et al. Laminin alterations after in vitro nonenzymatic glycosylation. *Diabetes* 1990;39:807-814.
- [16] Fujii N, Harada K, Momose Y, Ishii N, Akaboshi M. D-amino acid formation induced by a chiral field within a human lens protein during aging. *Biochem Biophys Res Commun* 1999;263:322-326.
- [17] Fujii N, Shimo-Oka T, Ogiso M, et al. Localization of biologically uncommon D-beta-aspartate-containing alphaA-crystallin in human eye lens. *Mol Vis* 2000;6:1-5.

- [18] Fujii N, Takemoto LJ, Momose Y, Matsumoto S, Hiroki K, Akaboshi M. Formation of four isomers at the asp-151 residue of aged human alphaA-crystallin by natural aging. *Biochem Biophys Res Commun* 1999;265:746-751.
- [19] Kaji Y, Oshika T, Takazawa Y, Fukayama M, Takata T, Fujii N. Localization of D-beta-aspartic acid-containing proteins in human eyes. *Invest Ophthalmol Vis Sci* 2007;48:3923-3927.
- [20] Masters PM. Stereochemically altered noncollagenous protein from human dentin. *Calcif Tissue Int* 1983;35:43-47.
- [21] Shapira R, Chou CH. Differential racemization of aspartate and serine in human myelin basic protein. *Biochem Biophys Res Commun* 1987;146:1342-1349.
- [22] Fujii N, Tajima S, Tanaka N, Fujimoto N, Takata T, Shimo-Oka T. The presence of D-beta-aspartic acid-containing peptides in elastic fibers of sun-damaged skin: a potent marker for ultraviolet-induced skin aging. *Biochem Biophys Res Commun* 2002;294:1047-1051.
- [23] Powell JT, Vine N, Crossman M. On the accumulation of D-aspartate in elastin and other proteins of the ageing aorta. *Atherosclerosis* 1992;97:201-208.
- [24] McFadden PN, Clarke S. Methylation at D-aspartyl residues in erythrocytes: possible step in the repair of aged membrane proteins. *Proc Natl Acad Sci U S A* 1982;79:2460-2464.
- [25] Shapiro SD, Endicott SK, Province MA, Pierce JA, Campbell EJ. Marked longevity of human lung parenchymal elastic fibers deduced from prevalence of D-aspartate and nuclear weapons-related radiocarbon. *J Clin Invest* 1991;87:1828-1834.
- [26] Ritz-Timme S, Laumeier I, Collins M. Age estimation based on aspartic acid racemization in elastin from the yellow ligaments. *Int J Legal Med* 2003;117:96-101.
- [27] Yang D, Fujii N, Takata T, et al. Immunological detection of D-beta-aspartate-containing protein in lens-derived cell lines. *Mol Vis* 2003;9:200-204.
- [28] Ishibashi T, Murata T, Hangai M, et al. Advanced glycation end products in age-related macular degeneration. *Arch Ophthalmol* 1998;116:1629-1632.
- [29] Kaji Y, Oshika T, Takazawa Y, Fukayama M, Fujii N. Accumulation of D-beta-aspartic acid-containing proteins in age-related ocular diseases. *Chem Biodivers* 2010;7:1364-1370.
- [30] Wautier JL, Wautier MP, Schmidt AM, et al. Advanced glycation end products (AGEs) on the surface of diabetic erythrocytes bind to the vessel wall via a specific receptor inducing oxidant stress in the vasculature: a link between surface-associated AGEs and diabetic complications. *Proc Natl Acad Sci U S A* 1994;91:7742-7746.
- [31] Vlassara H, Li YM, Imani F, et al. Identification of galectin-3 as a high-affinity binding protein for advanced glycation end products (AGE): a new member of the AGE-receptor complex. *Mol Med* 1995;1:634-646.
- [32] Ohgami N, Nagai R, Ikemoto M, et al. Cd36, a member of the class b scavenger receptor family, as a receptor for advanced glycation end products. *J Biol Chem* 2001;276:3195-3202.
- [33] Ma W, Lee SE, Guo J, et al. RAGE ligand upregulation of VEGF secretion in ARPE-19 cells. *Invest Ophthalmol Vis Sci* 2007;48:1355-1361.
- [34] Zhou J, Cai B, Jang YP, Pachydaki S, Schmidt AM, Sparrow JR. Mechanisms for the induction of HNE- MDA- and AGE-adducts, RAGE and VEGF in retinal pigment epithelial cells. *Exp Eye Res* 2005;80:567-580.

- [35] Shibayama R, Araki N, Nagai R, Horiuchi S. Autoantibody against N(epsilon)-(carboxymethyl)lysine: an advanced glycation end product of the Maillard reaction. *Diabetes* 1999;48:1842-1849.
- [36] Tai AW, Newkirk MM. An autoantibody targeting glycated IgG is associated with elevated serum immune complexes in rheumatoid arthritis (RA). *Clin Exp Immunol* 2000;120:188-193.
- [37] Klein R, Knudtson MD, Klein BE, et al. Inflammation, complement factor h, and age-related macular degeneration: the Multi-ethnic Study of Atherosclerosis. *Ophthalmology* 2008;115:1742-1749.
- [38] Becerra EM, Morescalchi F, Gandolfo F, et al. Clinical evidence of intravitreal triamcinolone acetonide in the management of age-related macular degeneration. *Curr Drug Targets* 2011;12:149-172.
- [39] Metz TO, Alderson NL, Thorpe SR, Baynes JW. Pyridoxamine, an inhibitor of advanced glycation and lipoxidation reactions: a novel therapy for treatment of diabetic complications. *Arch Biochem Biophys* 2003;419:41-49.
- [40] Williams ME, Bolton WK, Khalifah RG, Degenhardt TP, Schotzinger RJ, McGill JB. Effects of pyridoxamine in combined phase 2 studies of patients with type 1 and type 2 diabetes and overt nephropathy. *Am J Nephrol* 2007;27:605-614.
- [41] Kinouchi T, Nishio H, Nishiuchi Y, et al. Isolation and characterization of mammalian D-aspartyl endopeptidase. *Amino Acids* 2007;32:79-85.

IntechOpen



Age Related Macular Degeneration - The Recent Advances in Basic Research and Clinical Care

Edited by Dr. Gui-Shuang Ying

ISBN 978-953-307-864-9

Hard cover, 300 pages

Publisher InTech

Published online 20, January, 2012

Published in print edition January, 2012

Age-related Macular Degeneration (AMD) is the leading cause of vision loss and blindness in the developed countries. In the past decade, great progress has been made in understanding the pathobiology and genetics of this blinding disease, as well as in finding new therapies for its treatment. These include the discovery of several genes that are associated with the risk of AMD, new anti-VEGF treatments for wet AMD and new imaging techniques to diagnose and monitor the AMD. All chapters in this book were contributed by outstanding research scientists and clinicians in the area of AMD. I hope this timely book will provide the basic scientists and clinicians with an opportunity to learn about the recent advances in the field of AMD.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Yuichi Kaji, Tetsuro Oshika and Noriko Fujii (2012). Non-Enzymatic Post-Translational Modifications in the Development of Age-Related Macular Degeneration, *Age Related Macular Degeneration - The Recent Advances in Basic Research and Clinical Care*, Dr. Gui-Shuang Ying (Ed.), ISBN: 978-953-307-864-9, InTech, Available from: <http://www.intechopen.com/books/age-related-macular-degeneration-the-recent-advances-in-basic-research-and-clinical-care/non-enzymatic-post-translational-modifications-in-the-development-of-age-related-macular-degeneratio>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen