

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Myocardial Self-Repair and Congenital Heart Disease

Fariba Chalajour, Xiaoyuan Ma and R. Kirk Riemer
Stanford University, Stanford, CA
USA

1. Introduction

In this chapter we will explore the current understanding about the dynamics of myocardial cell populations after birth and how they may contribute to homeostasis and response to injury. Recent investigations of myocardial cell biology have revealed that the heart is not the terminally differentiated organ it was once thought to be. We now know that myocardium contains limited populations of cardiac progenitor cells (CPC) that are capable of generating all myocardial cell types. The changes in myocardial cell populations with time and the role these changes play in neonatal myocardial tissue expansion will be discussed. The contribution of CPC to heart growth and therapeutic strategies for congenital cardiac diseases will be explored.

2. Stem cells and heart development

2.1 Stem cells: The perspective has evolved

Pluripotent embryonic stem cells (ESC) are derived from the inner cell mass of an embryonic blastocyst (primitive ectoderm). The *pluripotency* of ESC defines that they are capable of differentiation into one of the three germ layers: ectoderm, mesoderm, or endoderm; reviewed in (Rossant, 2008; Bolli & Chaudhry, 2010). Embryonic mesoderm cells (identified by the expression of Brachyury T) in the developing primary heart field undergo committed differentiation into a pre-cardiac lineage (expressing Mesp-1) and have the potential to form committed cardiac progenitor cells (CPC, expressing Nkx2.5), which are capable of differentiation into cardiomyocytes, vascular smooth muscle cells, and endothelial cells; reviewed in (Sturzu & Wu, 2011). A cardiac progenitor cell is thus defined as a *multipotent* progenitor cell of the pre-cardiac lineage.

Originally it was thought that only ESC are pluripotent, and they could only be obtained from living embryos. Consequently, studies of human ESC have been impeded by ethical concerns. However, continued research has led to the realization that pluripotency is in fact a plastic multidirectional state. It has now been clearly demonstrated that a pluripotent state can be induced in adult somatic cells from humans, and other species; reviewed in Yamanaka (Yamanaka & Blau, 2010). The biochemical methods through which so-called induced pluripotent cells (iPSC) may be derived from adult cells are numerous and still expanding. Several recent reviews (Rossant, 2008; Yamanaka & Blau, 2010) of both the technology and the biology of iPSC may be consulted for the interested reader; the topic will not be further discussed here.

A variety of techniques have been used to induce the differentiation of ESC into a specific cell type of interest. The directed differentiation of ESC into cardiomyocytes has been achieved by several methods and has enabled the use of this approach to produce a large enough number of cells for repairing the injured heart to restore myocardial function (Gonzales & Pedrazzini, 2009; Laflamme & Murry, 2011). A large number of studies have been conducted and are still being conducted with the goal of restoring the function of infarcted myocardium using stem cell therapy with ESC-derived cardiomyocytes and also other cell types, with the expectation that they would continue to differentiate into cardiomyocytes after being delivered to the injured heart tissue (Murry et al., 2005). As this therapy is less relevant to *congenital* heart disease, it will not be further discussed here.

2.2 How cardiac progenitor cells form the developing heart

A brief review of cardiac development illustrates how cardiac stem cells are identified and how multiple distinct populations of pre-cardiac cells coalesce to form the distinct subregions of the heart.

2.2.1 Heart fields and cardiac progenitor cells

The development of the multi-chambered heart is complex and requires a precise regulation of cell migration, proliferation, and differentiation in a highly organized positional and temporal order. Several distinct CPC populations contribute to the formation of different heart fields. The first heart field (FHF) of cardiac progenitor cells is localized in the primitive streak and mostly contributes to formation of the left ventricle and part of the atria (Garcia-Martinez & Schoenwolf, 1993; Tam et al., 1997). A second heart field (SHF) of CPC from the pharyngeal mesoderm migrates into the arterial pole of the heart and is a major source of the cardiac progenitors that form the outflow tract (OFT), the right ventricle and the atria (Abu-Issa & Kirby, 2007). Continued development of the OFT and heart valves is achieved by migration of a population of non-mesodermal neural crest cells (NCC) from the neural fold into the arterial pole and endocardial cushion (Kirby & Waldo, 1995; Hutson & Kirby, 2007). Another population of mesenchymal CPC, the proepicardium, has been demonstrated to be an additional cell source contributing to the formation of coronary arteries and cardiac fibroblasts (Dettman et al., 1998). CPC from both FHF and SHF act in a close collaboration during the formation of the embryonic heart. During heart morphogenesis, CPC directed by different signaling pathways differentiate to mature cardiac cells. However, CPC of the postnatal heart have been shown to express markers common to both FHF and SHF. Whether these postnatal resident CPC contribute to remodeling and growth of the heart is considered likely but is still unsettled.

2.2.2 Second heart field and congenital heart disease

Studies of heart development in mice have shown that mutation of genes expressed by cells of the SHF causes congenital heart disease (CHD). Mutations in *Nkx2.5* cause a spectrum of congenital heart defects including cardiac conduction abnormalities and ventricular- and atrial- septal defects (VSD, ASD) (Basson et al., 1997). Deletion of *Tbx1* results in malformations of the cardiac outflow tract and VSDs due to failure in the migration of NCC to the heart (Jerome & Papaioannou, 2001; Merscher et al., 2001). Mutations in *GATA4*, some of which disrupt its interaction with *Tbx5*, cause ASDs and VSDs (Garg et al., 2003). *Hand2* is essential for survival of second heart field progenitors and loss of *Hand2* function in this cardiac progenitor population can cause a spectrum of congenital heart malformations

(Tsuchihashi et al., 2011). Cai (Cai et al., 2003) reported that *Isl1*⁺ cells are mostly localized in the SHF and give rise to the outflow tract. They demonstrated that disruption of *Isl1* results in a complete failure to form the outflow tract in the mutant mice. The role of *Fgf8* in early and late heart development has also been studied (Sun et al., 1999; Abu-Issa et al., 2002). *Fgf8* is required for migration of mesoderm out of the primitive streak (Sun et al., 1999). In addition *Fgf8* plays an important role in development of all the pharyngeal arches and in NCC survival (Abu-Issa et al., 2002). Cardiac NCC are multipotent and after migration to the SHF contribute to cardiovascular patterning (Kirby & Waldo, 1995). Ablation of the pre-migratory cardiac NCC causes numerous outflow tract septation defects: persistent truncus arteriosus, double-outlet right ventricle, tetralogy of Fallot, double-inlet left ventricle, tricuspid atresia, straddling tricuspid valve, and the absence of a varying combination of aortic arch arteries derived from pharyngeal arches 3, 4 and 6 (Nishibatake et al., 1987; Hutson & Kirby, 2007). Studies of interactions between cells of SHF origin with NCC in the pharyngeal region of mice (Vitelli et al., 2002; Moraes et al., 2005) have shown that mutation in *Tbx1* disrupts formation of the pharyngeal arch arteries and causes septation defects due to failure in NCC migration. Bradshaw (Bradshaw et al., 2009) showed that abnormal distribution of *Isl1*-expressing cells in a neural crest-deficient mutant mouse causes instability of posterior arch arteries and outflow tract septation defects, leading to a double outlet right ventricle. Discovery of the presence and role of CPC in the normal and abnormal development of heart fields and their possible role in postnatal life suggests that targeting these cells is a valuable approach for both understanding the ontology of CHD and developing therapeutic approaches.

The implications of resident CPC expressing neural markers in the heart are manifold for congenital heart disease repair. First, the secondary heart field of cells creates the structures most affected by congenital disease: the outflow tracts and pulmonary and aortic trunks (Dyer & Kirby, 2009; Jain et al., 2010). Second, the resident CPC-derived from this field (i.e. *Isl1*⁺ or *Nestin*⁺ CPC) may form the primary cardiac stem cell pool that would potentially be mobilized to participate in the repair (cell replacement) of structures derived from the anterior/secondary heart field such as the OFT. Thirdly, therapeutics aimed at CPC mobilization may act more effectively on the right heart if they target NCC-derived CPC.

2.2.3 Postnatal distribution of CPC in the heart

Resident CPC have been localized in small groups or clusters in a unique microenvironment known as a “stem cell niche” which provides conditions for stem cells to maintain their multipotency and renewal capacity (Morrison & Spradling, 2008). Niches comprise a specific arrangement of stem cells, supporting cells, and extracellular matrix. The niche environment provides a paracrine signaling influence that maintains stem cells in their quiescent state and also responds to conditions outside the niche to initiate activation of stem cell replication and/or differentiation, for example in response to tissue injury (Morrison & Spradling, 2008). Classically, three well-characterized stem cell niches have been identified: germline, hematopoietic, and epithelial (Lemischka, 1997; Xie & Spradling, 2000; Spradling et al., 2001). However in the past decade, identification of resident stem cells in most mammalian organs suggests local tissue-specific niches are a general rule. Discovery of CPC niches in the adult mouse heart strengthens the growing appreciation that the heart is not a terminally differentiated organ and resident CPC might play a role in the postnatal remodeling and repair of cardiac tissue (Urbanek et al., 2006). The early study of Messina (Messina et al., 2004) demonstrated that cardiac stem cells are present in both atrial and ventricular human samples and in a wide range of ages (1 to 80 years old). According to this

study and mouse studies by Beltrami (Beltrami et al., 2003) and Oh (Oh et al., 2003), cardiac stem cells can be isolated from adult myocardium and differentiated into different cardiac cell lineages. Our own studies (Amir et al., 2008) have investigated the number and characteristics of CPC in human neonatal myocardium from patients with congenital heart disease. We demonstrated that CPC comprise a high percentage relative to cardiomyocytes in the human neonate and that their fractional representation declines as the heart grows in size, suggesting a dilutional effect. In other words, an apparently fixed population of CPC resides in the heart postnatally. Pouly (Pouly et al., 2008) reported that CPC concentration in the right atrium is greater than that of the septum of transplanted hearts. Other myocardial niches have been reported. Schenke-Layland (Schenke-Layland et al., 2011) reported that in human and mouse heart endogenous, multipotent $Isl1^+/Flk1^+$ CPC reside within niche clusters in the right ventricular free wall, the atria and outflow tracts. They were tightly circumscribed by the basement membrane proteins collagen V and laminin. However, systematic mapping of the complete heart to localize major niches and determine the distribution of these regions of high CPC density has not been reported. A general concept is that cardiac regions protected from greatest mechanical stress (atria, septum, apex) appear to contain the highest density of CPC. A more useful perspective may be consideration of the cardiac developmental sequence in which fields of CPC undergo migration and strategically controlled distribution, since the niches may constitute remnants of specific precursor fields in the primordial heart anlagen.

The role of niche-associated cells in regulating the stability and activation of resident stem cells is still poorly understood, especially in the heart. Recently, it has been shown that Notch-1 of murine CPC interacts with surrounding cells via the Jagged ligand. Through a Jagged-mediated activation signaling process induced in vitro, notch signaling, down regulation of c-Kit and upregulation of $Nkx2.5$ were found to be associated with increased myocyte proliferation (Boni et al., 2008). These data support the concept that interaction of CPC with supporting cells and matrix regulates their commitment to the myocyte lineage.

2.2.4 Postnatal Characterization of CPC fates

2.2.4.1 Prenatal formation of the heart from different fields

During heart development CPC express field-specific markers (**Figure 1.**) for FHF, SHF, and neural crest (Vincent & Buckingham, 2010). $Mesp1$ is one of the earliest markers for cardiac primordial cells (Saga et al., 1999). Depending on the stage of cell differentiation, genetic markers will be upregulated or down regulated in the CPC, which complicates clear classification of CPC originating from FHF and SHF based on markers. $Nkx2.5$ is a critical cardiac transcription factor in the first lineage, and $Isl1$ along with $Foxh1$, GATA factors, and $Hand2$ are key regulators in the second heart progenitor field (Moretti et al., 2006). The earliest differentiated cells in the cardiac crescent express GATA4 and $Nkx2.5$ (Vincent & Buckingham, 2010). The final differentiation to cardiomyocytes is controlled by Tbx (Takeuchi & Bruneau, 2009). Known markers for SHF CPC are $Isl1$, $Tbx1$ and $Fgf10$ (Kelly et al., 2001; Cai et al., 2003; Xu et al., 2004). $Pax3$ is recognized as a major regulatory gene for NCC, having an important role in their migration (Bradshaw et al., 2009). There is considerable overlap between the heart field-specific genetic markers. This issue is more evident in the case of the proepicardial organ, which gives rise to coronary arteries and epicardium. Both $Isl1$ and $Nkx2.5$ are expressed in the CPC from the proepicardial organ (Dettman et al., 1998), the cells of which also express $Tbx18$ and $Wt1$ (Martinez-Estrada et al., 2010).

2.2.4.2 Markers of CPC in the postnatal heart

In the postnatal heart, identification of CPC is based primarily on the cardiac lineage markers Nkx2.5, Mef2c, and Isl1. Identification of c-Kit⁺ cells in adult mammalian heart suggests another postnatal source of CPC, however, the origin of these cells is unsettled. Although expression of the key CPC markers is preserved both in rodents and humans, species-specific markers limit translation of animal studies to human studies. Our studies of human infant RVOFT myocardium showed a mixed population of c-Kit⁺, Isl1⁺, and Nkx2.5⁺ CPC (Amir et al., 2008). The neonatal human heart is a rich source of CPC bearing markers such as SSEA4, Isl1, Nkx2.5, and c-Kit. In addition to our report of myocardial SSEA4 localization (Amir et al., 2008), SSEA4 expression has been previously only reported in adult human kidney (Ward et al., 2011). Isl1⁺ CPC have also been localized to the atrium (Laugwitz et al., 2005). Stem cell antigen-1 (Sca-1) is a marker for mouse resident CPC. In the mouse and human studies performed by Messina (Messina et al., 2004) and Beltrami (Beltrami et al., 2003) c-Kit⁺ cells are considered to be CPC, whereas Oh (Oh et al., 2003) used c-Kit⁻, Sca-1⁺ cells as a marker for CPC. This apparent controversy may be explained by the existence of heterogeneous pools of cardiac stem cells in different stages of differentiation rather than multiple populations of distinct resident CPC. In addition, the anatomical location (i.e., heart field of origin) of the heart regions studied may determine the markers expressed by postnatal CPC.




Fig. 1. Overview of myocardial CPC Markers in relation to their ontology.

Pre-cardiac mesoderm cells express Mesp1 until differentiation to cardiogenic mesoderm, marked by the expression of Nkx2.5. The cardiogenic mesoderm differentiates to form two heart fields. Cells in the first heart field (FHF) express GATA4 in addition to Nkx2.5. Cells of the secondary heart field (SHF) express Isl1 and Nkx2.5. The SHF also becomes populated by neural crest cells (NCC) expressing Pax3. Markers associated with the heart field cells but showing greater variation in their expression are indicated in blue type. In addition to cardiogenic mesoderm, cells from the pro-epicardial organ and also of hematopoietic lineage have been identified in the myocardium. Abbreviations: NCC, Neural crest cells; SHF, Second heart field; FHF, First heart field.

Mouse studies have demonstrated homing of bone marrow-derived stem cells into the infarcted myocardium, suggesting a hematopoietic origin for c-Kit⁺ CPC. However, transdifferentiation of these cells into cardiomyocytes is controversial (Orlic et al., 2001; Murry et al., 2004; Bearzi et al., 2007). On the other hand, Tallini (Tallini et al., 2009) reported that c-Kit is expressed by immature cardiomyocytes as well as endothelial cells during development of mouse heart and that the populations rapidly expanded during the first 2 days of postnatal life. Their study compared the difference between the multipotency potential of CPC derived from the neonatal heart and the adult injured heart. They concluded that the neonatal c-Kit⁺ cells showed an ability to differentiate into cardiomyocytes, smooth muscle cells, and endothelial cells. In contrast, they found that c-Kit⁺ cells in the adult injured mouse heart had no myogenic capacity. The co-expression of c-Kit and neuronal marker Nestin in neonatal CPC (Tallini et al., 2009) suggests a transitional status of NCC in postnatal life and the potential for NCC contributing to cardiac repair and remodeling (Drapeau et al., 2005).

2.3 Postnatal heart growth

The infant heart grows rapidly after birth. The pulmonary valve annular diameter doubles in the first 7 years of life and plateaus by age 14 (Sairanen & Louhimo, 1992) while left ventricular volume doubles over approximately the first 10 years of life (Nielsen et al., 2010). Although cardiomyocytes increase in volume (hypertrophic growth), in part by cellular fusion into very large multinucleate myocytes, there is also an overall increase in cellularity (cell number) in the heart during this rapid expansion phase of heart growth. That the number of myocytes continues to increase after birth has only recently been demonstrated. For decades, it was taken as fact that cardiomyocytes were incapable of dividing postnatally, if only because mitosis of a multinucleated cell is considered impossible.

One of the earliest reports of new cardiomyocytes appearing in the postnatal human heart was provided by studies of adult aortic stenosis patients, in which cells co-expressing cardiac and stem cell markers could be localized (Urbanek et al., 2003). Additionally, studies of transplanted adult donor hearts revealed the appearance of new cardiomyocytes in the donor heart by the presence of sex chromosomes opposite to that of the recipient host (Bayes-Genis et al., 2007). Highly compelling further evidence that new myocytes are added to the heart over time arises from carbon isotope data (Bergmann et al., 2009) showing that nearly 50% of myocytes are replaced over a lifetime. Studies of the hearts of dogs with advanced dilated cardiomyopathy performed by Leri (Leri et al., 2001) showed postnatal proliferation of cardiomyocytes, documented by the expression of cell proliferation marker Ki67 and telomerase. Telomerase activity is not only required for cardiac growth and survival but it also suppresses cardiomyocyte apoptosis (Oh & Schneider, 2002). We (Amir et al., 2008) also demonstrated that the neonatal human heart contains cardiomyocytes expressing Ki67. We also determined that the number of these proliferating myocytes declines nearly 6-fold in the first two months of life.

2.4 Summary: CPC in the postnatal myocardium

Work by a number of investigators has demonstrated that a population of multipotent cardiac lineage-determined cells (i.e., CPC) capable of further differentiation to all cardiac cell types is present in all hearts. Hence, continued cardiomyocyte renewal has slowly been gaining acceptance as both an important aspect of normal myocardial biology and a potential strategy to assist with repair of a diseased heart. The ability to expand the

population of cardiomyocytes presents a potentially vast opportunity for therapeutic intervention for congenital heart diseases.

3. CPC and Therapy for CHD

3.1 Cardiac repair using CPC

3.1.1 Role of CPC in homeostatic repair and postsurgical healing

If the heart can add new cells, then the question arises regarding why it doesn't always heal itself? The generally offered explanation for the fact that resident CPC are unable to naturally rescue a moderately infarcted adult heart is that the region of damaged tissue is too great for this mechanism to work rapidly enough to restore function. Although not widely recognized, "silent" repair by CPC has been observed (see section 3.2.1). It is probable that small lesions in the myocardium are actually self-repaired silently through expansion of the resident CPC population. Large ischemic lesions would be expected to also result in the loss of resident CPC in the infarcted region, further impairing self-repair. However, progress has recently been reported. Intra-myocardial injection of autologous bone marrow derived stem cells has been used in adults with chronic ischemic heart disease (i.e. not infarcted) to achieve functional recovery and reverse ventricular remodeling (Williams et al., 2011). Such reports of success strengthen the concept that endogenous stem cells can provide clinically significant benefits to heart disease patients of all ages.

3.1.2 Therapies using CPC for tissue engineering

3.1.2.1 Tissue engineered myocardial grafts

Presently, many investigators are attempting to use cardiac stem cells to produce engineered myocardial sheet grafts for myoplasty of larger regions of infarct-damaged adult myocardium. Success has been limited to moderate. Major limiting factors to this approach remain. For example, integration of the graft into the existing myocardium so that it provides clinically significant augmented force development has been problematic. Additionally, achieving sufficient revascularization of the grafts to sustain viability has been difficult; see review by Sui (Sui et al., 2011). Although the potential utility of such grafts in congenital heart disease is not expectedly large, marked progress in this approach could conceivably provide alternative therapies for patients with failing Fontan circulations or possibly for those with cardiomyopathies.

3.1.2.2 Tissue engineered myocardial vascular and valve grafts

One area of potentially beneficial therapy that has previously received little attention in *congenital* heart disease patients is the engineering of vascular and valve grafts that are capable of meeting the rapid growth rate typical of the neonatal heart and great vessels during the first decade of life. The availability of graft materials with the ability to grow along with the young patient is highly desirable, but thus far not available. This is an area of recently increasing research interest, and one in which further research could provide enormous benefit to CHD patients.

3.1.3 Mobilizing resident CPC

3.1.3.1 Alternatives to cell delivery: "Activation" of resident CPC

Given the apparent difficulty in achieving clinically valuable augmentation of cardiac performance through the delivery of cardiomyocytes to a damaged heart, therapeutic

approaches that are designed to mobilize resident CPC to expand the population of cardiomyocytes in situ are being given much more consideration. Therapeutic exploitation of the paracrine environments of the CPC niche and enhancing homing to sites of repair is a very attractive alternative approach to cell-based therapies. In concept, it is a matter of using biomolecules to mimic or enhance endogenous CPC “awakening” mechanisms to obtain greater quantities of cells to differentiate into functional cardiomyocytes. Recent reports reveal that this general approach has a high potential for success and is quite worthy of further investigation. High mobility group box protein 1 (HMGB1), an endogenous chromatin-associated protein, and Thymosine Beta4, a G-actin monomer binding protein, have been identified as paracrine factors potentially able to promote regeneration of myocardium. HMGB1 is released from necrotic cells and has been shown to stimulate the homing of fibroblasts and smooth muscle cells. It has been identified as a potential mediator of resident stem cell activation/mobilization (Palumbo & Bianchi, 2004). Limana (Limana et al., 2005) demonstrated that injection of HMGB1 into the infarcted region of mice induced the appearance of new myocytes and an increase in ventricular performance, leading these investigators to conclude that HMGB1 is a “potent inducer of myocardial regeneration.” They demonstrated that c-kit⁺ CPC express the receptor for HMGB1 and that treatment increased the number of c-kit⁺ cells in mouse heart. Thymosine Beta4 has been shown to stimulate epicardial-derived cells (EPDC) to migrate and potentially promote neovascularization in the infarcted mouse heart (Smart et al., 2010); see review by (Bollini et al., 2011). Although there was no proof that EPDC were a source of the new cardiomyocytes, they may facilitate collateral vessel growth and thereby support the cardiomyocyte regeneration process. Damaged myocardium has a different paracrine environment, which if properly understood and exploited, may provide unique approaches for therapy via resident CPC activation.

Growth factor treatment has also been used to activate resident CPC populations for myocardial repair. Linke (Linke et al., 2005) used a canine MI model to show that IGF-1 and HGF treatment (intra-myocardial injection) could increase the density of proliferating (Ki67⁺) CPC following MI. These same investigators recently extended their observations, showing that CPC aging is related to a decline in signaling through IGF-1 and HGF, which in turn reduces their effect to antagonize the aging effect that the local renin-angiotensin system induces on CPC. They found that IGF-1 and HGF were able to partially reverse age-related decline in cardiac function in rats (Gonzalez et al., 2008). Consistent with these reports of IGF-1 stimulation of CPC proliferation, D’Amario (D’Amario et al., 2011) has proposed that CPC senescence is regulated by paracrine and autocrine signaling: positive through IGF-1 & -2, and balanced by an opposing signal mediated primarily by angiotensin II, all acting via their cognate receptors. Accordingly, they propose that IGF-1 promotes CPC proliferation and survival via IGF-1 receptor signaling, CPC differentiation via action on both IGF-1 and -2 receptors, and increasing angiotensin signaling and reduced IGF-1 receptor signaling with aging promotes CPC and cardiomyocyte apoptosis.

CPC from the secondary/anterior heart field (i.e., Nestin⁺) are resident in the secondary heart field regions of the rat heart (Drapeau et al., 2005). The recent review by Di Felice (Di Felice & Zummo, 2009) surveyed the many mutations of secondary heart field cells that are associated with human Tetralogy of Fallot. They concluded that potentially improved approaches to therapy would be better informed by a clearer understanding of the behavior and patterns of migration of CPC of the secondary heart field. That neural stem cells (CPC

from the secondary heart field) may participate in recovery from myocardial injury in a region-specific manner has also been shown. These cells apparently migrate and home to infarcted region of rat hearts, although they differentiate into neuronal, not myocardial cells (Beguin et al., 2011). Tamura (Tamura et al., 2011) used a mouse transgenic approach to tag neural crest cells and show that these cells migrated to the ischemic border zone of an infarct and transdifferentiated into cardiomyocytes. Although evidence for Nestin⁺ cells in the human heart is lacking, it is worth considering the different heart field lineages in the context of devising strategies for therapeutic targeting of human CPC. The targeting of neural crest-derived CPC to correct late ventricular arrhythmias occurring in patients after surgical repair operations for TOF is a potential therapeutic strategy (Di Felice & Zummo, 2009).

3.2 Potential roles of CPC in therapy for CHD

3.2.1 CPC silently contribute to therapy

3.2.1.1 (Pre-) Failing Fontan rescue via activation-RV strengthening

In the setting of failing Fontan physiology the potential for boosting cardiac performance through manipulation of cell number represents a new horizon. Ventricular assist devices (VAD) are presently used as a bridge to transplant in pediatric Fontan patients (Fynn-Thompson & Almond, 2007). VAD are also used to reduce ventricular load with the objective of enhancing the ability of the ventricle to support a greater load, i.e., to “rest” or re-train the ventricle of patients with dilated cardiomyopathy (CMP), even restoring function to the point of enabling pump removal (Birks et al., 2011). It is conceivable that the use of VAD may increase cardiomyocyte number, and evidence of this comes from recent prospective studies of myocardial biopsies showing an increased number of diploid myocytes in end stage congestive heart failure patients supported as a bridge to transplant with LVAD (Wohlschlaeger et al., 2010). Manginas (Manginas et al., 2009) has shown that endothelial progenitor cells are also mobilized by VAD use in patients. EPC may be stimulated to home into myocardium supported by VAD and improve myocardial function. As reviewed by Tsiavou (Tsiavou & Manginas, 2010), rescue of myocardial function during cardiac support by VAD may be mediated by CD45⁺ EPC promoting neovascularization and transdifferentiating into or fusing with cardiomyocytes. However, the possible involvement of CPC fusion with existing myocytes as repair mechanism is not widely accepted. The implications of the above observations are that progenitor cells may at least be participating in, if not be a primary mechanism mediating these physiological changes observed with VAD therapy.

What physiological mechanisms might underlie the expansion of the myocardial cell population during VAD support? Current thinking is that paracrine mechanisms are involved. For example, mechanical stimulation by the VAD may induce the production of cytokines such as growth factors (see section 3.1.3.1), which then stimulate mitotic expansion of CPC. Additionally, the production of chemokines under the influence of the same mechanical stimulation may promote the homing of bone marrow-derived progenitor cells. Indeed, much of the success of cell delivery-based therapies, although still rather modest, are thought to be largely due to paracrine effects mediated through the injected cells, the carrier media or in some cases the physical-mechanical changes induced by injection of liquid boluses into the muscle wall.

3.2.2 Augmenting current therapy with CPC targeting

3.2.2.1 CPC activation as an adjuvant to VAD “resting” of ventricle

It has been reported that pharmacological therapy in conjunction with VAD support may enhance myocyte population expansion. Recent studies by Soppa used therapy with the beta-2 agonist clenbuterol (Soppa et al., 2008) in a murine heterotopic abdominal transplant of failing hearts. Testing the effects of clenbuterol on mechanical unloading, they were able to show that it improves LV function. Bhavsar demonstrated that clenbuterol positively affects cardiac physiology through myocyte hypertrophy, concluding that the effect was mediated by a paracrine action of fibroblast-derived IGF-1 (Bhavsar et al., 2010). At the present time, we can find no reports of an effect of clenbuterol on CPC recruitment/activation. It is conceivable that such combined pharmacological plus mechanical therapy approaches could provide a markedly re-strengthened ventricle capable of many more years of function if not for the rest of the patient’s life. CPC recruitment could be combined with VAD-mediated myocardial unloading to augment myocardial tissue while “resting” the ventricle, whether left or right sided. Another possible use of therapies designed to expand the population of CPC is in the area of “ventricular training” a potential approach to prepare the LV of, e.g., delayed repair TGA patients for greater force production once the arterial switch operation has been performed. If VAD use can augment cardiomyocyte populations in otherwise normal ventricles, this could potentially help prepare the TGA patient for the switch operation by enhancing ventricular adaptation to increased loads.

3.2.2.2 Cardiomyopathy

Is it realistic to expect the manipulation of resident CPC populations to achieve a reversal in the decline in myocardial function in the setting of cardiomyopathy? Given the genetic nature of known lesions in sarcomeric proteins in this disease (Frazier et al., 2011), one may anticipate that the progenitor cell population may also harbor the same mutant alleles and therefore the expansion of that population may provide no benefit. However, for cardiomyopathy induced by chemical injury such as doxorubicin/adriamycin therapy (Shi et al., 2011), strategies to promote expansion of the resident CPC population could be considered an adjuvant or co-therapy used to mitigate cardiotoxicity. It is important to consider that, although the resident CPC may have an advantage in already being present within the muscular wall of the heart, therapies designed to help recruit bone marrow-derived CPC (and EPC) (see section 3.1.3.1) are certainly worthy of exploration while investigators try to understand the possibly different significance of resident versus bone marrow-derived CPC. Indeed, given the mismatch between patients in need of transplantation and the availability of transplantable hearts there would seem to be little reason not to emphasize the exploration of multiple approaches designed to utilize endogenous progenitor cells whether they be resident in the tissue or delivered to the tissue from sites such as the bone marrow.

3.2.2.3 Summary: Potential therapeutic opportunities

At present, investigators have only begun to exploit the potentials of using small molecule (drugs, growth factors) based therapies to expand desirable cell populations. We know of no examples of demonstrated CPC-based therapies for congenital cardiac disease at this time. Since CPC likely undergo senescence and their number as a percentage of total cells declines

with expansion of the myocardial cell population, conceivably the benefits of such therapy may be far greater for the pediatric population than the adult.

4. Conclusions

The presence of resident CPC in myocardium is well supported through multiple studies. There is still much independent confirmation to be completed to clarify the promise of cardiac progenitor cell-mediated repair. Importantly, the major novel discoveries in the field of CPC biology have been made by only a small group of investigators and interpretation of some data is impaired by the lack of independent corroborating studies. Controversies also continue regarding the origin of CPC: during cardiac development, e.g., *Isl1*⁺ CPC, or from bone marrow, e.g., *c-Kit*⁺ CPC. Most likely, both sources are important but their therapeutic utilization may need to be approached with different strategies. Methods for activating resident CPC to realize their potential for effecting endogenous cardiac repair are still in the early discovery period. The fundamental question of CPC role in homeostatic maintenance of the myocardium throughout life has yet to be fully clarified, although an understanding of this highly significant role appears to be limited only by a lack of detection. Nonetheless, the potential applications of CPC-focused therapies in congenital heart disease treatments are likely manifold, awaiting only further investigation and implementation.

5. Acknowledgement

We thank Olaf Reinhartz, MD for reviewing this manuscript, and Laura Piechura for editorial assistance.

6. References

- Abu-Issa, R. & Kirby, M. L. (2007). Heart Field: From Mesoderm to Heart Tube. *Annual review of cell and developmental biology* Vol.23, (2007) pp. 45-68, ISBN 1081-0706 (Print) 1081-0706 (Linking)
- Abu-Issa, R., Smyth, G., Smoak, I., Yamamura, K. & Meyers, E. N. (2002). Fgf8 Is Required for Pharyngeal Arch and Cardiovascular Development in the Mouse. *Development* Vol.129, No.19, (Oct, 2002) pp. 4613-4625, ISBN 0950-1991 (Print) 0950-1991 (Linking)
- Amir, G., Ma, X., Reddy, V. M., Hanley, F. L., Reinhartz, O., Ramamoorthy, C. & Riemer, R. K. (2008). Dynamics of Human Myocardial Progenitor Cell Populations in the Neonatal Period. *The Annals of thoracic surgery* Vol.86, No.4, (2008) pp. 1311-1319, ISBN 1552-6259
- Basson, C. T., Bachinsky, D. R., Lin, R. C., Levi, T., Elkins, J. A., Soultz, J., Grayzel, D., Kroumpouzou, E., Traill, T. A., Leblanc-Straceski, J., Renault, B., Kucherlapati, R., Seidman, J. G. & Seidman, C. E. (1997). Mutations in Human *Tbx5* [Corrected] Cause Limb and Cardiac Malformation in Holt-Oram Syndrome. *Nature genetics* Vol.15, No.1, (Jan, 1997) pp. 30-35, ISBN 1061-4036 (Print) 1061-4036 (Linking)
- Bayes-Genis, A., Roura, S., Prat-Vidal, C., Farre, J., Soler-Botija, C., de Luna, A. B. & Cinca, J. (2007). Chimerism and Microchimerism of the Human Heart: Evidence for Cardiac Regeneration. *Nature clinical practice. Cardiovascular medicine* Vol.4 Suppl 1, (Feb, 2007) pp. S40-45, ISBN 1743-4300 (Electronic) 1743-4297 (Linking)

- Bearzi, C., Rota, M., Hosoda, T., Tillmanns, J., Nascimbene, A., De Angelis, A., Yasuzawa-Amano, S., Trofimova, I., Siggins, R. W., Lecapitaine, N., Cascapera, S., Beltrami, A. P., D'Alessandro, D. A., Zias, E., Quaini, F., Urbanek, K., Michler, R. E., Bolli, R., Kajstura, J., Leri, A. & Anversa, P. (2007). Human Cardiac Stem Cells. *Proc Natl Acad Sci U S A* Vol.104, No.35, (Aug 28, 2007) pp. 14068-14073, ISBN 0027-8424 (Print) 0027-8424 (Linking)
- Beguín, P. C., El-Helou, V., Gillis, M. A., Duquette, N., Gosselin, H., Brugada, R., Villeneuve, L., Lauzier, D., Tanguay, J. F., Ribuot, C. & Calderone, A. (2011). Nestin (+) Stem Cells Independently Contribute to Neural Remodelling of the Ischemic Heart. *Journal of cellular physiology* Vol.226, No.5, (May, 2011) pp. 1157-1165, ISBN 1097-4652 (Electronic) 0021-9541 (Linking)
- Beltrami, A. P., Barlucchi, L., Torella, D., Baker, M., Limana, F., Chimenti, S., Kasahara, H., Rota, M., Musso, E., Urbanek, K., Leri, A., Kajstura, J., Nadal-Ginard, B. & Anversa, P. (2003). Adult Cardiac Stem Cells Are Multipotent and Support Myocardial Regeneration. *Cell* Vol.114, No.6, (Sep 19, 2003) pp. 763-776, ISBN 0092-8674 (Print) 0092-8674 (Linking)
- Bergmann, O., Bhardwaj, R. D., Bernard, S., Zdunek, S., Barnabe-Heider, F., Walsh, S., Zupicich, J., Alkass, K., Buchholz, B. A., Druid, H., Jovinge, S. & Frisen, J. (2009). Evidence for Cardiomyocyte Renewal in Humans. *Science* Vol.324, No.5923, (Apr 3, 2009) pp. 98-102, ISBN 1095-9203 (Electronic) 0036-8075 (Linking)
- Bhavsar, P. K., Brand, N. J., Felkin, L. E., Luther, P. K., Cullen, M. E., Yacoub, M. H. & Barton, P. J. (2010). Clenbuterol Induces Cardiac Myocyte Hypertrophy Via Paracrine Signalling and Fibroblast-Derived Igf-1. *Journal of cardiovascular translational research* Vol.3, No.6, (Dec, 2010) pp. 688-695, ISBN 1937-5395 (Electronic)
- Birks, E. J., George, R. S., Hedger, M., Bahrami, T., Wilton, P., Bowles, C. T., Webb, C., Bougard, R., Amrani, M., Yacoub, M. H., Dreyfus, G. & Khaghani, A. (2011). Reversal of Severe Heart Failure with a Continuous-Flow Left Ventricular Assist Device and Pharmacological Therapy: A Prospective Study. *Circulation* Vol.123, No.4, (Feb 1, 2011) pp. 381-390, ISBN 1524-4539 (Electronic) 0009-7322 (Linking)
- Bolli, P. & Chaudhry, H. W. (2010). Molecular Physiology of Cardiac Regeneration. *Annals of the New York Academy of Sciences* Vol.1211, (Nov, 2010) pp. 113-126, ISBN 1749-6632 (Electronic) 0077-8923 (Linking)
- Bollini, S., Smart, N. & Riley, P. R. (2011). Resident Cardiac Progenitor Cells: At the Heart of Regeneration. *Journal of molecular and cellular cardiology* Vol.50, No.2, (Feb, 2011) pp. 296-303, ISBN 1095-8584 (Electronic) 0022-2828 (Linking)
- Boni, A., Urbanek, K., Nascimbene, A., Hosoda, T., Zheng, H., Delucchi, F., Amano, K., Gonzalez, A., Vitale, S., Ojaimi, C., Rizzi, R., Bolli, R., Yutzey, K. E., Rota, M., Kajstura, J., Anversa, P. & Leri, A. (2008). Notch1 Regulates the Fate of Cardiac Progenitor Cells. *Proc Natl Acad Sci U S A* Vol.105, No.40, (Oct 7, 2008) pp. 15529-15534, ISBN 1091-6490 (Electronic) 0027-8424 (Linking)
- Bradshaw, L., Chaudhry, B., Hildreth, V., Webb, S. & Henderson, D. J. (2009). Dual Role for Neural Crest Cells During Outflow Tract Septation in the Neural Crest-Deficient Mutant *Splotch*(2h). *Journal of anatomy* Vol.214, No.2, (Feb, 2009) pp. 245-257, ISBN 1469-7580 (Electronic) 0021-8782 (Linking)
- Cai, C. L., Liang, X., Shi, Y., Chu, P. H., Pfaff, S. L., Chen, J. & Evans, S. (2003). *Isl1* Identifies a Cardiac Progenitor Population That Proliferates Prior to Differentiation and

- Contributes a Majority of Cells to the Heart. *Dev Cell* Vol.5, No.6, (Dec, 2003) pp. 877-889, ISBN 1534-5807 (Print) 1534-5807 (Linking)
- D'Amario, D., Cabral-Da-Silva, M. C., Zheng, H., Fiorini, C., Goichberg, P., Steadman, E., Ferreira-Martins, J., Sanada, F., Piccoli, M., Cappetta, D., D'Alessandro, D. A., Michler, R. E., Hosoda, T., Anastasia, L., Rota, M., Leri, A., Anversa, P. & Kajstura, J. (2011). Insulin-Like Growth Factor-1 Receptor Identifies a Pool of Human Cardiac Stem Cells with Superior Therapeutic Potential for Myocardial Regeneration. *Circulation research* Vol.108, No.12, (Jun 10, 2011) pp. 1467-1481, ISBN 1524-4571 (Electronic) 0009-7330 (Linking)
- Dettman, R. W., Denetclaw, W., Jr., Ordahl, C. P. & Bristow, J. (1998). Common Epicardial Origin of Coronary Vascular Smooth Muscle, Perivascular Fibroblasts, and Intermycardial Fibroblasts in the Avian Heart. *Developmental biology* Vol.193, No.2, (Jan 15, 1998) pp. 169-181, ISBN 0012-1606 (Print) 0012-1606 (Linking)
- Di Felice, V. & Zummo, G. (2009). Tetralogy of Fallot as a Model to Study Cardiac Progenitor Cell Migration and Differentiation During Heart Development. *Trends Cardiovasc Med* Vol.19, No.4, (May, 2009) pp. 130-135, ISBN 1873-2615 (Electronic) 1050-1738 (Linking)
- Drapeau, J., El-Helou, V., Clement, R., Bel-Hadj, S., Gosselin, H., Trudeau, L. E., Villeneuve, L. & Calderone, A. (2005). Nestin-Expressing Neural Stem Cells Identified in the Scar Following Myocardial Infarction. *Journal of cellular physiology* Vol.204, No.1, (Jul, 2005) pp. 51-62, ISBN 0021-9541 (Print) 0021-9541 (Linking)
- Dyer, L. A. & Kirby, M. L. (2009). The Role of Secondary Heart Field in Cardiac Development. *Developmental biology* Vol.336, No.2, (Dec 15, 2009) pp. 137-144, ISBN 1095-564X (Electronic) 0012-1606 (Linking)
- Frazier, A. H., Ramirez-Correa, G. A. & Murphy, A. M. (2011). Molecular Mechanisms of Sarcomere Dysfunction in Dilated and Hypertrophic Cardiomyopathy. *Progress in pediatric cardiology* Vol.31, No.1, (Jan 1, 2011) pp. 29-33, ISBN 1058-9813 (Print) 1058-9813 (Linking)
- Fynn-Thompson, F. & Almond, C. (2007). Pediatric Ventricular Assist Devices. *Pediatric cardiology* Vol.28, No.2, (Mar-Apr, 2007) pp. 149-155, ISBN 0172-0643 (Print) 0172-0643 (Linking)
- Garcia-Martinez, V. & Schoenwolf, G. C. (1993). Primitive-Streak Origin of the Cardiovascular System in Avian Embryos.Pdf. *Developmental Biology* Vol.159, 1993) pp. 706-719, ISBN
- Garg, V., Kathiriyai, I. S., Barnes, R., Schluterman, M. K., King, I. N., Butler, C. A., Rothrock, C. R., Eapen, R. S., Hirayama-Yamada, K., Joo, K., Matsuoka, R., Cohen, J. C. & Srivastava, D. (2003). Gata4 Mutations Cause Human Congenital Heart Defects and Reveal an Interaction with Tbx5. *Nature* Vol.424, No.6947, (Jul 24, 2003) pp. 443-447, ISBN 1476-4687 (Electronic) 0028-0836 (Linking)
- Gonzales, C. & Pedrazzini, T. (2009). Progenitor Cell Therapy for Heart Disease. *Exp Cell Res* Vol.315, No.18, (Nov 1, 2009) pp. 3077-3085, ISBN 1090-2422 (Electronic) 0014-4827 (Linking)
- Gonzalez, A., Rota, M., Nurzynska, D., Misao, Y., Tillmanns, J., Ojaimi, C., Padin-Iruegas, M. E., Muller, P., Esposito, G., Bearzi, C., Vitale, S., Dawn, B., Sanganalmath, S. K., Baker, M., Hintze, T. H., Bolli, R., Urbanek, K., Hosoda, T., Anversa, P., Kajstura, J. & Leri, A. (2008). Activation of Cardiac Progenitor Cells Reverses the Failing Heart


- Senescent Phenotype and Prolongs Lifespan. *Circ Res* Vol.102, No.5, (Mar 14, 2008) pp. 597-606, ISBN
- Hutson, M. R. & Kirby, M. L. (2007). Model Systems for the Study of Heart Development and Disease. Cardiac Neural Crest and Conotruncal Malformations. *Seminars in cell & developmental biology* Vol.18, No.1, (Feb, 2007) pp. 101-110, ISBN 1084-9521 (Print) 1084-9521 (Linking)
- Jain, R., Rentschler, S. & Epstein, J. A. (2010). Notch and Cardiac Outflow Tract Development. *Annals of the New York Academy of Sciences* Vol.1188, (Feb, 2010) pp. 184-190, ISBN 1749-6632 (Electronic) 0077-8923 (Linking)
- Jerome, L. A. & Papaioannou, V. E. (2001). Digeorge Syndrome Phenotype in Mice Mutant for the T-Box Gene, *Tbx1*. *Nature genetics* Vol.27, No.3, (Mar, 2001) pp. 286-291, ISBN 1061-4036 (Print) 1061-4036 (Linking)
- Kelly, R. G., Brown, N. A. & Buckingham, M. E. (2001). The Arterial Pole of the Mouse Heart Forms from *Fgf10*-Expressing Cells in Pharyngeal Mesoderm. *Developmental cell* Vol.1, No.3, (Sep, 2001) pp. 435-440, ISBN 1534-5807 (Print) 1534-5807 (Linking)
- Kirby, M. L. & Waldo, K. L. (1995). Neural Crest and Cardiovascular Patterning. *Circulation research* Vol.77, No.2, (Aug, 1995) pp. 211-215, ISBN 0009-7330 (Print) 0009-7330 (Linking)
- Laflamme, M. A. & Murry, C. E. (2011). Heart Regeneration. *Nature* Vol.473, No.7347, (May 19, 2011) pp. 326-335, ISBN 1476-4687 (Electronic) 0028-0836 (Linking)
- Laugwitz, K. L., Moretti, A., Lam, J., Gruber, P., Chen, Y., Woodard, S., Lin, L. Z., Cai, C. L., Lu, M. M., Reth, M., Platoshyn, O., Yuan, J. X., Evans, S. & Chien, K. R. (2005). Postnatal *Isl1*⁺ Cardioblasts Enter Fully Differentiated Cardiomyocyte Lineages. *Nature* Vol.433, No.7026, (Feb 10, 2005) pp. 647-653, ISBN 1476-4687 (Electronic) 0028-0836 (Linking)
- Lemischka, I. R. (1997). Microenvironmental Regulation of Hematopoietic Stem Cells. *Stem cells* Vol.15 Suppl 1, 1997) pp. 63-68, ISBN 1066-5099 (Print) 1066-5099 (Linking)
- Leri, A., Barlucchi, L., Limana, F., Deptala, A., Darzynkiewicz, Z., Hintze, T. H., Kajstura, J., Nadal-Ginard, B. & Anversa, P. (2001). Telomerase Expression and Activity Are Coupled with Myocyte Proliferation and Preservation of Telomeric Length in the Failing Heart. *Proceedings of the National Academy of Sciences of the United States of America* Vol.98, No.15, (Jul 17, 2001) pp. 8626-8631, ISBN 0027-8424 (Print) 0027-8424 (Linking)
- Limana, F., Germani, A., Zacheo, A., Kajstura, J., Di Carlo, A., Borsellino, G., Leoni, O., Palumbo, R., Battistini, L., Rastaldo, R., Muller, S., Pompilio, G., Anversa, P., Bianchi, M. E. & Capogrossi, M. C. (2005). Exogenous High-Mobility Group Box 1 Protein Induces Myocardial Regeneration after Infarction Via Enhanced Cardiac C-Kit⁺ Cell Proliferation and Differentiation. *Circ Res* Vol.97, No.8, (Oct 14, 2005) pp. e73-83, ISBN 1524-4571 (Electronic) 0009-7330 (Linking)
- Linke, A., Muller, P., Nurzynska, D., Casarsa, C., Torella, D., Nascimbene, A., Castaldo, C., Cascapera, S., Bohm, M., Quaini, F., Urbanek, K., Leri, A., Hintze, T. H., Kajstura, J. & Anversa, P. (2005). Stem Cells in the Dog Heart Are Self-Renewing, Clonogenic, and Multipotent and Regenerate Infarcted Myocardium, Improving Cardiac Function. *Proc Natl Acad Sci U S A* Vol.102, No.25, (Jun 21, 2005) pp. 8966-8971, ISBN 0027-8424 (Print) 0027-8424 (Linking)

- Manginas, A., Tsiavou, A., Sfyraakis, P., Giamouzis, G., Tsourelis, L., Leontiadis, E., Degiannis, D., Cokkinos, D. V. & Alivizatos, P. A. (2009). Increased Number of Circulating Progenitor Cells after Implantation of Ventricular Assist Devices. *The Journal of heart and lung transplantation : the official publication of the International Society for Heart Transplantation* Vol.28, No.7, (Jul, 2009) pp. 710-717, ISBN 1557-3117 (Electronic) 1053-2498 (Linking)
- Martinez-Estrada, O. M., Lettice, L. A., Essafi, A., Guadix, J. A., Slight, J., Velecela, V., Hall, E., Reichmann, J., Devenney, P. S., Hohenstein, P., Hosen, N., Hill, R. E., Munoz-Chapuli, R. & Hastie, N. D. (2010). Wt1 Is Required for Cardiovascular Progenitor Cell Formation through Transcriptional Control of Snail and E-Cadherin. *Nat Genet* Vol.42, No.1, (Jan, 2010) pp. 89-93, ISBN 1546-1718 (Electronic) 1061-4036 (Linking)
- Merscher, S., Funke, B., Epstein, J. A., Heyer, J., Puech, A., Lu, M. M., Xavier, R. J., Demay, M. B., Russell, R. G., Factor, S., Tokooya, K., Jore, B. S., Lopez, M., Pandita, R. K., Lia, M., Carrion, D., Xu, H., Schorle, H., Kobler, J. B., Scambler, P., Wynshaw-Boris, A., Skoultschi, A. I., Morrow, B. E. & Kucherlapati, R. (2001). Tbx1 Is Responsible for Cardiovascular Defects in Velo-Cardio-Facial/Digeorge Syndrome. *Cell* Vol.104, No.4, (Feb 23, 2001) pp. 619-629, ISBN 0092-8674 (Print) 0092-8674 (Linking)
- Messina, E., De Angelis, L., Frati, G., Morrone, S., Chimenti, S., Fiordaliso, F., Salio, M., Battaglia, M., Latronico, M. V., Coletta, M., Vivarelli, E., Frati, L., Cossu, G. & Giacomello, A. (2004). Isolation and Expansion of Adult Cardiac Stem Cells from Human and Murine Heart. *Circulation research* Vol.95, No.9, (Oct 29, 2004) pp. 911-921, ISBN 1524-4571 (Electronic) 0009-7330 (Linking)
- Moraes, F., Novoa, A., Jerome-Majewska, L. A., Papaioannou, V. E. & Mallo, M. (2005). Tbx1 Is Required for Proper Neural Crest Migration and to Stabilize Spatial Patterns During Middle and Inner Ear Development. *Mechanisms of development* Vol.122, No.2, (Feb, 2005) pp. 199-212, ISBN 0925-4773 (Print) 0925-4773 (Linking)
- Moretti, A., Caron, L., Nakano, A., Lam, J. T., Bernshausen, A., Chen, Y., Qyang, Y., Bu, L., Sasaki, M., Martin-Puig, S., Sun, Y., Evans, S. M., Laugwitz, K. L. & Chien, K. R. (2006). Multipotent Embryonic Isl1+ Progenitor Cells Lead to Cardiac, Smooth Muscle, and Endothelial Cell Diversification. *Cell* Vol.127, No.6, (Dec 15, 2006) pp. 1151-1165, ISBN 0092-8674 (Print) 0092-8674 (Linking)
- Morrison, S. J. & Spradling, A. C. (2008). Stem Cells and Niches: Mechanisms That Promote Stem Cell Maintenance Throughout Life. *Cell* Vol.132, No.4, (Feb 22, 2008) pp. 598-611, ISBN 1097-4172 (Electronic) 0092-8674 (Linking)
- Murry, C. E., Field, L. J. & Menasche, P. (2005). Cell-Based Cardiac Repair: Reflections at the 10-Year Point. *Circulation* Vol.112, No.20, 2005) pp. 3174-3183, ISBN 1524-4539
- Murry, C. E., Soonpaa, M. H., Reinecke, H., Nakajima, H., Nakajima, H. O., Rubart, M., Pasumarthi, K. B. S., Virag, J. I., Bartelmez, S. H., Poppa, V., Bradford, G., Dowell, J. D., Williams, D. A. & Field, L. J. (2004). Haematopoietic Stem Cells Do Not Transdifferentiate into Cardiac Myocytes in Myocardial Infarcts. *Nature* Vol.428, No.6983, 2004) pp. 664-668, ISBN 1476-4687
- Nielsen, J. C., Lytrivi, I. D., Ko, H. H., Yau, J., Bhatla, P., Parness, I. A. & Srivastava, S. (2010). The Accuracy of Echocardiographic Assessment of Left Ventricular Size in Children by the 5/6 Area X Length (Bullet) Method. *Echocardiography* Vol.27, No.6, (Jul, 2010) pp. 691-695, ISBN 1540-8175 (Electronic) 0742-2822 (Linking)

- Nishibatake, M., Kirby, M. L. & Van Mierop, L. H. (1987). Pathogenesis of Persistent Truncus Arteriosus and Dextroposed Aorta in the Chick Embryo after Neural Crest Ablation. *Circulation* Vol.75, No.1, (Jan, 1987) pp. 255-264, ISBN 0009-7322 (Print) 0009-7322 (Linking)
- Oh, H., Bradfute, S. B., Gallardo, T. D., Nakamura, T., Gaussin, V., Mishina, Y., Pocius, J., Michael, L. H., Behringer, R. R., Garry, D. J., Entman, M. L. & Schneider, M. D. (2003). Cardiac Progenitor Cells from Adult Myocardium: Homing, Differentiation, and Fusion after Infarction. *Proc Natl Acad Sci U S A* Vol.100, No.21, (Oct 14, 2003) pp. 12313-12318, ISBN 0027-8424 (Print) 0027-8424 (Linking)
- Oh, H. & Schneider, M. D. (2002). The Emerging Role of Telomerase in Cardiac Muscle Cell Growth and Survival. *Journal of molecular and cellular cardiology* Vol.34, No.7, (Jul, 2002) pp. 717-724, ISBN 0022-2828 (Print) 0022-2828 (Linking)
- Orlic, D., Kajstura, J., Chimenti, S., Jakoniuk, I., Anderson, S. M., Li, B., Pickel, J., McKay, R., Nadal-Ginard, B., Bodine, D. M., Leri, A. & Anversa, P. (2001). Bone Marrow Cells Regenerate Infarcted Myocardium. *Nature* Vol.410, No.6829, (Apr 5, 2001) pp. 701-705, ISBN 0028-0836 (Print) 0028-0836 (Linking)
- Palumbo, R. & Bianchi, M. E. (2004). High Mobility Group Box 1 Protein, a Cue for Stem Cell Recruitment. *Biochemical pharmacology* Vol.68, No.6, (Sep 15, 2004) pp. 1165-1170, ISBN 0006-2952 (Print) 0006-2952 (Linking)
- Pouly, J., Bruneval, P., Mandet, C., Proksch, S., Peyrard, S., Amrein, C., Bousseaux, V., Guillemain, R., Deloche, A., Fabiani, J. N. & Menasche, P. (2008). Cardiac Stem Cells in the Real World. *J Thorac Cardiovasc Surg* Vol.135, No.3, (Mar, 2008) pp. 673-678, ISBN 1097-685X (Electronic) 0022-5223 (Linking)
- Rossant, J. (2008). Stem Cells and Early Lineage Development. *Cell* Vol.132, No.4, (Feb 22, 2008) pp. 527-531, ISBN 1097-4172 (Electronic) 0092-8674 (Linking)
- Saga, Y., Miyagawa-Tomita, S., Takagi, A., Kitajima, S., Miyazaki, J. & Inoue, T. (1999). Mesp1 Is Expressed in the Heart Precursor Cells and Required for the Formation of a Single Heart Tube. *Development* Vol.126, No.15, (Aug, 1999) pp. 3437-3447, ISBN 0950-1991 (Print) 0950-1991 (Linking)
- Sairanen, H. & Louhimo, I. (1992). Dimensions of the Heart and Great Vessels in Normal Children. A Postmortem Study of Cardiac Ventricles, Valves and Great Vessels. *Scandinavian journal of thoracic and cardiovascular surgery* Vol.26, No.2, 1992) pp. 83-92, ISBN 0036-5580 (Print) 0036-5580 (Linking)
- Schenke-Layland, K., Nsair, A., Van Handel, B., Angelis, E., Gluck, J. M., Votteler, M., Goldhaber, J. I., Mikkola, H. K., Kahn, M. & Maclellan, W. R. (2011). Recapitulation of the Embryonic Cardiovascular Progenitor Cell Niche. *Biomaterials* Vol.32, No.11, (Apr, 2011) pp. 2748-2756, ISBN 1878-5905 (Electronic) 0142-9612 (Linking)
- Shi, Y., Moon, M., Dawood, S., McManus, B. & Liu, P. P. (2011). Mechanisms and Management of Doxorubicin Cardiotoxicity. *Herz* (Jun 9, 2011) pp. ISBN 1615-6692 (Electronic) 0340-9937 (Linking)
- Smart, N., Risebro, C. A., Clark, J. E., Ehler, E., Miquerol, L., Rossdeutsch, A., Marber, M. S. & Riley, P. R. (2010). Thymosin Beta4 Facilitates Epicardial Neovascularization of the Injured Adult Heart. *Annals of the New York Academy of Sciences* Vol.1194, (Apr, 2010) pp. 97-104, ISBN 1749-6632 (Electronic) 0077-8923 (Linking)
- Soppa, G. K., Lee, J., Stagg, M. A., Felkin, L. E., Barton, P. J., Siedlecka, U., Youssef, S., Yacoub, M. H. & Terracciano, C. M. (2008). Role and Possible Mechanisms of

- Clenbuterol in Enhancing Reverse Remodelling During Mechanical Unloading in Murine Heart Failure. *Cardiovascular research* Vol.77, No.4, (Mar 1, 2008) pp. 695-706, ISBN 0008-6363 (Print) 0008-6363 (Linking)
- Spradling, A., Drummond-Barbosa, D. & Kai, T. (2001). Stem Cells Find Their Niche. *Nature* Vol.414, No.6859, (Nov 1, 2001) pp. 98-104, ISBN 0028-0836 (Print) 0028-0836 (Linking)
- Sturzu, A. C. & Wu, S. M. (2011). Developmental and Regenerative Biology of Multipotent Cardiovascular Progenitor Cells. *Circ Res* Vol.108, No.3, (Feb 4, 2011) pp. 353-364, ISBN 1524-4571 (Electronic) 0009-7330 (Linking)
- Sui, R., Liao, X., Zhou, X. & Tan, Q. (2011). The Current Status of Engineering Myocardial Tissue. *Stem cell reviews* Vol.7, No.1, (Mar, 2011) pp. 172-180, ISBN 1558-6804 (Electronic) 1550-8943 (Linking)
- Sun, X., Meyers, E. N., Lewandoski, M. & Martin, G. R. (1999). Targeted Disruption of Fgf8 Causes Failure of Cell Migration in the Gastrulating Mouse Embryo. *Genes & development* Vol.13, No.14, (Jul 15, 1999) pp. 1834-1846, ISBN 0890-9369 (Print) 0890-9369 (Linking)
- Takeuchi, J. K. & Bruneau, B. G. (2009). Directed Transdifferentiation of Mouse Mesoderm to Heart Tissue by Defined Factors. *Nature* Vol.459, No.7247, (Jun 4, 2009) pp. 708-711, ISBN 1476-4687 (Electronic) 0028-0836 (Linking)
- Tallini, Y. N., Greene, K. S., Craven, M., Spealman, A., Breitbach, M., Smith, J., Fisher, P. J., Steffey, M., Hesse, M., Doran, R. M., Woods, A., Singh, B., Yen, A., Fleischmann, B. K. & Kotlikoff, M. I. (2009). C-Kit Expression Identifies Cardiovascular Precursors in the Neonatal Heart. *Proceedings of the National Academy of Sciences of the United States of America* Vol.106, No.6, (Feb 10, 2009) pp. 1808-1813, ISBN 1091-6490 (Electronic) 0027-8424 (Linking)
- Tam, P. P., Parameswaran, M., Kinder, S. J. & Weinberger, R. P. (1997). The Allocation of Epiblast Cells to the Embryonic Heart and Other Mesodermal Lineages: The Role of Ingression and Tissue Movement During Gastrulation. *Development* Vol.124, No.9, (May, 1997) pp. 1631-1642, ISBN 0950-1991 (Print) 0950-1991 (Linking)
- Tamura, Y., Matsumura, K., Sano, M., Tabata, H., Kimura, K., Ieda, M., Arai, T., Ohno, Y., Kanazawa, H., Yuasa, S., Kaneda, R., Makino, S., Nakajima, K., Okano, H. & Fukuda, K. (2011). Neural Crest-Derived Stem Cells Migrate and Differentiate into Cardiomyocytes after Myocardial Infarction. *Arterioscler Thromb Vasc Biol* Vol.31, No.3, (Mar, 2011) pp. 582-589, ISBN 1524-4636 (Electronic) 1079-5642 (Linking)
- Tsiavou, A. & Manginas, A. (2010). Dynamics of Progenitor Cells and Ventricular Assist Device Intervention. *Journal of cardiovascular translational research* Vol.3, No.2, (Apr, 2010) pp. 147-152, ISBN 1937-5395 (Electronic)
- Tsuchihashi, T., Maeda, J., Shin, C. H., Ivey, K. N., Black, B. L., Olson, E. N., Yamagishi, H. & Srivastava, D. (2011). Hand2 Function in Second Heart Field Progenitors Is Essential for Cardiogenesis. *Developmental biology* Vol.351, No.1, (Mar 1, 2011) pp. 62-69, ISBN 1095-564X (Electronic) 0012-1606 (Linking)
- Urbanek, K., Cesselli, D., Rota, M., Nascimbene, A., De Angelis, A., Hosoda, T., Bearzi, C., Boni, A., Bolli, R., Kajstura, J., Anversa, P. & Leri, A. (2006). Stem Cell Niches in the Adult Mouse Heart. *Proc Natl Acad Sci U S A* Vol.103, No.24, (Jun 13, 2006) pp. 9226-9231, ISBN 0027-8424 (Print) 0027-8424 (Linking)

- Urbanek, K., Quaini, F., Tasca, G., Torella, D., Castaldo, C., Nadal-Ginard, B., Leri, A., Kajstura, J., Quaini, E. & Anversa, P. (2003). Intense Myocyte Formation from Cardiac Stem Cells in Human Cardiac Hypertrophy. *Proc Natl Acad Sci U S A* Vol.100, No.18, (Sep 2, 2003) pp. 10440-10445, ISBN 0027-8424 (Print) 0027-8424 (Linking)
- Vincent, S. D. & Buckingham, M. E. (2010). How to Make a Heart: The Origin and Regulation of Cardiac Progenitor Cells. *Curr Top Dev Biol* Vol.90, 2010) pp. 1-41, ISBN 1557-8933 (Electronic) 0070-2153 (Linking)
- Vitelli, F., Morishima, M., Taddei, I., Lindsay, E. A. & Baldini, A. (2002). Tbx1 Mutation Causes Multiple Cardiovascular Defects and Disrupts Neural Crest and Cranial Nerve Migratory Pathways. *Human molecular genetics* Vol.11, No.8, (Apr 15, 2002) pp. 915-922, ISBN 0964-6906 (Print) 0964-6906 (Linking)
- Ward, H. H., Romero, E., Welford, A., Pickett, G., Bacallao, R., Gattone, V. H., 2nd, Ness, S. A., Wandinger-Ness, A. & Roitbak, T. (2011). Adult Human Cd133/1(+) Kidney Cells Isolated from Papilla Integrate into Developing Kidney Tubules. *Biochimica et biophysica acta*(Jan 19, 2011) pp. ISBN 0006-3002 (Print) 0006-3002 (Linking)
- Williams, A. R., Trachtenberg, B., Velazquez, D. L., McNiece, I., Altman, P., Rouy, D., Mendizabal, A. M., Pattany, P. M., Lopera, G. A., Fishman, J., Zambrano, J. P., Heldman, A. W. & Hare, J. M. (2011). Intramyocardial Stem Cell Injection in Patients with Ischemic Cardiomyopathy: Functional Recovery and Reverse Remodeling. *Circulation research* Vol.108, No.7, (Apr 1, 2011) pp. 792-796, ISBN 1524-4571 (Electronic) 0009-7330 (Linking)
- Wohlschlaeger, J., Levkau, B., Brockhoff, G., Schmitz, K. J., von Winterfeld, M., Takeda, A., Takeda, N., Stypmann, J., Vahlhaus, C., Schmid, C., Pomjanski, N., Bocking, A. & Baba, H. A. (2010). Hemodynamic Support by Left Ventricular Assist Devices Reduces Cardiomyocyte DNA Content in the Failing Human Heart. *Circulation* Vol.121, No.8, (Mar 2, 2010) pp. 989-996, ISBN 1524-4539 (Electronic) 0009-7322 (Linking)
- Xie, T. & Spradling, A. C. (2000). A Niche Maintaining Germ Line Stem Cells in the Drosophila Ovary. *Science* Vol.290, No.5490, (Oct 13, 2000) pp. 328-330, ISBN 0036-8075 (Print) 0036-8075 (Linking)
- Xu, H., Morishima, M., Wylie, J. N., Schwartz, R. J., Bruneau, B. G., Lindsay, E. A. & Baldini, A. (2004). Tbx1 Has a Dual Role in the Morphogenesis of the Cardiac Outflow Tract. *Development* Vol.131, No.13, (Jul, 2004) pp. 3217-3227, ISBN 0950-1991 (Print) 0950-1991 (Linking)
- Yamanaka, S. & Blau, H. M. (2010). Nuclear Reprogramming to a Pluripotent State by Three Approaches. *Nature* Vol.465, No.7299, (Jun 10, 2010) pp. 704-712, ISBN 1476-4687 (Electronic) 0028-0836 (Linking)



Congenital Heart Disease - Selected Aspects

Edited by Prof. P. Syamasundar Rao

ISBN 978-953-307-472-6

Hard cover, 348 pages

Publisher InTech

Published online 18, January, 2012

Published in print edition January, 2012

There are significant advances in the understanding of the molecular mechanisms of cardiac development and the etiology of congenital heart disease (CHD). However, these have not yet evolved to such a degree so as to be useful in preventing CHD at this time. Developments such as early detection of the neonates with serious heart disease and their rapid transport to tertiary care centers, availability of highly sensitive noninvasive diagnostic tools, advances in neonatal care and anesthesia, progress in transcatheter interventional procedures and extension of complicated surgical procedures to the neonate and infant have advanced to such a degree that almost all congenital cardiac defects can be diagnosed and "corrected". Treatment of the majority of acyanotic and simpler cyanotic heart defects with currently available transcatheter and surgical techniques is feasible, effective and safe. The application of staged total cavo-pulmonary connection (Fontan) has markedly improved the long-term outlook of children who have one functioning ventricle. This book, I hope, will serve as a rich source of information to the physician caring for infants, children and adults with CHD which may help them provide optimal care for their patients.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Fariba Chalajour, Xiaoyuan Ma and R. Kirk Riemer (2012). Myocardial Self-Repair and Congenital Heart Disease, Congenital Heart Disease - Selected Aspects, Prof. P. Syamasundar Rao (Ed.), ISBN: 978-953-307-472-6, InTech, Available from: <http://www.intechopen.com/books/congenital-heart-disease-selected-aspects/myocardial-self-repair-and-congenital-heart-disease>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen