

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

**WEB OF SCIENCE™**Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us? Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com

Metabolic Optimization by Enzyme-Enzyme and Enzyme-Cytoskeleton Associations

Daniela Araiza-Olivera et al.*

*Instituto de Fisiología Celular, Universidad Nacional Autónoma de México,
Mexico*

1. Introduction

Probably enzymes are not dispersed in the cytoplasm, but are bound to each other and to specific cytoskeleton proteins. Associations result in substrate channeling from one enzyme to another. Multienzymatic complexes, or metabolons have been detected in glycolysis, the Krebs cycle and oxidative phosphorylation. Also, some glycolytic enzymes interact with mitochondria. Metabolons may associate with actin or tubulin, gaining stability. Metabolons resist inhibition mediated by the accumulation of compatible solutes observed during the stress response. Compatible solutes protect membranes and proteins against stress. However, when stress is over, compatible solutes inhibit growth, probably due to the high viscosity they promote. Viscosity inhibits protein movements. Enzymes that undergo large conformational changes during catalysis are more sensitive to viscosity. Enzyme association seems to protect the more sensitive enzymes from viscosity-mediated inhibition. The association-mediated protection of the enzymes in a given metabolic pathway would constitute a new property of metabolons: that is, to enhance survival during stress. It is proposed that resistance to inhibition is due to elimination of non-productive conformations in highly motile enzymes.

2. Metabolons: Enzyme complexes that channel substrates

The cytoplasm should not be regarded as a liquid phase containing a large number of soluble enzymes and particles. Instead, it has become evident that there is a high degree of organization where different lipid and protein structures associate among themselves and with other molecules. The high molecule concentration found in the cytoplasm promotes macromolecule associations such as protein-protein, protein-membrane, protein-nucleic acid, protein-polysaccharide and thus is a control factor for all biological processes (Srere & Ovadi, 1990). Indeed, the classical studies by Green (Green *et al.*, 1965), Clegg (Clegg, 1964)

* Salvador Uribe-Carvajal^{1,**}, Natalia Chiquete-Félix¹, Mónica Rosas-Lemus¹, Gisela Ruíz-Granados¹, José G. Sampedro², Adela Mújica³ and Antonio Peña¹

¹*Instituto de Fisiología Celular, Universidad Nacional Autónoma de México,*

²*Instituto de Física, Universidad Autónoma de San Luís Potosí and*

³*CINVESTAV, Instituto Politécnico Nacional*

Mexico

** Corresponding Author

and Fulton (Fulton, 1982) have suggested that enzymes are not dispersed in the cytoplasm. Instead, enzymes are localized at specific sites where they are associated between them and with the cytoskeleton. The cytoskeleton is a trabecular network of fibrous proteins that micro-compartmentalizes the cytoplasm (Porter *et al.*, 1983). Associated enzymes channel substrates from one to another preventing their diffusion to the aqueous phase (Gaertner *et al.*, 1978; Minton & Wilf, 1981; Ovadi *et al.*, 1996).

In a multienzyme complex, intermediaries can be channeled more than once from the active site of an enzyme to the next to obtain the final product (Al-Habori, 2000; Robinson *et al.*, 1987). Channeling requires stable interactions of the multienzymatic metabolons (Al-Habori, 2000; Cascante *et al.*, 1994; Ovadi & Srere, 1996; Ovadi & Saks, 2004; Srere & Ovadi, 1990; Srere, 1987). The metabolon stability is facilitated by the compartmentalization of the cell in different organelles and structures (Jorgensen *et al.*, 2005).

There are many advantages inherent to metabolons (Jorgensen *et al.*, 2005) (Fig. 1): I) Improved catalytic efficiency of the enzymes. This is obtained by channeling an intermediary from the active site of an enzyme directly to the active site of the next. II) Channeling optimizes kinetic constants. III) Labile or toxic intermediates are retained within the metabolon. IV) Inhibitors are excluded from the active site of enzymes. V) Control and coordination of the enzymes in a given pathway is enhanced. VI) Finally, alternative metabolons may favor different pathways (Fig. 1). Most metabolons seem to be transient, opening the possibility for a quick change in some elements that allows them to redirect metabolism (Jorgensen *et al.*, 2005).




Fig. 1. Advantages of Metabolons. (A) In isolated enzymes the substrate (green), intermediaries (red and yellow) and product (orange) diffuse into the aqueous phase (little arrows). Toxic intermediaries and inhibitors (grey) are free to exit/enter the active site in each enzyme. (B) In metabolons (we show filamentous actin in red and white) channeling allows transfer of the substrate (green) from the active site of an enzyme direct to the next to obtain a final product (orange) without diffusion to the cytoplasm of intermediaries (not-depicted) are prevented, while inhibitors (grey) are excluded from the active sites.

The enzymes from the Krebs cycle are attached to the mitochondrial membrane in an enzymatic complex; this was the first “metabolon” described (Srere, 1987). In oxidative

phosphorylation, multiprotein complexes seem to associate in supercomplexes and eventually in respiratory chains resulting in controlled electron channeling and proton-pumping stoichiometry (Guerrero-Castillo *et al.*, 2011). It has been proposed that these supercomplexes constitute an exquisite mechanism to regulate the yield of ATP (Guerrero-Castillo *et al.*, 2009; 2011; Schägger *et al.*, 2001). In addition, in some organisms such as trypanosomes, glycolytic enzymes are contained in small organelles called glycosomes, where channeling is highly efficient (Aman *et al.*, 1985). Tumor cells also produce aggregates containing glycolytic enzymes (Coe & Greenhouse, 1973). Interactions between organelles such as the endoplasmic reticulum and mitochondria have been described (Dorn & Scorrano, 2010; Kornmann *et al.*, 2009; Lebedzinska *et al.*, 2009). Mitochondria are both, the main source of ATP and inducers of cellular death (Anesti & Scorrano, 2006). Mitochondrial functions are regulated by interactions with other organelles and cytoplasmic proteins (Kostal & Arriaga, 2011). Cytoskeletal proteins such as actin and tubulin, direct mitochondria to specific sites in the cell (Senning & Marcus, 2010) and control coupling of phosphorylation by interacting with mitochondrial porin (Xu *et al.*, 2001; Lemasters & Holmuhamedov, 2006; Rostovtseva *et al.*, 2008; Rostovtseva *et al.*, 2004; Xu *et al.*, 2001). In addition to cytoskeletal proteins, hexokinase, a glycolytic enzyme binds mitochondria in mammals (Pastorino & Hoek, 2008), yeast and plants (Balasubramanian *et al.*, 2008) regulating the energy yield of mitochondria as well as the induction of programmed cell death (Kroemer *et al.*, 2005; Pastorino & Hoek, 2008; Xie & Wilson, 1988). All the above data suggest that enzymes are highly organized (Clegg & Jackson, 1989) and the cytoskeleton plays an important role (Minaschek *et al.*, 1992; Keleti *et al.*, 1989; Porter *et al.*, 1983).

3. The cytoskeleton: A scaffold where metabolons are bound

The eukaryotic cytoplasm is supported by the cytoskeleton, a network of structural proteins that shapes the cell and has binding sites for different enzymes. Such sites have been identified in filamentous actin (F-actin), in microtubules and in the cytoplasmic domain of the erythrocyte band 3, which is also an anion exchanger. Glycolytic enzyme binding to actin usually results in stimulation, whereas binding to microtubules or to band 3 inhibits activity (Real-Hohn *et al.*, 2010). Actin is involved in a variety of cell functions that include contractility, cytokinesis, maintenance of cell shape, cell locomotion and organelle localization. In addition, glycolytic enzymes and F-actin co-localize in muscle cells, probably reflecting compartmentation of the glycolytic pathway (Waingeh *et al.*, 2006).

Actin is highly conserved in eukaryotic cells. It may be found as a monomer (G-actin) or as a polymeric filament (F-actin) that is interconverted in an extremely dynamic, highly controlled process. The polar actin monomers polymerize head-to-tail to yield a polar filament. Actin filaments are constituted by 8 nm diameter, double-helical structures formed by assemblies of monomeric actin with a barbed end (or plus end) and a pointed end (or minus end). The spontaneous polymerization of actin monomers occurs in three phases: nucleation, elongation and maintenance. Nucleation consists in the formation of a dimer, followed by the addition of a third monomer to yield a trimer; this process is very slow. Further monomer addition becomes thermodynamically favorable and the filament elongates rapidly: much faster at the plus end than at the minus end. In the maintenance phase, there is no net filament growth and the concentration of ATP-G-actin is kept stationary (Fig. 2).

Upon incorporation to a filament, G-actin-bound ATP is hydrolyzed. ADP and Pi remain non-covalently bound. Then Pi is released slowly. Thus, the elongating filaments contain: the barbed end, rich in ATP-actin, the center, rich in ADP-Pi-actin and the pointed end containing ADP-actin. Many actin-binding proteins regulate actin polymerization. Profilin is an actin monomer-binding protein; Arp 2/3 complex are nucleation proteins; CapZ and gelsolin regulate the length of the actin filament and the cofilin/ADP family cuts F-actin and accelerates depolymerization (Kustermans *et al.*, 2008). However, protein functions may vary; in *Dictyostelium*, CapZ prevents filament elongation and increases the concentration of unpolymerized actin; in contrast, in yeast this same protein prevents depolymerization increasing F-actin concentration (Welch *et al.*, 1997). The cytoskeleton can be rapidly remodeled by the small RhoGTPases (Rho, Rac and Cdc42), which act in response to extracellular stimuli (Kustermans *et al.*, 2008). There are exogenous natural compounds that can disturb actin dynamics (Kustermans *et al.*, 2008).

4. The glycolytic metabolon

The association of enzymes with the cytoskeleton probably stabilizes metabolons. In this regard, glycolytic enzymes such as fructose 1,6-bisphosphate aldolase (aldolase), glyceraldehyde 3-phosphate dehydrogenase (GAPDH), pyruvate kinase (PK), glucose phosphate isomerase (GPI), and lactate dehydrogenase (LDH) associate with actin. Other glycolytic enzymes such as triose phosphate isomerase and phosphoglycerate mutase bind indirectly through interactions with other enzymes. Enzyme-enzyme-actin complexes are called piggy-back interactions. Also, aldolase and GAPDH compete for binding sites (Knull & Walsh, 1992; Waingeh *et al.*, 2006).




Fig. 2. **Actin polymerization.** During nucleation, actin monomers aggregate to form a trimer. Then during elongation actin filaments grow actively at both ends. Growth stops in the maintenance phase, also known as stationary phase. (Modified from Kustermans *et al.*, 2008)

Enzyme/actin interaction is regulated by ionic strength (Waingeh *et al.*, 2006). In homogenates of muscle tissue suspended in isosmotic sucrose, proteins such as F-actin, myosin, troponin and tropomyosin associate with glycolytic enzymes (Brooks & Storey, 1991). Glycolytic enzyme association to actin is not accepted universally, for instance, the F-actin/glycolytic enzyme interaction has been modeled mathematically at physiological ionic strength and protein concentrations. The results suggest that under cellular conditions only a small percentage of TPI, GAPDH, PGK and LDH would be associated with F-actin (Brooks & Storey, 1991).

Protein dynamics seem important for their interactions. Brownian dynamics (BD) simulations detect that rabbit F-actin has different binding modes/affinities for aldolase and GAPDH (Forlemu *et al.*, 2006). Some metabolites such as ATP and ADP modulate enzyme interactions and the resulting substrate channeling (Forlemu *et al.*, 2006).

A barely explored effect of the association of enzymes with the cytoskeleton is the modulation of the dynamics of actin polymerization. Such an effect has been reported for aldolase (Chiquete-Felix *et al.*, 2009; Schindler *et al.*, 2001). An interesting finding is that some growth factors, such as PGF and EGF enhance the GAPDH/cytoskeleton interaction, possibly increasing keratinocyte migration (Tochio *et al.*, 2010). Indeed, GAPDH seems to participate in cytoskeleton dynamics processes such as endocytosis, membrane fusion, vesicular transport and nuclear tRNA transport (Cueille *et al.*, 2007).

In red blood cell membranes, GAPDH, aldolase and PFK interact with an acidic sequence at the amino-terminal extreme of band 3 with high affinity (Campanella *et al.*, 2005). Under physiological conditions, the binding of glycolytic enzymes to band 3 results in inhibition of the glycolytic flux (Real-Hohn *et al.*, 2010).

Association to microtubules regulates the energetic metabolism (Keleti *et al.*, 1989; Keller *et al.*, 2007; Walsh *et al.*, 1989) at the level of some glycolytic enzymes such as pyruvate kinase, phosphofructokinase (Kovács *et al.*, 2003) and enolase (Keller *et al.*, 2007). When the glycolytic enzymes are associated and anchored to the sarcomere, ATP is produced more efficiently (Keller *et al.*, 2007). The interaction of enzymes with themselves and with the cytoskeleton confers more stability to the enzyme activity and to the whole network (Keleti *et al.*, 1989; Volker *et al.*, 1995; Walsh *et al.*, 1989). F-actin stabilizes some glycolytic enzymes of muscle and sperm (Walsh & Knull, 1988; Ovadi & Saks, 2004). That is the case of the phosphofructokinase (PFK) and aldolase where the dilution-mediated inactivation of PFK is stopped upon aldolase addition. If PFK is associated with microtubules, it still loses activity when diluted, however, in these conditions it recovers the lost activity upon aldolase addition (Raïs *et al.*, 2000; Vértessy *et al.*, 1997). All this evidence supports the existence of a cytoskeleton-bound glycolytic metabolon.

5. Compatible solutes protect cellular structures during stress

Compatible solutes are defined as molecules that reach high concentrations in the cell without interfering with metabolic functions (Brown & Simpson, 1972). These are mostly amino acids and amino acid derivatives, polyols, sugars and methylamines. Compatible solutes are typically small and harbor chemical groups that interact with protein surfaces. Indeed, some authors have proposed to call them “chemical or pharmacological chaperones” as they stabilize native structures (Loo & Clarke, 2007; Romisch, 2004). Some compatible solutes are:

glycine betaine, a thermoprotectant in *B. subtilis* (Chen & Murata, 2011; Holtmann & Bremer, 2004). Ectoine, that in halophile microorganisms confers resistance to salt and temperature stress (Pastor *et al.*, 2010). Glycerol is accumulated in yeast under high osmotic pressure (Blomberg, 2000). Glycerol stabilizes thermolabile enzymes preventing their inactivation (Zancan & Sola-Penna, 2005). The disaccharide trehalose protects against environmental injuries (heat, cold, desiccation, and anoxia) and nutritional limitations (Argüelles, 2000; Crowe *et al.*, 1984) in bacteria, yeast, fungi, plants and invertebrates. In biotechnology, trehalose is one of the best protein stabilizing known (Jain & Roy, 2008; Sampedro *et al.*, 2001).

6. Effect of compatible solutes on the activity of enzymes

Compatible solute synthesis and accumulation is triggered by harsh conditions and results in protein stabilization and enhanced survival. Proteins may be unfolded, partially unfolded or native (Chilson & Chilson, 2003). In the absence of stress, high compatible solute concentrations inhibit cellular growth, metabolism and division (Wera *et al.*, 1999), e.g. a trehalase-deficient mutant of *S. cerevisiae* subjected to heat or saline stress accumulated high amounts of trehalose and survived. However, when these mutants were returned to normal conditions they are unable to grow or sustain metabolic activity (Garre & Matallana, 2009; Wera *et al.*, 1999).

6.1 Inhibition of isolated enzymes; possible role of viscosity

Under stress, high compatible solutes change the physicochemical properties of the cytoplasm. However, the effect of the high viscosity generated by molar concentrations of compatible solutes on enzyme activity has drawn little attention. Trehalose and other polyols protect proteins from thermal unfolding via indirect interactions (Liu *et al.*, 2010). Therefore the stabilizing mechanism must rely in the modified physicochemical properties of aqueous media.

Large-scale conformational changes in proteins involve the physical displacement of associated solvent molecules and solutes. The resistance to the movement or displacement of solvent molecules is a frictional process. Kramers theory provides the mathematical basis to understand and analyze reactions at high viscosity (Kramers, 1940). The application of Kramer's theory to proteins indicates that the movements involved in folding or in enzyme-substrate association and processing must be highly sensitive to viscosity (Jacob and Schmid, 1999; Jacob *et al.*, 1999; Sampedro and Uribe, 2004).

Studies on cellular viscosity in yeast cytoplasm showed a value of 2 cP at 30°C (Williams *et al.*, 1997). Also, *in vitro* determinations for 0.6 M trehalose solutions showed a viscosity of 1.5 cP at 30°C (Table 1). Therefore, one may infer that yeast cytoplasm viscosity with 0.6 M trehalose should be in the vicinity of 2.5-3 cP.

The plasma membrane H⁺-ATPase from yeast depends on large domain motion for catalysis (Kulbrandt, 2004), was inhibited at all trehalose concentrations tested (Sampedro *et al.*, 2002). The rate constant for the ATPase reaction ($V_{max} = k_{cat} [E_t]$) was inversely dependent on solution viscosity; as higher the viscosity lower the reaction rate of catalysis (Sampedro *et al.*, 2002). Notably, when temperature was raised inhibition disappeared, in agreement with the fact that viscosity decreases when temperature increases (Table 1). Similar results have been obtained with Na⁺/K⁺-ATPase and Na⁺-ATPase in the presence of polyethylene glycol and

glycerol (Esmann *et al.*, 2008). In glucose oxidase, activity inhibition by varying concentrations of trehalose was due to the promotion of a highly compact state, which correlated with the increased viscosity of the medium (Paz-Alfaro *et al.*, 2009).

TREHALOSE (M)	0.2	0.4	0.5	0.6	0.8
TEMP (°C)	VISCOSITY (cP)				
20	1.35	1.59	1.81	2.04	2.58
25	1.20	1.37	1.51	1.74	2.20
30	1.08	1.18	1.33	1.50	1.91
35	0.94	1.03	1.18	1.31	1.67
40	0.86	0.94	1.04	1.13	1.49
45	0.75	0.81	0.90	1.04	1.31

Data modified from Sampedro *et al.*, 2002.

Table 1. Viscosity of trehalose solutions at different concentrations and temperatures.




Fig. 3. Reaction coordinate diagram, comparing an enzyme reaction at normal viscosity (blue) and at high viscosity (η ; red). When a diffusive protein domain process is present in the catalytic cycle, it becomes rate limiting when viscosity is high. Therefore the overall activation energy (E_a) increases.

Many enzymes are inhibited by viscosity. Glutathione reductase is inhibited at 25°C, by trehalose (70% inhibition at 1.5 M trehalose) although inhibition disappears at 40°C (Sebollela *et al.*, 2004). Also pyrophosphatase and glucose 6-phosphate dehydrogenase show temperature dependence of trehalose-mediated inhibition (Sebollela *et al.*, 2004).

Aminoglycoside nucleotidyltransferase 2"-I is inhibited by glycerol in a temperature-dependent way (Gates & Northrop, 1988). The hyaluronan-synthase from *Streptococcus equisimilisis* is inhibited by PEG, ethylene glycol, glycerol or sucrose (Tlapak-Simmons *et al.*, 2004). At high viscosities (greater than 4 mPa s⁻¹) different carbohydrates inhibit egg-white lysozyme (Lamy *et al.*, 1990; Monkos, 1997).

Detailed studies on diffusive protein-structural components demonstrated that for β -lactam synthase a conformational change is rate-limiting on k_{cat} . Therefore, the rate for catalysis shows a high inhibition by medium viscosity (Raber *et al.*, 2009). Crystallographic analysis of adhesion kinase-1 shows a large conformational motion of the activation loop upon ATP binding. This is an essential step during catalysis and explains the viscosity inhibitory effect (Schneck *et al.*, 2010). In the plasma membrane H⁺-ATPase, the enzyme fluctuates between two structural conformations (E1 \leftrightarrow E2) during catalysis. The N-domain (nucleotide binding) rotates 73° towards the phosphorylation site to deliver ATP to the phosphorylation site (Kuhlbrandt, 2004). In all cases, the rate-limiting step is a conformational change that seems to be the one inhibited by viscosity (Fig. 3).

6.2 Enzyme association results in protection against inhibition

Compatible solute-mediated inhibition does not seem to uniformly affect all enzymes. Furthermore, in the face of both the stress condition and the compatible solute, catabolic pathways seem to resist inhibition, thus providing the energy needed for survival (Hoffmann & Holzhütter, 2009; Hounsa *et al.*, 1998). In our hands, in a yeast cytoplasmic extract, compatible solutes inhibit the whole glycolytic pathway much less than many of its individual, isolated enzymes (Araiza-Olivera *et al.*, 2010). In contrast, anabolism seems to be shot both during the stress situation and later (Attfield, 1987). Inhibition of anabolism would explain the inability of cells to reproduce (Wera *et al.*, 1999). The mechanism for resistance to inhibition, exhibited by the catabolic enzymes is a matter of study (Marcondes *et al.*, 2011; Raïs *et al.*, 2000).

The effect of a compatible solute (trehalose) on the activity of some yeast glycolytic enzymes such as GAPDH, HXK, ALD and PGK has been analyzed. These enzymes were tested individually or in mixtures (Araiza-Olivera *et al.*, 2010). When isolated, GAPDH and HXK were inhibited by trehalose while others, such as ALD and PGK were resistant. Probably GAPDH and HXK are more motile than ALD and PGK. Remarkably, when the sensitive enzymes were mixed with the resistant enzymes a protection effect was observed. This led to analyze the whole glycolytic pathway and again, inhibition was minimal in comparison with the individual, isolated enzymes (Araiza-Olivera *et al.*, 2010). Thus, it was decided to explore the possible mechanisms underlying this effect, i.e, why some metabolic pathways, such as glycolysis resist the viscosity-mediated inhibition promoted by compatible solutes, even if they contain several viscosity-sensitive enzymes.

The protection effect was specific for each protein couple, as GAPDH was not protected by neither HXK, albumin or lactate-dehydrogenase. Also, the pentose pathway enzyme glucose 6-phosphate dehydrogenase (G6PDH) was not protected by ALD against inhibition by trehalose. Once in the complexes, probably the more flexible enzymes that are more sensitive to viscosity (Sampedro & Uribe 2004) are stabilized by the more resistant, more rigid enzymes forming a less motile, more resistant complex.

The proposal that enzyme association favors a more stable folded state would require the motile enzymes to eliminate some non-productive conformations (Villali & Kern, 2010). These associations are probably further stabilized by some elements of the cytoskeleton, such as tubulin (Raïs *et al.*, 2000; Walsh *et al.*, 1989) or F-actin (Minaschek *et al.*, 1992; Waingeh *et al.*, 2006). Thus, it is proposed that another function of enzyme association into metabolons, in addition to substrate channeling and metabolic control might be to resist compatible solute-mediated inhibition.

7. Concluding remarks

Under stress, compatible solutes accumulate to very high levels in the cytoplasm. This results in enhanced viscosity. As revised in section 6.1, viscosity is known to inhibit diverse enzymes. Indeed, high viscosity may be the mechanism by which diverse cell functions are inhibited in the presence of high compatible solute concentrations, e.g. cells are unable to. In contrast, catabolism remains active even in the presence of compatible solutes. One possible mechanism for this resistance to inhibition is probably the specific association of glycolytic enzymes among themselves and probably with the cytoskeleton. Resistance to viscosity-mediated inhibition is proposed as a novel, important property of enzyme association into metabolons. The mechanism of protection that association confers against viscosity still has to be defined. Protection of activity is needed for survival during stress.

8. References

- Al-Habori M. Microcompartmentation, metabolic channelling and carbohydrate metabolism. *Int. J. Biochem. Cell. Biol.* 1995; 27(2):123-32.
- Aman RA & Wang CC. An improved purification of glycosomes from the procyclic trypomastigotes of *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* 1986; 21(3):211-20.
- Anesti V & Scorrano L. The relationship between mitochondrial shape and function and the cytoskeleton. *Biochim Biophys Acta.* 2006; 1757(5-6): 692-9.
- Araiza-Olivera D, Sampedro JG, Mújica A, Peña A & Uribe-Carvajal S. The association of glycolytic enzymes from yeast confers resistance against inhibition by trehalose. *FEMS Yeast Res.* 2010; 10 (3):282-9.
- Argüelles JC. Physiological roles of trehalose in bacteria and yeasts: a comparative analysis. *Arch Microbiol.* 2000; 174(4):217-24.
- Attfield PV. Trehalose accumulates in *Saccharomyces cerevisiae* during exposure to agents that induce heat shock response. *FEBS Lett.* 1987; 225(1-2):259-63.
- Balasubramanian R, Karve A & Moore BD. Actin-based cellular framework for glucose signaling by *Arabidopsis hexokinase1*. *Plant Signal Behav.* 2008; 3(5):322-4.
- Blomberg A. Metabolic surprises in *Saccharomyces cerevisiae* during adaptation to saline conditions: questions, some answers and a model. *FEMS Microbiol Lett.* 2000; 82(1):1-8.
- Brooks SP & Storey KB. Where is the glycolytic complex? A critical evaluation of present data from muscle tissue. *FEBS Lett.* 1991; 278(2):135-8.
- Brown AD & Simpson JR. Water relations on sugar-tolerant yeast: the role of intracellular polyols. *J. Gen Microbiol.* 1972; 72:589-591.

- Campanella ME, Chu H & Low PS. Assembly and regulation of a glycolytic enzyme complex on the human erythrocyte membrane. *Proc Natl Acad Sci.* 2005; 102(7):2402-7.
- Cascante M, Sorribas A & Canela EI. Enzyme-enzyme interactions and metabolite channelling: alternative mechanisms and their evolutionary significance. *Biochem. J.* 1994; 298 (Pt 2):313-20
- Chen TH & Murata N. Glycinebetaine protects plants against abiotic stress: mechanisms and biotechnological applications. *Plant Cell Environ.* 2011; 34(1):1-20.
- Chilson OP & Chilson AE. Perturbation of folding and reassociation of lactate dehydrogenase by proline and trimethyl amine oxide. *Eur.J.Biochem.* 2003; 270,4823-4834.
- Chiquete-Felix N, Hernández JM, Méndez JA, Zepeda-Bastida A, Chagolla-López A & Mújica A. In guinea pig sperm, aldolase A forms a complex with actin, WAS, and Arp2/3 that plays a role in actin polymerization. *Reproduction.* 2009; 137(4):669-78.
- Clegg JS. The control of emergence and metabolism by external osmotic pressure and the role of free glycerol in developing cysts of *Artemia salina*. *J. Exp. Biol.* 1964; 41:879-92.
- Clegg JS & Jackson SA. Evidence for intermediate channelling in the glycolytic pathway of permeabilized L-929 cells. *Biochem. Biophys. Res. Commun.* 1989; 160(3):1409-14.
- Coe EL & Greenhouse WV. Possible regulatory interactions between compartmentalized glycolytic systems during initiation of glycolysis in ascites tumor cells. *Biochim. Biophys. Acta.* 1973; 329(2):171-82.
- Crowe JH & Crowe LM, Chapman D. Preservation of Membranes in Anhydrobiotic Organisms: The Role of Trehalose. *Science.* 1984; 223(4637):701-703.
- Cueille N, Blanc CT, Riederer IM & Riederer BM. Microtubule-associated protein 1B binds glyceraldehyde-3-phosphate dehydrogenase. *J Proteome Res.* 2007; 6(7):2640-7.
- Dorn GW 2nd & Scorrano L. Two close, too close: sarcoplasmic reticulum-mitochondrial crosstalk and cardiomyocyte fate. *Circ Res.* 2010; 107(6):689-99. Review.
- Esmann M, Fedosova NU & Marsh D. Osmotic Stress and Viscous Retardation of the Na,K-ATPase Ion Pump. *Biophysical J.* 2008; 94:2767-2776.
- Forlemu NY, Waingeh VF, Ouporov IV, Lowe SL & Thomasson KA. Theoretical study of interactions between muscle aldolase and F-actin: insight into different species. *Biopolymers.* 2007; 85(1):60-71.
- Fulton AB. How crowded is the cytoplasm? *Cell.* 1982; 30(2):345-7.
- Gaertner FH. Unique catalytic properties of enzyme clusters. *Trends Biochem. Sci.* 1978; 3, 63.
- Garre E & Matallana E. The three trehalases Nth1p, Nth2p and Ath1p participate in the mobilization of intracellular trehalose required for recovery from saline stress in *Saccharomyces cerevisiae*. *Microbiology.* 2009; 155:3092-3099.
- Gates CA & Northrop DB. Determination of the rate-limiting segment of aminoglycoside nucleotidyltransferase 2"-I by pH and viscosity-dependent kinetics. *Biochemistry.* 1988; 27(10):3834-3842.
- Green DE, Murer E, Hultin HO, Richardson SH, Salmon B, Brierley GP & Baum H. Association of integrated metabolic pathways with membranes. I. Glycolytic enzymes of the red blood corpuscle and yeast. *Arch. Biochem. Biophys.* 1965; 112(3):635-47.

- Guerrero-Castillo S, Vázquez-Acevedo M, González-Halphen D & Uribe-Carvajal S. In *Yarrowia lipolytica* mitochondria, the alternative NADH dehydrogenase interacts specifically with the cytochrome complexes of the classic respiratory pathway. *Biochim Biophys Acta*. 2009; 1787 (2):75-85.
- Guerrero-Castillo S, Araiza-Olivera D, Cabrera-Orefice A, Espinasa-Jaramillo J, Gutiérrez-Aguilar M, Luévano-Martínez LA, Zepeda-Bastida A & Uribe-Carvajal S. Physiological uncoupling of mitochondrial oxidative phosphorylation. Studies in different yeast species. *J Bioenerg Biomembr*. 2011; 43(3):323-31.
- Hoffmann S & Holzhütter HG. Uncovering metabolic objectives pursued by changes of enzyme levels. *Ann. N. Y. Acad. Sci*. 2009; 1158:57-70
- Holtmann G & Bremer E. Thermoprotection of *Bacillus subtilis* by exogenously provided glycine betaine and structurally related compatible solutes: involvement of Opu transporters. *J Bacteriol*. 2004; 186(6):1683-93.
- Hounsa CG, Brandt EV, Thevelein J, Hohmann S & Prior BA. Role of trehalose in survival of *Saccharomyces cerevisiae* under osmotic stress. *Microbiology* 1998; 144:671-80.
- Jacob M, Geeves M, Holterman G, & Schmid FX. Diffusional crossing in a two-state protein folding reaction. 1999 *Nat. Struct. Biol*. 1999; 6:923-926.
- Jacob M & Schmid FX. Protein folding as a diffusional process. *Biochemistry* 1999; 38:13773-13779.
- Jain NK & Roy I. Role of trehalose in moisture-induced aggregation of bovine serum albumin. *Eur J Pharm Biopharm*. 2008; 69(3):824-34.
- Jørgensen K, Rasmussen AV, Morant M, Nielsen AH, Bjarnholt N, Zagrobelny M, Bak S & Møller BL. Metabolon formation and metabolic channeling in the biosynthesis of plant natural products. *Curr. Opin. Plant Biol*. 2005; 8(3):280-91.
- Keleti T, Ovádi J & Batke J. Kinetic and physico-chemical analysis of enzyme complexes and their possible role in the control of metabolism. *Prog. Biophys. Mol. Biol*. 1989; 53(2):105-52.
- Keller A, Peltzer J, Carpentier G, Horváth I, Oláh J, Duchesnay A, Orosz F & Ovádi J. Interactions of enolase isoforms with tubulin and microtubules during myogenesis. *Biochim. Biophys. Acta*. 2007;1770 (6):919-26.
- Knull HR & Walsh JL. Association of glycolytic enzymes with the cytoskeleton. *Curr Top Cell Regul*. 1992; 33:15-30. Review.
- Kornmann B, Currie E, Collins SR, Schuldiner M, Nunnari J, Weissman JS & Walter P. An ER-mitochondria tethering complex revealed by a synthetic biology screen. *Science*. 2009; 325(5939):477-81.
- Kostal V & Arriaga EA. Capillary Electrophoretic Analysis Reveals Subcellular Binding between Individual Mitochondria and Cytoskeleton. *Anal Chem*. 2011.
- Kovacs J, Low P, Pacz A, Horvath I, Olah J & Ovadi J. Phosphoenolpyruvate-dependent tubulin-pyruvate kinase interaction at different organizational levels. *J. Biol. Chem*. 2003; 278(9):7126-30.
- Kramers HA. Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica*. 1940; 7:284-304.
- Kroemer G, El-Deiry WS, Golstein P, Peter ME, Vaux D, Vandenabeele P, Zhivotovsky B, Blagosklonny MV, Malorni W, Knight RA, Piacentini M, Nagata S & Melino G. Classification of cell death: recommendations of the Nomenclature Committee on Cell Death. *Cell Death Differ*. 2005, Suppl 2: 1463-7.

- Kühlbrandt W. Biology, structure and mechanism of p-type ATPases. *Nature*. 2004; 5:282-295.
- Kustermans G, Piette J & Legrand-Poels S. Actin-targeting natural compounds as tools to study the role of actin cytoskeleton in signal transduction. *Biochem Pharmacol*. 2008; 76(11):1310-22.
- Lamy L, Portmann MO, Mathlouthi M & Larreta-Garde V. Modulation of egg-white lysozyme activity by viscosity intensifier additives. *Biophys Chem*. 1990; 36(1):71-76.
- Lebiedzinska M, Szabadkai G, Jones AW, Duszynski J & Wieckowski MR. Interactions between the endoplasmic reticulum, mitochondria, plasma membrane and other subcellular organelles. *Int J Biochem Cell Biol*. 2009; 41(10):1805-16. Review.
- Lemasters JJ & Holmuhamedov E. Voltage-dependent anion channel (VDAC) as mitochondrial governor--thinking outside the box. *Biochim Biophys Acta*. 2006; 1762(2): 181-90.
- Liu FF, Ji L, Zhang L, Dong XY & Sun Y . Molecular basis for polyol-induced protein stability revealed by molecular dynamics simulations. *J Chem Phys*. 2010; 132(22):225103.
- Loo TW & Clarke DM. Chemical and pharmacological chaperones as new therapeutic agents. *Expert Rev Mol Med*. 2007; 9(16):1-18.
- Marcondes MC, Sola-Penna M, Torres Rda S & Zancan P. Muscle-type 6-phosphofructo-1-kinase and aldolase associate conferring catalytic advantages for both enzymes. *IUBMB Life*. 2011; 63(6):435-45.
- Minaschek G, Gröschel-Stewart U, Blum S & Bereiter-Hahn J. Microcompartmentation of glycolytic enzymes in cultured cells. *Eur J Cell Biol*. 1992; 58(2):418-28.
- Minton AP & Wilf J. Effect of macromolecular crowding upon the structure and function of an enzyme: glyceraldehyde-3-phosphate dehydrogenase. *Biochemistry* 1981; 20(17):4821-6.
- Monkos K. Concentration and temperature dependence of viscosity in lysozyme aqueous solutions. *Biochim Biophys Acta*. 1997; 1339(2):304-10.
- Ovádi J & Srere PA. Metabolic consequences of enzyme interactions. *Cell. Biochem. Funct*. 1996; 14(4):249-58.
- Ovádi J & Saks V. On the origin of intracellular compartmentation and organized metabolic systems. *Mol. Cell Biochem*. 2004; 256-257(1-2):5-12.
- Pastor JM, Salvador M, Argandoña M, Bernal V, Reina-Bueno M, Csonka LN, Iborra JL, Vargas C, Nieto JJ & Cánovas M. Ectoines in cell stress protection: uses and biotechnological production. *Biotechnol Adv*. 2010; 28(6):782-801. Review.
- Pastorino JG & Hoek JB. Regulation of hexokinase binding to VDAC. *J Bioenerg Biomembr*. 2008; 40(3): 171-82.
- Paz-Alfaro K, Ruiz-Granados YG, Uribe-Carvajal S & Sampedro JG. Trehalose-mediated stabilization of glucose oxidase from *Aspergillus niger*. *J Biotechnol*. 2009; 141:130-136.
- Porter ME & Johnson KA. Transient state kinetic analysis of the ATP-induced dissociation of the dynein-microtubule complex. *J. Biol. Chem*. 1983; 258(10):6582-7.
- Raber ML, Freeman MF & Townsend CA. Dissection of the stepwise mechanism to beta-lactam formation and elucidation of a rate-determining conformational change in beta-lactam synthetase. *J Biol Chem*. 2009; 284(1):207-217.

- Rais B, Ortega F, Puigjaner J, Comin B, Orosz F, Ovádi J & Cascante M. Quantitative characterization of homo- and heteroassociations of muscle phosphofructokinase with aldolase. *Biochim Biophys Acta*. 2000; 1479(1-2):303-14.
- Real-Hohn A, Zancan P, Da Silva D, Martins ER, Salgado LT, Mermelstein CS, Gomes AM & Sola-Penna M. Filamentous actin and its associated binding proteins are the stimulatory site for 6-phosphofructo-1-kinase association within the membrane of human erythrocytes. *Biochimie*. 2010; 92(5):538-44.
- Robinson JB Jr, Inman L, Sumegi B & Srere PA. Further characterization of the Krebs tricarboxylic acid cycle metabolon. *J. Biol. Chem*. 1987; 262 (4):1786-90.
- Römisch K. A cure for traffic jams: small molecule chaperones in the endoplasmic reticulum. *Traffic*. 2004; 5(11):815-820.
- Rostovtseva TK, Antonsson B, Suzuki M, Youle RJ, Colombini M & Bezrukov S. M. Bid, but not Bax, regulates VDAC channels. *J Biol Chem*. 2004; 279(14): 13575-83.
- Rostovtseva TK, Sheldon KL, Hassanzadeh E, Monge C, Saks V, Bezrukov SM & Sackett DL. Tubulin binding blocks mitochondrial voltage-dependent anion channel and regulates respiration. *Proc Natl Acad Sci U S A*. 2008; 105(48): 18746-51.
- Sampedro JG, Muñoz-Clares RA & Uribe S. Trehalose-mediated inhibition of the plasma membrane H⁺-ATPase from *Kluyveromyces lactis*: dependence on viscosity and temperature. *J Bacteriol*. 2002; 184(16):4384-91.
- Sampedro JG, Cortés P, Muñoz-Clares RA, Fernández A & Uribe S. Thermal inactivation of the plasma membrane H⁺-ATPase from *Kluyveromyces lactis*. Protection by trehalose. *Biochim. Biophys. Acta* 2001; 1544(1-2):64-73.
- Sampedro JG & Uribe S. Trehalose-enzyme interactions result in structure stabilization and activity inhibition. The role of viscosity. *Mol. Cell. Biochem*. 2004; 256-257(1-2):319-27.
- Schägger H & Pfeiffer K. The ratio of oxidative phosphorylation complexes I-V in bovine heart mitochondria and the composition of respiratory chain supercomplexes. *J Biol Chem*. 2001; 276 (41):37861-7.
- Schindler R, Weichselsdorfer E, Wagner O & Bereiter-Hahn J. Aldolase-localization in cultured cells: cell-type and substrate-specific regulation of cytoskeletal associations. *Biochem Cell Biol*. 2001; 79(6):719-28.
- Schneck JL, Briand J, Chen S, Lehr R, McDevitt P, Zhao B, Smallwood A, Concha N, Oza K, Kirkpatrick R, Yan K, Villa JP, Meek TD & Thrall SH. Kinetic mechanism and rate-limiting steps of focal adhesion kinase-1. *Biochemistry*. 2010; 49(33):7151-7163.
- Sebollela A, Louzada PR, Sola-Penna M, Sarone-Williams V, Coelho-Sampaio T & Ferreira ST. Inhibition of yeast glutathione reductase by trehalose: possible implications in yeast survival and recovery from stress. *Int J Biochem Cell Biol*. 2004; 36(5):900-908.
- Senning EN & Marcus AH. Actin polymerization driven mitochondrial transport in mating *S. cerevisiae*. *Proc Natl Acad Sci*. 2010; 107(2):721-5.
- Srere PA. Complexes of sequential metabolic enzymes. *Annu. Rev. Biochem*. 1987; 56:89-124.
- Srere PA & Ovadi J. Enzyme-enzyme interactions and their metabolic role. *FEBS Lett*. 1990; 268(2):360-4.
- Tlapak-Simmons VL, Baron CA & Weigel PH. Characterization of the purified hyaluronan synthase from *Streptococcus equisimilis*. *Biochemistry*. 2004; 43(28):9234-9242.

- Tochio T, Tanaka H, Nakata S & Hosoya H. Fructose 1,6-bisphosphate aldolase A is involved in HaCaT cell migration by inducing lamellipodia formation. *J Dermatol Sci.* 2010; 58(2):123-9.
- Villali J & Kern D. Choreographing an enzyme's dance. *Curr Opin Chem Biol.* 2010; 14(5):636-43.
- Vértessy BG, Orosz F, Kovács J & Ovádi J. Alternative binding of two sequential glycolytic enzymes to microtubules. Molecular studies in the phosphofructokinase/ aldolase/ microtubule system. *J. Biol. Chem.* 1997; 272(41): 25542-6.
- Volker KW, Reinitz CA & Knull HR. Glycolytic enzymes and assembly of microtubule networks. *Comp Biochem Physiol B Biochem Mol Biol.* 1995;112(3):503-14.
- Waingeh VF, Gustafson CD, Kozliak EI, Lowe SL, Knull HR & Thomasson KA. Glycolytic enzyme interactions with yeast and skeletal muscle F-actin. *Biophys J.* 2006; 90(4):1371-84.
- Walsh JL & Knull HR. Heteromeric interactions among glycolytic enzymes and of glycolytic enzymes with F-actin: effects of poly(ethylene glycol). *Biochim Biophys Acta.* 1988; 952(1):83-91.
- Walsh JL, Keith TJ & Knull HR. Glycolytic enzyme interactions with tubulin and microtubules. *Biochim. Biophys. Acta* 1989; 999(1):64-70.
- Welch MD, Mallavarapu A, Rosenblatt J & Mitchison TJ. Actin dynamics in vivo. *Curr Opin Cell Biol.* 1997; 9(1):54-61.
- Wera S, De Schrijver E, Geyskens I, Nwaka S & Thevelein JM. Opposite roles of trehalase activity in heat-shock recovery and heat-shock survival in *Saccharomyces cerevisiae*. *Biochem J.* 1999; 343Pt 3:621-626.
- Williams SP, Haggie PM & Brindle KM. 19F-NMR Measurements of the rotational mobility of proteins in vivo. *Biophys J.* 1997; 72:490-498.
- Xie GC & Wilson JE. Rat brain hexokinase: the hydrophobic N-terminus of the mitochondrially bound enzyme is inserted in the lipid bilayer. *Arch Biochem Biophys* 1988; 267(2): 803-10.
- Xu X, Forbes JG & Colombini M. Actin modulates the gating of *Neurospora crassa* VDAC. *J Membr Biol.* 2001; 180(1): 73-81.
- Zancan P & Sola-Penna M. Trehalose and glycerol stabilize and renature yeast inorganic pyrophosphatase inactivated by very high temperatures. *Arch. Biochem. Biophys.* 2005; 444(1):52-60.



Cell Metabolism - Cell Homeostasis and Stress Response

Edited by Dr. Paula Bubulya

ISBN 978-953-307-978-3

Hard cover, 208 pages

Publisher InTech

Published online 25, January, 2012

Published in print edition January, 2012

A global research community of scientists is teasing out the biochemical mechanisms that regulate normal cellular physiology in a variety of organisms. Much of current research aims to understand the network of molecular reactions that regulate cellular homeostasis, and to learn what allows cells to sense stress and activate appropriate biochemical responses. Advanced molecular tools and state-of-the-art imaging techniques discussed in this book continue to provide novel insights into how environmental changes impact organisms, as well as to develop therapeutic interventions for correcting aberrant pathways in human disease.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Daniela Araiza-Olivera, Salvador Uribe-Carvajal, Natalia Chiquete-Félix, Mónica Rosas-Lemus, Gisela Ruíz-Granados, José G. Sampedro, Adela Mújica and Antonio Peña (2012). Metabolic Optimization by Enzyme-Enzyme and Enzyme-Cytoskeleton Associations, Cell Metabolism - Cell Homeostasis and Stress Response, Dr. Paula Bubulya (Ed.), ISBN: 978-953-307-978-3, InTech, Available from:
<http://www.intechopen.com/books/cell-metabolism-cell-homeostasis-and-stress-response/metabolic-optimization-by-enzyme-enzyme-and-enzyme-cytoskeleton-associations->

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen