

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Identification of the Novel Plasminogen Receptor, Plg-R_{KT}

Lindsey A. Miles¹, Nicholas M. Andronicos², Emily I. Chen³,
Nagyung Baik¹, Hongdong Bai⁴, Caitlin M. Parmer⁵,
Shahzad Lighvani¹, Samir Nangia^{4,6}, William B. Kiosses¹,
Mark P. Kamps⁶, John R. Yates III¹ and Robert J. Parmer^{4,6}

1. Introduction

1.1 The plasminogen activation system

Initiation of the plasminogen activation system results in generation of the broad spectrum serine protease, plasmin, from the circulating zymogen, plasminogen. Plasminogen is activated to plasmin by plasminogen activators (PA's), either urokinase type-plasminogen activator (u-PA) or tissue-type plasminogen activator (t-PA), via specific proteolytic cleavage (Castellino & Ploplis, 2005). Plasmin is the major enzyme responsible for degradation of fibrin clots (fibrinolysis) to maintain normal blood homeostasis (Bugge et al., 1995; Ploplis et al., 1995). Dysregulation of the plasminogen activation system can result in hemorrhage (excess fibrinolysis) or thrombosis (insufficient fibrinolysis). The plasminogen activation system is regulated by direct inhibition of plasmin (by the circulating serpin, α_2 -antiplasmin) and by synthesis and secretion of plasminogen activators and the serpin, plasminogen activator inhibitor 1 (PAI-1)] (Collen, 1999). In a key regulatory step, plasminogen activation is promoted when plasminogen and its activator, t-PA, bind concomitantly to lysine residues on the surface of fibrin clots, resulting in a marked reduction in the K_m for activation of plasminogen compared with the reaction in solution (Hoylaerts et al., 1982).

1.2 Functions of plasminogen receptors

In the past 25 years an additional mechanism for positive regulation of plasminogen activation has been recognized: co-localization of plasminogen and PA's on cell surfaces markedly decreases the K_m for plasminogen activation in a mechanism analogous to assembly of components of the plasminogen activation system on fibrin (Miles et al., 2005). The plasmin produced is retained on the cell surface where it is protected from its inhibitor, α_2 -antiplasmin (Figure 1) (Hall et al., 1991; Plow et al., 1986). Thus, the cell surface becomes

¹The Scripps Research Institute, USA

²CSIRO Livestock Industries, Armidale, NSW, Australia


³Stony Brook University, USA

⁴Veterans Administration San Diego Healthcare System, USA

⁵Yale University, USA

⁶University of California, San Diego, USA

armed with the broad spectrum proteolytic activity of plasmin. Cell surface plasmin plays a key role in processes in which cells must degrade an extracellular matrix in order to migrate, including inflammation (Busuttill et al., 2004; Ploplis et al., 1998; Plow et al., 1999), wound healing (Creemers et al., 2000; Romer et al., 1996), metastasis (Palumbo et al., 2003; Ranson et al., 1998) and neurite outgrowth (Gutierrez-Fernandez et al., 2009; Jacovina et al., 2001). Cell surface plasmin also plays a key role in myogenesis (Lopez-Aleman et al., 2003) and prohormone processing (Jiang et al., 2001, 2002).



Activation of cell-associated plasminogen (Plg) to plasmin (Pm) by cell-associated plasminogen activators (PA) is markedly enhanced compared to the reaction in solution. The Pm formed remains on the cell surface where it is relatively protected from its inhibitor, α_2 -antiplasmin (α_2 -AP).


Fig. 1. Enhancement of plasminogen activation on the cell surface.

1.3 Characteristics of plasminogen receptors

Cellular plasminogen binding sites are very broadly distributed on both prokaryotic and eukaryotic cells. Of the many cell types examined to date, only red cells do not exhibit detectable plasminogen binding ability (Miles et al., 2005). The interactions of plasminogen with cells are of very high capacity, reaching 3×10^7 molecules/cell on lung fibroblasts (Plow et al., 1986), for example. Thus, the plasminogen binding capacity of a cell is made up of contributions from a set of distinct cell surface proteins.

An important aspect of the mechanism of promotion of plasminogen activation on cell surfaces is that a subset of carboxypeptidase-B-sensitive plasminogen binding proteins is responsible for enhancement of plasminogen activation on eukaryotic cells. When cells are treated with carboxypeptidase B, the ability to stimulate plasminogen activation is lost (Félez et al., 1996). Since carboxypeptidase B removes C-terminal basic residues, these results imply that proteins exposing C-terminal basic residues on cell surfaces are responsible for stimulation of plasminogen activation (Figure 2,A). Known carboxypeptidase-B-sensitive cell surface plasminogen receptors could previously be divided into two classes: 1) proteins synthesized with C-terminal lysines and having

additional known intracellular functions, including α -enolase (Miles et al., 1991; Redlitz et al., 1995), cytokeratin 8 (Hembrough et al., 1995, 1996), S100A10 (Choi et al., 2003; Kassam et al., 1998), TIP49a (Hawley et al., 2001) and histone H2B (Herren et al., 2006) and; 2) proteins requiring proteolytic processing in order to reveal a C-terminal basic residue (lysine), including actin (Dudani & Ganz, 1996; Miles et al., 2006) and annexin 2 (Hajjar et al., 1994). However, until recently, no integral membrane plasminogen binding proteins synthesized with a C-terminal basic residue had been identified. The existence of a plasminogen receptor with the latter characteristics would reveal a novel mechanism for stimulation of plasminogen activation because release and rebinding of intracellular proteins or proteolytic cleavage of membrane proteins to expose C-terminal basic residues would not be required.



Plasminogen binding to carboxyl terminal lysines on the cell surface. Panel A) The binding of plasminogen to cell surface proteins occurs via receptors exposing carboxyl terminal lysines to the extracellular environment. Cell surface proteins with carboxyl terminal lysines that are masked or in other inaccessible orientations on the cell surface, or membrane-associated proteins with carboxyl terminal lysines that are located on the inner face of the membrane, cannot serve as plasminogen receptors. Panel B) CpB treatment of intact cells removes carboxyl terminal lysines from plasminogen receptors, and plasminogen binding to the cell surface is reduced. Reprinted with permission from (Hawley, Green, and Miles 2000, 84:882-890).

Fig. 2. Plasminogen binding to carboxyl terminal lysines on the cell surface.

1.4 Need for a proteomics approach to identify integral membrane plasminogen receptor(s) with C-terminal basic residues

Previous methodologies and characteristics of plasminogen binding proteins may have precluded identification of an integral membrane plasminogen binding protein with a C-terminal basic residue. The identification of plasminogen receptors has relied previously on cell surface labeling followed by affinity chromatography on plasminogen-Sepharose columns and N-terminal sequencing of fractions eluted from SDS gels. Thus, many

intracellular proteins that are also present on the cell surface were readily identified because protein fractions that bound to plasminogen-Sepharose included the labeled, surface-associated protein, as well as nonlabeled intracellular protein. Using these methods, a lower abundance integral membrane plasminogen binding protein might not have been detectable.

Previously, we used a proteomics approach to examine monocytoïd cell membranes for the presence of proteins exposing carboxyl terminal lysines on the extracellular face of the cell membrane (Hawley et al., 2000). We compared plasminogen ligand blots of 2-D gels of membrane fractions of intact cells treated with carboxypeptidase B with untreated membranes (e.g. Figure 2,B). We eluted a prominent carboxypeptidase B-sensitive protein from the 2-D gels and obtained two peptide sequences using tandem mass spectrometry. These peptide sequences corresponded to TATA-binding protein-interacting protein (TIP49a) (Hawley et al., 2001). However, TIP49a is a member of the class of cell surface plasminogen receptors synthesized with a C-terminal lysine and also having intracellular functions and is not an integral membrane protein.

The methodology used to identify TIP49a and other plasminogen receptors has required elution of candidate proteins from 2-D SDS polyacrylamide gels. However, many membrane proteins are not well resolved on SDS polyacrylamide gels. Therefore, we used an isolation method that used column chromatography instead of SDS polyacrylamide gel analysis: We took advantage of the exquisite sensitivity of multidimensional protein identification technology (MudPIT) to search for integral membrane plasminogen receptor(s) exposing a C-terminal basic residue on the cell surface and present on viable cells.

2. Methods

2.1 Plasminogen receptor isolation

Plasminogen receptor isolation was performed as described (Andronicos et al., 2010). Briefly, progenitor and M-CSF-differentiated Hoxa9-ER4 cells were separately biotinylated, using EZ-Link Amine-PEO₃-Biotin. The cells were then subjected to dead cell removal on annexin V-coated magnetic microspheres that resulted in a 99% enrichment of viable cells. Membrane fractions were prepared from the viable cells by dounce homogenization in the presence of Complete Protease Inhibitor Cocktail in Invitrosol, followed by centrifugation steps as used in our laboratory (Hawley et al., 2000, 2001) and were applied to a plasminogen-Sepharose affinity column as described (Miles et al., 1991). The column was washed in phosphate buffered saline containing 1 X Invitrosol until no protein was detected at 280 nm followed by elution with the washing buffer containing 0.2 M ϵ -aminocaproic acid (EACA). The eluant from the plasminogen-Sepharose column was incubated with 50 μ l of immobilized avidin for 30 minutes at 4°C. The proteins bound to the immobilized avidin were resuspended in Invitrosol and heated at 60°C. Then, 80% acetonitrile was added and the samples were digested with trypsin. After 24 h, the solvent was evaporated in a speedvac, and peptides were dissolved in buffer A (95% H₂O, 5% acetonitrile, and 0.1% formic acid).

2.2 Multidimensional chromatography and tandem mass spectrometry

Multidimensional chromatography and tandem mass spectrometry were performed as described (Andronicos et al., 2010). Briefly, the protein digest was subjected to MudPIT [reviewed in (Eng et al., 1994)]. Peptide mixtures were resolved by strong cation exchange

liquid chromatography upstream of reversed phase liquid chromatography (Larmann, Jr. et al., 1993; Link et al., 1999; Opiteck & Jorgenson, 1997; Wolters et al., 2001). Eluting peptides were electrosprayed onto an LTQ ion trap mass spectrometer equipped with a nano-LC electrospray ionization source. Full MS spectra were recorded over a 400–1600 m/z range, followed by three tandem mass (MS/MS) events sequentially generated in a data-dependent manner on the first, second, and third most intense ions selected from the full MS spectrum (at 35% collision energy). Mass spectrometer scan functions and HPLC solvent gradients were controlled by the Xcalibur data system.

2.3 Database search and interpretation of MS/MS datasets

Database searching and interpretation of MS/MS Datasets were performed as described (Andronicos et al., 2010). Briefly, tandem mass spectra were extracted from raw files, and a binary classifier (Bern et al., 2004), previously trained on a manually validated data set, was used to remove low quality MS/MS spectra. Remaining spectra were searched against a mouse protein database containing 50,370 protein sequences downloaded as FASTA-formatted sequences from EBI-IPI and 124 common contaminant proteins, for a total of 66,743 target database sequences (Peng et al., 2003). To calculate confidence levels and false positive rates, a decoy database containing the reverse sequences of the 66,743 proteins appended to the target database and the SEQUEST algorithm (Yates, III, 1998) were used to find the best matching sequences from the combined database.

SEQUEST searches were done on an Intel Xeon 80-processor cluster running under the Linux operating system. The peptide mass search tolerance was set to 3 Da. No differential modifications were considered. No enzymatic cleavage conditions were imposed on the database search, so the search space included all candidate peptides whose theoretical mass fell within the 3 Da mass tolerance window, despite their tryptic status.


The validity of peptide/spectrum matches was assessed in DTASelect2 (Tabb et al., 2002) using SEQUEST-defined parameters, the cross-correlation score (XCorr) and normalized difference in cross-correlation scores (DeltaCN). The search results were grouped by charge state (+1, +2, and +3) and tryptic status (fully tryptic, half-tryptic, and non-tryptic), resulting in 9 distinct sub-groups. In each one of the sub-groups, the distribution of XCorr and DeltaCN values for direct and decoy database hits was obtained, and the two subsets were separated by quadratic discriminant analysis. Outlier points in the two distributions (for example, matches with very low Xcorr but very high DeltaCN) were discarded. Full separation of the direct and decoy subsets is not generally possible; therefore, the discriminant score was set such that a false positive rate of 5% was determined based on the number of accepted decoy database peptides. This procedure was independently performed on each data subset, resulting in a false positive rate independent of tryptic status or charge state.

3. Results

3.1 Isolation of an integral membrane plasminogen receptor exposing a C-terminal lysine on the cell surface

We used specific proteolysis followed by MudPIT to probe the membrane proteome of differentiated mouse monocyte progenitor cells (Hoxa9-ER4) for the presence of integral membrane plasminogen receptor(s) exposing a C-terminal basic residue on the cell surface, as outlined in Figure 3. [The Hoxa9-ER4 cell line was derived from primary murine bone

marrow myeloid precursors immortalized with an estrogen regulated conditional oncoprotein, HoxA9-ER4 (Wang et al., 2006). The Hoxa9-ER4 line is factor-dependent and differentiates to monocytes when estrogen is removed from the medium, thereby inactivating the Hoxa9-ER protein. The mature monocytes respond to M-CSF (Odegaard et al., 2007)]. First, the Hoxa9-ER4 monocyte progenitor cells were differentiated with macrophage colony stimulating factor (M-CSF), which induces plasminogen receptors on these cells (Andronicos et al., 2010). Then intact cells were biotinylated. Because early apoptotic and non-viable/necrotic cells exhibit markedly enhanced plasminogen binding ability (Mitchell et al., 2006; O'Mullane & Baker, 1998, 1999) we passed the biotinylated cells over a dead cell removal column to enrich for live cells. Cells were then lysed and membrane fractions prepared and passed over a plasminogen-Sepharose affinity column and specifically eluted with ϵ -aminocaproic acid (EACA), a lysine analog that blocks the binding of plasminogen to cells (Miles & Plow, 1985). Biotinylated proteins bound to the avidin column and were digested with trypsin while still on the column. The peptide digest was then subjected to MudPIT.



Monocyte (Hoxa9-ER4) progenitor cells were differentiated with macrophage colony stimulating factor (M-CSF), which induces plasminogen receptors (\blacktriangle) on these cells. Then intact cells were biotinylated (\bullet) and passed over a dead cell removal column. Live cells were then lysed and membrane fractions prepared and passed over a plasminogen-Sepharose affinity column and specifically eluted. Biotinylated plasminogen receptors ($\blacktriangle\bullet$) were then bound to an avidin column and digested with trypsin.

Fig. 3. Isolation of plasminogen receptors.

In MudPIT, the peptide mixtures were first resolved by strong cation exchange liquid chromatography followed by reversed phase liquid chromatography. Eluting peptides were electrosprayed onto an LTQ ion trap mass spectrometer and full MS spectra were recorded over a 400-1600 m/z range, followed by three tandem mass events. The spectra obtained were searched against a mouse protein database. Using this method, only one protein with a predicted transmembrane sequence and a C-terminal basic residue was identified: the hypothetical protein, C9orf46 homolog (IPI00136293), homologous to the protein predicted to be encoded by human chromosome 9, open reading frame 46. The peptides corresponding to C9orf46 homolog that were obtained in the MudPIT analysis are shown in Table 1. We have designated the protein, Plg-R_{KT}, to indicate a plasminogen receptor with a C-terminal lysine and having a transmembrane domain (see below). [A limitation of shotgun proteomics, such as MudPIT, is that they typically under sample a proteome because they use data dependent data acquisition (a computer-driven data acquisition approach). This can lead to variations in the proteins identified, particularly amongst the lower abundant proteins. Thus, we cannot exclude the possibility that other membrane proteins exposing C-terminal lysines were present in the membrane proteome.]

Xcorr	DeltCN	Conf%	ObsM+H+	CalcM+H+	Peptide Sequences
3.8378	0.2884	99.8%	2195.8743	2196.4788	K.SMNENMKNQQEFMVTHAR.L (3+)
2.6749	0.1167	95.2%	1359.5521	1361.5181	K.NQQEFMVTHAR.L (2+)
2.6771	0.2534	99.8%	1160.4321	1160.3514	R.HLTMQNEMR.E (2+)
4.7468	0.3052	100%	1523.5322	1523.6954	R.MKSEAEDILETEK.T (2+)
5.1774	0.3788	100%	2335.5544	2333.6997	R.MKSEAEDILETEKTKLELPK.G (3+)
3.775	0.3164	100%	1264.0922	1264.3287	K.SEAEDILETEK.T (2+)
2.995	0.0655	96.8%	1137.2722	1137.3184	K.GLITFESLEK.A (2+)
2.893	0.2591	99.7%	1364.4922	1364.5848	K.GLITFESLEKAR.R (2+)

SEQUEST-defined parameters (Xcorr, DeltCN, and Conf%) are shown for each peptide. (core: cross-correlation score; DeltCN: normalized difference in cross-correlation scores; Conf%: confidence level of the peptide; ObsM+H+: observed peptide mass; CalcM+H+: theoretical peptide mass). Observed peptide mass, theoretical peptide mass, and charges of the peptide identified (3+ or 2+) are also shown to demonstrate accurate peptide identification. This research was originally published in Blood, Andronicos, N.M., Chen, E.I., Baik, N., Bai, H., Parmer, C.M., Kiosses, W.B., Kamps, M.P., Yates, J.R., III, Parmer, R.J., Miles, L.A., Proteomics-based discovery of a novel, structurally unique, and developmentally regulated plasminogen receptor, Plg-R_{KT}, a major regulator of cell surface plasminogen activation, Blood. 2010, 115: 1319-30.

Table 1. Peptides obtained corresponding to C9orf46 homolog

A key advantage of MudPIT is that proteins in a given proteome can be identified simultaneously. As proof of principle of our isolation method, peptides corresponding to other proteins previously identified as plasminogen binding proteins on monocytes were also detected in the membrane preparations: α -enolase, gamma actin, S100A10, histone H2B, annexin 2, and β_2 integrin.

	10	20	30	40	50	60
Mouse	MGFIFSKSMN	ENMKNQQEFM	VTHARLQLER	HLTMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Human	MGFIFSKSMN	ESMKNQKEFM	LMNARLQLER	QLIMQSEMRE	RQMAMQIAWS	REFLKYFGTF
Rat	MGFIFSKSMN	ENMKNQQEFM	VMHARLQLER	QLIMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Dog	MGFIFSKSMN	ENMKNQQEFM	LMNARLQMER	QLMMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Cow	MGFIFSKSMN	ENLKSQQEFM	LMNSRLQLER	QLIMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Alpaca	MGFIFSKSMN	ENMKSQQEFM	LMNARLQLER	QLMMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Chimpanzee	MGFIFSKSMN	ESMKNQKEFM	LMNARLQLER	QLIMQSEMRE	RQMAMQIAWS	REFLKYFGTF
Dolphin	MGFIFSKSMN	ENMKSQQEFM	LMNARLQLER	QLMMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Gibbon	MGFIFSKSMN	ESMKNQKEFM	LMNARLQLER	QLIMQSEMRE	RQMAMQIAWS	REFLKYFGTF
Guinea Pig	MGFIFSKSMN	ENMKNQQEFM	LMNARLQLER	QLLLQNEMRE	RQMAMQIAWS	REFLKYFGTF
Horse	MGFIFSKSMN	ENMKNQQEFM	LMNARLQLER	QLTMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Lemur	MGFIFSKSMK	ENMQNQQEFM	LMNARLQLER	QLTMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Opossum	MGFLFSKHMN	ENMKQQEFM	LMNARLQMER	QLTIQNEMRE	RQMAMQIAWT	REFLKYFGTF
Orangutan	MGFIFSKSMN	ESMKNQKEFM	LMNARLQLER	QLIMQSEMRE	RQMAMQIAWS	REFLKYFGTF
Panda (Giant)	MGFIFSKSMS	ENMKNQQEFM	LMNARLQLER	QLMMQNEMRE	RQMALQIAWS	REFLKYFGTF
Pig	MGFIFSKSMN	ENMKRQQEFM	LMNTRLQLER	QLIMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Rabbit	MGFIFSKSMN	ENLKNQQEFM	LMNARLQLER	QLMLQNEMRE	RQMAMQIAWS	REFLKYFGTF
Rhesus Monkey	MGFIFSKSMN	ESMKNQKEFM	LMSARLQLER	QLIMQSEMRE	RQMAMQIAWS	REFLKYFGTF
Tarsier	MGFIFSKSMN	ENMKHQEFM	LMNAQLQLER	QLTMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Tree Shrew	MGFIFSKSMN	ENMKNQQEFM	LMNARLQLER	QLMMQNEMRE	RQMAMQIAWS	REFLKYFGTF
Lizard (arboreal)	MGFIFSKSMN	ENLKNQQEFM	IMNSRLQLER	QLLMQNEMRE	RQMAMQIAWT	REFLKYFGAF
Frog (xenopus)	MGSLSKATE	TQMKKQQELM	QMNAQIQLER	QITIMQNMRE	RQMAMQIAWS	REFLKYGFSF
	70	80	90	100	110	120
Mouse	FGIATISLAT	GALKRKKPAF	LVPIVPLSFI	FTYQYDLGYG	TLLQRMKSEA	EDILETEKTK
Human	FGLAAISLTA	GAIKKKKPAF	LVPIVPLSFI	LTYQYDLGYG	TLLERMKGEA	EDILETEKSK
Rat	FGIATISLAA	GAIKRKKPAF	LPIVPLSFI	FTYQYDLGYG	TLLQRMKSEA	EDILETEKTK
Dog	FGIAAISLTA	GAIRKKKPAF	LFPIIPLSFI	FTYQYDLGYG	TLLQRMKGEA	ENILETEKSK
Cow	FGTAAISLTA	GAIKGKPLV	IFPIVPLGFV	LAYQYDMGYG	TLLHRMKGEA	ENILETEKSK
Alpaca	FGIAAISLTA	GAIKRKKPAF	IFPIVPLGFV	LTYQFDLGYG	TLLQRMKGEA	ENILETEKSK
Chimpanzee	FGLAAISLTA	GAIKKKKPAF	LVPIVPLSFI	LTYQYDLGYG	TLLERMKGEA	EDILETEKSK
Dolphin	FGIAAISLTA	GAIKKKKPAF	VFPIVPLGFV	LAYQYDMGYG	TLLIQRMKGEA	DNILETEKSK
Gibbon	FGTAAISLTA	GAIKKKKPAF	LVPIVPLSFI	LTYQYDLGYG	TLLERMKGEA	ENILETEKSK
Guinea Pig	FGISAISLTA	RAIKQKKPAF	FPIVPLSFI	LAYQYDLGYG	TLLQRMKGEA	EDILETEKSK
Horse	FGIAAISLTA	GALKRKKPAF	LFPIVPLGFV	LTYQYDLGYG	TLLQRMKGEA	ENILETEKSK
Lemur	FGITAISLTA	GAIKSKKPGF	LFPIVPLSFI	LAYQYDLGYG	TLLQRMKGEA	EDILETEKSK
Opossum	FGIAAISLTA	GAIKKKKPAF	FFPIVPLSFI	LAYQYDMGYG	TLLQRMKGEA	ENILETEKSK
Orangutan	FGLAAISLTA	GAIKKKKPAF	LVPIVPLSFI	LTYQYDLGYG	TLLERMKGEA	EDILETEKSK
Panda (Giant)	FGITAISLTA	GAIRKKKPAF	LFPIIPLSFI	FTYQYDLGYG	TLLQRMKGEA	ENILETEKSK
Pig	FGIASVALTA	GAIKRKKPAF	FLPIIPLGFV	FTYQYDLGYG	TLLQRMKGEA	ENILETEKSK
Rabbit	FGVATISLTA	GAMRRKKPAF	LLPIVPLSFI	FTYQYDLGYG	TLLQRMKGEA	ENILETEKSK
Rhesus Monkey	FGFAAISLTA	GAIKKKKPAF	LVPIVPLSFI	LTYQYDLGYG	TLLERMKGEA	EDILETEKSK
Tarsier	FGITAISLTA	GAIKRKKPAL	LLPIVPLSFI	FTYQYDLGYG	TLLERMKGEA	EEILBAEKNM
Tree Shrew	FGIAAISLTA	GAIKKKNPAF	FFPIVPLSFI	LTYQYDLGYG	TLLPRMKSEA	EDILETEKSK
Lizard (arboreal)	SGLAAVGLTV	GAIKRKKPAF	FLPMVPLSFI	LAYQYDMGYG	SLLKRMKSEA	ESILDTESTT
Frog (xenopus)	FSLAVIGLTV	GAVKNKKPAL	FTPVIPLTFV	FAYQFDMGYG	TLVTRMKGEA	ENILEKEHIL
	130	140	147			
Mouse	LELPKGLITF	ESLEKARREQ	SKLFSK			
Human	LQLPRGMITF	ESIEKARKEQ	SRFFIDK			
Rat	LELPKGLITF	ESLEKARREQ	SKFFSDK			
Dog	LQLPRGMITF	ESLEKARREQ	SKFFIDK			
Cow	LQLPKGMITF	ESLEKARKEQ	SKFFIDK			
Alpaca	LQLPKGIITF	ESLEKARKEQ	SKFFIDK			
Chimpanzee	LQLPRGMITF	ESIEKARKEQ	SRFFIDK			
Dolphin	LQLPKGMITF	ENLEKARREQ	SKFFIDK			
Gibbon	LQLPRGMITF	ESIEKARKEQ	SKFFIDK			
Guinea Pig	LELPKGVITF	ESLEKARREQ	SKFFLGK			
Horse	LQLPKGMITF	ESLEKARREQ	SKFFIDK			
Lemur	LQLPKGMITF	ESLEKARREQ	SKFFIEK			
Opossum	LQLPRGSITF	ETLEKARKAQ	SKFFIEK			
Orangutan	LQLPRGMITF	ESIEKARKEQ	SRFFIDK			
Panda (Giant)	LQLPRGMITF	ENLEKARREQ	SKFFIDK			
Pig	LQLPKGMITF	EGLEKARREQ	SKFFIDK			
Rabbit	LQLPGGMITF	ESLEKARREQ	SKFFIDK			
Rhesus Monkey	LQLPRGMITF	ESIEKARKEQ	SKFFIDK			
Tarsier	LQLPKGMITF	ESLEKTRREQ	SKFFTDK			
Tree Shrew	LELPKGMITF	ESLEKARREQ	SKFFVVK			
Lizard (arboreal)	LEMPKGPLTF	ESIEKARRAQ	SKFFIEK			
Frog (xenopus)	LEMPQGLPTF	EGIEKTRKAH	RSLLL-K			

Table 2. Alignment of Orthologs of Plg-R_{KT}

3.2 Conservation of Plg-R_{KT} across species

The C9orf46 homolog/Plg-R_{KT} murine DNA sequence encodes a protein of 147 amino acids with a molecular mass of 17,261 Da and a C-terminal lysine (Table 2, first line). We blasted the C9orf46 homolog/Plg-R_{KT} sequence against all species using NCBI Blast and obtained unique human, rat, dog, cow, dog, giant panda, gibbon, horse, pig, rabbit, and rhesus monkey predicted orthologs, with high identity and homology (e.g. human versus rhesus monkey = 99% similarity), high identity (e.g. human vs rhesus monkey = 98% identity) and no gaps in the sequence (Table 2). Of key importance, a C-terminal lysine was predicted for all of the mammalian orthologs obtained in the blast search. In a query of the Ensembl Gene Report, DNA sequences of all 10 other sequenced mammalian orthologs encoded C-terminal lysines (Table 2).

In addition, the DNA sequences of xenopus and the green lizard also encoded C-terminal lysines (Table 2). Furthermore, Plg-R_{KT} orthologs with 149 amino acids with a C-terminal lysine were encoded in bony fish (salmon and zebrafish) and the high similarity with a mammalian ortholog is illustrated in the alignment of the mouse and zebrafish proteins in Table 3.

The Plg-R_{KT} sequence also encodes a putative conserved DUF2368 domain (encompassing amino acids 1-135), an uncharacterized protein with unknown function conserved from nematodes to humans. Notably, Plg-R_{KT} orthologs of lower organisms were of different predicted lengths and did not consistently predict C-terminal lysines. It is interesting to note that the evolutionary origin of plasminogen is currently believed to originate with protochordates (Liu & Zhang, 2009), so that lower organisms without plasminogen would not utilize the C-terminal lysine of Plg-R_{KT} to bind plasminogen.


Mouse	1	MGFIFSKSMNENMKNQQEFMVTHARLQLERHLTMQNEMRERQMAMQIAWSREFLKYFGTF	60
Zebrafish	1	MGFVLSKGMQNFQKQQEFMLLNARLQLERQLAMQNQMRRERQMAMQLAWSREFLKYFGSF	60
Mouse	61	FGIATISLATGALKRKKPAFLVPIVPLSFIFTYQYDLGYGTLQRMKSEAEDILETEKTK	120
Zebrafish	61	FGLATLGLTVGAVKKRKPALLAPVIPLSFILVYQMDAAYGTMLQRMRAEAESIMVSECER	120
Mouse	121	LLEPKGLITTFESLEKARREQSKL--FSDK	147
Zebrafish	121	LDVPHGMPTFESIEKSRRAKAHLTTLTEK	149

Table 3. Alignment of Mouse and Zebrafish Plg-R_{KT} Sequences

It is also noteworthy that the primary sequence of C9orf46 homolog/Plg-R_{KT} is apparently tightly conserved in humans, with no validated polymorphisms (cSNPs) within the 6 exons encoded by the gene (on chromosome 9p24.1) in the NCBI human genome sequence variation database (dbSNP, <http://www.ncbi.nlm.nih.gov/SNP>).

3.3 Topology of Plg-R_{KT}

The C9orf46 homolog/Plg-R_{KT} sequence was analyzed in the TMpred site (www.ch.embnet.org/cgi-bin/TMPRED). The model predicted two transmembrane helices extending from F₅₃-L₇₃ (secondary helix, oriented from outside the cell to inside the cell) and P₇₈-Y₉₉ (primary helix, oriented from inside the cell to outside the cell) (Figure 4). Hence a 52 amino acid N-terminal region and a 48 amino acid C-terminal tail with a C-terminal lysine were predicted to be exposed on the cell surface.




A. Membrane fractions or cytoplasmic fractions from either undifferentiated or M-CSF-treated Hoxa9-ER4 cells were electrophoresed on 12% sodium dodecyl sulfate polyacrylamide gels under reducing conditions and western blotted with either anti-Plg-R_{KT} mAb, anti-α-enolase mAb as a loading control, or isotype control mAb. B. M-CSF-treated Hoxa9-ER4 cell membranes were solubilized in 3% Triton X-114. After heating at 37°C and separation of the phases by centrifugation, an aliquot of both phases was electrophoresed and western blotted with anti-Plg-R_{KT} mAb. This research was originally published in *Blood*, Andronicos, N.M., Chen, E.I., Baik, N., Bai, H., Parmer, C.M., Kiosses, W.B., Kamps, M.P., Yates, J.R., III, Parmer, R.J., Miles, L.A., Proteomics-based discovery of a novel, structurally unique, and developmentally regulated plasminogen receptor, Plg-R_{KT}, a major regulator of cell surface plasminogen activation, *Blood*. 2010, 115: 1319-30.

Fig. 5. Plg-R_{KT} behaves as a regulated integral membrane protein.

removed but intracellular C-terminal lysines are protected (see Figure 2,B). Under this condition, no peptides corresponding to Plg-R_{KT} were obtained in the MudPIT analysis, consistent with cell surface exposure of the C-terminal lysine of Plg-R_{KT}.

In order to experimentally evaluate whether the N-terminus of Plg-R_{KT} was exposed on the cell surface, PC12 (rat pheochromocytoma) cells were stably transfected with V5-pCIneo-Plg-R_{KT} that expressed a V5 tag at the N-terminus of Plg-R_{KT}. (The V5 sequence was added in front of the mammalian expression vector, pCIneo using PCR and then the full-length 443 bp Plg-R_{KT} cDNA was subcloned into the V5-pCIneo vector using the Xho1 and Sma1 cloning sites. Constructs were transfected into cells using Lipofectamine 2000 and stable transfectants were selected with G418.)

A specific band migrating with a M_{rapp} of 17,000 was detected in cell membranes of the stably transfected cells with both anti-V5 mAb and anti-Plg-R_{KT} mAb (Figure 6, lane 1). The band was not detected by either mAb after trypsin digestion of the isolated membrane fraction (lane 2). When intact cells were incubated with trypsin and the trypsin neutralized with SBTI prior to preparation of the membrane fraction, the majority of the band detectable with either anti-V5 or anti-Plg-R_{KT} was lost (lane 4). In controls, treatment with soybean trypsin inhibitor (SBTI) fully neutralized the ability of trypsin to degrade the V5-tagged Plg-R_{KT} in purified membrane fractions (lane 3), demonstrating that the trypsin had been neutralized prior to membrane fractionation of the treated cells. These results suggest that the N-terminus of Plg-R_{KT} is accessible to trypsin proteolysis of intact cells and is, therefore,




Membrane fractions of PC12 cells stably transfected with V5-pCIneo-Plg-R_{KT} were incubated with either buffer (lane 1), trypsin (1mg/ml) (lane 2) or trypsin 1 mg/ml + soybean trypsin inhibitor (SBTI) (2 mg/ml) (lane 3) for 30 minutes at 37°C or intact PC12 cells were incubated with 1 mg/ml trypsin for 2 hr at 37°C, followed by 2 mg/ml SBTI for 15 min. Following neutralization of trypsin with SBTI, the membrane fraction was prepared from the treated, intact cells (lane 4). 30 µg/lane of membrane fractions were electrophoresed on 18% SDS PAGE under reducing conditions and western blotted with either anti-V5, anti-Plg-R_{KT} mAb or isotype control.

Fig. 6. The N-termini and C-termini of Plg-R_{KT} are exposed on the cell surface.

exposed on the extracellular face. Furthermore, because the anti-Plg-R_{KT} mAb reacts with the C-terminus of Plg-R_{KT}, these data also confirm the exposure of the C-terminus on the extracellular face of the cell membrane.

3.4 Role of the C-terminal lysine of Plg-R_{KT} in plasminogen binding

We further addressed the exposure of the C-terminus of Plg-R_{KT} on the cell surface using confocal microscopy with a mAb raised against the Plg-R_{KT} C-terminal peptide. (The mAb reacted with the C-terminal peptide of murine Plg-R_{KT} and blocked plasminogen binding to CEQSKLFSDK). When cells were incubated with anti-Plg-R_{KT} mAb and a polyclonal anti-plasminogen antibody, Plg-R_{KT} and plasminogen were both immunodetected in small aggregates dispersed over the cell surface (Figure 7,A), in a similar distribution to that published for confocal analyses of monocyte-associated plasminogen (Das et al., 2007). Most importantly, after preincubation of monocytes with plasminogen, immunodetection of Plg-R_{KT} was reduced by half (Figure 7,A,B). These results demonstrate that the C-terminus of Plg-R_{KT} is exposed on the cell surface. Furthermore, these results show that plasminogen binds to the C-terminal domain of Plg-R_{KT} on the cell surface.




A. M-CSF-differentiated (Hoxa9-ER4) cells were grown on coverslips and preincubated with either phosphate buffered saline (- plasminogen) or 2 μ M plasminogen (+ plasminogen), then fixed in 1% formaldehyde, washed and stained with polyclonal anti-plasminogen IgG or anti-Plg-R_{KT} mAb and stained with a combination of Alexa 488- F(ab')₂ of goat anti-rabbit IgG and Alexa 568- F(ab')₂ fragment of goat anti-mouse IgG.

B. The number and size of each labeled aggregate was determined. The results reflect counts from over 40 cells in 2 independent experiments. Data represent mean \pm SEM. * $p < 0.001$. This research was originally published in *Blood*, Andronicos, N.M., Chen, E.I., Baik, N., Bai, H., Parmer, C.M., Kiosses, W.B., Kamps, M.P., Yates, J.R., III, Parmer, R.J., Miles, L.A., Proteomics-based discovery of a novel, structurally unique, and developmentally regulated plasminogen receptor, Plg-R_{KT}, a major regulator of cell surface plasminogen activation, *Blood*. 2010, 115: 1319-30.

Fig. 7. Plg-R_{KT} binds plasminogen on the cell surface.

To further address the plasminogen binding function of the C-terminus of Plg-R_{KT}, we tested whether the synthetic peptide, corresponding to the C-terminus of Plg-R_{KT} could bind plasminogen. The peptide, CEQSKLFSDK, was coupled to BSA and then coated onto wells of microtiter plates. Biotinylated Glu-plasminogen was incubated with the wells and specific binding was detected with HRP-streptavidin (Figure 8). We tested the ability of the soluble C-terminal peptide to inhibit Glu-plasminogen binding under solution phase equilibrium

conditions. The soluble peptide competed for Glu-plasminogen binding in a dose-dependent manner with an IC_{50} of 2 μ M (Figure 8), similar to the K_d values we have previously determined for Glu-plasminogen binding to cells (Miles et al., 2005). In addition, a mutated peptide with the C-terminal lysine substituted with alanine did not compete for plasminogen binding at concentrations up to 1 mM (Figure 8), further supporting the role of the C-terminal lysine in the interaction of Plg- R_{KT} with plasminogen.



The peptide, CEQSKLFSDK, was coupled to BSA and immobilized on microtiter wells. Biotinylated-Glu-plasminogen (25 nM) was incubated with immobilized CEQSKLFSDK in the presence of increasing concentrations of CEQSKLFSDK (●) or a K147A mutant peptide, CEQSKLFSDA (○). Biotinylated Glu-plasminogen binding was detected with HRP-streptavidin. Data are as mean \pm SEM, $n=3$, for each determination. This research was originally published in *Blood*, Andronicos, N.M., Chen, E.I., Baik, N., Bai, H., Parmer, C.M., Kiosses, W.B., Kamps, M.P., Yates, J.R., III, Parmer, R.J., Miles, L.A., Proteomics-based discovery of a novel, structurally unique, and developmentally regulated plasminogen receptor, Plg- R_{KT} , a major regulator of cell surface plasminogen activation, *Blood*. 2010, 115: 1319-30.


Fig. 8. Plasminogen binds to the C-terminal peptide of Plg- R_{KT} .

3.5 Plg- R_{KT} regulates cell surface plasminogen activation

We verified that plasminogen activation was promoted in the presence of differentiated Hoxa9-ER4 cells. Plasminogen activation was stimulated 12.7-fold in the presence of differentiated monocyte progenitor cells, compared to the reaction in the absence of cells (Figure 9). In order to test the role of Plg- R_{KT} in plasminogen activation, we tested the effect of anti-Plg- R_{KT} mAb raised against the synthetic peptide, CEQSKLFSDK. Anti-Plg- R_{KT} mAb substantially suppressed cell-dependent plasminogen activation (Figure 9). In controls, plasminogen activation in the absence of cells was not affected by anti-Plg- R_{KT} mAb.

3.6 Tissue and cellular distribution and regulation of the Plg- R_{KT} transcript

We searched results of gene expression array analyses for expression of the C9orf46 homolog/Plg- R_{KT} transcript. The transcript is broadly expressed in normal human and mouse tissues [as determined in high-throughput gene expression profiling in which RNA samples from human and murine tissues were hybridized to high-density gene expression arrays (Su et al., 2002; Su et al., 2004)]. The C9orf46 homolog/Plg- R_{KT} transcript has been



Plasminogen activation was determined in either the presence or absence of differentiated Hoxa9-ER4 cells and in the presence of either anti-Plg-R_{KT} mAb (filled bars) or isotype control rat IgG2a (open bars). ***p < 0.001, compared to the corresponding isotype control. This research was originally published in Blood, Andronicos, N.M., Chen, E.I., Baik, N., Bai, H., Parmer, C.M., Kiosses, W.B., Kamps, M.P., Yates, J.R., III, Parmer, R.J., Miles, L.A., Proteomics-based discovery of a novel, structurally unique, and developmentally regulated plasminogen receptor, Plg-R_{KT}, a major regulator of cell surface plasminogen activation, Blood. 2010, 115: 1319-30.

Fig. 9. Plg-R_{KT} regulates cell surface plasminogen activation.

detected in spleen, lymph node, thymus, bone marrow, lung, intestine, adrenal, pituitary, and other endocrine tissues, vascular tissue, kidney, liver, stomach, bladder, and neuronal tissue (hippocampus, hypothalamus, cerebellum, cerebral cortex, olfactory bulb and dorsal root ganglion) (Table 4).

We also searched for C9orf46 homolog/Plg-R_{KT} mRNA microarray expression data at <http://www.ebi.ac.uk/microarray-as/aew/>. C9orf46 homolog/Plg-R_{KT} mRNA is present in monocytes, leukocytes, natural killer (NK) cells, T cells, myeloid, dendritic, and plasmacytoid cells, breast cancer, acute lymphoblastic leukemia and Molt-4 acute lymphoblastic leukemia cells (Table 5).

• Spleen	• Liver
• Thymus	• Stomach
• Lymph Node	• Bladder
• Lung	• Hippocampus
• Intestine	• Hypothalamus
• Bone Marrow	• Cerebellum
• Adrenal	• Cerebral Cortex
• Pituitary	• Olfactory Bulb
• Vascular Tissue	• Dorsal Root Ganglion
• Kidney	

Results of high-throughput gene expression profiling (54).

Table 4. Tissue Distribution of Plg-R_{KT} mRNA

-
- Monocytes
 - NK cells
 - T cells
 - Myeloid cells
 - Dendritic cells
 - Plasmacytoid cells
 - B-cell precursor cells
 - Acute lymphoblastic leukemia cells
 - Molt-4 acute lymphoblastic leukemia cells
 - Breast cancer cells, SKBR3, MDA468, BT474, T47D
-

www.ebi.ac.uk/microarray-as/aew/

Table 5. Microarray Expression Data for Plg-R_{KT} mRNA

These data are consistent with previous reports documenting expression of plasminogen binding sites on peripheral blood leukocytes (Miles & Plow, 1987), breast cancer cells (Correc et al., 1990; Ranson et al., 1998) and other tissues [reviewed in (Miles et al., 2005)]. In addition, results obtained by searching the ArrayExpress Warehouse (<http://www.ebi.ac.uk/microarray>) indicated that the C9orf46 homolog gene is also regulated in other tissues by lipopolysaccharide, aldosterone, canrenoate, H₂O₂, and dexamethasone (Table 6).

In a previously published genome-scale quantitative image analysis, overexpression of a cDNA that we now recognize to be the Plg-R_{KT} cDNA, resulted in dramatic increases in cell proliferation whereas knockdown of the corresponding mRNA resulted in apoptosis (Harada et al., 2005). Consistent with an anti-apoptotic role of Plg-R_{KT}, we have shown that cell-bound plasminogen inhibits TNF α -induced apoptosis (Mitchell et al., 2006). In microarray studies, C9orf46 homolog mRNA expression has a high power to predict cervical lymph node metastasis in oral squamous cell carcinoma (Nguyen et al., 2007).

Experiment	Tissue	Agonist	Effect*
E-MEXP-420	Hippocampal microglial cells	lipopolysaccharide	↓
E-TABM-229	Kidney	aldosterone	↓
E-TABM-229	Kidney	canrenoate	↓
E-MEXP-710	Cholinergic cells	H ₂ O ₂	↓
E-MEXP-774	Preadipocytes	Dexamethasone	↓

Data were obtained from ArrayExpress Warehouse (<http://www.ebi.ac.uk/microarray>).

* ↓ = downregulation. This research was originally published in Blood, Andronicos, N.M., Chen, E.I., Baik, N., Bai, H., Parmer, C.M., Kiosses, W.B., Kamps, M.P., Yates, J.R., III, Parmer, R.J., Miles, L.A., Proteomics-based discovery of a novel, structurally unique, and developmentally regulated plasminogen receptor, Plg-R_{KT}, a major regulator of cell surface plasminogen activation, Blood. 2010, 115: 1319-30.

Table 6. Regulation of C9orf46 homolog/Plg-R_{KT} mRNA in Tissues

4. Conclusions

In conclusion, MudPIT has allowed us to identify a new protein, Plg-R_{KT}, a novel plasminogen receptor with unique characteristics: integral to the cell membrane and exposing a C-terminal lysine on the cell surface in an orientation to bind plasminogen and promote plasminogen activation. Thus, Plg-R_{KT} is likely to play a key role in plasminogen-dependent functions of cells including inflammation, wound healing, development, metastasis, neurite outgrowth, fibrinolysis, myogenesis and prohormone processing. The

broad distribution of the Plg-R_{KT} transcript and its regulation in tissues that have been demonstrated to express plasminogen binding sites, suggest that Plg-R_{KT} provides plasminogen receptor function that may serve to modulate plasmin proteolytic functions (both physiologic and pathologic) specific to a large number of tissues. Furthermore, the potential function of Plg-R_{KT} in the regulation of apoptosis and proliferation may play a key role in cancer and metastasis. Future studies with knockout mice should build on our initial results using MudPIT to elucidate the role of Plg-R_{KT}.

5. Acknowledgments

Supported by National Institutes of Health Grants (HL38272, HL45934, and HL081046 to L.A.M., HL50398 to R.J.P., NIH P41 RR011823 to J.R.Y., NIAID sub-contract grant UCSD/MCB0237059 to E.I.C.) and Department of Veterans Affairs to R.J.P. S.L was supported by NIH training grant, T32 HL007195. We thank Dr. Ray Stevens at The Scripps Research Institute and Dr. Nuala Booth and Dr. Ian Booth, University of Aberdeen, for helpful discussions. We thank Ms. Linda Bonafede for manuscript preparation. This is publication #21378 from The Scripps Research Institute.


6. References

- Andronicos, N. M., E. I. Chen, N. Baik, H. Bai, C. M. Parmer, W. B. Kiosses, M. P. Kamps, J. R. Yates, III, R. J. Parmer, and L. A. Miles. 2010. Proteomics-based discovery of a novel, structurally unique, and developmentally regulated plasminogen receptor, Plg-RKT, a major regulator of cell surface plasminogen activation. *Blood* 115, no. 7:1319-1330.
- Bern, M., D. Goldberg, W. H. McDonald, and J. R. 3rd Yates. 2004. Automatic quality assessment of peptide tandem mass spectra. *Bioinformatics* 20, Suppl 1:149-154.
- Bordier, C. 1981. Phase separation of integral membrane proteins in Triton X-114 solution. *Journal of Biological Chemistry* 256, no. 4:1604-1607.
- Bugge, T. H., M. J. Flick, C. C. Daugherty, and J. L. Degen. 1995. Plasminogen deficiency causes severe thrombosis but is compatible with development and reproduction. *Genes and Development* 9:794-807.
- Busuttill, S. J., V. A. Ploplis, F. J. Castellino, L. Tang, J. W. Eaton, and E. F. Plow. 2004. A central role for plasminogen in the inflammatory response to biomaterials. *J.Thromb.Haemost.* 2, no. 10:1798-1805.
- Castellino, F. J. and V. A. Ploplis. 2005. Structure and function of the plasminogen/plasmin system. *Thrombosis and Haemostasis* 93, no. 4:647-654.
- Choi, K. S., D. K. Fogg, C. S. Yoon, and D. M. Waisman. 2003. p11 regulates extracellular plasmin production and invasiveness of HT1080 fibrosarcoma cells. *FASEB J* 17, no. 2:235-246.
- Collen, D. 1999. The plasminogen (fibrinolytic) system. *Thrombosis and Haemostasis* 82:259-270.
- Correc, P., M.-C. Fondanèche, M. Bracke, and P. Burtin. 1990. The presence of plasmin receptors on three mammary carcinoma MCF-7 sublines. *International Journal of Cancer* 46:745-750.
- Creemers, E., J. Cleutjens, J. Smits, S. Heymans, L. Moons, D. Collen, M. Daemen, and P. Carmeliet. 2000. Disruption of the plasminogen gene in mice abolishes wound healing after myocardial infarction. *American Journal of Pathology* 156, no. 6:1865-1873.
- Das, R., T. Burke, and E. F. Plow. 2007. Histone H2B as a functionally important plasminogen receptor on macrophages. *Blood* 110, no. 10:3763-3772.

- Dudani, A. K. and P. R. Ganz. 1996. Endothelial cell surface actin serves as a binding site for plasminogen, tissue plasminogen activator and lipoprotein(a). *British Journal of Haematology* 95:168-178.
- Eng, J. K., A. L. McCormick, and J. R. III Yates. 1994. An approach to correlate tandem mass spectral data of peptides with amino acid sequences in a protein database. *J.Am.Soc.Mass Spectrom.* 5:976-989.
- Estreicher, A., A. Wohlowend, D. Belin, W-D. Scluning, and J. D. Vassalli. 1989. Characterization of the cellular binding site for the urokinase-type plasminogen activator. *Journal of Biological Chemistry* 264:1180-1189.
- Félez, J., L. A. Miles, P. Fábregas, M. Jardi, E. F. Plow, and R. H. Lijnen. 1996. Characterization of cellular binding sites and interactive regions within reactants required for enhancement of plasminogen activation by tPA on the surface of leukocytic cells. *Thrombosis and Haemostasis* 76:577-584.
- Gutierrez-Fernandez, A., N. A. Gingles, H. Bai, F. J. Castellino, R. J. Parmer, and L. A. Miles. 2009. Plasminogen enhances neuritogenesis on laminin-1. *Journal of Neuroscience* 29, no. 40:12393-12400.
- Hajjar, K. A., A. T. Jacovina, and J. Chacko. 1994. An endothelial cell receptor for plasminogen/tissue plasminogen activator. I. Identity with annexin II. *Journal of Biological Chemistry* 269:21191-21197.
- Hall, S. W., J. E. Humphries, and S. L. Gonias. 1991. Inhibition of cell surface receptor-bound plasmin by α_2 -antiplasmin and α_2 -macroglobulin. *Journal of Biological Chemistry* 266:12329-12336.
- Harada, J. N., K. E. Bower, A. P. Orth, S. Callaway, C. G. Nelson, C. Laris, J. B. Hogenesch, P. K. Vogt, and S. K. Chanda. 2005. Identification of novel mammalian growth regulatory factors by genome-scale quantitative image analysis. *Genome Res.* 15, no. 8:1136-1144.
- Hawley, S. B., M. A. Green, and L. A. Miles. 2000. Discriminating between cell surface and intracellular plasminogen-binding proteins: heterogeneity in profibrinolytic plasminogen-binding proteins on monocytoid cells. *Thrombosis and Haemostasis* 84, no. 5:882-890.
- Hawley, S. B., T. Tamura, and L. A. Miles. 2001. Purification, cloning, and characterization of a profibrinolytic plasminogen-binding protein, TIP49a. *Journal of Biological Chemistry* 276:179-186.
- Hembrough, T. A., K. R. Kralovich, L. Li, and S. L. Gonias. 1996. Cytokeratin 8 released by breast carcinoma cells *in vitro* binds plasminogen and tissue-type plasminogen activator and promotes plasminogen activation. *Biochemical Journal* 317:763-769.
- Hembrough, T. A., J. Vasudevan, M. M. Allietta, W. F. Glass, and S. L. Gonias. 1995. A cytokeratin 8-like protein with plasminogen-binding activity is present on the external surfaces of hepatocytes, HepG2 cells and breast carcinoma cell lines. *J.Cell Sci.* 108 (Pt 3):1071-1082.
- Herren, T., T. A. Burke, R. Das, and E. F. Plow. 2006. Identification of histone H2B as a regulated plasminogen receptor. *Biochemistry* 45, no. 31:9463-9474.
- Hoylaerts, M., D. C. Rijken, H. R. Lijnen, and D. Collen. 1982. Kinetics of the activation of plasminogen by human tissue plasminogen activator. *Journal of Biological Chemistry* 257:2912-2919.
- Jacovina, A. T., F. Zhong, E. Khazanova, E. Lev, A. B. Deora, and K. A. Hajjar. 2001. Neuritogenesis and the nerve growth factor-induced differentiation of PC-12 cells requires annexin II-mediated plasmin generation. *Journal of Biological Chemistry* 276, no. 52:49350-49358.

- Jiang, Q., L. Taupenot, S. K. Mahata, M. Mahata, D. T. O'Connor, L. A. Miles, and R. J. Parmer. 2001. Proteolytic cleavage of chromogranin A (CgA) by plasmin. Selective liberation of a specific bioactive CgA fragment that regulates catecholamine release. *J Biol.Chem.* 276, no. 27:25022-25029.
- Jiang, Q., S. Yasothornsrikul, L. Taupenot, L. A. Miles, and R. J. Parmer. 2002. The local chromaffin cell plasminogen/plasmin system and the regulation of catecholamine secretion. *Annals of the New York Academy of Sciences* 971:445-449.
- Kassam, G., B. H. Le, K. S. Choi, H. M. Kang, S. L. Fitzpatrick, P. Louie, and D. M. Waisman. 1998. The p11 subunit of the annexin II tetramer plays a key role in the stimulation of t-PA-dependent plasminogen activation. *Biochemistry* 37, no. 48:16958-16966.
- Larmann, J. P., Jr., A. V. Lemmo, A. W. Moore, Jr., and J. W. Jorgenson. 1993. Two-dimensional separations of peptides and proteins by comprehensive liquid chromatography-capillary electrophoresis. *Electrophoresis* 14, no. 5-6:439-447.
- Link, A. J., J. Eng, D. M. Schieltz, E. Carmack, G. J. Mize, D. R. Morris, B. M. Garvik, and J. R. Yates, III. 1999. Direct analysis of protein complexes using mass spectrometry. *Nat.Biotechnol.* 17, no. 7:676-682.
- Liu, M. and S. Zhang. 2009. A kringle-containing protease with plasminogen-like activity in the basal chordate *Branchiostoma belcheri*. *Biosci.Rep.* 29, no. 6:385-395.
- Lopez-Aleman, R., M. Suelves, and P. Munoz-Canoves. 2003. Plasmin generation dependent on alpha-enolase-type plasminogen receptor is required for myogenesis. *Thrombosis and Haemostasis* 90, no. 4:724-733.
- Miles, L. A., N. M. Andronicos, N. Baik, and R. J. Parmer. 2006. Cell-surface actin binds plasminogen and modulates neurotransmitter release from catecholaminergic cells. *Journal of Neuroscience* 26, no. 50:13017-13024.
- Miles, L. A., C. M. Dahlberg, J. Plescia, J. Felez, K. Kato, and E. F. Plow. 1991. Role of cell-surface lysines in plasminogen binding to cells: Identification of alpha-Enolase as a candidate plasminogen receptor. *Biochemistry* 30:1682-1691.
- Miles, L. A., S. B. Hawley, N. Baik, N. M. Andronicos, F. J. Castellino, and R. J. Parmer. 2005. Plasminogen receptors: the sine qua non of cell surface plasminogen activation. *Front Biosci.* 10:1754-1762.
- Miles, L. A. and E. F. Plow. 1985. Binding and activation of plasminogen on the platelet surface. *Journal of Biological Chemistry* 260:4303-4311.
- Miles, L. A. and E. F. Plow. 1987. Receptor mediated binding of the fibrinolytic components, plasminogen and urokinase, to peripheral blood cells. *Thrombosis and Haemostasis* 58:936-942.
- Mitchell, J. W., N. Baik, F. J. Castellino, and L. A. Miles. 2006. Plasminogen inhibits TNF{alpha}-induced apoptosis in monocytes. *Blood* 107, no. 11:4383-4390.
- Nguyen, S. T., S. Hasegawa, H. Tsuda, H. Tomioka, M. Ushijima, M. Noda, K. Omura, and Y. Miki. 2007. Identification of a predictive gene expression signature of cervical lymph node metastasis in oral squamous cell carcinoma. *Cancer Sci.* 98, no. 5:740-746.
- O'Mullane, M. J. and M. S. Baker. 1998. Loss of cell viability dramatically elevates cell surface plasminogen binding and activation. *Experimental Cell Research* 242:153-164.
- O'Mullane, M. J. and M. S. Baker. 1999. Elevated plasminogen receptor expression occurs as a degradative phase event in cellular apoptosis. *Immunol.Cell Biol.* 77, no. 3:249-255.
- Odegaard, J. I., D. Vats, L. Zhang, R. Ricardo-Gonzalez, K. L. Smith, D. B. Sykes, M. P. Kamps, and A. Chawla. 2007. Quantitative expansion of ES cell-derived myeloid progenitors capable of differentiating into macrophages. *Journal of Leukocyte Biology* 81, no. 3:711-719.

- Opiteck, G. J. and J. W. Jorgenson. 1997. Two-dimensional SEC/RPLC coupled to mass spectrometry for the analysis of peptides. *Anal.Chem.* 69, no. 13:2283-2291.
- Palumbo, J. S., K. E. Talmage, H. Liu, C. M. La Jeunesse, D. P. Witte, and J. L. Degen. 2003. Plasminogen supports tumor growth through a fibrinogen-dependent mechanism linked to vascular patency. *Blood* 102, no. 8:2819-2827.
- Peng, J., J. E. Elias, C. C. Thoreen, L. J. Licklider, and S. P. Gygi. 2003. Evaluation of multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-MS/MS) for large-scale protein analysis: the yeast proteome. *J.Proteome.Res.* 2, no. 1:43-50.
- Ploplis, V. A., P. Carmeliet, S. Vazirzadeh, Vlaenderen Van, I, L. Moons, E. F. Plow, and D. Collen. 1995. Effects of disruption of the plasminogen gene on thrombosis, growth, and health in mice. *Circulation* 92, no. 9:2585-2593.
- Ploplis, V. A., E. L. French, P. Carmeliet, D. Collen, and E. F. Plow. 1998. Plasminogen deficiency differentially affects recruitment of inflammatory cell populations in mice. *Blood* 91:2005-2009.
- Plow, E. F., D. E. Freaney, J. Plescia, and L. A. Miles. 1986. The plasminogen system and cell surfaces: Evidence for plasminogen and urokinase receptors on the same cell type. *Journal of Cell Biology* 103:2411-2420.
- Plow, E. F., V. A. Ploplis, S. Busuttill, P. Carmeliet, and D. Collen. 1999. A role of plasminogen in atherosclerosis and restenosis models in mice. *Thrombosis and Haemostasis* 82 Suppl 1:4-7.
- Ranson, M., N. M. Andronicos, M. J. O'Mullane, and M. S. Baker. 1998. Increased plasminogen binding is associated with metastatic breast cancer cells: differential expression of plasminogen binding proteins. *British Journal of Cancer* 77:1586-1597.
- Redlitz, A., B. J. Fowler, E. F. Plow, and L. A. Miles. 1995. The role of an enolase-related molecule in plasminogen binding to cells. *European Journal of Biochemistry* 227:407-415.
- Romer, J., T. H. Bugge, C. Pyke, L. R. Lund, M. J. Flick, J. L. Degen, and K. Dano. 1996. Impaired wound healing in mice with a disrupted plasminogen gene. *Nat.Med.* 2, no. 3:287-292.
- Su, A. I., M. P. Cooke, K. A. Ching, Y. Hakak, J. R. Walker, T. Wiltshire, A. P. Orth, R. G. Vega, L. M. Sapinoso, A. Moqrich, A. Patapoutian, G. M. Hampton, P. G. Schultz, and J. B. Hogenesch. 2002. Large-scale analysis of the human and mouse transcriptomes. *Proc.Natl.Acad.Sci.U.S.A* 99, no. 7:4465-4470.
- Su, A. I., T. Wiltshire, S. Batalov, H. Lapp, K. A. Ching, D. Block, J. Zhang, R. Soden, M. Hayakawa, G. Kreiman, M. P. Cooke, J. R. Walker, and J. B. Hogenesch. 2004. A gene atlas of the mouse and human protein-encoding transcriptomes. *Proc.Natl.Acad.Sci.U.S.A* 101, no. 16:6062-6067.
- Tabb, D. L., W. H. McDonald, and J. R. Yates, III. 2002. DTASelect and Contrast: tools for assembling and comparing protein identifications from shotgun proteomics. *J.Proteome.Res.* 1, no. 1:21-26.
- Wang, G. G., K. R. Calvo, M. P. Pasillas, D. B. Sykes, H. Hacker, and M. P. Kamps. 2006. Quantitative production of macrophages or neutrophils ex vivo using conditional Hoxb8. *Nat.Methods* 3, no. 4:287-293.
- Wolters, D. A., M. P. Washburn, and J. R. Yates, III. 2001. An automated multidimensional protein identification technology for shotgun proteomics. *Anal.Chem.* 73, no. 23:5683-5690.
- Yates, J. R., III. 1998. Database searching using mass spectrometry data. *Electrophoresis* 19, no. 6:893-900.



Proteomics - Human Diseases and Protein Functions

Edited by Prof. Tsz Kwong Man

ISBN 978-953-307-832-8

Hard cover, 438 pages

Publisher InTech

Published online 10, February, 2012

Published in print edition February, 2012

Biomedical research has entered a new era of characterizing a disease or a protein on a global scale. In the post-genomic era, Proteomics now plays an increasingly important role in dissecting molecular functions of proteins and discovering biomarkers in human diseases. Mass spectrometry, two-dimensional gel electrophoresis, and high-density antibody and protein arrays are some of the most commonly used methods in the Proteomics field. This book covers four important and diverse areas of current proteomic research: Proteomic Discovery of Disease Biomarkers, Proteomic Analysis of Protein Functions, Proteomic Approaches to Dissecting Disease Processes, and Organelles and Secretome Proteomics. We believe that clinicians, students and laboratory researchers who are interested in Proteomics and its applications in the biomedical field will find this book useful and enlightening. The use of proteomic methods in studying proteins in various human diseases has become an essential part of biomedical research.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Lindsey A. Miles, Nicholas M. Andronicos, Emily I. Chen, Nagyung Baik, Hongdong Bai, Caitlin M. Parmer, Shahrzad Lighvani, Samir Nangia, William B. Kiosses, Mark P. Kamps, John R. Yates III and Robert J. Parmer (2012). Identification of the Novel Plasminogen Receptor, Plg-RKT, Proteomics - Human Diseases and Protein Functions, Prof. Tsz Kwong Man (Ed.), ISBN: 978-953-307-832-8, InTech, Available from: <http://www.intechopen.com/books/proteomics-human-diseases-and-protein-functions/identification-of-the-novel-plasminogen-receptor-plg-rkt>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen