

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Thoracic Critical Care

Seyed Mohammad Reza Hashemian and Seyed Amir Mohajerani
*Chronic Respiratory Disease Research Center (CRDRC),
NRITLD,
Masih Daneshvari Hospital,
Shahid Beheshti University of Medical Sciences, Tehran,
Iran*

1. Introduction

1.1 Ventilator associated pneumonia

Ventilator associated pneumonia (VAP) is one of the sub-types of nosocomial acquired pneumonia occur in patients admitted to ICU who are under ventilator assistant mechanical ventilation occurring more than 48 h after patients have been intubated and received mechanical ventilation. Between 250,000 and 300,000 cases per year occur in the United States solely, which is an incidence rate of 5 to 10 cases per 1,000 hospital admissions¹. The incidence of VAP increases with the duration of mechanical ventilation higher in day 10 compare to day five², and it is associated with high mortality rates (0-50%) in ICU patients, and pneumonia accounts for second cause of death in ICU patients³, although Using various scoring systems for the mortality prediction along with the guideline-based medicine have helped decrease in VAP mortality rates^{4,5}. Although mortality of viral VAP is not determined in ICU patients, the scoring systems provide an acceptable clinical index for such cases [46]. The misuse of insufficient dose or inappropriate antibiotic will lead to outgrow multi drug resistant serotypes of bacterial VAP and induce higher mortality⁶. This high mortality rate also depends on the type of underlying disease, with highest mortality attributable to VAP in patients with trauma or acute respiratory distress syndrome⁷, and the type of organism affecting the patient. Higher mortality rates have been explored in VAP caused by *Pseudomonas aeruginosa* (in patients with underlying respiratory problems) ⁸, *Acinetobacter*, and *Stenotrophomonas maltophilia* than those associated with other organisms⁹. Bacterial VAP can be due to colonization and spread of organisms from oropharynx, sinus cavities, nares, dental plaque, gastrointestinal tract, patient-to-patient contact, and the ventilator circuit to the lungs¹⁰. Essentially, each ICU should have an established protocol in place to initial empirical therapy based on previously accepted guidelines modified by local knowledge of prevalence of resistant serotypes unique to that ICU. Notably, empiric therapy should be both appropriate by using more specific antibiotics and adequate by using correct dose and good penetration to the site of infection¹¹. Duration of antibiotics are also been a point of controversy; although 8 days of therapy has been effective in non-resistant organism, but duration of antibiotic therapy for multiple drug resistant (MDR) organism such as *P aeruginosa* and *Acinetobacter* spp, is unknown¹².

On the other hand, a key point in management of MDR VAP is rapid diagnosis of VAP and providing culture and anti-biogram in detecting the responsible organism. The antibiotic duration for patients with MDR VAP remains a controversial issue. Several serum biomarkers have been applied as potential biomarker contributing to guide antibiotic use in patients with VAP caused by MDR pathogens. Previously, pro-calcitonin (PCT) has been broadly used as a marker for community acquired pneumonia (CAP) and VAP^{13,14}, however it does not incorporate into hospital acquired pneumonia(HAP). Using PCT has shortened duration of anti-microbial treatment in VAP¹⁵ in which patients with MDR VAP whose serum PCT concentrations are less than 0.5 ng/mL or decreased by 80% or more, compared with the first peak concentration, antibiotics may be terminated 3 days after initiation¹⁶. On the other hand usefulness of other bio-markers such as C reactive proteins(CRP) have yielded to conflicting data's¹⁷, probably due to acute phase reactant release in ICU patients such as IL-6 and TNF- α which stimulate CRP release to surmountable amounts.

Our current data on epidemiology, pathogenesis, clinical importance, and risk factors of viral VAP and viral pneumonia in ICU has many pitfalls due to some challenges¹⁸. First, the diagnosis of viral VAP in critically ill patients requires a high clinical suspicion combined with bedside examination, radiographic examination, and microbiologic analysis of respiratory secretions. Besides, the presence of indolent viral VAP in a critically ill patient makes diagnosis more challenging and increases the mortality rate of patients. It is important to investigate these viral markers in VAP to probe them earlier and estimate their role in increasing mortality rate. ICU patients assumed as immunocompetent are also at risk for Herpes simplex virus (HSV) and cytomegalovirus (CMV) VAP¹⁹. Although CMV reactivation assumed to increase morbidities like increased length of stay in the ICU but impact on mortality particularly in patients with low CMV-DNA plasma levels is in doubt²⁰. Currently there is increasing tendency to CMV infection among ICU patients. One simple explanation is that any bacterial colonization in ICU patients promotes the release of immunomodulatory cytokines and lead to reactivation of CMV²¹, and reactivation from the latency induces CMV infection. Because of nonspecific signs and symptoms ICU patients are rarely monitored routinely for active CMV infection, the development of active CMV infection could remain under-diagnosed in critically ill patients.

The 2009 H1N1 influenza virus pandemic has highlighted another challenge faced by intensivists in managing severe influenza A, especially for those at high risk of severe respiratory disease in ICUs⁴⁴. Thus, intensivists should consider performing rapid diagnostic tests and specific scoring for drug resistance genotyping for high risk patients especially in tertiary care centers⁴⁵.

2. Hyperglycemia and hypoglycemia in ICU

Essentially, ICU admitted patients encounter a profound hyperglycemia due to stress hormone surge, corticosteroid usage, and inhibition of insulin release of sepsis or trauma induced mediators²². Hyperglycemia could harm ICU patients by increase susceptibility to sepsis and increase mortality of critically ill patients²³. It is supposed to have a tight control of hyperglycemia in ICU patients although threshold of blood glucose is still

controversial. As a matter of fact, glucose monitoring and route of insulin injection is mainstay of hyperglycemia control in ICU patients. Nonetheless, different studies have suggested various blood glucose levels but majorly 180 mg/dl has considered the safe treatment threshold and 140-180 for target glucose level^{24,25,26,27,28}. It is widely assumed that sampling in ICU patients could also be a bottleneck in glucose tight control as catheter sampling is easy but have danger of contamination with IV fluids, whether fingerprints sampling may be inaccurate in patients with edema or anemia²⁹. On the other hand, insulin therapy induced hyperglycemia may cause severe neurologic damages while neurologic symptoms of hypoglycemia are difficult to detect in critically ill patients, but they are a real concern³⁰. Severe hypoglycemia <40 mg/dl could occur in almost high proportion of patients in intensive insulin therapy and could majorly increase mortality rates of patients^{31,32}. Besides, all the studies with target glucose concentration of 80 to 110 mg/dl showed increased rates of hypoglycemia³³. Notably, even a blood glucose target of 180 mg/dl or less resulted in lower mortality than did a target of 81 to 108 mg/dl³⁴.

3. Venous thromboembolism (VTE) and Pulmonary Embolism (PE)

Venous thromboembolism (VTE) is a common and lethal complication in critically ill patients, due to several predisposing factors such as pre-morbid conditions (e.g., trauma, major surgeries, malignancy, sepsis), invasive interventions like central venous catheterization, and prolonged immobility³⁵. The incidence of VTE is reported variously in different studies based on the population, prophylactic interventions and screening methods. Patients in intensive care unit have a higher risk of lower limb deep venous thrombosis (DVT) in comparison with other hospitalized patients which may be undiagnosed in considerable number of cases³⁶. On the other hand, VTE could remain unrecognized in the intensive care unit because of the difficulty in eliciting signs and symptoms from intubated, sedated patients. It is likely that quite large number of patients under mechanical ventilation with unexpected episodes of tachycardia, hypotension, or hypoxia may have unnoticed pulmonary embolism (PE)³⁷ which could be diagnosed based on Geneva score with an acceptable predictive accuracy in low and intermediate-probability groups³⁸. Undiagnosed or barely suspected PE may also lead to delay weaning patients from mechanical ventilation. Intensive care unit patients, who have reduced cardiopulmonary reserve, are prone to have significant complications of PE. Recent data suggest that the duration of therapy and recurrence rate is associated with persistently elevated levels of d-dimer³⁹. Long-term treatment of thrombosis with the low-molecular-weight heparin has been shown to be associated with fewer thromboembolic recurrences in ICU patients⁴⁰.

4. Management of ICU-associated agitation by dexmedetomidine

Agitated delirium is a common complication that occurs commonly in patients undergoing mechanical ventilation in ICU, and is often treated with haloperidol despite concerns about its adverse effects such as unpredictable hepatic toxicity and cardiotoxicity⁴¹. Inadequate sedation in ICU particularly in intubated patients adversely affects their

morbidity and mortality. Use of sedatives during agitation while patient under mechanical intubation precludes further extubation. Other than that agitation in intensive care may be associated with self-extubation, removal of vascular catheters, increased oxygen consumption and failure to cooperate with treatment. Drug of choice should be effective, safe, titrable, rapidly acting agent that has both sedative and analgesic activity, and could prevent anxiety and unpleasant memories. The newly introduced drug, Dexmedetomidine, a novel selective α_2 -agonist with sedative and anxiolytic properties, have showed particular utility in ICU-associated delirious agitated patients under mechanical ventilation without inducing excessive sedation, with fewer side effects than haloperidol, little interaction with other drugs and easily titrable. Studies have reported the successful use of dexmedetomidine in this context although multi-centric clinical trials may needed to establish those assumptions. Dexmedetomidine is more effective than conventional haloperidol therapy for the treatment of combined agitation and delirium in intubated patients in the ICU; besides Dexmedetomidine reduce the need of extra sedative known to cause agitation. Its administration superiority to traditional sedatives is reliably shown that has reduced ICU length of stay, hastened liberation from mechanical restraint, reduced the need for supplementary sedation, reduced QTc interval prolongation and possibly reduced the need for tracheostomy⁴². In a study in comparing dexmedetomidine and propofol in patients requiring sedation in ICU, dexmedetomidine revealed safer and protect against myocardial infarction⁴³. In addition, some studies have proposed that patients who receive a dexmedetomidine bolous have no clinically significant hypotension or increased epinephrine requirement but others not. Dexmedetomidine induce no respiratory depression and should be considered for patient failing spontaneous breathing due to agitation or anxiety. On the other hand, in dose related hypotension and bradycardia bolus dexmedetomidine is not recommended. Besides, clinicians should consider higher starting dose of dexmedetomidine if used as monotherapy.

Medication	Dose	Consideration
Dexmedetomidine	0.2-1.5 mcg/kg/hr	patients failing spontaneous breathing trials secondary to agitation
Haloperidol	2.5-5mg IV q 15 min prn/ 2.5-5 mg PO q6 hr	Caution if baseline QTc >440 msec
Aripiprazole	10-15 mg po daily	Consider when baseline QTc>440 msec
Quetiapine	50-200 mg po q12 hr	Consider if sedative properties desired
Risperidone	0.5-1 mg po q12 hr	Caution baseline QTc >440 msec
Propofol	1 mg/kg loding+1-3mg/kg/h	Consider vasodilation risk

Table 1. Pharmacologic Treatment for Delirium.

5. References

- [1] McEachern, R., and G. D. Campbell, Jr. 1998. Hospital-acquired pneumonia: epidemiology, etiology, and treatment. *Infect. Dis. Clin. N. Am.* 12:761-779.
- [2] Cook DJ, Walter SD, Cook RJ, Griffith LE, Guyatt GH, Leasa D, et al. Incidence of and risk factors for ventilator-associated pneumonia in critically ill patients. *Ann Intern Med.* Sep 15 1998;129(6):433-40.
- [3] Baker, A. M., J. W. Meredith, and E. F. Haponik. 1996. Pneumonia in intubated trauma patients. Microbiology and outcomes. *Am. J. Respir. Crit. Care Med.* 153:343-349.
- [4] Hashemian SM, Jamaati HR, Malekmohammad M, Ehteshami Afshar E, Alish O, Radmand G, Fakharian A. Assessing the Performance of Two Clinical Severity Scoring Systems in the ICU of a Tertiary Respiratory Disease Center. *Tanaffos* 2010; 9(3): 58-64.
- [5] Nooraei N, Hashemian SM, Golfam A, Saghebi A, G Radmand. Preoperative Assessment of Mechanical Ventilation Requirement after Surgical Treatment of Esophageal Cancer. *Tanaffos* 2010; 9(1): 34-41.
- [6] Niederman MS, Craven DE, Bonten MJ, et al. Guidelines for the management of adults with hospital-acquired, ventilator-associated, and healthcare-associated pneumonia. *Am J Respir Crit Care Med* 2005;171:388-416.
- [7] Melsen WG, Rovers MM, Bonten MJ. Ventilator-associated pneumonia and mortality: a systematic review of observational studies. *Crit Care Med.* Oct 2009;37(10):2709-18.
- [8] Jamaati HR, Malekmohammad M, Hashemian SM, Nayebi M, Bashar zad N. Ventilator-Associated Pneumonia: Evaluation of Etiology, Microbiology and Resistance Patterns in a Tertiary Respiratory Center. *Tanaffos* 2010; 9(1): 21-27.
- [9] Kunis KA, Puntillo KA. Ventilator-associated pneumonia in the ICU: its pathophysiology, risk factors, and prevention. *Am J Nurs.* 2003;133(8):64AA-64GG.
- [10] Staudinger T, Bojic A, Holzinger U, Meyer B, Rohwer M, Mallner F, et al. Continuous lateral rotation therapy to prevent ventilator-associated pneumonia. *Crit Care Med.* Feb 2010;38(2):486-90
- [11] Nicasio AM, Eagye KJ, Nicolau DP, et al. Pharmacodynamic-based clinical pathway for empiric antibiotic choice in patients with ventilator-associated pneumonia. *J Crit Care* 2010;25:69-77.
- [12] Chastre J, Wolff M, Fagon JY, et al. Comparison of 8 vs 15 days of antibiotic therapy for ventilator associated pneumonia in adults: a randomized trial. *JAMA* 2003;290:2588-98.
- [13] Stolz D, Smyrniotis N, Eggimann P, et al. Procalcitonin for reduced antibiotic exposure in ventilator-associated pneumonia: a randomised study. *Eur Respir J* 2009; 34:1364-75.
- [14] Kopterides P, Siempos II, Tsangaris I, et al. Procalcitonin-guided algorithms of antibiotic therapy in the intensive care unit: a systematic review and metaanalysis of randomized controlled trials. *Crit Care Med* 2010;38:2229-41.


- [15] Bouadma L, Luyt CE, Tubach F, et al. Use of procalcitonin to reduce patients' exposure to antibiotics in intensive care units (PRORATA trial): a multicentre randomised controlled trial. *Lancet* 2010;375:463-74.
- [16] Hochreiter M, Kohler T, Schweiger AM, et al. Procalcitonin to guide duration of antibiotic therapy in intensive care patients: a randomized prospective controlled trial. *Crit Care* 2009;13:R83.
- [17] Povoia P, Coelho L, Almeida E, et al. Early identification of intensive care unit-acquired infections with daily monitoring of C-reactive protein: a prospective observational study. *Crit Care* 2006;10:R63.
- [18] Moradi A, Nadji SA, Tabarsi P, Hashemian SM, Marjani M, Sigaroodi A, Mansouri D, Masjedi MR, Velayati AA. Prevalence of Oseltamivir-Resistant 2009 H1N1 Influenza Virus among Patients with Pandemic 2009 H1N1 Influenza infection in NRITLD. *Tanaffos* 2011; 10(1): 8-11
- [19] Hodson EM, Craig JC, Strippoli GF, Webster AC. Antiviral medications for preventing cytomegalovirus disease in solid organ transplant recipients. *Cochrane Database Syst Rev.* 2008 Apr 16;(2):CD003774.
- [20] Heininger A, Haeberle H, Fischer I, Beck R, Riessen R, Rohde F, Meisner C, Jahn G, Koenigsrainer A, Unertl K, Hamprecht K. Cytomegalovirus reactivation and associated outcome of critically ill patients with severe sepsis. *Crit Care.* 2011 Mar 1;15(2):R77.
- [21] Döcke WD, Prösch S, Fietze E, Kimel V, Zuckermann H, Klug C, Syrbe U, Krüger DH, von Baehr R, Volk HD: Cytomegalovirus reactivation and tumour necrosis factor. *Lancet* 1994, 343(8892):268-269.
- [22] Dungan KM, Braithwaite SS, Preiser JC. Stress hyperglycaemia. *Lancet* 2009; 373:1798-807.
- [23] van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med* 2001; 345:1359-67.
- [24] De La Rosa GDC, Donado JH, Restrepo AH, et al. Strict glycaemic control in patients hospitalised in a mixed medical and surgical intensive care unit: a randomized clinical trial. *Crit Care* 2008; 12(5):R120.
- [25] Preiser JC, Devos P, Ruiz-Santana S, et al. A prospective randomised multi-centre controlled trial on tight glucose control by intensive insulin therapy in adult intensive care units: the Glucontrol study. *Intensive Care Med* 2009;35:1738-48.
- [26] Griesdale DE, de Souza RJ, van Dam RM, et al. Intensive insulin therapy and mortality among critically ill patients: a meta-analysis including NICE-SUGAR study data. *CMAJ* 2009;180:821-7.
- [27] Marik PE, Preiser JC. Toward understanding tight glycemic control in the ICU: a systematic review and metaanalysis. *Chest* 2010;137:544-51.
- [28] Wiener RS, Wiener DC, Larson RJ. Benefits and risks of tight glucose control in critically ill adults: a meta-analysis. *JAMA* 2008;300:933-44.
- [29] Corstjens AM, Ligtenberg JJ, van der Horst IC, et al. Accuracy and feasibility of point-of-care and continuous blood glucose analysis in critically ill ICU patients. *Crit Care* 2006;10(5):R135.

- [30] Kosiborod M, Inzucchi SE, Goyal A, et al. Relationship between spontaneous and iatrogenic hypoglycemia and mortality in patients hospitalized with acute myocardial infarction. *JAMA* 2009; 301:1556-64.
- [31] Van den Berghe G, Wilmer A, Hermans G, et al. Intensive insulin therapy in the medical ICU. *N Engl J Med* 2006; 354:449-61.
- [32] Egi M, Bellomo R, Stachowski E, et al. Hypoglycemia and outcome in critically ill patients. *Mayo Clin Proc* 2010;85:217-24.
- [33] Kavanagh BP, McCowen KC. Glycemic control in ICU. *N Engl J Med* 2010;363:2540-6.
- [34] Chittock D. R., Yu-Shuo S., Blair D., Dhingra V. Intensive versus Conventional Glucose Control in Critically Ill Patients. *N Engl J Med* 2009; 360: 1283 - 97.
- [35] Tapson V. F. Acute Pulmonary Embolism. *N Engl J Med* 2008;358:1037-52.
- [36] Segal JB, Streiff MB, Hofmann LV, Thornton K, Bass EB.. Management of venous thromboembolism: a systematic review for a practice guideline. *Ann Intern Med* 2007;146:211-22.
- [37] Goldhaber SZ, Visani L, De Rosa M. Acute pulmonary embolism: clinical outcomes in the International Cooperative Pulmonary Embolism Registry (ICOPER). *Lancet* 1999;353:1386-9.
- [38] Jamaati H. R., Hashemian S. M., Malekmohammad M., Bagheri Moghadam A., Kahkouee S., Miri M., Radmand G., Masjedi M. R. Predictive Accuracy of Revised Geneva Score in the Diagnosis of Pulmonary Embolism. *Tanaffos* 2009; 8 (4): 7 - 1
- [39] Palareti G, Cosmi B, Legnani C, et al. d-Dimer testing to determine the duration of anticoagulation therapy. *N Engl J Med* 2006;355:1780-9
- [40] Lee AYY, Levine MN, Baker RI, et al. Low-molecular-weight heparin versus a coumarin for the prevention of recurrent venous thromboembolism in patients with cancer. *N Engl J Med* 2003;349:146-53.
- [41] Fraser GL, Prato BS, Riker RR, Berthiaume D, Wilkins ML: Frequency, severity, and treatment of agitation in young versus elderly patients in the ICU. *Pharmacotherapy* 2000, 20:75-82.
- [42] Dotson B, Peeters MJ: Sedation with dexmedetomidine vs lorazepam in mechanically ventilated patients. *JAMA* 2008;299:1540.
- [43] Venn RM, Grounds RM. Comparison between dexmedetomidine and propofol for sedation in the ICU: patient and clinician perception. *British Journal of Anesthesia*.2001;87(5):648-90
- [44] Moradi A, Nadji SA, Tabarsi P, Hashemian SM, Marjani M, Sigaroodi A, Mansouri D, Masjedi MR, Velayati AA. Prevalence of Oseltamivir-Resistant 2009 H1N1 Influenza Virus among Patients with Pandemic 2009 H1N1 Influenza infection in NRITLD, Tehran, Iran. *Tanaffos* 2011; 10(1): 8-11.
- [45] Pereira JM, Moreno RP, Matos R, Rhodes A, Martin-Loeches A, Cecconi M, Lisboa T, Rello J. Severity assessment tools in ICU patients with 2009 Influenza A (H1N1) pneumonia.. *Clinical Microbiology and Infection* 2010. DOI: 10.1111/j.1469-0691.2011.03736.x

- [46] Hashemian SM, Jamaati HR, Malekmohammad M, Ehteshami Afshar E, Alish O, Radmand G, Fakharian A. Assessing the Performance of Two Clinical Severity Scoring Systems in the ICU of a Tertiary Respiratory Disease Center. *Tanaffos* 2010; 9(3): 58-64.

IntechOpen

IntechOpen



Topics in Thoracic Surgery

Edited by Prof. Paulo Cardoso

ISBN 978-953-51-0010-2

Hard cover, 486 pages

Publisher InTech

Published online 15, February, 2012

Published in print edition February, 2012

Thoracic Surgery congregates topics and articles from many renowned authors around the world covering several different topics. Unlike the usual textbooks, Thoracic Surgery is a conglomerate of different topics from Pre-operative Assessment, to Pulmonary Resection for Lung Cancer, chest wall procedures, lung cancer topics featuring aspects of VATS major pulmonary resections along with traditional topics such as Pancoast tumors and recurrence patterns of stage I lung disease, hyperhidrosis, bronchiectasis, lung transplantation and much more. This Open Access format is a novel method of sharing thoracic surgical information provided by authors worldwide and it is made accessible to everyone in an expedite way and with an excellent publishing quality.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Seyed Mohammad Reza Hashemian and Seyed Amir Mohajerani (2012). Thoracic Critical Care, Topics in Thoracic Surgery, Prof. Paulo Cardoso (Ed.), ISBN: 978-953-51-0010-2, InTech, Available from: <http://www.intechopen.com/books/topics-in-thoracic-surgery/thoracic-critical-care>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen