

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Anion Channels in Osteoarthritic Chondrocytes

Elizabeth Perez-Hernandez¹, Nury Perez-Hernandez²,
Fidel de la C. Hernandez-Hernandez³ and Juan B. Kouri-Flores³

¹Hospital de Ortopedia Dr. Victorio de la Fuente Narváez –IMSS,

²Departamento de Biomedicina Molecular, Escuela Nacional de
Medicina y Homeopatía- IPN,

³Departamento de Infectómica y Patogénesis Molecular, CINVESTAV-IPN,
México

1. Introduction

Osteoarthritis (OA) is the most common form of arthritis and represents a global health problem. OA is estimated to affect 40% of the population over 70 years of age and is a major cause of pain and physical disability (Lawrence, 2008). This is a condition that predominantly involves hip, knee, spine, foot and hands. Several risk factors have been associated with the initiation and progression of OA, increasing age, sex, obesity, occupational loading, malalignment, articular trauma and crystal deposition.

OA is a chronic degenerative joint disorder characterized primarily by destruction of articular cartilage, formation of reparative fibrocartilage and subchondral bone remodelling. However, not only the cartilage and bone are affected, also the synovium and the joint-stabilizing structures such as ligaments and meniscus (Baker-LePain, 2010). Apparently, all the tissues of the joint respond to mechanical stress with the consequent loss of function and clinical deterioration.

The application of mechanical forces under physiological conditions is a preponderant factor in cartilage homeostasis. It is now well known, that environmental factors, such as compressive and tensile forces, load and shear stress have a significant influence on the chondrocyte metabolism. Also, conditions that alter the load distribution on the articular surface can induce the development of OA (Roos, 2005).

This degradative process of the cartilage in OA is a consequence of an imbalance between anabolism and catabolism of chondrocytes. Generally, chondrocytes respond with increased expression of inflammatory mediators and matrix-degrading proteinases (Kurz, 2005).

Other effect observed in chondrocytes under mechanical stimulation is the change in membrane and osmotic potential (Wright, 1992, Bush, 2001). In OA, chondrocytes undergo depolarization instead of hyperpolarization (Millward-Sadler, 2000), and the decrease of the osmotic potential has been associated with loss of volume control in early stages of OA (Stockwell, 1991, Bush, 2003). However, the sequence of mechanobiological events necessary for the maintenance of extracellular matrix (ECM) homeostasis and its involvement in the pathogenesis of OA is still poorly understood.

2. Articular cartilage structure

Articular cartilage is a specialized connective tissue that contains a single cell type, the chondrocytes, embedded in an ECM which is composed of water and macromolecules. Collagens, proteoglycans and noncollagenous proteins are the main components of matrix. The chondrocytes are highly differentiated cells and constitute 1% of the total cartilage volume (Stockwell, 1967). These cells are responsible of the production and organization of ECM. Cartilage is divided into four horizontal zones: superficial, transitional or middle, radial or deep, and calcified zones. Each of these layers has specific morphological and functional characteristics related to the metabolic activity.

In the superficial zone, chondrocytes are flattened and oriented parallel to the surface. This layer consists of a low proportion of proteoglycans, thin collagen fibers and higher water content (Weiss, 1968). Functionally, this zone is responsible of the highest resistance to compressive forces during joint movement.

The transitional or middle zone consist of rounded chondrocytes, a network of collagen fibers arranged radially and an increased proteoglycan content.

In the deep zone, the chondrocytes are grouped in perpendicular columns to the articular surface. This layer contains the highest proportion of aggrecan and long collagen fibers, approximately of 55 μm across (Minns, 1977).

The calcified zone links and anchors the articular cartilage to subchondral bone through collagen fibers arranged perpendicularly. The hypertrophic chondrocytes are scarce and located in uncalcified lacunae.

Also, the matrix is subdivided into the pericellular, territorial and interterritorial regions which are arranged around the chondrocytes. The pericellular matrix is composed for sulfated proteoglycans and glycoproteins. They provide protection to chondrocytes when exposed to chondrocyte load and maintain water homeostasis. Together, the chondrocyte and pericellular matrix constitute the chondron (Poole, 1997). Adjacent to this region, the territorial matrix contains a dense meshwork of collagen fibers that provide mechanical protection to the chondrocytes. The concentration of proteoglycans rich in chondroitin sulfate is higher in this region. In the interterritorial region predominates proteoglycans rich in keratan sulfate and the collagen fibers of largest diameter (Stockwell, 1990).

The biochemical properties of cartilage are dependent of the integrity of the matrix. In the ECM of mature mammalian articular cartilage, the collagen represents 50-60% dry weight. The network consists of collagen IX (1%), collagen XI (3%) and collagen II ($\geq 90\%$) (Eyre, 1987). Type II collagen, a fibrillar protein, is composed of three identical α -1 chains in form triple helix and constitutes the basic structure of cartilage. Type XI collagen is probably copolymerized and type IX collagen is covalently linked to the type II collagen fibrils (Mendler, 1989). Among other types of collagen found in cartilage, type X collagen is described in the calcified zone (Gannon, 1991), type VI collagen promotes the chondrocyte-matrix attachment (Wu, 1987) and type III collagen is copolymerized and linked to collagen II (Wu, 1996). Generally, these collagen fibers provided to the cartilage the tensile stiffness and strength.

The second largest component of the ECM are the proteoglycans, which constitute 5 to 10% of the tissue wet weight. These are molecules consisting of glycosaminoglycans (GAGs) chains bound to a protein core. Also, the aggrecan are composed of proteoglycan aggregates in association with hyaluronic acid (HA) and link protein (Hardingham, 1974). The keratan sulfate and chondroitin sulfate are the main types of GAGs. The anionic groups of aggrecan

attract cations such as Na^+ , this allows an osmotic imbalance and the accumulation of fluid in the ECM, which is critical for the biomechanical properties of cartilage. Collagen fibrils interact with aggrecan monomers in the keratan sulfate-rich regions (Hedlund, 1999). Other proteoglycans include decorin, biglycan, lumican and fibromodulin, consisting of small leucine-rich repeat. They bind to fibrillar collagen via its core protein regulating the fibril diameter and fibril-fibril interaction in the ECM.

In addition, there are nocollagenous proteins such as fibronectin and tenascin favoring chondrocyte-matrix interaction.

3. Chondrocytes mechanotransduction

The articular cartilage is continuously exposed to mechanical stress which significantly affects the response of chondrocytes to their environment (Grodzinsky, 2000). Under conditions of continuous compression, the chondrocytes undergo changes in potential membrane (Wright, 1992), matrix water content, ion concentrations and pH (Mobasher, 1998; Mow, 1999). The process of converting physical forces into biochemical signals and the subsequent transduction into cellular responses is referred to as mechanotransduction (Huang, 2004).

Miscellaneous "mechanoreceptors" including ion channels and cell adhesion molecules, such as integrins have been described in chondrocytes. Thus far, $\alpha1\beta1$ integrin, a receptor for collagen, $\alpha5\beta1$ integrin, a receptor for fibronectin and $\alphaV\beta5$ integrin, a receptor for vitronectin have been implicated in the regulation of chondrocyte behaviour (Ostergaard, 1998; Loeser, 1995). Voltage-gated Na^+ and K^+ channels (Sugimoto, 1996), epithelial sodium channels (ENaC) (Mobasher, 1999) and N-/L-type voltage-gated Ca^{2+} channels (Wang, 2000) have been considered as potential mechanosensitive ion channels.

3.1 Membrane potential

The negative resting membrane potential (RMP) in most cells, is the result from activity of K^+ (Wilson, 2004) and Cl^- channels (CIC) (Tsuga, 2002; Funabashi, 2010). In chondrocytes, the RMP ranging from -10.6 mV to -46 mV (Wright, 1992; Clark, 2010; Lewis 2011), and those values are determined by the influx and efflux of cations and anions in the cell membrane. Consequently, the stability of the ECM is linked to the ionic changes and the RMP in chondrocytes. Several studies related the use of ionic blockers such as lidocaine and verapamil, showed a decrease in the synthesis of GAGs (Wu, 2000; Mouw, 2007), inhibition of chondrocytes proliferation (Wohlrab, 2002) and induction of apoptosis (Wohlrab, 2005). The permeability of ion channels is regulated by chondrocyte channelome (Barret-Jolley, 2010).

The ion channels so far identified in chondrocytes are summarized in Table 1.

The membrane potential in articular chondrocytes is primarily related to the Cl^- conductance greater than K^+ conductance (Tsuga, 2002; Funabashi, 2010). More recently, it was showed the involvement of TRPV5 (gadolinium-sensitive cation channels) in the relatively positive RMP of the chondrocytes. Furthermore, has been suggested that the positive membrane potential is due to TRVP5 higher than that generated by potassium ions, which would allow efflux of potassium ions into the cell limiting the increase in cell volume in situations of reduced osmolarity (Lewis, 2011).

Ion channels	Reference
Voltage-gated K ⁺ channels	Walsh, 1992 Sugimoto, 1996 Wilson, 2004 Mobasheri, 2005 Ponce, 2006 Clark, 2010
Voltage-gated Na ⁺ channels	Sugimoto, 1996 Ramage, 2008
Voltage-gated Ca ²⁺ channels	Shakibaei, 2003 Sanchez, 2004 Mancilla, 2007 Xu, 2009
Voltage-activated H ⁺ channels	Sanchez, 2006
Cl ⁻ channels (ClC)	Sugimoto, 1996 Tsuga, 2002 Isoya, 2009 Okumura, 2009 Funabashi, 2010 Perez, 2010
ATP-dependent K ⁺ channels	Mobasheri, 2007
Ca ²⁺ -dependent K ⁺ channels	Grandolfo, 1992 Long, 1994 Martina, 1997 Mozrzymas, 1997 Mobasheri, 2010
Transient receptor potential (TRP) channel	Sanchez, 2003 Phan, 2009 Lewis, 2010
Epithelial sodium channels (ENaC)	Trujillo, 1999 Shakibaei, 2003 Lewis, 2008

Table 1. Ion channels described in chondrocytes

Therefore, it has been proposed that the ClCs functions in chondrocytes could be involved in setting of the membrane potential or anionic osmolyte channels (Barret-Jolley, 2010).

3.2 Mechanotransduction in osteoarthritic chondrocytes

In vivo, the extracellular environment surrounding the chondrocytes is negatively charged and provides a high osmolarity, from 480 mOsm to 550 mOsm under load (Urban, 1993; 1994; Xu, 2010). Under physiological conditions, chondrocytes are able to regulate their volume through osmotic pressure cycles (Bush, 2001). However, in OA, the chondrocytes have decreased osmotic potential and increased water content (Stockwell, 1991). Furthermore, in osteoarthritic chondrocytes described a high recovery of the increased volume, which promotes the progression of the disease (Jones, 1999; Bush, 2005).

In this matter, in the superficial and middle zones of normal cartilage and the zone of fibrillated osteoarthritic cartilage has been demonstrated the expression of ATP-dependent K⁺ channels (Mobasher, 2007). Under normal conditions, mechanical stimulation produces hyperpolarization of the membrane of chondrocytes, but in OA the depolarization is induced. According to studies performed with sodium channel blockers, this response could be due to the involvement of Ca²⁺-dependent K⁺ channels (stretch-activated channels) in the process of chondrocyte mechanotransduction (Wright, 1996).

In osteoarthritic chondrocytes has been suggested that the response of membrane depolarization is due to the autocrine / paracrine activity of soluble factors. *In vitro*, mechanical stimulation of chondrocytes at 0.33 Hz induces membrane hyperpolarization due to IL-4 (Millward-Sadler, 1999) with the consequent increase in aggrecan mRNA levels. Interestingly, in osteoarthritic chondrocytes, membrane depolarization is induced by proinflammatory cytokine IL-1 (Salter, 2002). Altered mechanotransduction in OA may contribute to the response of the chondrocyte favoring ECM degradation and disease progression.

Moreover, osmotic fluctuations, in addition to the volume change in chondrocytes (Bush, 2005; Kerrigan, 2006; Lewis, 2011) involve the filamentous actin restructuring (Chao, 2006; Erickson, 2003). Similarly, the osmotic stimulation of chondrocytes increases cytosolic calcium concentrations with the consequent regulation of metabolism, cell volume, and gene expression (Liu, 2006; Hardingham, 1999). In this case, a possible candidate in the chondrocyte mechanotransduction is the transient receptor potential vanilloid 4 channel (TRPV4), a Ca²⁺-permeable, nonspecific cation channel (Liedtke, 2000; Phan, 2009).

4. Anion channels

The anion channels are a membranal class of porin ion channel that allow the passive diffusion of negatively charged ions along their electrochemical gradient. These channels allow the flow of monovalent anions such as I⁻, NO₃⁻, Br⁻, and Cl⁻, however, these are known as CIC (Fahlke, 2001). The CIC constitute a large family of Cl⁻ selective channels (Jentsch, 2002). In humans, have been described nine isoforms of CICs. The first branch of this gene family encodes plasma membrane channels (CIC-1, -2, -Ka and -Kb) and others two branches (CIC-3, -4 and -5; CIC-6/-7) which encode for proteins found on intracellular vesicles (Jentsch, 2005).

In the plasma membrane, the main functions of CICs are regulation cell volume and ionic homeostasis, transepithelial transport and excitability. Channels in intracellular organelles facilitate the exchange of anionic substrates between the biosynthetic compartments involved in acidification of vesicles in the endosomal pathway (Jentsch, 2002; 2007).

Different diseases have been associated with anionic channels. The loss of CIC-5 is related to Dent's disease (Lloyd, 1996), while the disruption of CIC-3, -6, or -7 in mice promotes the development of neurodegenerative disorders in the central nervous system (Kasper, 2005; Poët, 2006), and CIC-7 mutations have been related with osteopetrosis and lysosomal storage disease (Kasper, 2005; Poët, 2006).

4.1 CIC and chondrocytes proliferation

OA is a complex morphology reflect the severity of damage to articular structures. The histopathological parameters in OA include cartilage degradation with formation of

fibrocartilage, fissures, denudation with exposure, repair and remodeling of subchondral bone. In addition, chondrocytes suffer proliferation and apoptosis (Pritzker, 2006) (Fig. 1).




Fig. 1. OA cartilage morphology. A-B fibrillation zone, C apoptotic chondrocytes, D vertical fissures, E-F "clones" or chondrocytes aggregates. Hematoxylin and eosin staining. 10x, 20x and 40x.

This is a condition in which it was reported that the proliferative activity of chondrocytes is low compared with its absence in normal cartilage (Rothwell, 1973; Hulth, 1972). Apparently, the osteoarthritic chondrocytes have a greater access to proliferation factors due to alterations in the ECM (Lee, 1993). It is also possible that aggregates of chondrocytes or clones are a manifestation of the proliferative activity (Horton, 2005).

Moreover, apoptotic cell death is a mechanism widely described in association with OA (Blanco, 1998; Hashimoto, 1998a, 1998b; Kim, 2000; Kouri, 2000). The classic morphologic appearance of this process is characterized by fragmentation and nuclear condensation, and formation of apoptotic bodies (Aigner, 2002). In general, the response of osteoarthritic chondrocytes to the metabolic imbalance (Sandell, 2001), phenotypic modulation (Aigner, 1997) and cell death induces proliferation that compensate the cell loss and the demand of synthetic activity (Aigner, 2001; Rothwell, 1973; Hulth, 1972).

Processes such as proliferation and cell death have been linked to the involvement of ion channels. In this regard, it has been described the association between the activity of K⁺ channels and the proliferation of Schwann cells and B lymphocytes (Deutsch, 1990), keratinocytes (Harmon, 1993), melanoma (Nilius, 1994), neuroblastoma and astrocytoma cells (Lee YS, 1993). It also has been associated intracellular free Ca²⁺ concentration and cell proliferation (Wohlrab, 1998).

In chondrocytes, there are changes in the membrane potential of human chondrocytes in response to modulators of ion channels such as tetraethylammonium (TEA), 4-aminopyridine (4-AP), 4',4'-diisothiocyanato-stilbene-2,2'-disulfonic acid (DIDS), 4-acetamido-4'-isothiocyano-2,2'-disulfonic acid (SITS), verapamil and lidocaine. About this, it was reported an increase in DNA synthesis with lidocaine and 4-AP after 12 days in chondrocytes culture (Wohlrab, 2002). Moreover, the exposure of chondrocytes to high concentrations of SITS induced necrosis while that the use of 4-AP caused cytotoxic effects and suppression of proliferation on the same system (Wohlrab, 2004).

4.2 CLIC proteins

The chloride intracellular channel (CLIC) proteins belong to the glutathione S-transferase (GST) superfamily and these are group of proteins with possible role of anion channels. However, they differ from classical GSTs because they contain an active site with cysteine residue, reactive for the protein itself and not through a thiol group (Littler, 2010). The CLICs are proteins highly conserved in vertebrates and these are referred to as CLIC1–CLIC6.

These proteins are present in soluble and membrane-inserted form. CLICs have been reported in membranous organelles, cytoplasmic and vesicular compartments and the nucleus (Valenzuela, 1997; Duncan, 1997; Chuang, 1999). The functions of CLICs include pH-dependent ion channel activity (Tulk, 2002; Littler, 2005) and enzymatic which likely to involve a glutathione (GSH) related cofactor (Littler, 2010). They could also be involved in maintaining the structure of intracellular organelles and their interaction with cytoskeleton proteins (Singh, 2007).

Functions of CLICs reported in the musculoskeletal system involve these proteins in the acidification processes in bone resorption (Edwards, 2006; Schlesinger, 1997). Despite their involvement in different systems with activities of CLIC, has not been possible to attribute the channel protein function permanently.

CLICs proteins are implicated in cell cycle regulation, cell differentiation, and apoptosis. Specifically, CLIC4 has been linked to apoptosis and differentiation of fibroblasts into myofibroblasts (Fernandez-Salas, 2002; Ronnov-Jessen, 2002). In addition, it was reported that CLIC3 interacts with ERK7, a mitogen-activated protein kinase, allowing the regulation of the phosphatases or kinases (Qian, 1999).

4.3 CLIC6 protein

CLIC6 consist of 704 aminoacids with decapeptide repeats (Friedli, 2003) and and reportedly bind to the dopamine D(2)-like receptors (Griffon, 2003). It is also suggested that CLIC6 could be involved in the regulation of water and secretion of hormones (Nishizawa, 2000).

In chondrocytes, recently, we reported the identification of CLIC6 protein from proteomic analysis of rat normal articular cartilage (Perez, 2010). In human cartilage obtained from total knee arthroplasty, we found immunoreactivity for CLIC6 protein largely restricted to the aggregates of chondrocytes (clones or clusters) in the superficial zone (Fig. 2).




Fig. 2. CLIC6 immunostaining of NoAC and OAC. A-B Surface zone (NoAC), C-E Chondrocytes clusters (OAC). The anti-CLIC6 antibody was coupled to FITC (fluorescein-5-isothiocyanate) and the nucleus were counterstained with propidium iodide.

Immunolabelling was arranged in a coarse granular pattern located in the cytoplasm of most chondrocytes. Comparatively, chondrocytes of non osteoarthritic human articular cartilage (NoAC) showed scarce immunoreactivity predominantly in cells of superficial zones (unpublished data).

Although controversial, the formation of cell clusters in osteoarthritic cartilage (OAC) has been considered an event of repair and regeneration (Horton, 2005). However, electrophysiological and molecular investigations are required to define role of the CLIC6 protein on healthy and osteoarthritic chondrocytes.

5. Conclusion

The pathogenesis of OA includes homeostatic alterations that induce imbalance in the anabolic and catabolic processes. These have been associated with changes in the viability and chondrocytes proliferation. Previous studies have showed a low proliferative activity of OAC. However, the arrangement of the chondrocytes in groups (clones), a hallmark of OA, could possibly be considered a sign of proliferation.

CLIC6 protein expression in normal articular cartilage of rat, and immunolocalization in chondrocytes clusters removed from patients with OA, could suggest the involvement of this anion channel in the pathophysiological processes of disease.

6. Acknowledgment

We wish to thank Jose C. Luna Muñoz, PhD, for image capture support at the Departamento de Fisiología y Unidad de Microscopia Confocal, CINVESTAV-IPN, México.

7. References

- Aigner, T.; Dudhia, J. (1997). Phenotypic modulation of chondrocytes as a potential therapeutic target in osteoarthritis: a hypothesis. *Ann Rheum Dis*. Vol. 56, pp. 287-291.
- Aigner, T.; Hemmel, M.; Neureiter, D.; Gebhard, P.M.; Zeiler, G.; Kirchner, T.; McKenna, L. (2001). Apoptotic cell death is not a widespread phenomenon in normal aging and osteoarthritis human articular knee cartilage: a study of proliferation, programmed cell death (apoptosis), and viability of chondrocytes in normal and osteoarthritic human knee cartilage. *Arthritis Rheum*. Vol. 44, pp. 1304-1312.
- Aigner, T.; Kim, H.A. (2002). Apoptosis and cellular vitality: issues in osteoarthritic cartilage degeneration. *Arthritis Rheum*. Vol. 46, pp. 1986-1996.
- Baker-LePain, J.C.; Lane, N.E. (2010). Relationship between joint shape and the development of osteoarthritis. *Curr Opin Rheumatol* Vol. 22, pp. 538-543.
- Barrett-Jolley, R.; Lewis, R.; Fallman, R.; Mobasher, A. (2010). The emerging chondrocyte channelome. *Front Physiol*. Vol. 1, pp. 135.
- Blanco, F.J.; Guitian, R.; Vazquez-Martul, E.; de Toro, F.J.; Galdo, F. (1998). Osteoarthritis chondrocytes die by apoptosis. A possible pathway for osteoarthritis pathology. *Arthritis Rheum* Vol. 41, pp. 284-289.
- Bush, P.G.; Hall, A.C. (2001). The osmotic sensitivity of isolated and in situ bovine articular chondrocytes. *J Orthop Res* Vol. 19, pp. 768-778.
- Bush, P.G.; Hall, A.C. (2003). The volume and morphology of chondrocytes within non-degenerate and degenerate human articular cartilage. *Osteoarthritis Cartilage* Vol. 11, pp. 242-251.
- Bush, P.G.; Hodkinson, P.D.; Hamilton, G.L.; Hall, A.C. (2005). Viability and volume of in situ bovine articular chondrocytes--changes following a single impact and effects of medium osmolarity. *Osteoarthritis Cartilage* Vol. 13, pp. 54-65.
- Chao, P.H.; West, A.C.; Hung, C.T. (2006). Chondrocyte intracellular calcium, cytoskeletal organization, and gene expression responses to dynamic osmotic loading. *Am J Physiol Cell Physiol* Vol. 291, pp. C718-725.
- Chuang, J.Z.; Milner, T.A.; Zhu, M.; Sung, C.H. (1999). A 29 kDa intracellular chloride channel p64H1 is associated with large dense-core vesicles in rat hippocampal neurons. *J Neurosci* Vol. 19, pp. 2919-2928.
- Clark, R.B.; Hatano, N.; Kondo, C.; Belke, D.D.; Brown, B.S.; Kumar, S.; Votta, B.J.; Giles, W.R. (2010). Voltage-gated k⁺ currents in mouse articular chondrocytes regulate membrane potential. *Channels* Vol. 4, pp. 179-191.
- Deutsch, C. (1990). K⁺ channels and mitogenesis, *Prog Clin Biol Res*. Vol. 334, pp. 251-271.
- Duncan, R.R.; Westwood, P.K.; Boyd, A.; Ashley, R.H. (1997). Rat brain p64H1, expression of a new member of the p64 chloride channel protein family in endoplasmic reticulum. *J Biol Chem* Vol. 272, pp. 23880-23886.

- Edwards, J.C.; Cohen, C.; Xu, W.; Schlesinger, P.H. (2006). c-Src control of chloride channel support for osteoclast HCl transport and bone resorption. *J Biol Chem* Vol. 281, pp. 28011–28022.
- Erickson, G.R.; Northrup, D.L.; Guilak, F. (2003). Hypo-osmotic stress induces calcium-dependent actin reorganization in articular chondrocytes. *Osteoarthritis Cartilage* Vol. 11, pp. 187–197.
- Eyre, D.R.; Wu, J.J. (1987). Type XI or 1 α 2 α 3 α collagen. In *Structure and Function of Collagen Types*. R. Mayne, R.E. Burgeson (Ed), New York: Academic Press, pp. 261–281.
- Fahlke, C. (2001) Ion permeation and selectivity in ClC-type chloride channels. *Am J Physiol Renal Physiol*. Vol. 280, pp. F748–F757.
- Fernández-Salas, E.; Suh, K.S.; Speransky, V.V.; Bowers, W.L.; Levy, J.M.; Adams, T.; Pathak, K.R.; Edwards, L.E.; Hayes, D.D.; Cheng, C.; Steven, A.C.; Weinberg, W.C.; Yuspa, S.H. (2002). mtCLIC/CLIC4, an organellar chloride channel protein, is increased by DNA damage and participates in the apoptotic response to p53. *Mol Cell Biol* Vol. 22, pp. 3610–3620.
- Friedli, M.; Guipponi, M.; Bertrand, S.; Bertrand, D.; Neerman-Arbez, M.; Scott, H.S.; Antonarakis, S.E.; Reymond, A. (2003). Identification of a novel member of the CLIC family, CLIC6, mapping to 21q22.12. *Gene*. Vol. 320, pp. 31–40.
- Funabashi, K.; Fujii, M.; Yamamura, H.; Ohya, S.; Imaizumi, Y. (2010). Contribution of chloride channel conductance to the regulation of resting membrane potential in chondrocytes. *J Pharmacol Sci*. Vol. 113, pp. 94–99.
- Gannon, J.M.; Walker, G.; Fischer, M.; Carpenter, R.; Thompson, R.C. Jr.; Oegema, T.R. Jr. (1991). Localization of type X collagen in canine growth plate and adult canine articular cartilage. *J Orthop Res* Vol. 9, pp. 485–494.
- Grandolfo, M.; D'Andrea, P.; Martina, M.; Ruzzier, F.; Vittur, F. (1992). Calcium-activated potassium channels in chondrocytes. *Biochem Biophys Res Commun*. Vol. 182, pp. 1429–1434.
- Griffon, N., Jeanneteau, F., Prieur, F., Diaz, J. and Sokoloff, P. (2003) CLIC6, a member of the intracellular chloride channel family, interacts with dopamine D(2)-like receptors. *Brain Res Mol Brain Res*. Vol. 117, pp. 47–57.
- Grodzinsky AJ, Levenston ME, Jin M, Frank EH. (2000). Cartilage tissue remodeling in response to mechanical forces. *Annu Rev Biomed Eng*. Vol. 2, pp. 691–713.
- Hardingham, T.E.; Muir, H. (1974). Hyaluronic acid in cartilage and proteoglycan aggregation. *Biochem J*. Vol. 139, pp. 565–581.
- Hardingham, G.E.; Bading, H. (1999). Calcium as a versatile second messenger in the control of gene expression. *Microsc Res Tech* Vol. 46, pp. 348–355.
- Harmon SC, Lutz D, Ducote J. (1993). Potassium channel openers stimulate DNA synthesis in mouse epidermal keratinocyte and whole hair follicle cultures. *Skin Pharmacol*. Vol. 6, pp. 170–178.
- Hashimoto, S.; Ochs, R.L.; Komiya, S.; Lotz, M. (1998a). Linkage of chondrocyte apoptosis and cartilage degradation in human osteoarthritis. *Arthritis Rheum* Vol. 41, pp. 1632–1638.
- Hashimoto, S.; Takahashi, K.; Amiel, D.; Coutts, R.D.; Lotz, M. (1998b). Chondrocyte apoptosis and nitric oxide production during experimentally induced osteoarthritis. *Arthritis Rheum* Vol. 41, pp. 1266–1274.

- Hedlund, H.; Hedbom, E.; Heinegard, D.; Mengarelli-Widholm, S.; Reinholt, F.P.; Svensson, O. (1999). Association of the aggrecan keratan sulfate-rich region with collagen in bovine articular cartilage. *J Biol Chem* Vol. 274, pp. 5777–5781.
- Horton, W.E. Jr.; Yagi, R.; Lavery, D.; Weiner, S. (2005). Overview of studies comparing human normal cartilage with minimal and advanced osteoarthritic cartilage. *Clin Exp Rheumatol* Vol. 23, pp. 103–112.
- Huang, H.; Kamm, R.D.; Lee, R.T. (2004). Cell mechanics and mechanotransduction: pathways, probes, and physiology. *Am J Physiol Cell Physiol*. Vol. 287, pp. C1-11.
- Hulth, A.; Lindberg, L.; Telhag, H. (1972). Mitosis in human osteoarthritic cartilage. *Clin Orthop Rel Res* Vol. 84, pp. 197–199.
- Isoya, E.; Toyoda, F.; Imai, S.; Okumura, N.; Kumagai, K.; Omatsu-Kanbe, M.; Kubo, M.; Matsuura, H.; Matsusue, Y. (2009). Swelling-activated Cl⁻ current in isolated rabbit articular chondrocytes: inhibition by arachidonic acid. *J Pharmacol Sci*. Vol. 109, pp. 293–304.
- Jentsch, T.J.; Stein, V.; Weinreich, F.; Zdebik, A.A. (2002). Molecular structure and physiological function of chloride channels. *Physiol Rev*. Vol. 82, pp. 503–568.
- Jentsch, T.J.; Poët, M.; Fuhrmann, J.C.; Zdebik, A.A. (2005). Physiological functions of CLC Cl⁻-channels gleaned from human genetic disease and mouse models. *Annu Rev Physiol*. Vol. 67, pp. 779–807.
- Jentsch, T.J. (2007) Chloride and the endosomal-lysosomal pathway: emerging roles of CLC chloride transporters. *J Physiol*. Vol. 578, pp. 633–40.
- Jones, W.R.; Ting-Beall, H.P.; Lee, G.M.; Kelley, S.S.; Hochmuth, R.M.; Guilak, F. (1999). Alterations in the young's modulus and volumetric properties of chondrocytes isolated from normal and osteoarthritic human cartilage. *J Biomech* Vol. 32, pp. 119–127.
- Kasper, D.; Planells-Cases, R.; Fuhrmann, J.C.; Scheel, O.; Zeitz, O.; Ruether, K.; Schmitt, A.; Poët, M.; Steinfeld, R.; Schweizer, M.; Kornak, U.; Jentsch, T.J. (2005). Loss of the chloride channel CLC-7 leads to lysosomal storage disease and neurodegeneration. *EMBO J*. Vol. 24, pp. 1079–1091.
- Kerrigan, M.J.; Hook, C.S.; Qusous, A.; Hall, A.C. (2006). Regulatory volume increase (RVI) by in situ and isolated bovine articular chondrocytes. *J Cell Physiol* Vol. 209, pp. 481–492.
- Kim, H.A.; Lee, Y.J.; Seong, S.C.; Choe, K.W.; Song, Y.W. (2000). Apoptotic chondrocyte death in human osteoarthritis. *J Rheum* Vol. 27, pp. 455–462.
- Kouri, J.B.; Aguilera, J.M.; Reyes, J.; Lozoya, K.A.; Gonzalez S. (2000). Apoptotic chondrocytes from osteoarthrotic human articular cartilage and abnormal calcification of subchondral bone. *J Rheum* Vol. 27, pp. 1005–1019.
- Kurz, B.; Lemke, A.K.; Fay, J.; Pufe, T.; Grodzinsky, A.J.; Schünke, M. (2005). Pathomechanisms of cartilage destruction by mechanical injury. *Ann Anat* Vol. 187, pp. 473–485.
- Lawrence, R.C.; Felson, D.T.; Helmick, C.G.; Arnold, L.M.; Choi, H.; Deyo, R.A.; Gabriel, S.; Hirsch, R.; Hochberg, M.C.; Hunder, G.G.; Jordan, J.M.; Katz, J.N.; Kremers, H.M.; Wolfe, F. (2008). Estimates of the prevalence of arthritis and other rheumatic conditions in the United States. Part II. *Arthritis Rheum*. Vol. 58, pp. 26–35.
- Lee, D.A.; Bentley, G.; Archer, C.W. (1993). The control of cell division in articular chondrocytes. *Osteoarthritis Cartilage* Vol. 1, pp. 137–146.


- Lee, Y.S.; Sayeed, M.M.; Wurster, R.D. (1993). Inhibition of cell growth by K1 channel modulators is due to interference with agonist induced Ca²⁺ release. *Cell Signal* Vol. 5, pp. 803–809.
- Lewis, R., Mobasheri, A., and Barrett-Jolley, R. (2008). Electrophysiological identification of epithelial sodium channels in canine articular chondrocytes. *Proc Physiol Soc.* Vol. 11, pp. C47.
- Lewis, R.; Purves, G.; Crossley, J.; Barrett-Jolley, R. (2010). Modelling the membrane potential dependence on non-specific cation channels in canine articular chondrocytes. *Biophys J.* Vol. 98. (Suppl. 1), pp. 340A.
- Lewis, R.; Asplin, K.E.; Bruce, G.; Dart, C.; Mobasheri, A.; Barrett-Jolley R. (2011). The role of the membrane potential in chondrocyte volume regulation. *J Cell Physiol.* Feb 15.
- Liedtke W, Choe Y, Martí-Renom MA, Bell AM, Denis CS, Sali A, Hudspeth AJ, Friedman JM, Heller S. (2000). Vanilloid receptor-related osmotically activated channel (VR-OAC), a candidate vertebrate osmoreceptor. *Cell* Vol. 103, pp. 525–535.
- Littler, D.R.; Assaad, N.N.; Harrop, S.J.; Brown, L.J.; Pankhurst, G.J.; Luciani, P.; Aguilar, M.I.; Mazzanti, M.; Berryman, M.A.; Breit, S.N.; Curmi, P.M. (2005). Crystal structure of the soluble form of the redox-regulated chloride ion channel protein CLIC4. *FEBS J.* Vol. 272, pp. 4996–5007.
- Littler, D.R.; Harrop, S.J.; Goodchild, S.C.; Phang, J.M.; Mynott, A.V.; Jiang, L.; Valenzuela, S.M.; Mazzanti, M.; Brown, L.J.; Breit, S.N.; Curmi, P.M. (2010). The enigma of the CLIC proteins: Ion channels, redox proteins, enzymes, scaffolding proteins? *FEBS Lett.* Vol. 584, pp. 2093–2101
- Liu, X.; Bandyopadhyay, B.C.; Nakamoto, T.; Singh, B.; Liedtke, W.; Melvin, J.E.; Ambudkar I. (2006). A role for AQP5 in activation of TRPV4 by hypotonicity: concerted involvement of AQP5 and TRPV4 in regulation of cell volume recovery. *J Biol Chem* Vol. 281, pp. 15485–15495.
- Lloyd, S.E.; Pearce, S.H.; Fisher, S.E.; Steinmeyer, K.; Schwappach, B.; Scheinman, S.J.; Harding, B.; Bolino, A.; Devoto, M.; Goodyer, P.; Rigden, S.P.; Wrong, O.; Jentsch, T.J.; Craig, I.W.; Thakker, R.V. (1996). A common molecular basis for three inherited kidney stone diseases. *Nature* Vol. 379, pp. 445–449.
- Loeser, R.F.; Carlson, C.S.; McGee, M.P. (1995). Expression of beta 1 integrins by cultured articular chondrocytes and in osteoarthritic cartilage. *Exp Cell Res* Vol. 217, pp. 248–257.
- Long, K.J.; Walsh, K.B. (1994). Calcium-activated potassium channel in growth-plate chondrocytes: regulation by protein-kinase A. *Biochem Biophys Res Commun.* Vol. 201, pp. 776–781.
- Mancilla, E.E.; Galindo, M.; Fertilio, B.; Herrera, M.; Salas, K.; Gatica, H.; Goecke, A. (2007). L-type calcium channels in growth plate chondrocytes participate in endochondral ossification. *J Cell Biochem.* Vol. 101, pp. 389–398.
- Martina, M.; Mozrzymas, J.W.; Vittur, F. (1997). Membrane stretch activates a potassium channel in pig articular chondrocytes. *Biochim Biophys Acta.* Vol. 1329, pp. 205–210.
- Mendler, M.; Eich-Bender, S.V.; Vaughan, L.; Winterhalter, K.M.; Bruckner P. (1989). Cartilage contains mixed fibrils of collagen types II, IX, and XI. *J Cell Biol* Vol. 108, pp. 191–197.
- Millward-Sadler, S.J.; Wright, M.O.; Lee, H.S.; Nishida, K.; Caldwell, H.; Nuki, G.; Salter, D.M. (1999). Integrin-regulated secretion of interleukin 4: a novel pathway of

- mechanotransduction in human articular chondrocytes. *J Cell Biol* Vol. 145, pp. 183-189.
- Millward-Sadler, S.J.; Wright, M.O.; Lee, H.S.; Caldwell, H.; Nuki, G.; Salter, D.M. (2000). Altered electrophysiological responses to mechanical stimulation and abnormal signaling through alpha5beta1 integrin in chondrocytes from osteoarthritic cartilage. *Osteoarthritis Cartilage* Vol. 8, pp. 272-278.
- Minns, R.J.; Steven, F.S. (1977). The collagen fibril organization in human articular cartilage. *J Anat* Vol. 123, pp. 437- 457.
- Mobasheri, A.; Mobasheri, R.; Francis, M.J.; Trujillo, E.; Alvarez de la Rosa, D.; Martín-Vasallo, P. (1998). Ion transport in chondrocytes: membrane transporters involved in intracellular ion homeostasis and the regulation of cell volume, free $[Ca^{2+}]$ and pH. *Histol Histopathol.* Vol. 13, pp. 893-910.
- Mobasheri, A.; Martín-Vasallo, P. (1999). Epithelial sodium channels in skeletal cells; a role in mechanotransduction? *Cell Biol Int.* Vol. 23, pp. 237-240.
- Mobasheri, A.; Gent, T.C.; Womack, M.D.; Carter, S.D.; Clegg, P.D.; Barrett-Jolley, R. (2005). Quantitative analysis of voltage-gated potassium currents from primary equine (*Equus caballus*) and elephant (*Loxodonta africana*) articular chondrocytes. *Am J Physiol Regul Integr Comp Physiol.* Vol. 289, pp. R172-R180.
- Mobasheri, A.; Gent, T.C.; Nash, A.I.; Womack, M.D.; Moskaluk, C.A.; Barrett-Jolley, R. (2007). Evidence for functional ATP-sensitive (K(ATP)) potassium channels in human and equine articular chondrocytes. *Osteoarthritis Cartilage.* Vol. 15, pp. 1-8.
- Mobasheri, A.; Lewis, R.; Maxwell, J.E.; Hill, C.; Womack, M.; Barrett-Jolley, R. (2010). Characterization of a stretch-activated potassium channel in chondrocytes. *J Cell Physiol.* Vol. 223, pp. 511-518.
- Mouw, J.K.; Imler, S.M.; Levenston, M.E. (2007). Ion-channel regulation of chondrocyte matrix synthesis in 3D culture under static and dynamic compression. *Biomech Model Mechanobiol* Vol. 6, pp. 33-41.
- Mow, V.C.; Wang, C.C.; Hung, C.T. (1999). The extracellular matrix, interstitial fluid and ions as a mechanical signal transducer in articular cartilage. *Osteoarthritis Cartilage.* Vol. 7, pp. 41-58.
- Mozzrymas, J.W.; Martina, M.; Ruzzier, F. (1997). A large-conductance voltage-dependent potassium channel in cultured pig articular chondrocytes. *Pflugers Arch.* Vol. 433, pp. 413-427.
- Nilius, B.; Droogmans, G. (1994). A role for potassium channels in cell proliferation?. *News Physiol Sci.* Vol. 16:1-12.
- Nishizawa, T., Nagao, T., Iwatsubo, T., Forte, J.G. and Urushidani, T. (2000) Molecular cloning and characterization of a novel chloride intracellular channel-related protein, parchorin, expressed in water-secreting cells. *J Biol Chem.* Vol. 275, pp. 11164-11173.
- Okumura, N.; Imai, S.; Toyoda, F.; Isoya, E.; Kumagai, K.; Matsuura, H.; Matsusue, Y. (2009). Regulatory role of tyrosine phosphorylation in the swelling-activated chloride current in isolated rabbit articular chondrocytes. *J Physiol* Vol. 587, pp. 3761-3776.
- Ostergaard, K.; Salter, D.M.; Petersen, J.; Bendtzen, K.; Hvolris, J.; Andersen, C.B. (1998). Expression of alpha and beta subunits of the integrin superfamily in articular cartilage from macroscopically normal and osteoarthritic human femoral heads. *Ann Rheum Dis* Vol. 57, pp. 303-308.

- Pérez, E.; Gallegos, J.L.; Cortés, L.; Calderón, K.G.; Luna, J.C.; Cázares, F.E.; Velasquillo, M.C.; Kouri, J.B.; Hernández, F.C. (2010). Identification of latexin by a proteomic analysis in rat normal articular cartilage. *Proteome Sci.* Vol. 5, pp. 27.
- Phan, M.N.; Leddy, H.A.; Votta, B.J.; Kumar, S.; Levy, D.S.; Lipshutz, D.B.; Lee, S.H.; Liedtke, W.; Guilak, F. (2009). Functional characterization of TRPV4 as an osmotically sensitive ion channel in porcine articular chondrocytes. *Arthritis Rheum.* Vol. 60, pp. 3028–3037.
- Poët, M.; Kornak, U.; Schweizer, M.; Zdebik, A.A.; Scheel, O.; Hoelter, S.; Wurst, W.; Schmitt, A.; Fuhrmann, J.C.; Planells-Cases, R.; Mole, S.E.; Hübner, C.A.; Jentsch, T.J. (2006). Lysosomal storage disease upon disruption of the neuronal chloride transport protein CIC-6. *Proc Natl Acad Sci USA* Vol. 103, pp. 13854–13859.
- Ponce, A. (2006). Expression of voltage dependent potassium currents in freshly dissociated rat articular chondrocytes. *Cell Physiol Biochem.* Vol. 18, pp. 35–46.
- Poole, C.A. (1997). Articular cartilage chondrons: form, function and failure. *J Anat* Vol. 191, pp. 1–13.
- Pritzker, K.P.; Gay, S.; Jimenez, S.A.; Ostergaard, K.; Pelletier, J.P.; Revell, P.A.; Salter, D.; van den Berg WB. (2006). Osteoarthritis cartilage histopathology: grading and staging. *Osteoarthritis Cartilage.* Vol. 14, pp. 13–29.
- Qian, Z.; Okuhara, D.; Abe, M.K.; Rosner, M.R. (1999). Molecular cloning and characterization of a mitogen activated protein kinase-associated intracellular chloride channel. *J Biol Chem.* Vol. 274, pp. 1621–1627.
- Ramage, L.; Martel, M.A.; Hardingham, G.E.; Salter, D.M. (2008). NMDA receptor expression and activity in osteoarthritic human articular chondrocytes. *Osteoarthritis Cartilage.* Vol. 16, pp. 1576–1584.
- Ronnov-Jessen, L.; Villadsen, R.; Edwards, J.C.; Petersen, O.W. (2002). Differential expression of a chloride intracellular channel gene, CLIC4, in transforming growth factor-beta1-mediated conversion of fibroblasts to myofibroblasts. *Am J Pathol* Vol. 161, pp. 471–480.
- Roos, E.M. (2005). Joint injury causes knee osteoarthritis in young adults. *Curr Opin Rheumatol* Vol. 17, pp. 195–200.
- Rothwell, A.G.; Bentley, G. Chondrocyte multiplication in osteoarthritic articular cartilage. (1973). *J Bone Joint Surg Br.* Vol. 55, pp. 588–594.
- Salter, D.M.; Millward-Sadler, S.J.; Nuki, G.; Wright, M.O. (2002). Differential responses of chondrocytes from normal and osteoarthritic human articular cartilage to mechanical stimulation. *Biorheology* Vol. 39, pp. 97–108.
- Sanchez, J.C.; Danks, T.A.; Wilkins, R.J. (2003). Mechanisms involved in the increase in intracellular calcium following hypotonic shock in bovine articular chondrocytes. *Gen Physiol Biophys.* Vol. 22, pp. 487–500.
- Sanchez, J. C., and Wilkins, R. J. (2004). Changes in intracellular calcium concentration in response to hypertonicity in bovine articular chondrocytes. *Comp Biochem Physiol A Mol Integr Physiol.* Vol. 137, pp. 173–182.
- Sanchez, J.C.; Powell, T.; Staines, H.M.; Wilkins, R.J. (2006). Electrophysiological demonstration of voltage-activated H⁺ channels in bovine articular chondrocytes. *Cell Physiol Biochem.* Vol. 18, pp. 85–90.
- Sandell, L.J.; Aigner, T. (2001). Articular cartilage and changes in arthritis. An introduction: cell biology of osteoarthritis. *Arthritis Res.* Vol. 3, pp. 107–113.

- Schlesinger, P.H.; Blair, H.C.; Teitelbaum, S.L.; Edwards, J.C. (1997). Characterization of the osteoclast ruffled border chloride channel and its role in bone resorption. *J Biol Chem* Vol. 272, pp. 18636–18643.
- Shakibaei, M.; Mobasher, A. (2003). Beta1-integrins co-localize with Na, K-ATPase, epithelial sodium channels (ENaC) and voltage activated calcium channels (VACC) in mechanoreceptor complexes of mouse limb-bud chondrocytes. *Histol Histopathol.* Vol. 18, pp. 343–351.
- Singh, H.; Cousin, M.A.; Ashley, R.H. (2007). Functional reconstitution of mammalian 'chloride intracellular channels' CLIC1, CLIC4 and CLIC5 reveals differential regulation by cytoskeletal actin. *FEBS J* Vol. 274, pp. 6306–6316.
- Stockwell, R.A. (1967). The cell density of human articular and costal cartilage. *J Anat* Vol. 101, pp. 753–763.
- Stockwell, R.A. (1990). Morphology of cartilage. In *Maroudas A & Keuttner K* (eds.). *Methods in cartilage research*. San Diego: Academic Press, pp. 61–63.
- Stockwell, R.A. (1991). Cartilage failure in osteoarthritis: Relevance of normal structure and function. *Clinical Anatomy* Vol. 4, pp. 161–191.
- Sugimoto, T.; Yoshino, M.; Nagao, M.; Ishii, S.; Yabu, H. (1996). Voltage-gated ionic channels in cultured rabbit articular chondrocytes. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol.* Vol. 115, pp. 223–232.
- Trujillo, E., Alvarez de la Rosa, D., Mobasher, A., Gonzalez, T., Canessa, C. M., and Martin-Vasallo, P. (1999). Sodium transport systems in human chondrocytes. II. Expression of ENaC, Na⁺/K⁺/2Cl⁻ cotransporter and Na⁺/H⁺ exchangers in healthy and arthritic chondrocytes. *Histol Histopathol.* Vol. 14, pp. 1023–1031.
- Tsuga, K.; Tohse, N.; Yoshino, M.; Sugimoto, T.; Yamashita, T.; Ishii, S.; Yabu, H. (2002). Chloride conductance determining membrane potential of rabbit articular chondrocytes. *J Membr Biol.* Vol. 185, pp. 75–81.
- Tulk, B.M.; Kapadia, S.; Edwards, J.C. (2002). CLIC1 inserts from the aqueous phase into phospholipid membranes, where it functions as an anion channel. *Am J Physiol Cell Physiol.* Vol. 282, pp. C1103–1112.
- Urban, J.P.; Hall, A.C.; Gehl, K.A. (1993). Regulation of matrix synthesis rates by the ionic and osmotic environment of articular chondrocytes. *J Cell Physiol.* Vol. 154, pp. 262–270.
- Urban, J.P. (1994). The chondrocyte: a cell under pressure. *Br J Rheumatol.* Vol. 33, pp. 901–908.
- Valenzuela SM, Martin DK, Por SB, Robbins JM, Warton K, Bootcov MR, Schofield PR, Campbell TJ, Breit SN. (1997). Molecular cloning and expression of a chloride ion channel of cell nuclei. *J Biol Chem* Vol. 272, pp. 12575–12582.
- Walsh, K.B.; Cannon, S.D.; Wuthier, R.E. (1992). Characterization of a delayed rectifier potassium current in chicken growth plate chondrocytes. *Am J Physiol.* Vol. 262, pp. C1335–C1340.
- Wang, X.T.; Nagaba, S.; Nagaba, Y.; Leung, S.W.; Wang, J.; Qiu, W.; Zhao, P.L.; Guggino, S.E. (2000). Cardiac L-type calcium channel alpha 1-subunit is increased by cyclic adenosine monophosphate: messenger RNA and protein expression in intact bone. *J Bone Miner Res.* Vol. 15, pp. 1275–1285.
- Weiss, C.; Rosenberg, L.; Helfet, A.J. (1968). An ultrastructural study of normal young adult human articular cartilage. *J Bone Joint Surg Am* Vol. 50, pp. 663–674.

- Wilson, J.R.; Duncan, N.A.; Giles, W.R.; Clark, R.B. (2004). A voltage dependent K⁺ current contributes to membrane potential of acutely isolated canine articular chondrocytes. *J Physiol*. Vol. 557, pp. 93-104.
- Wohlrab, D. (1998). Der Einfluß elektrophysiologischer Membraneigenschaften auf die Proliferation humaner Keratinozyten. Tectum, Marburg, pp. 1-89
- Wohlrab, D.; Lebek, S.; Krüger, T.; Reichel, H. (2002). Influence of ion channels on the proliferation of human chondrocytes. *Biorheology* Vol. 39, pp. 55-61.
- Wohlrab, D.; Vocke, M.; Klapperstück, T.; Hein, W. (2004). Effects of potassium and anion channel blockers on the cellular response of human osteoarthritic chondrocytes. *J Orthop Sci*. Vol. 9, pp. 364-371.
- Wohlrab, D.; Vocke, M.; Klapperstück, T.; Hein, W. (2005). The influence of lidocaine and verapamil on the proliferation, CD44 expression and apoptosis behavior of human chondrocytes. *Int J Mol Med*. Vol. 16, pp. 149-57.
- Wright, M.O.; Stockwell, R.A.; Nuki, G. (1992). Response of the plasma membrane to applied hydrostatic pressure in chondrocytes and fibroblasts. *Connect Tissue Res* Vol. 28, pp. 49-70.
- Wright, M.O.; Jobanputra, P.; Bavington, C.; Salter, D.M.; Nuki, G. (1996). The effects of intermittent pressurisation on the electrophysiology of cultured human articular chondrocytes: evidence for the presence of stretch activated membrane ion channels. *Clin Sci (Lond)*. Vol. 90, pp. 61-71.
- Wu, J.J.; Eyre, D.R.; Slayter, H.S. (1987). Type VI collagen of the intervertebral disc. Biochemical and electron microscopic characterization of the native protein. *Biochem J* Vol. 248, pp. 373-381.
- Wu, J.J.; Murray, J.; Eyre, D.R. (1996). Evidence for copolymeric crosslinking between types II and III collagens in human articular cartilage. *Trans Orthop Res Soc* Vol. 21, pp. 42.
- Wu, Q. Q., and Chen, Q. (2000). Mechanoregulation of chondrocyte proliferation, maturation, and hypertrophy: ion-channel dependent transduction of matrix deformation signals. *Exp Cell Res* Vol. 256, pp. 383-391.
- Xu, J.; Wang, W.; Clark, C.C.; Brighton, C.T. (2009). Signal transduction in electrically stimulated articular chondrocytes involves translocation of extracellular calcium through voltage-gated channels. *Osteoarthritis Cartilage*. Vol. 17, pp. 397-405.
- Xu, X.; Urban, J.P.; Tirlapur, U.K.; Cui, Z. (2010). Osmolarity effects on bovine articular chondrocytes during three-dimensional culture in alginate beads. *Osteoarthritis Cartilage* Vol. 18, pp. 433-439.



Principles of Osteoarthritis- Its Definition, Character, Derivation and Modality-Related Recognition

Edited by Dr. Bruce M. Rothschild

ISBN 978-953-51-0063-8

Hard cover, 590 pages

Publisher InTech

Published online 22, February, 2012

Published in print edition February, 2012

This volume addresses the nature of the most common form of arthritis in humans. If osteoarthritis is inevitable (only premature death prevents all of us from being afflicted), it seems essential to facilitate its recognition, prevention, options, and indications for treatment. Progress in understanding this disease has occurred with recognition that it is not simply a degenerative joint disease. Causative factors, such as joint malalignment, ligamentous abnormalities, overuse, and biomechanical and metabolic factors have been recognized as amenable to intervention; genetic factors, less so; with metabolic diseases, intermediate. Its diagnosis is based on recognition of overgrowth of bone at joint margins. This contrasts with overgrowth of bone at vertebral margins, which is not a symptomatic phenomenon and has been renamed spondylosis deformans. Osteoarthritis describes an abnormality of joints, but the severity does not necessarily produce pain. The patient and his/her symptoms need to be treated, not the x-ray.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Elizabeth Perez-Hernandez, Nury Perez-Hernandez, Fidel de la C. Hernandez-Hernandez and Juan B. Kouri-Flores (2012). Anion Channels in Osteoarthritic Chondrocytes, Principles of Osteoarthritis- Its Definition, Character, Derivation and Modality-Related Recognition, Dr. Bruce M. Rothschild (Ed.), ISBN: 978-953-51-0063-8, InTech, Available from: <http://www.intechopen.com/books/principles-of-osteoarthritis-its-definition-character-derivation-and-modality-related-recognition/anion-channels-in-osteoarthritic-chondrocytes>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen